

# Biodiesel from microalgae lipids: from inorganic carbon to energy production

Marc Veillette, Anne Giroir-Fendler, Nathalie Faucheux & Michèle Heitz

To cite this article: Marc Veillette, Anne Giroir-Fendler, Nathalie Faucheux & Michèle Heitz (2018) Biodiesel from microalgae lipids: from inorganic carbon to energy production, *Biofuels*, 9:2, 175-202, DOI: [10.1080/17597269.2017.1289667](https://doi.org/10.1080/17597269.2017.1289667)

To link to this article: <https://doi.org/10.1080/17597269.2017.1289667>



Published online: 16 Mar 2017.



Submit your article to this journal [↗](#)



Article views: 447



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 11 View citing articles [↗](#)



# Biodiesel from microalgae lipids: from inorganic carbon to energy production

Marc Veillette<sup>a,b</sup>, Anne Giroir-Fendler<sup>b</sup>, Nathalie Fauchoux<sup>a</sup> and Michèle Heitz<sup>a</sup>

<sup>a</sup>Department of Chemical Engineering and Biotechnological Engineering, Engineering Faculty, Université de Sherbrooke, Canada;

<sup>b</sup>Université Lyon 1, CNRS, UMR 5256, IRCELYON, Institut de Recherches sur la Catalyse et l'Environnement de Lyon, 2 avenue Albert Einstein, 69626 Villeurbanne Cedex, France

## ABSTRACT

Following the United Nations Conference on Climate Change, COP21 (Paris, France), several countries have attempted to reduce their greenhouse gas emissions. In order to reach this objective, microalgae could be used to capture carbon dioxide and transform it into a biomass composed essentially of lipids, carbohydrates and proteins. Moreover, cultivating microalgae does not require arable land, in opposition to several oleaginous plants used to produce biofuels. Despite the fact that microalgae could be transformed into several biofuels such as bioethanol (by fermentation of hydrocarbons) and biomethane (by anaerobic digestion), transforming lipids into biodiesel could allow the reduction of oil-based diesel consumption. However, microalgae biodiesel production costs remain high for a short-term commercialization. The microalgae lipids can be transesterified into biodiesel in the presence of catalysts (homogeneous or heterogeneous). In order to commercialize biodiesel from microalgae, biodiesel physicochemical properties must respect the American Society for Testing and Materials (ASTM) standards. The aim of the study was to describe the current technologies available to produce biodiesel from microalgae.

## ARTICLE HISTORY

Received 30 March 2016

Accepted 6 May 2016

## KEYWORDS

Microalgae; culture; lipids; extraction; transesterification; biodiesel

## Introduction

Between 1980 and 2011, the world carbon dioxide (CO<sub>2</sub>) emissions increased from 18.5 to 31.5 billion metric tons per year, which represented an increase of 420 million metric tons per year [1]. In order to reduce these emissions, several countries signed the United Nations Framework Convention on Climate Change at COP21 (Paris, France) in December 2015. Among them, by 2030, the Canadian government is committed to reduce greenhouse gases (GHG) emissions by 30% under the 2005 GHG emission level of 607 Mt CO<sub>2</sub> eq [2]. In Canada (2013), near 80% of all GHG, around 581 Mt CO<sub>2</sub> eq, was attributable to CO<sub>2</sub> emissions [3,4]. The Canadian CO<sub>2</sub> emissions related to energy consumption increased from 457 to 581 million metric tons between 1980 and 2013, which represents around 16 metric tons of CO<sub>2</sub> per person in 2013 [1,4]. The main sectors responsible for CO<sub>2</sub> emissions are the energy sector (93%) and industrial processes (6.7%) [5].

The exploitation of fossil fuel in western Canada has, among other things, contributed to increasing CO<sub>2</sub> emissions linked to the energy sector from 467 to 588 Mt CO<sub>2</sub> eq, an increase of 25% between 1990 and 2013 [4,5]. Indeed, Canada owns the second most important petroleum deposit (all forms included) in the world with 178 billion barrels, which is the 4<sup>th</sup> most important oil producer in the world with 3.3 million barrels per

day (2012); most of this oil is exported to the United States of America [6].

Between 1992 and 2012, the world proven petroleum deposit evaluated by the BP petroleum company seems to have increased from 1040 to 1690 billion barrels [7]. The world petroleum deposit was evaluated at 1526 billion barrels in 2012 [8]. Some authors predict that these petroleum deposits will be sufficient to fulfill the demand for petroleum-based products until 2042 [9]. This could be attributable to the fact that the crude petroleum consumption increased from 78 to 90 million barrels per day from 2002 to 2012 [7]. In 2012, the world's most important petroleum oil consumers were the United States, China and Japan with 18.5, 10.3 and 4.7 million barrels per day, respectively [10]. According to the same source, Canada was the 10<sup>th</sup> most important oil consumer with 2.3 million barrels per day in 2012.

Alternatively to non-renewable energy sources, energy can be produced from biomass such as lignocellulosic plants, herbaceous plants, water plants and animal manure [11]. Indeed, in 2011, 82% of all the world's non-renewable energy was produced from biomass, followed by hydropower, wind, solar and other with 15, 2.0, 0.2 and 0.4%, respectively [12].

In order to find more sustainable solutions to replace oil-based fuel, other solutions are currently being developed such as liquid biofuels, gaseous

biofuels (hydrogen (H<sub>2</sub>), methane (CH<sub>4</sub>), etc.), solar energy, fuel cell and hybrid technologies [13]. Consequently, several countries (mostly European countries) have voted for laws stating that the fuel used by transportation must contain at least 10% (v/v) biofuel such as bioethanol or biodiesel by 2010 [14]. However, since most biofuels are produced from plants cultivated on arable lands, the European countries reduced, in 2013, their target of biofuels in fuels to 6% (v/v) [15].

Despite the fact that the biodiesel production is around 4 times lower than the bioethanol production [16], the interest in biodiesel has constantly been rising since the fluctuation of oil prices [17]. In April 2011, the average price of a barrel of oil reached a peak value exceeding 121 US\$ [18,19] while in 2015 it decreases to less than 40 US\$ [20]. Furthermore, the world biodiesel production increased from 77 to  $403 \times 10^3$  barrels per day between 2005 and 2011, but the Canadian biodiesel production remains low at 3600 barrels per day in 2012 [10].

Among the vegetable oleaginous species used as a source of biomass, microalgae are an interesting source because these microorganisms, which do not require being cultivated on arable lands, can be utilized to produce biodiesel [17].

Microalgae can also be utilized as a raw material for the production of food supplements, medical chemicals and food [21,22]. Moreover, microalgae can be transformed into energy using gasification (synthesis gas), torrefaction, anaerobic digestion, direct combustion (heat), liquefaction, pyrolysis, hydrogenation and transesterification [23].

Biodiesel from vegetable oil can produce the required energy to be used in transportation and generating low air pollutants such as sulfur and polycyclic aromatic compounds [24]. For example, increasing the biodiesel content in petrodiesel from 0 to 20% (v/v) reduces the emissions of particulate matter (−10%), carbon monoxide (−11%) and unburned hydrocarbons (−21%), but slightly increases the nitrogen oxide emissions (+2%) [25,26]. Moreover, with a 20% (v/v) biodiesel-petrodiesel blend (B20), mercaptan emissions (bad smells) can be reduced up to 18% [27]. In addition, increasing the biodiesel content from 20 to 100% (v/v) in petrodiesel can increase the distance traveled from 0.38 to 4.5 km/L [28]. According to the same source, the main problem with biodiesel is the feedstock cost, which represents 80% of the operating costs. However, the research on microalgae biodiesel must continue because the biodiesel production costs remain high between 2.4 and 6.6 \$US/L (microalgae culture in ponds) [29]. Reducing the cost associated with microalgae biodiesel is an important challenge for the replacement of first-generation biodiesel, produced from oleaginous plants such as soybean.

This literature review will describe microalgae culture, composition, harvesting, lipid extraction and lipid

transesterification. Moreover, the possibilities of using microalgae as a raw material for biodiesel production will be discussed through biodiesel properties, by-products, cost calculation and Canadian biodiesel production.

## Microalgae: a source of sustainable development

Currently, first-generation biofuels are mainly produced from agricultural biomass such as soybean, rapeseed, palm, etc. Most of the biodiesel produced in the United States comes from soybean culture [17]. However, using arable lands to produce biodiesel can have negative effects on the economy, the environment and the society.

On an economic level, producing biofuels such as biodiesel on a large scale using arable lands in order to cultivate the biomass could lead to an increase of food prices. As an example, between 2000 and 2011, the FAO Food Price Index and FAO Oils/Fats Price Index of several food and oil prices increased by 250% [30]. This situation would, as a consequence, increase the disparity between rich and poor countries, threatening the food security of the poorest countries [31]. At an environmental level, producing biofuels on a large scale using arable land could create problems of soil impoverishment, land pollution and deforestation [32–34].

Using microalgae as a raw material for biofuel production could solve these problems as microalgae culture does not compete with human and animal food production [35]. Furthermore, this type of culture does not require heavy equipment for chemical product spreading or for watering because nutrients and water are added to the bioreactor.

One of the main advantages of microalgae is their lipid content. As an example, palm oil can contain up to 36% (dry weight) of lipids (triglycerides) with a maximal culture yield of 5.4 m<sup>3</sup> lipids/ha/year [36], while microalgae can contain up to 75% (dry weight) lipids with a culture yield of 137 m<sup>3</sup> lipids/ha/year [17].

## Microalgae composition

Microalgae are composed of four main components: lipids, carbohydrates, proteins and nucleic acids [37].

### Carbohydrates

Carbohydrates are structural material, with a metabolic function, of the microalga, which can be found as starch, polysaccharides, monosaccharides (like glucose) and more [38,39]. Concerning biofuels, one of the main interests of carbohydrates is to use microalgal glucose and transform it by fermentation into bioethanol with a yield up to 50 wt% (compared to the total carbohydrates) for *Chlorococcum humicola* microalgae (a 100%

conversion based on glucose). [40]. Another study stated that cyanobacteria could synthesize directly bioethanol from CO<sub>2</sub> with an annual production of 20 m<sup>3</sup>/ha [41].

### Proteins

Like carbohydrates, proteins contained in microalgae have metabolic and structural functions [42]. Depending on the type of microalgae, the protein content ranges from 30 to 50 wt% of the dry biomass [43,44]. Several companies (such as Cyanotech, Martek Biosciences Corporation, Mera Pharmaceutical, etc.) have cultivated microalgae for their protein content (dietary supplements) or other components like pigments [43].

### Lipids

Lipids are also structural elements with an energy storage function [42]. Despite the fact that lipids are a complex group of 14 categories, such as hydrocarbons, fatty acids, phospholipids, glycolipids and sulfated lipids [45], several of these lipids cannot be transformed into fatty acids methyl esters (FAME) and are called unsaponified lipids.

Some polyunsaturated lipids (omega 3 and omega 6) such as eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA) showed a potential reduction of some types of cancer such as colorectal cancer [46], and have antiangiogenic, antivasoproliferative and neuroprotective properties [47].

Depending on the type of microalgae, the lipid yield obtained from microalgae can reach more than 75% (g lipid/g dry biomass) [48–50]. The microalgae with the most important lipid contents are (g lipid/g dry biomass): *Botryococcus braunii* (25–75%), *Schizochytrium* sp. (50–77%), *Dunaliella tertiolecta* (16–71%), *Dunaliella* sp. (17–67%) and *Neochloris oleoabundans* (29–65%).

Several microalgae species have a high carotenoid content, like the microalga *Dunaliella salina* that contains 10% (g/g dry biomass) of  $\beta$ -carotene (a kind of carotenoid) [51]. Some factors affect the carotenoid content in microalgae such as temperature, pH, and dissolved oxygen level in the culture medium [52]. As an example, using a stream of air containing 2% (v/v) of CO<sub>2</sub> for the growth of the microalga *Chlorococcum* sp. for 10 days at an irradiance of 200  $\mu$ mol photons/m<sup>2</sup>/s, Liu *et al.* [51] increased the initial pH from 5 to 9 and found that a pH of 8 was the best pH tested for carotenoid growth with a total carotenoid content of 7.4 mg/g.

Other studies tried to recover extracellular polymeric substances (EPSs; extracellular biomass) of microalgae [53,54]. For example, a recent study used solvents to extract a maximum concentration of EPSs in an aqueous phase of 0.944 g/L of EPSs from the microalga *Dunaliella salina* and obtained a maximal emulsifying activity of 86% EPSs [53].

### Nucleic acids and chlorophyll

In microalgae, the content of nucleic acids generally varies from 4 to 6% (g/g dry biomass) [55]. On the other hand, a high level of nucleic acids (> 0.3 g microalgae/kg body weight/day) induces oral toxicity for the human body [56], which means that microalgae must be processed before human consumption. For animal food, unprocessed microalgae do not seem to be toxic [55].

Chlorophyll is a green pigment. Industrially, this pigment is mainly used as an additive by the pharmaceutical industry (cosmetic products) and also acts as a natural dying agent [57]. Moreover, according to the same source, these compounds have antioxidant and antimutagenic properties. The solvents used to extract chlorophyll are methanol, ethanol, dimethyl formamide, acetone and supercritical CO<sub>2</sub> [57]. As an example, Bai *et al.* [58] studied microalgae pigments (chlorophyll and carotenoids). They obtained chlorophyll contents ranging from 1.7 to 5.6% (g lipid/g dry biomass) for a few species of microalgae (including *Chlorella* sp., *Chlorella minutissima*, *Dunaliella promolecta*, *Isochrysis galbana* and *Nannochloropsis oculata*) by using methanol as an extraction solvent.

### Microalgae metabolisms

There are more than 30,000 species of algae known on Earth [59], that can be classified as a function of their taxonomy, size and metabolism [34,42,60]. Based on their metabolism, there are four main classifications to identify microalgae: photoautotrophic, heterotrophic, mixotrophic and photoheterotrophic.

#### Photoautotrophic

Inorganic carbon (CO<sub>2</sub> or bicarbonate) is transformed into biomass (microalgae) and oxygen following an input of light energy, water and nutrients [57,61]. Carbon dioxide (CO<sub>2</sub>) can be supplied by several sources. A common source of CO<sub>2</sub> often cited in the literature comes from thermal power plants [62].

Despite the fact that the photoautotrophic metabolism seems to be a sustainable solution to reduce the CO<sub>2</sub> emissions, this process has some limits. Even if the lipid content of some microalgae (*Botryococcus braunii*, *Dunaliella tertiolecta*, *Neochloris oleoabundans*) is relatively high (around 75% (g lipid/g dry biomass)), the concentrations of microalgae biomass obtained by these photoautotrophic metabolisms are low to medium (0.5–6.7 g/L) which makes microalgae biomass harvesting problematic compared to other oleaginous species because microalgae require a high volume of culture medium.

#### Heterotrophic

Heterotrophic microalgae transform organic carbon sources without light within fermenters [63]. Some

microalgae species with heterotrophic metabolisms were studied because they can have high lipid contents (up to 56% (g lipid/dry biomass)) [64,65]. It should be noted that heterotrophic microalgae, such as *Chlorella pyrenoidosa*, can reach high biomass concentrations (up to 216 g/L), but rheological problems in fermenters (mass transfer and mixing problems) were observed at biomass concentrations higher than 50 g/L [66,67].

Despite the fact that glucose is the most common carbon source [65], other carbon sources were tested for microalgae growth, such as fructose, galactose, acetic acid, sodium acetate, glycerol and various hydrolysis products [64,68]. Among the hydrolysis products are sugarcane [69], artichoke [70,71], tapioca [72] and rice straw [65]. A study demonstrated that hydrolysis products have a higher potential than glucose does. For example, growing the microalga *Chlorella pyrenoidosa* with glucose and rice straw hydrolysate, as carbon sources, at a concentration of 10 g/L, Li *et al.* [65] obtained a higher lipid content of 56% (g lipid/g dry biomass) using rice straw hydrolysate than glucose 50% (g lipid/g dry biomass). Furthermore, these authors obtained respective biomass concentrations of 2.83 g/L and 0.92 g/L for rice straw hydrolysate and glucose as carbon sources.

Despite the potential of heterotrophic microalgae for biodiesel production, no commercial scale plant of microalgae for biodiesel purpose has been built [64] up to now. This could be explained by the fact that the first objective of microalgae culture is to decrease CO<sub>2</sub> emissions. Moreover, heterotrophic culture requires a cheap organic source such as organic matter hydrolysate [73], which is directly in competition with bioethanol production.

### Mixotrophic and photoheterotrophic

Microalgae with a mixotrophic metabolism can have a photoautotrophic or heterotrophic metabolism. They can use inorganic or organic carbon with or without light energy [68,74]. When the light is absent, these types of microalgae can continue to grow by using inorganic or organic carbon. Liang *et al.* [75] used glucose and CO<sub>2</sub> (from air) in the absence of light and obtained respective biomass productivities of 151 and 10 mg/L/day. When they used glucose in similar but dark conditions, the biomass productivity increased to 254 mg/L/day. The term 'photoheterotrophic' is kept for microalgae that are able to use organic carbon and light [74].

### Microalgae production

Industrial microalgae culture for lipid extraction was considered a long time ago. As an example, some studies were performed in the 1950s in order to produce the *Chlorella* algae on a wide scale as a food source

[76]. On the other hand, the interest in a specific microalgae production for a biofuel purpose is linked to the increase of oil prices and to the reduction of CO<sub>2</sub> mentioned previously. An interesting fact is that microalgae can mitigate CO<sub>2</sub> emissions of several industries such as thermal power plants that burned oil fuel [77,78]. Moreover, microalgae can also be cultivated using wastewaters (as a source of nutrients) despite the abundance of heavy metals, pathogenic microorganisms, nitrogen or phosphorus [79,80].

A major problem with microalgae production is the fact that unrealistic biomass production rates reported in research articles have led to the cancellation of several microalgae production scaling-up projects [81].

On an industrial scale, four modes of culture can be considered: open ponds, photobioreactors, hybrid modes and fermenters.

### Open ponds

Open ponds are divided into three categories [82]: (1) lakes, (2) lagoons and (3) ponds. Most industrial microalgae cultures are performed in raceway type ponds, shallow (less than 30 cm deep), provided with a paddle wheel, which creates a flow to avoid microalgae settling [17]. Nutrients and water are added in front of the paddle wheel and microalgae are harvested at the back of the wheel [17,83].

The main advantage of open culture systems (outside) is to use free sunlight energy [84]. However, these systems have the disadvantage of producing low biomass density (lower than 5 g/L) [85], which complicates the dewatering process. Cultivating the microalga *Chlorella* sp. in an open pond, Doucha and Lívanský [86] obtained a high biomass concentration (40 g/L) after 18 days of production. On the other hand, to our best knowledge of the literature, no other studies have reported such a high biomass concentration in open ponds. The main advantage of open systems is linked to their facility of scaling up and their lower investment costs, even if these systems have non-negligible disadvantages compared to closed systems: risk of contamination, higher harvesting costs, loss of CO<sub>2</sub> (no CO<sub>2</sub> recycled), higher water evaporation rates and bad use of light energy [87].

Despite the fact that open ponds are used for industrial microalgae production for food and cosmetic purposes [88–90], production costs (around 3.65 US\$/kg biomass) make this technology hard to apply industrially for the production of biodiesel [91].

### Photobioreactors

In order to solve the problems linked to microalgae culture in ponds (biomass concentration, contamination, etc.), microalgae culture in photobioreactors is often preferred [92]. There are several types of photobioreactors [87, 93]: (1) tubular, (2) flat plate, (3) airlift, (4) bubbling column and (5) stirred tank.



Tubular photobioreactors, made of plastic or glass, can have different configurations: spiral [94,95], external looped [96] and straight [87]. These systems can be paired with a pump or an airlift system [87,96].

The flat plate photobioreactors have parallel glass plates between which microalgae grows with sunlight. Kurano *et al.* [97] cultivated the microalgae *Chlorococcum littorale* with this type of reactor and obtained biomass doubling every 2 h at a temperature of 40°C. Sometimes, this type of reactor is used with only two parallel glass plates in which the gas is fed like in a bubbling column [98].

The airlift bioreactors are a group of two vertical concentric tubes [99] in which the inside tube is perforated at the inlet and the outlet [100] or is shorter than the outside tube [87,99]. This configuration must allow microalgae to go up in the inlet tube and go down between the two tubes. Air is fed in at the bottom of the reactor and makes microalgae flow from the bottom to the top to avoid settling.

A bubbling photobioreactor works similarly to an airlift photobioreactor. The bubbles are produced with a perforated flat plate in which air and CO<sub>2</sub> form bubbles at the bottom of the reactor. Some of these reactors have produced interesting biomass yields. For example, Ranjbar *et al.* [101] cultivated the microalga *Hoematococcus pluvialis* in a bubbling air column of 1.1 L and obtained a biomass concentration of 6.7 g/L after 33 days of culture (illumination at 21.5 μmol/m<sup>2</sup>/s; temperature of 20°C; pH of 7.5 by adjusting the CO<sub>2</sub> concentration).

Stirred tanks are photobioreactors rarely used despite a biomass productivity of 47.8 g/m<sup>2</sup>/day (in pseudo steady-state regime) for a configuration of bioreactor in series [102]. The little interest in this type of reactor could be explained by the fact that some studies obtained low growth rates and/or low biomass concentrations. For example, using a stirred tank photobioreactor to cultivate the marine microalgae *Dunaliella salina*, Li *et al.* [103] obtained a maximum growth rate and biomass concentration of 0.119 1/h and 0.55 g/L, respectively, after around 116 h of culture (temperature of 29°C; pH of around 7.4).

Other configurations of dome-shaped photobioreactors have been tested. For example, Sato *et al.* [104] grew the microalga *Chlorococcum littorale* and reached a biomass concentration of 2.5 g/L (temperature of 25°C; pH of 7.5; time of 288 h).

### Hybrid culture

As technical problems occurred with both of the previous systems (costs, scaling up, contamination, biomass concentration, etc.), some researchers paired ponds and photobioreactors [105,106]. The first step is a photobioreactor in which the culture conditions are controlled to minimize the risks of contamination [105]. The temperature is controlled with a water bath (16–

18°C) [106]. Then, microalgae are transferred into a pond until the harvesting with nutrient deprivation [91].

### Fermenters

Fermenters are stirred reactors in which microalgae with heterotrophic metabolisms are grown. For this type of reactor, biomass concentrations and lipid content can reach 116 g/L and 56% (g lipid/g biomass), respectively [107]. On the other hand, a higher biomass concentration can be reached (up to 150 g/L) without rheological problems [66]. According to Cantin *et al.* [64], a few companies such as Solazyme Inc. (South San Francisco, USA) and Fermenlg (Libourne, France) produced biodiesel within fermenters with organic carbon and without light. Solazyme Inc. affirmed that some of their biodiesels, Soladiesel<sub>RD</sub> et Soladiesel<sub>BD</sub>, could have a higher cetane number (74) or better cold properties than conventional biodiesel [108]. However, no specification is given on the production capacity or the costs of the biodiesel produced.

### Microalgae culture parameters

As there are few projects of industrial scale for microalgae with heterotrophic metabolisms [64] and the production costs are high [72, 109], the present study will focus on the autotrophic culture metabolism. The main factors that influence microalgae productivity are: (1) inorganic carbon concentration, (2) light energy, (3) oxygen concentration, (4) temperature, (5) pH and (6) stirring.

#### • Inorganic carbon concentration

Inorganic carbon (such as CO<sub>2</sub>) is the main source of carbon for autotrophic microalgae. Despite the fact that CO<sub>2</sub> of surrounding air can be used for microalgae growth, CO<sub>2</sub> concentration ranging from 2 to 15% (v/v) have been tested [84,110,111]. Some microalgae species such as the strain ZY-1 (*Chlorella*) can stand high CO<sub>2</sub> concentrations (up to 70% (v/v)) [112]. According to our best knowledge of the literature, the optimal CO<sub>2</sub> concentration is between 4 and 10% (v/v) [110,112,113].

#### • Energy light intake

One of the most important factors is the light energy intake, which is an essential factor for microalgae growth. In the laboratory, the microalga uses an absorption spectrum between 400 to 700 nm with an efficiency that depends on the photosynthetic efficiency [84,114,115].

As the sun is a free and an unlimited energy resource, microalgae culture is performed outdoors. Consequently, the biomass productivity can be influenced depending of the time of the year. For example,

Blanco *et al.* [116] cultivated the microalgae *Muriellopsis* sp. in open ponds (Isla de la Cartuja, Sevilla, Spain) and obtained a higher biomass productivity during the months of May, June and July (20 g/m<sup>3</sup>/day), while the lowest biomass productivity was obtained during the month of November (8 g/m<sup>3</sup>/day).

#### • Oxygen concentration (O<sub>2</sub>)

Oxygen is a microalgae production by-product, but the O<sub>2</sub> concentration must not be too high because it has an inhibitory effect on microalgae growth [117]. Kitaya *et al.* [110] increased the O<sub>2</sub> concentration from 10 to 30% (v/v) during the growth of the microalga *Euglena gracilis* and noticed that an increase of O<sub>2</sub> concentration decreased the maximum specific growth rate from 0.45 to 0.14 1/h.

#### • Nutrients intake

At a laboratory scale, culture medium with water and mineral salts [101,114,115] was often used because it allows a better control of nutrient intake. The main macronutrients are nitrogen, phosphorus, hydrogen, sulfur, calcium, magnesium, sodium and potassium [84]. A modification of the nutrients has different effects depending on the microalga. The main macronutrients used for the lipid synthesis are carbon, hydrogen, oxygen, phosphorus and sulfur [42]. Carbon, oxygen and hydrogen are supplied by the transformation of CO<sub>2</sub> and water as discussed above.

The macronutrient that has the most important effect (except carbon, hydrogen and oxygen) is nitrogen. Some microalgae species are more sensitive to the type of nitrogen that increases their lipid content [118,119]. As an example, using different sources of nitrogen (ammonium nitrate, ammonium chloride and urea) at a concentration of 0.64 mmol-N/L with the microalga *Ellipsoidion* sp., Xu *et al.* [118] obtained similar maximum growth rates (0.29 ± 0.02 1/h), but with lipid contents of 28, 33 and 22% (g lipid/g dry biomass) for ammonium nitrate, ammonium chloride and urea, respectively. In comparison, the absence of nitrogen gave a maximum growth rate of 0.17 1/h with a lipid content of 8% (g lipid/g dry biomass). In another study, nitrogen deprivation of a particular species (NAVIC1) increased the lipid content from 22 to 58% (g lipid/g dry biomass) [120]. A more recent study showed an enhancement of lipid content after nitrogen deprivation [121].

Some studies showed that increasing the phosphorus concentration has a positive effect on biomass productivity but few effects on lipid productivity. For example, by increasing the phosphorus concentration from 0.1 to 2 mg/L in the culture medium of the microalga *Scenedesmus* sp., Xin *et al.* [119] obtained an

increase of the biomass concentration from 0.14 to 0.37 g/L, while the lipid concentration remained stable at 0.08 ± 0.01 g/L. Consequently, for the range of phosphorus concentrations, the lipid content decreased from 53 to 24% (g lipid/g dry biomass).

Despite the fact that sulfur is an essential nutrient, few studies have been performed, to our best knowledge of the literature, in comparison to phosphorus or nitrogen. Giorgano *et al.* [122] tested the effect of sulfur concentrations from 6 μM to around 1.1 mM and obtained an increase of the maximum specific growth rate of the microalga *Dunaliella salina* from 0.006 to 0.02 1/h [123]. However, sulfur concentration can have an inhibitory effect on microalgae growth for concentrations higher than 1.1 mM [122,123].

Some microalgae species are also sensitive to salinity [124]. For example, by using a culture medium with salinity from 1 to 3 wt%, Renaud *et al.* [125] obtained an increase of the lipid content from 28 to 33% (g lipid/g dry biomass) for the microalga *Nannochloropsis oculata*. On the other hand, for the alga *Nitzschia (frustulwn)*, a similar increase of the salinity had an opposite effect, decreasing the lipid content from 33 to 29% (g lipid/g dry biomass).

Some other micronutrients are also important, such as iron, bore, manganese, copper, molybdenum, vanadium, cobalt, nickel, silica and selenium [84]. In order to decrease the production costs, some microalgae can be grown using waste water as a source of nutrients. Some species such as *Chlamydomonas globosa*, *Chlorella minutissima* or *Scenedesmus bijuga* can metabolize nitrogen, phosphorus, sulfur and metals (Al, Fe, Mn, Zn) [113,126,127].

#### • Temperature

Despite the weather conditions, the optimal conditions of temperature must remain between 20 and 30°C [17,91,128]. As the nutrient intake, the optimum temperature depends on the microalgae species [91,128]. For example, for the microalgae species *Nannochloropsis* sp., *Tetraselmis* sp. and *Isochrysis* sp., Abu-Rezq *et al.* [129] obtained maximum biomass yields with an ambient air CO<sub>2</sub> concentration (0.035% v/v) at respective optimal temperature ranges of 19–21, 19–21 and 24–26°C.

As the sun inevitably heats the culture medium, an external source of water must be provided to cool down the culture medium. For microalgae cultivated in ponds, cooling down is mainly done by evaporation [17]. As to the photobioreactors, cooling is performed with external input of water [130,131]. Consequently, culturing microalgae requires a non-costly source of water (river, sea, groundwater, etc.) whatever the type of bioreactor used.

## • pH

The optimum pH varies as a function of the microalgae species. An increase of CO<sub>2</sub> in the gas phase has the effect of decreasing the pH of the reaction medium, as CO<sub>2</sub> produces carbonic acid in aqueous phase. Consequently, when microalgae consume CO<sub>2</sub>, there is an increase of pH [84]. Using sodium hydroxide (NaOH) or hydrochloric acid for pH regulation and a temperature of 20°C, Bitaubé Pérez *et al.* [132] obtained an optimal pH of the culture medium between 8 and 9 for a specific growth rate of 7.2 1/h for the microalga *Phaeodactylum tricornutum*. However, the optimum pH varies as a function of the microalga. For example, by using microalgae strain ZY-1 (*Chlorella* sp.), Yue and Chen [112] obtained an optimal pH of 5 for a biomass growth rate of 0.08 g/L/h. They also observed few variations of the optimal biomass growth rate for pH variations between 4 and 6.

## • Stirring

Stirring is an important parameter, as the latter prevents microalgae from settling. Moreover, stirring improves the light energy and the gas transfer, and standardizes the temperature and the concentration of

nutrients [84,133]. Stirring can be performed mechanically or pneumatically. Another study tried to demonstrate (without success) that the superficial gas velocity could harm the microalgae *Dunaliella tertiolecta* and *Chlamydomonas reinhardtii* wild-type [134]. However, according to this study, the superficial velocity (0.076 m/s) alone cannot explain the death of microalgae.

## Harvesting and drying methods

Figure 1 presents the different methods of biodiesel production from microalgae. The different methods will be described in the following sections. When the microalgae are harvested, the water content may vary between 0.5 g/L for open systems for photoautotrophic microalgae [135] and 116 g/L for fermenters with heterotrophic microalgae [107]. In order to reduce the production costs, water must be reduced by using microsieve, filtration, centrifugation or flocculation (aluminium chloride, iron chloride (FeCl<sub>3</sub>), chitosan, organoclays etc.) or by sedimentation [82,91,136,137]. Despite the fact that some authors mention flotation as a harvesting method, this method remains technically and economically limited [91]. Otherwise, the harvesting costs can represent between 20 and 30% of the total production costs.




Figure 1. Process diagram of a biodiesel production process with all possible extraction methods.



Some companies stated that they have made major breakthroughs for harvesting and dehydration microalgae. As an example, Algaeventure Inc. states that they can increase microalgae solid content from 20 to 40 wt% of solid matter without forced drying by using a porous belt filtration system [138]. This company affirms that their process could reduce up to 90% of the cost linked to microalgae harvesting [139]. Evodos Inc. developed a harvesting process using a spiral plate technology (SPT), which could concentrate microalgae up to 32 wt% of solids, but few details have been released [140]. Other patents have been filed for the membrane used to concentrate the microalgae biomass [141].

### Drying

In order to increase the contact between the lipids and the solvent, microalgae are sometimes dried. Several techniques have been tested including freeze [142], oven [143], spray [142] and vacuum dryings [144,145]. Furthermore, the techniques with low pressures (less than 10 kPa) such as freeze-drying are often seen as too costly at an industrial scale. These techniques allow direct transesterification because water reduces the FAME yield, as water hydrolyzes the FAME [146]. Some patents state that hot air, between 60 and 90°C, could be used to dry microalgae [147]. On the other hand, drying microalgae at these temperatures for a long period of time could induce the degradation of some microalgae components such as the peroxydation of unsaturated lipid double bounds [45]. Consequently, this type of drying is more appropriate for microalgae applications where drying is essential, such as gasification, pyrolysis or thermal oxidation. Drying from solar energy could be energetically interesting, but few studies to our best knowledge have tested this kind of drying, which could only be used during hot, dry, sunny days.

On the other hand, removing water from microalgae by evaporation implies higher costs as the lipid extraction process require 2.5 times more energy than does a process without water removal [148]. According to these authors, the biodiesel production process implying a drying step would mean a negative energy balance (−2.6 MJ/kg biodiesel) with 82 MJ/kg biodiesel linked to the drying of microalgae biomass. As the energy produced by the biodiesel is around 41 MJ/kg biodiesel [149], using drying techniques seems to be impossible for now.

## Lipid extraction methods

### Methods using solvents

There are two main types of solvents used to extract lipids from microalgae: organic solvents and supercritical fluids.

- Organic solvents

Extraction methods using chloroform have been known to extract and evaluate the total lipids for a long time [150]. The 'Bligh and Dyer' method [151] using a blend of chloroform-methanol-water (1:1:0.5 v/v) is one of the most used methods to determine the total lipids in biological samples [152]. This extraction method is used directly [153] or with mechanical or thermal pretreatments [142,154] in order to extract the total lipids.

The order of blending solvents (chloroform, methanol or water) with microalgae biomass also has an influence on the extracted lipids yield [143,154]. As an example, Lewis *et al.* [153] obtained a FAME yield that increased from 26 to 33% (g lipid/g dry biomass) by inverting water and chloroform in the sequence water-methanol-chloroform. They explained this difference by the fact that a non-polar solvent added first could weaken the link between the lipids and the cell walls of microalgae.

Nevertheless, using chloroform is risky because this solvent could have carcinogenic effects on the human being [155]. With fewer toxic effects, hexane is sometimes preferred to chloroform [156]. However, because of the non-polar nature of hexane, the lipid yields are lower, since microalgae often have a high content in more polar lipids such as glycolipids and phospholipids [42]. Moreover, using toxic organic solvents to extract lipids (chloroform, hexane, etc.) prevents or makes more difficult the valorization of microalgae for food purposes. For microalgae lipid extraction, some studies have attempted to replace the chloroform-methanol extraction (2:1, v/v) by different blends such as hexane-isopropanol (3:2, v/v), dichloroethane-methanol (1:1, v/v), dichloroethane-ethanol (1:1, v/v) and acetone-dichloromethane (1:1, v/v), but the lipid yield obtained was significantly lower [157].

Samorì *et al.* [158] used polar solvents such as ethanol, octanol or 1,8-diazabicyclo-[5,4,0]-undec-7-ene (DBU). In this study, despite the fact that the lipid yield obtained was almost twice as high (16 g lipid/g dry biomass), the biodiesel yield was 5 times lower (0.65% g biodiesel/dry biomass) than for hexane extraction. A patent describes a process for proteins and sugar extraction from microalgae in aqueous phase using organic solvents (acetone, methanol, ethanol, isopropanol, methyl ethyl ketone, dimethylether or propionic aldehyde) which make the lipid extraction easier [159]. This process includes: (1) centrifugation of the aqueous microalgae, (2) blending microalgae with an organic solvent, (3) centrifugation of the organic solvent/humid microalgae blend, (4) heating the organic solvent/humid microalgae blend, (5) organic solvent separation, (6) evaporation of the organic solvent and (7) separation of the lipids from the aqueous phase. On the other hand, the heating phase includes high temperatures (50 to 150°C) and pressures (0.5 to 3 MPa),

which considerably increase the extraction costs. However, in this patent, no yields were specified.

- Supercritical carbon dioxide

As some organic solvents imply risks to human health, CO<sub>2</sub> supercritical extraction was considered to recover lipids from microalgae such as polyunsaturated fatty acids for food and pharmaceutical purposes [160–162] and for biodiesel [156,163,164]. Otherwise, a patent describing a process using supercritical CO<sub>2</sub> to extract lipids from microalgae to produce biodiesel was filed by the China Green Oil Co Ltd (Hong Kong, China) [165]. Several companies still hesitate to use supercritical extraction at an industrial scale based on the high investment costs [166] and the high pressures required [156,160,167]. Still, some studies indicate that this technology has a high potential to extract lipids from microalgae, even when the latter are still wet [156]. Using supercritical CO<sub>2</sub> to extract lipids from the microalga *Nannochloropsis* sp., Andrich *et al.* [160] obtained a lipid yield of 26% (g lipid/g dry biomass) at a temperature of 40°C and a pressure of 70 MPa.

### Physicochemical methods

- Organic solvents

As microalgae cell wall is a barrier to lipid extraction, physicochemical methods are used to increase the lipid content and the biodiesel obtained. Tables 1 and 2 present the lipid and the biodiesel yields, respectively, for different microalgae lipid extraction studies. Among the pre-treatments used, the main methods are microwave, ultrasound, bead-beating, freeze-drying, homogenization, French press, crushing, autoclave and osmotic shock. Other researchers own a patent for a cavitation system implying pressures from 0.3 to 30 MPa allowing microalgae cell walls to break up even if no extraction yield was specified in the patent filed [168,169].

Some pre-treatments such as microwave and bead-beating seem to be effective to increase the lipids extracted from microalgae when the latter is in aqueous phase [142,154,157]. As an example, after a pre-treatment by bead beating of the microalga *Botryococcus braunii* UTEX 572 in aqueous phase, Lee *et al.* [157] obtained an increase in the lipids extracted from 14 (without pre-treatment) to 29% (g lipid/g dry biomass) (bead beating). At last, the company Solazyme Inc. (South San Francisco, USA) owns a patent for a process based on lysis of microalgae cell walls and the use of an organic solvent for the lipid separation [170]. For example, the patent mentioned cell lysis with heating at temperatures ranging from 50 to 130°C, the addition of a strong alkali such as potassium hydroxide (KOH) or NaOH, a strong acid such as sulfuric acid (H<sub>2</sub>SO<sub>4</sub>), a

mechanical lysis (1500 bar with a shrinkage), an osmotic shock, and the use of ultrasound and a pressure oscillation. Among others, they used KOH and the H<sub>2</sub>SO<sub>4</sub> concentrations that varied from 0 to 160 mN with temperatures ranging from 25 to 130°C. Furthermore, Dillon *et al.* [170] also used several surfactants such as NINOL<sup>®</sup> 201 (Oleamide DEA from Stepan Company) to extract lipids of the heterotrophic microalgae *Phlorella protothecoides* at a temperature of 55°C for 5 h and obtained lipid yields of up to 92% (based on the theoretical mass of lipids).

Lyophilization is mainly used in studies focusing on solvent extraction in order to obtain a better contact between a non-polar solvent and the lipids. Other studies tried to show that lyophilization could break the cell walls of microalgae and increase the lipid yield. For example, Lewis *et al.* [153] obtained an increase of the lipid yield extracted from the microalga *Botryococcus braunii* UTEX 572 from 14 to 19% (g lipid/g dry biomass) after lyophilization.

Some techniques used with solvent extraction allowed an increase in the amount of lipids extracted. As an example, using ultrasounds on a blend of chloroform-methanol (3:1, v/v), the lipid yield obtained from the microalga *Scenedesmus* sp. was increased from 2 to 6% (g lipid/g dry biomass) compared with no treatment [171]. However, another study used ultrasounds and observed no significant effect on the lipid yield with a blend of chloroform-methanol-water (1:2:0.8, v/v) for a heterotrophic microalga (strains ACEM 6063 and ACEM A) [153].

Despite the fact there are few studies using electric energy to extract lipids from microalgae, some patents affirm that this technique could be used to reduce microalgae extraction cost. As an example, Diversified Technologies, Inc. (Bedford, MA, USA) stated that their technology of pulsed electric field (PEF) could reduce the extraction costs from 0.46 US\$/L oil for a conventional drying to 0.03 US\$/L oil [172].

- Solvent-free and ionic liquid extraction

In order to eliminate some steps in the lipid recovery, a separation of lipids in aqueous phase could be performed based on the difference of specific gravities and solubilities of both lipids and water. The main advantage of the lipid extraction in aqueous phase is to allow the valorization of residual biomass in agriculture. To our best knowledge of the literature, no studies have demonstrated the feasibility of this type of extraction for microalgae lipids. On the other hand, a patent from Solazyme Inc. (South San Francisco, USA) affirms that it is possible to recover lipids directly in aqueous phase following cell wall lysis due to heating (higher than ambient temperature) and an acid pre-treatment, and then a recovery of lysate with a decrease of the temperature (20°C, 24 h) and a

Table 1. Physico-chemical pre-treatments effect on lipid yield for several studies.

Microalgae species	Initial state	Pre-treatment	Parameters	Solvents	Max lipid yield (% g dry biomass)	FAME yield (% g FAME/ g dry biomass)	Ref.
<i>Botryococcus</i> sp.	Lyophilized	None Autoclave Bead beating Microwave Ultrasound Osmotic shock	None 125 °C – 1.5 MPa 0.1 mm – 2800 rpm 100 °C – 2450 MHz – 5 min 10 kHz – 5 min Sodium chloride (NaCl) 10% (m/v) – vortex (1 min) – 48 h	Water-chloroform-methanol (2:1:1 v/v/v)	7.7 12 28 29 8.5 11	1.2	[142]
<i>Chlorella vulgaris</i>		None Autoclave Bead beating Microwave Ultrasound Osmotic shock	None 125 °C – 1.5 MPa 0.1 mm – 2800 rpm 100 °C – 2450 MHz – 5 min 10 kHz – 5 min NaCl 10% (m/v) – vortex (1 min) – 48 h		5.1 9.6 7.9 9.8 5.9 8.0	2.9	
<i>Scenedesmus</i> sp.		None Autoclave Bead beating Microwave Ultrasound Osmotic shock	None 125 °C – 1.5 MPa 0.1 mm – 2800 rpm 100 °C – 2450 MHz – 5 min 10 kHz – 5 min NaCl 10% (m/v) – vortex (1 min) – 48 h		1.8 5.5 8.3 10 7.3 6.8	1.9	
<i>Botryococcus braunii</i> UTEX 572	Refrigerated (4 °C)	None Ultrasound Homogenization French press Lyophilization Bead beating	None 1 min 1 min 170 atm Not specified 1 min – 1 mm	Chloroform-methanol (2:1 v/v)	14 16 15 23 19 29 20 19 19 18	Unspecified	[157]
<i>Scenedesmus</i> sp.	Oven dried at 45 °C (4–5 days)	None Ultrasound None Ultrasound (chloroform-methanol)	None 100 W – 30 min None 100 W – 30 min	Hexane-isopropanol (3:2 v/v) Dichloroethane: methanol (1:1 v/v) Dichloroethane: ethanol (1:1 v/v) Acetone: dichloromethane (1:1 v/v) Soxhlet with n-hexane Chloroform-methanol-water (3:1:1.2 v/v/v)	0.70 0.80 2.0 6.0	Unspecified	[171]
<i>Phaeodactylum tricomutum</i>	Frozen	Extraction time	1 <sup>st</sup> extraction: 600 min 1 <sup>st</sup> extraction: 600 min – 2 <sup>nd</sup> extraction: 90 min 1 <sup>st</sup> extraction: 600 min – 2 <sup>nd</sup> extraction: 90 min – 3 <sup>rd</sup> extraction: 80 min	Ethanol-water-hexane (1.6: 1.9:1 v/v/v) Hexane (4 × 32 mL)	5.0 6.2 6.3	Unspecified	[258]
<i>Botryococcus braunii</i>	Refrigerated	None Temperature (10 min)	None 75 °C 80 °C 85 °C 90 °C 120 °C	Hexane and methanol: chloroform (1:1 v/v) Hexane	37 0.15 1.2 32 36 36	Unspecified	[259]
<i>Botryococcus braunii</i>	Lyophilized	None	None	Hexane chloroform-methanol (2:1 v/v) 3 mL 1,8-diazabicyclo[5.4.0]-undec-7-ene (DBU) DBU-ethanol (1:0.39 v/v) 2 mL DBU/octanol (1:1 v/v) 3 mL octanol	8.0 15 16 13 1.2 4.3 0.64	2.7 0.54 0.65 0.59 Unspecified	[158]
	Fresh	Centrifugation	300 rpm – 2 h 300 rpm – 24 h 3000 rpm – 4 h 300 rpm – 2 h	3 mL n-hexane			(continued)

Table 1. (Continued)

Microalgae species	Initial state	Pre-treatment	Parameters	Solvents	Max lipid yield (% g lipid/ g dry biomass)	FAME yield (% g FAME/ g dry biomass)	Ref.		
<i>Nannochloropsis</i> sp.	Lyophilized	None	300 rpm – 24 h	2 mL DBU/octanol (1:1 v/v)	5.6	Unspecified	[160]		
			3000 rpm – 4 h		5.1				
			300 rpm – 2 h		2.4				
			300 rpm – 24 h		8.2				
			3000 rpm – 4 h	Supercritical CO <sub>2</sub>	6.9				
			40°C – 70 MPa		26				
			40°C – 55 MPa		25				
			40°C – 40 MPa		24				
			55°C – 70 MPa		26				
			55°C – 70 MPa		25				
<i>Chlorococcum</i> sp.	Oven dried (85°C) for 16 h and ground	None	55°C – 70 MPa	Hexane	25	Unspecified	[156]		
			55°C – 70 MPa		24				
			Soxhlet	Hexane-isopropanol (3:2, v/v)	4.8				
			Agitation						
			Agitation	Hexane	1.5				
			Soxhlet (5.5 h)		3.2				
			Soxhlet (7.5 h)	Supercritical CO <sub>2</sub>	5.7				
			80°C – 30 MPa		4.8				
			60°C – 30 MPa		5.8				
			60°C – 40 MPa		5.9				
60°C – 50 MPa	6.0								
60°C – 30 MPa	7.1								
Fresh									

FAME: fatty acid methyl esters.

FAME: fatty acid methyl esters.

centrifugation (4400 rpm) [170]. Moreover, they also noticed that the size of the emulsion increased from 0 to 30% when the temperature increased from 25 to 130°C. Even if they stated that a small amount of organic solvent (as small as 5% (v/v)) could be used, the patent does not give the lipid extraction yield. In the same sort of idea, other researchers filed to patent a process using acid hydrolysis (using H<sub>2</sub>SO<sub>4</sub>) of microalgae (100°C, 200 kPa) in aqueous phase including mechanical and chemical techniques in order to hydrolyze the lipids into free fatty acids (FFAs) and they esterified the FFAs in aqueous phase with methanol and H<sub>2</sub>SO<sub>4</sub> [173]. The problem with this kind of technique is that the excess water could give really low lipid (triglycerides, FFAs and others) conversions because water favors the reverse reaction (hydrolysis) [174–176].

The company Alfa Laval Lund AB (Lund, Sweden) filed for a patent for a centrifugation separator with disks, which is operated using a centrifuge force between 4500 and 5000 *g* that allows separating oil and microalgae in aqueous phase [177].

Following the pre-treatments mentioned previously, ionic liquids could be used to recover lipids [178,179]. As an example, a patent described a separation process using ionic solvents such as hydronium (H<sub>3</sub>O<sup>+</sup>), ammonium (NH<sub>4</sub><sup>+</sup>), alkyl ammonium, imidazolium and heterocycles (cycle of carbon with another element in the cycle) [178]. The ionic solvent is then recovered with an antisolvent such as inorganic salts (like calcium chloride (CaCl<sub>2</sub>)). However, the process allows recovering at least 80% of the ionic liquid, which might predict problems with the recovery of the ionic liquid. Moreover, the antisolvent has to be immiscible with the lipids.

Other patents reported the use of resins for the recovery of microalgae lipids. For example, Poenie *et al.* [180] filed a patent concerning a technique that extracts lipid from the microalga *Chlorella* sp. using resins (CHLOR 13A, NANN 13A, CHLOR 14B, CHLOR JB21, NANN JB21). The types of resins in this patent are made from polymers such as polystyrene, polyethylene, etc. After the saturation of the resin exchange column, water is evaporated using, for example, air streams. Then, an eluent, an alcohol such as methanol containing an acid (H<sub>2</sub>SO<sub>4</sub>) or an alkali (NaOH, KOH, sodium ethoxide (C<sub>2</sub>H<sub>5</sub>ONa), potassium methoxide (KCH<sub>3</sub>O)), is fed to the column to recover microalgae lipids.

Other researchers have developed processes for breaking microalgae cell walls by mechanical techniques. For example, Thomas *et al.* [181] own a patent for a vertical ram-type extractor which involves a mass of 5.6 kg dropped with a final velocity of 8 m/s in order to break microalgae cell walls. Echevaría Parres *et al.* [182] filed a patent in which microalgae are dragged into a turbine and then crushed between a rotor and a

Table 2. Physico-chemical pre-treatments effect on the FAME yield for several studies.

Microalgae	Initial state	Pre-treatment	Parameters	Solvent	FAME yield (% g FAME/g dry)	Reference
<i>Nannochloropsis</i>	Fresh	Oven dried and ground	Extraction time: 20h	Chloroform-methanol-water (1:2:0.8 v/v/v)	21.5 ± 4.3	[143]
		Oven dried and ground	Extraction time: 2h		21.3 ± 2.3	
		Oven dried	Extraction time: 2h		14.8 ± 6.8	
		Oven dried and frozen with liquid nitrogen	Extraction time: 2h	Water-methanol-chloroform-(0.8:2:1 v/v/v)	20.1 ± 0.5	
		Oven dried and frozen with dry ice	Extraction time: 2h		18.7 ± 3.2	
		Oven dried and ground	Extraction time: 2h		18.5 ± 0.5	
<i>Chlorella protothecoides</i>	Fresh	No	No	Water-methanol-chloroform-(0.8:2:1 v/v/v)	16.4 ± 2.1	[153]
		No	No		18.5 ± 0.5	
		Oven dried	No		21 ± 1.2	
		None	None	Chloroform-methanol-water (1:2:0.8 v/v/v)	25.9	
		Ultrasound in the solvent	100 W - 2 × 1 min		26.3	
		None	None		35	
Strain ACEM 6063	Lyophilized	Ultrasound in the solvent	100 W - 2 × 1 min	Chloroform-methanol-water (1:2:0.8 v/v/v)	34.3	[203]
		None	None		33.2	
		Ultrasound in the solvent	100 W - 2 × 1 min		34.6	
		None	None	Bligh and Dyer n-Hexane	19.9	
		Ground	No details		6.9	
		Ground and supercritical CO <sub>2</sub>	20 MPa - 313 K		6.3	
<i>Cryptocodinium cohnii</i>	Lyophilized	None	25 MPa - 313 K	Dichloromethane-methanol (2:1)	5.7	[260]
		None	30 MPa - 313 K		5.6	
		None	20 MPa - 323 K		7.1	
		None	25 MPa - 323 K	6.5 mg EMIM - methanol (1.2:1 w/w)	8.6	
		None	30 MPa - 323 K		17	
		None	None		33.5	
<i>Scenedesmus obliquus</i>	Lyophilized mortar	None Quartz sand - 0.2 g (10-30 mm) ultrasound - 35 kHz, 80 W, -4°C, 90 min	None			
<i>Chlorella</i> sp. <i>Dunaliella</i> sp.	Lyophilized	None	No	6.5 mg EMIM - dimethyl sulfoxide 6.5 mg EMIM - acetic acid 6.5 mg EMIM-acetone 6.5 mg EMIM - chloroform 6.5 mg EMIM - isopropyl alcohol	38	[261]
		1-ethyl-3-methyl imidazolium, methyl sulfate (EMIM) and a co-solvent	65°C - 18 h - stirring - molar excess of EMIM		8.6	
			co-solvent of 10 compared to EMIM		6.0	
					5.6	
					9.2	
					8.4	
<i>Chlorella</i> sp.		Enzymatic hydrolysis solution 20 g dry/L	Immobilized polyacrylonitrile cellulase	Hexane (28°C, 1-6 h)	53	[184]
		None	None		52	

FAME: fatty acid methyl esters.



stator. Zeiler *et al.* [183] filed a patent for a laser ultra short pulse (USP) producing ultraviolet of short wavelength (less than 400 nm) able to break microalgae cell walls. On the other hand, most of these patents practically did not report any lipid yield.

### Biochemical methods

In order to break the link between lipids and microalgae, biological pre-treatments can be used. As an example, using a hydrolysis pre-treatment of microalgae cell wall with cellulase, Fu *et al.* [184] obtained an increase of 70% of microalgae cell wall hydrolysis (measured by monosaccharide analysis) despite the fact that the lipid yield increased only from 52 to 53% (g lipid/g dry biomass). The second problem is that hydrolysis is performed in water phase and lipids must be extracted. In order to break microalgae cell wall, the company Solazyme Inc. (South San Francisco, USA) filed for a patent which mentions using enzymes such as hemicellulase, pectinase, cellulase and driselase, and also infections of microalgae by viruses (e.g. *Paramecium bursaria* chlorella virus) and autolysis (expression of a lysis gene) in order to degrade microalgae cell walls [170]. Another patent was filed using autolysis [185] or enzymes [186] to break the microorganism cell walls of microalgae and recover the lipids with, among other things, an organic solvent.

### Lipid purification

When the lipids are extracted, they contain various lipids, which result in a lower biodiesel yield during transesterification rather than the triglycerides whose yield can reach near 100% [187]. As an example, Nagle and Lemke [188] have determined that biodiesel yield for phospholipids and glycolipids could not reach more than 65 and 56% (g FAME/g lipid), respectively. These are important data because microalgae can contain up to 93 wt% of these two components [42]. Furthermore, a study shows that microalgae could contain also added-value compounds such as chlorophyll and carotenoids [58].

According to our best knowledge of the literature, few lab-scale studies have separated the different classes of lipids present in microalgae in order to recover the triglycerides. However, Lin *et al.* [189] filed a patent for a process that allows the selective isolation of some components of microalgae. The process in batch mode could be described as follows: After an extraction and a recovery of lipids with an organic solvent, the organic phase is blended with a mesoporous solid (nanoparticles of mesoporous carbon, activated carbon, nanoparticles of mesoporous silicon (MSN) or silicium gel) and filtration allows the separation of the organic phase. For example, using MSN, the process allows the recovery of 90 wt% of the fatty acid oleate (C18:1) from lipids of the microalga *Neocloris* sp.

### Direct transesterification

Direct transesterification is a process in which the lipids are simultaneously extracted and transesterified from microalgae cell walls. Direct transesterification is often used for heterotrophic microalgae [65,146,153,190]. Some authors obtained higher biodiesel yields using direct transesterification than using extraction followed by transesterification [146,187]. This process could have the advantage of reducing the amount of solvent used to transesterify the lipids that could never be extracted. However, most of these studies used an organic co-solvent (hexane, chloroform, etc.) during the direct transesterification in order to increase the biodiesel yield [146,187]. For example, using a homogeneous acid catalyst ( $H_2SO_4$ ) with methanol, Johnson and Wen [146] transformed lipids of the microalga *Schizochytrium limacinum* into biodiesel with and without a co-solvent (hexane). After 40 min of reaction at a temperature of 90°C, they respectively obtained biodiesel yields of 66 and 7% (g biodiesel/g dry biomass). Then, some studies compared hexane and chloroform as a co-solvent and found out that these two co-solvents had similar biodiesel yields [65,146].

In order to increase the biodiesel yield, some studies tested direct transesterification with physicochemical methods such as microwave and ultrasound. For example, replacing traditional heating with microwave heating, Koberg *et al.* [187] performed a transesterification with strontium oxide (SrO) as a catalyst and obtained an increase of the biodiesel yield from 7 to 37% (g biodiesel/g dry biomass). Despite the advantage of direct transesterification, the energetic costs from the mandatory drying of microalgae make this technique difficult to use at an industrial scale [148].

### Transesterification

#### Reaction

Vegetable oils or microalgae lipids cannot be used directly in diesel engines as their viscosity is relatively high (28–40 mm<sup>2</sup>/s) which causes formation of coke deposits in the engine and the injectors [191]. Furthermore, using vegetable crude oils can have negative effects on the biofuel combustion. For example, Altin *et al.* [192], tested the behavior of colza oil at a revolution velocity of 1300 rpm. Despite the fact that they obtained a similar power (97%) compared to petrodiesel, they noticed an increase of (1) the fuel consumption from 246 to 288 g/kWh, (2) the CO concentration from 2225 to 4000 ppmv and (3) the fume density (particular matter) from 29 to 49%. Transesterification of lipids allows reducing oil viscosity to a level similar to that of petrodiesel [191,193]. Despite the fact that most studies consider only a reaction between an oil (triglycerides) and an alcohol, biodiesel can be

**Table 3.** Studies on biodiesel production from microalgae and the yield obtained.

Microalgae species	Catalyst	Reaction time (min)	Temperature (°C)	Quantitative analysis	Biodiesel yield (% g FAME/g lipid)	Biodiesel yield (% g FAME/g dry biomass)	Ref.
<i>Chlorococcum</i> sp.	H <sub>2</sub> SO <sub>4</sub> /KCH <sub>3</sub> O	120/120	50/55	Gas chromatography (GC)	44	1.5	[156]
<i>Nochloropsis</i>	Strontium oxide (SrO)	2	60	<sup>1</sup> H NMR Spectroscopy	99.9	37.1	[187]
<i>Schizochytrium limacinum</i>	H <sub>2</sub> SO <sub>4</sub>	40	90	GC	73	42	[146]
<i>Chlorella mulleri</i>	HCl NaOH	6	70	Thin layer chromatography (latroscan)	68 1.3	Unspecified	[188]
<i>Chlorella protothecoides</i>	H <sub>2</sub> SO <sub>4</sub>	Unspecified	Unspecified	Elemental analyzer CE-440	68	37.4	[149]
<i>Botryococcus</i> sp.	KOH/HCl	10/10	75	GC	4.2	1.2	[142]
<i>Chlorella vulgaris</i>					29.6	2.9	
<i>Scenedesmus</i> sp.					19.8	1.9	
Microheterotrophic (strains ACEM 6063)	HCl	60	90	GC	Unspecified	41.3	[153]
<i>Botryococcus braunii</i>	NaOH/bore trifluoride (BF <sub>3</sub> )	10/2	Reflux	GC	21.6	2.7	[158]
<i>Nannochloropsis</i>	HCl	Unspecified	Unspecified	GC and <sup>1</sup> H NMR spectroscopy	Unspecified	8.4	[143]
<i>Nannochloropsis</i>	H <sub>2</sub> SO <sub>4</sub>					6.6	
<i>Chlorella protothecoides</i>	HCl					36.6	
<i>Chlorella protothecoides</i>	Acetyl chloride					56	
<i>Cryptocodinium cohnii</i>	Acetyl chloride	60	80	GC	Unspecified	19.9	[203]
<i>Scenedesmus obliquus</i>	H <sub>2</sub> SO <sub>4</sub>	240	70	GC	Unspecified	33.5	[260]
<i>Chlorella</i> sp.	BF <sub>3</sub>	15	100	GC	Unspecified	53	[184]
<i>Botryococcus braunii</i>	HCl	60	90	GC	Unspecified	5.5	[262]
	Acetyl chloride	60	100			12.1	
<i>Synechocystis</i> sp.	HCl	60	90			7.3	
	Acetyl chloride	60	100			6.0	
<i>Nochloropsis occulata</i>	Calcium oxide (CaO)/aluminium oxide (Al <sub>2</sub> O <sub>3</sub> ) CaO/Al <sub>2</sub> O <sub>3</sub>	240	50	GC	23 16	Unspecified	[144]
<i>Chlorella pyrenoidosa</i>	H <sub>2</sub> SO <sub>4</sub>	120	90	GC	93	52	[65]
<i>Chlorella protothecoides</i>	<i>Candida</i> sp.	720	Unspecified	GC	98	Unspecified	[145]
<i>Chlorella protothecoides</i>	Lipase (triacylglycerol acylhydrolase)	720	38	GC	98	Unspecified	[194]

FAME: fatty acid methyl esters.

produced from several classes of lipids including glycolipids, phospholipids, FFAs, etc.

Table 3 presents FAME yields and experimental conditions of transesterification of microalgae lipids into biodiesel. According to Table 3, the biodiesel yields (FAME) and the transesterification methods vary. The microalga *Chlorella pyrenoidosa* cultivated in heterotrophic conditions resulted in the highest FAME yield (52% (g FAME/g dry biomass)), which represents a FAME yield of 93% (g FAME/g lipid). Some studies mention biodiesel yields (compared to lipid) of 98% (g FAME/g lipid) using diverse heterotrophic metabolism microalgae [145,194], which can be compared to the yields obtained with vegetable oil transesterification [195].

### Alcohol

The most used alcohol for transesterification is methanol because of its lower cost [17], but other alcohols such as ethanol, propanol and butanol can produce similar biodiesel yields [196,197]. Some studies have tested the effect of the methanol-triglyceride molar

ratio (between 56:1 and 524:1) for different microalgae on the density of the biodiesel produced [149,198]. For example, by transforming the triglycerides of the microalga *Chlorella* sp. into FAME for 8 h at a reaction temperature of 25°C and increasing the methanol-triglyceride molar ratio from 105:1 to 524:1, Ehimen *et al.* [198] observed a decrease of the density of the biodiesel produced from 889 to 886 kg/m<sup>3</sup>. On the other hand, the authors did not specify whether this increase was significant. Methanol-triglyceride molar ratio also has an impact on the biodiesel yield because a molar excess of the alcohol (methanol) reduces the reverse reaction and shifts the equilibrium of FAME production [199]. Performing an acid transesterification (H<sub>2</sub>SO<sub>4</sub>) of triglycerides of the microalga *Chlorella protothecoides* at 50°C for 4h, Miao and Wu [149] varied the methanol-triglycerides molar ratio from 25:1 to 84:1 and obtained an optimum methanol-triglycerides molar ratio at 45:1 for a FAME yield of 69% (g FAME/g triglycerides). The molar ratios of methanol-triglyceride used for microalgae lipid transesterification are generally higher than in the case of vegetable oil which is

usually around 6:1 [200] (the stoichiometric molar ratio is 3:1 [201]).

### Reaction time

Generally, increasing the reaction time of transesterification has a positive influence on the FAME yield. As an example, using a homogeneous enzymatic transesterification of lipids of the microalga *Chlorella pyrenoidosa* at a temperature of 90°C, Li *et al.* [65] obtained a FAME yield ranging from 52 to 94% (g FAME/g lipid) following an increase of the reaction time from 0.5 to 2 h. Reaction time has an influence on the density of the biodiesel. As an example, using a homogeneous acid catalyst (H<sub>2</sub>SO<sub>4</sub>), a stirring rate of 160 rpm, and a methanol-triglyceride molar ratio of 56:1, Miao and Wu [149] observed a decrease in the density of the biodiesel produced from 0.919 to 0.864 kg/m<sup>3</sup> for respective reaction times of 0 and 4 h.

### Homogeneous catalysis

Despite the fact that homogeneous alkali catalysis using NaOH is widely used for vegetable oil transesterification (because it is 4000 times faster than homogeneous acid catalysis), homogeneous alkali transesterification is rarely used for microalgae lipid transesterification. This catalyst gives a biodiesel yield 50 times lower than that of a homogeneous acid catalyst (like HCl) [188]. Sometimes, homogeneous alkali catalysts such as KCH<sub>3</sub>O or NaOH are coupled to acid catalysts [142,156,202].

As shown in Table 3, homogeneous acid catalysts are mostly used in the transesterification of microalgae lipids. The main catalysts are HCl [143,149], H<sub>2</sub>SO<sub>4</sub> [143,146,202], acetyl chloride (CH<sub>3</sub>COCl) [154,203] and boric trifluoride (BF<sub>3</sub>) [158,184]. Some studies have shown that CH<sub>3</sub>COCl in methanol (5% v/v) is the catalyst which resulted in the highest FAME yield of 56% (g FAME/g dry biomass) for the heterotrophic microalgae *Chlorella protothecoides* compared to a yield of 37% (g FAME/g dry biomass) for HCl [143]. Catalyst concentration also has an effect on the transesterification yield of microalgae lipids. As an example, using four concentrations of H<sub>2</sub>SO<sub>4</sub> in methanol (0.56, 1.13, 1.35 and 2.25 mol/L), a temperature of 30°C and a reaction time of 5 h, Miao and Wu [149] mentioned that a catalyst concentration in methanol of 2.25 mol/L produces biodiesel with the lowest density (863 kg/m<sup>3</sup>) but with the lowest FAME yield at 38% (g FAME/g dry biomass).

Finally, other techniques that imply the use of catalysts with supercritical solvents such as methanol were tested, but the costs of this technology are relatively high [167]. To our best knowledge of the literature, this technique has not been tested on microalgae lipids.

### Heterogeneous catalysis

In order to reduce the amount of chemicals used, some studies consider using heterogeneous catalysts (solids). Moreover, some of these catalysts have less sensitivity to water [204] and can potentially be reused [205]. As for homogeneous catalysts, there are heterogeneous alkali and acid catalysts.

#### • Alkali

Microalgae lipids have been transformed into biodiesel by using heterogeneous alkali catalysts such as calcium oxide (CaO) [144], strontium oxide (SrO) [187] and mixed magnesium-zirconium oxides [206]. Some of these catalysts led to high biodiesel yields. As an example, performing a direct transesterification with a heterogeneous catalyst of SrO, Koberg *et al.* [187] achieved a conversion of 100% of *Nannochloropsis* sp. microalgae triglycerides.

Table 4 presents the results of studies for vegetable oil using methanol (as an alcohol) and heterogeneous catalysts.

#### • Acid

Few studies have used acid heterogeneous catalysts to transform microalgae lipid into biodiesel. As an example, Carrero *et al.* [207] compared two types of zeolites ( $\beta$  and ZSM-5) to transform microalgae (*Nannochloropsis gaditana*) lipids into biodiesel. However, at a temperature of 115°C (without specified reaction time, methanol-lipid ratio or catalyst-to-oil ratio) under reflux, they obtained a FAME content of 25% (g FAME/g lipid) compared to 90% (g FAME/g lipid) with H<sub>2</sub>SO<sub>4</sub> under the same reaction conditions.

As shown in Table 4, the most utilized heterogeneous acid catalysts used in transesterification studies are zeolites, heteropoly acids, functionalized (sulfate and tungsten) silica or zirconia, ion-exchanging resins, organo-sulfated acids (sulfonic, sulfonates and sulfonamides) functionalized silica [208]. Even if, to our best knowledge of the literature, most of these catalysts have not been tested with microalgae lipids, they show a great potential for microalgae biodiesel. However, heteropoly acids could be interesting as catalysts to transform microalgae lipids (high levels of FFAs) into biodiesel, as these catalysts showed almost no sensitivity to high FFA contents (up to 10 wt%), but are sensitive to water contents (water content higher than 0.1 wt%) [209].

On the other hand, heterogeneous acid catalysts have the disadvantage of ending up in lower lipid conversions or longer reaction times. As an example, by using a catalyst of sulfated zirconium oxide (0.86 wt%) and a molar ratio of 2-ethyl hexanol-dodecanoic acid of 25:1 at a temperature of 80°C, Omota *et al.* [210]

**Table 4.** Heterogeneous catalysts used for the transesterification of several oils.

Oil	Catalyst	Catalyst/oil (% w/w)	Temperature (°C)	Time (h)	Molar ratio (alcohol/oil)	Biodiesel yield (% g FAME/g oil)	Reference
Cotton seed	Magnesium oxide (MgO)/MgAl <sub>2</sub> O <sub>4</sub> /γ-Alumina (Al <sub>2</sub> O <sub>3</sub> )	1	180	15	6:1	36	[263]
	Mg–Al–CO <sub>3</sub> hydrotalcite					98	
	Carbon-based (acid)	0.3	220	3	21	89	[264]
<i>Jatropha curcas</i>	KSF clay	5	160	6	12:1	68	[265]
Palm	Potassium fluoride (KF)/hydrotalcite	3	65	5	12:1	92	[266]
	KF/Ca–Al hydrotalcite	5				98	[267]
	K–Mg–Al hydrotalcite	7	100	6	30:1	87	[268]
	CaO–Zinc oxide (ZnO)	10	60	1	20:1	94	[269]
Rapeseed	Mg–Al-hydrotalcites	1.5	65	4	6:1	91	[270]
Soybean	KF/CaO	3	60–65	1	12:1	90	[271]
	Al–Mg-hydrotalcite	5	180	1	12:1	93	[272]
	MgO					91	[273]
	NaX zeolite/KOH	3	65	8	10:1	86	[274]
	ZnO /strontium nitrate (Sr(NO <sub>3</sub> ) <sub>2</sub> )	5	65	1	12:1	87	[275]
	Potassium iodine (KI)/Al <sub>2</sub> O <sub>3</sub>	2	Reflux	6	15:1	87	[276]
	KI/Zirconium oxide (ZrO <sub>2</sub> )					78	
	KI/ZnO					73	
	KI/zeolite NaX					13	
	KI/zeolite KL					18	
	Al–Mg-hydrotalcite	7.5	Reflux	9	15:1	67	[277]
	Potassium nitrate (KNO <sub>3</sub> )/Al <sub>2</sub> O <sub>3</sub>	6.5	Reflux	7	15:1	87	[278]
	CaO/chitosan	13.8	60	3	13.4:1	97	[279]
	CaO/ eggshell calcinated	1	70	5	6.9:1	97	[280]
	MgO	5	70	7	55:1	64	[281]
	Mg-hydrotalcite oxide					14	
	ZnMg-hydrotalcite oxide					7	
	ZnO					0	
	Al <sub>2</sub> O <sub>3</sub>					0	
	Ca/mesoporous silica	5	60	8	16:1	95	[282]
	KI/mesoporous silica	5	70	8	16:1	90	[283]
	CaO/lanthanum oxide (La <sub>2</sub> O <sub>3</sub> )	8	65	1	10:1	75	[284]
	KF/Al <sub>2</sub> O <sub>3</sub>	2	72.7	3	12:1	99	[285]
	Mg –Al- hydrotalcite	5	230	1	13:1	90	[286]
	CaO	0.8	Reflux	2	11.7:1	90	[287]
	CaO/La <sub>2</sub> O <sub>3</sub>	8	65	3	7.8:1	97	[288]
	CaO/Cerium oxide (CeO <sub>2</sub> )	5	58	1	20:1	91	[289]
	La <sub>2</sub> O <sub>3</sub> /zirconium oxide (ZrO <sub>2</sub> )					8.7	
	CaO/La <sub>2</sub> O <sub>3</sub>					94	
	MgO/MgAl <sub>2</sub> O <sub>4</sub>	3	65	10	15:1	57	[290]
	La/Zeolite β	1.1	60	4	14:5	49	[291]
	CuVOP	3	60	5	6:1	65	[292]
	Calcined sodium silicate	3	60	1	7.5:1	100	[293]
	Phosphazene hydroxide/SiO <sub>2</sub>	4.5	75	12	60:1	90	[294]
Sunflower	Zeolite NaX	10	60	7	6:1	95	[295]
	Al–Si/Potassium carbonate (K <sub>2</sub> CO <sub>3</sub> )	2	120	1	15:1	97	[296]
	CaO/SBA15	1	60	5	12:1	95	[297]
	Magnesium (Mg)/Calcium (Ca)	2.5	60	3	12:1	92	[298]
	Mg/Aluminium (Al)					66	
Vegetable	KF/CaO	3	65	1	12:1	99.6	[299]
	Aluminium hydrogen sulphate	0.5	220	0.83	16:1	81	[300]
	Carbon-based (acide)	0.2	220	4.5	17	81–95 (FFA)	[301]
Vegetable (ethanol as an alcohol)	None	2	220	5	6.5:1	37	[302]
	MCM-41 (mesoporus silica)					46	
	KIT-41					36	
	SBA-15					39	
	MgO/MCM-41					68	
	MgO/KIT-41					82	
	K/MgO (A)					96	
<i>Zanthoxylum bungeanum</i>	CaO	2.5	95	2.5	11.7:1	96	[303]

FAME: fatty acid methyl esters.

achieved a conversion of dodecanoic acid of 100% for a 70-h reaction time.

### Enzymatic catalysis

The main advantage of enzymatic catalysis is the relatively high lipid conversion. A study has shown that a conversion of 98% [194] can be reached by using a lipase (triacylglycerol acylhydrolase, EC 3.1.1.3, 1,200 U,

30 wt%) immobilized on a macroporous resin, a methanol-oil molar ratio of 3:1, a temperature of 38°C, a pH of 7 and a reaction time of 12 h. Despite the fact that some studies used enzymatic catalysis to transesterify heterotrophic microalgae lipids, this type of catalysis has not been tested up to now at industrial scale, perhaps because of the enzymes' high costs [17,211].

### Temperature and stirring

An increase of the temperature can have a positive effect on the biodiesel yield. As an example, using an enzymatic transesterification (homogeneous) of the microalga *Chlorella pyrenoidosa* lipids, for a reaction time of 2 h, Li *et al.* [65] reached biodiesel yields of 43 to 95% (g FAME/g lipid) at temperatures ranging from 20 to 110°C, respectively. However, for acid homogeneous catalysis, the temperature has less significant effects. As an example, testing temperatures of 30, 50 and 90°C for the acid catalysis of the heterotrophic microalga *Chlorella protothecoides*, Miao and Wu [149] obtained biodiesel yields of 56, 58 and 38% (g FAME/g lipid), respectively.

The stirring rate has a positive effect on the quality of the biodiesel produced. As an example, for the homogeneous acid ( $H_2SO_4$ ) transesterification of microalga *Chlorella* sp. for 2 h at 60°C by using no stirring and stirring at 500 rpm, Ehimen *et al.* [198] observed a decrease of the biodiesel density from 903 to 883 kg/m<sup>3</sup>.

### Biodiesel yield and purification

For the microalgae triglyceride transesterification, a conversion of 100% (g FAME/g lipid) can be reached [187], while for other classes of lipids, the yield is lower. For example, phospholipids and glycolipids are converted into biodiesel with yields of 65 and 56% (g FAME/g lipid), respectively [188]. These data are important because some microalgae can have an important amount of phospholipids and glycolipids of 40 and 25% (g/g lipid), respectively [42]. As a consequence, producing biodiesel from microalgae results in a lower biodiesel yield than with vegetable oil (mainly composed of triglycerides), and other compounds have to be separated (chlorophyll, carotenoids etc.) [58].

Biodiesel produced from microalgae must be separated from glycerol, chlorophyll and other by-products. Sometimes, when the process uses organic solvents, water is added to favor the biodiesel separation [146,212], as non-polar solvents are insoluble in water. In cases where there is no organic solvent already present, the main separation processes would be hot water [145], organic solvents such as hexane [156,213] and a blend of water and organic solvents [115,158,203]. Other techniques of purification, based on vegetable oil processes, that could be used for large-scale microalgae biodiesel purification would be: (1) water washing, (2) dry washing and (3) membrane separation.

### Production standards and properties

Currently, there are two main biodiesel production standards: American standard (ASTM D6751-10) and European standard (UNE-EN 14214:2010) [214–216].

The following standards are inspired from the previous two standards: Australia (Acte 2000), Brazil (ANP Act No. 42/2003), China (GB/T 2008-2007), India (IS 15607-2005), Indonesia (SNI 04-7182-2006), Japan, Korea, New Zealand (NZS 7500-2005) and Thailand [217]. In Canada, there is a law that allows adding up to 5% (v/v) of biodiesel into petrodiesel [218].

There are important differences between standards UNE-EN 14214:2010 and ASTM D6751-10 used to produce biodiesel from microalgae. The European standard stipulates that the content of linolenic FAME (C18:3) and polyunsaturated FAME (number of double bonds > 3) must not exceed 12 and 1 wt%, respectively [216]. These data are important since microalgae often have a high content of polyunsaturated FAMES [146,187] at levels that can reach 56 wt% [153]. Some requirements of UNE-EN 14214:2010 are not found in ASTM 6756-10 standards such as the minimum FAME content (96.5% mol/mol min), the density (860–900 kg/m<sup>3</sup>, 15°C) and the iodine value (120 g I<sub>2</sub>/100 g max). The important biodiesel properties are cetane index, heating value, cold properties, viscosity, oxidation stability and lubricity [219,220].

### Cetane index

The cetane index is the capacity for a fuel to flame up during combustion [221]. A higher cetane index means that the shorter time a fuel takes to flame up, the better the quality of the combustion [221,222]. The minimal cetane index for diesel is 40, while the latter for different commercial biodiesels varies between 48 and 65 [223]. For FAME, the cetane index increases with the increase in the number of carbon and decreases with the number of double bonds on the carbon chain. To our best knowledge of the literature, no study has experimentally measured the cetane index of microalgae biodiesel. However, a study has evaluated the cetane index of microalgae biodiesel from their FAME content: varying between 39 and 54 for different species of microalgae such as *Isochrysis galbana* (51.7), *Chaetoceros* sp. (50.6) and *Ondotella weissflogii* (39.3) [224].

According to our knowledge of the literature, no study has tried to increase the cetane number of microalgae biodiesel. On the other hand, some techniques applied to first-generation biodiesel could be used to increase the cetane number of microalgae biodiesel. In addition to hydrogenation, a study has tested nitration of biodiesel with nitric acid (25 wt%) and ethanoic anhydride (63 wt%). Using frying oil, they obtained an increase in the cetane index of the biodiesel from 55 to 61 [225].

### Heating value

Heating value is the amount of energy released during biodiesel combustion. For commercial biodiesel, lower heating values vary between 36 and 41 MJ/kg, while



the lower heating value of microalgae biodiesel produced from acid transesterification ( $\text{H}_2\text{SO}_4$ ) was measured at 41 MJ/kg by Miao and Wu [149].

### Cold properties

Cold properties are defined as the biodiesel behavior when temperature is decreasing. These properties include cloud point and pour point. Cloud point is the temperature at which crystals ( $d \geq 0.5 \mu\text{m}$ ) begin to form, while pour point is the temperature where biodiesel stops flowing [34,226]. As biodiesel has generally poorer cold behavior compared to petrodiesel (higher cloud and pour points), it is recommended to consider these properties for a biodiesel-petrodiesel blend [214]. Based on our knowledge of the literature, no study has measured cloud and pour points for microalgae biodiesel. However, since microalgae contain a high amount of unsaturated FAMES that can reach 56 wt% [153], microalgae biodiesel would have better cold properties than biodiesel produced from oleaginous vegetable species [34].

### Viscosity

Viscosity is a measure of flowing resistance. The lowest viscosity possible is required in order to reduce the flowing resistance. Despite the fact that the transesterification allows a reduction of the viscosity of lipids, the kinematic viscosity of biodiesel (2.8 to 5.7  $\text{mm}^2/\text{s}$ ) is higher than that of petrodiesel (1.8 to 3.8  $\text{mm}^2/\text{s}$ ) [223]. A biodiesel with a high viscosity will cause deposit problems in the combustion chamber and will also increase most pollutant emissions [227,228]. The kinematic viscosity of microalgae biodiesel ranges from about 3.9 to 5.2  $\text{mm}^2/\text{s}$  [146,149]. Furthermore, like the cetane index, the viscosity increases with the length of the carbon chain and decreases as a function of the unsaturation degree of the FAMES.

### Oxidation stability and biodegradability

Oxidation stability is a very important parameter when biodiesel is stored. Indeed, when the oxygen contained in the air comes into contact with biodiesel, the latter may be transformed into deposits which consist of hydrogen peroxides, aldehydes, acids and oxygenated products [229]. Generally speaking, oxidation increases as a function of the unsaturation degree. Consequently, some authors have evaluated that biodiesel produced from several species of microalgae would have a low oxidation stability because the polyunsaturated FAME content can reach 56 wt% [153], which represents a real problem for storage [224]. Otherwise, if biodiesel is stored for more than a few months, adding antioxidants is recommended to improve its oxidation stability [223,230,231].

Biodegradability is another important factor during storage because microorganisms have the ability to

transform biodiesel into end products such as  $\text{O}_2$ , water and  $\text{CO}_2$  [232].

### Lubricity

Lubricity of a fuel is defined as the capacity of a fuel to reduce friction between the moving parts of an engine [233–235]. Blending biodiesel with petrodiesel can improve the lubricity inside an engine. As an example, at a temperature of 25°C, Knothe and Steidley [236] obtained friction values (without indicated units) for petrodiesel (without additives) and vegetable based biodiesel of 0.238 and 0.117, respectively. Several factors influence the lubricity of petrodiesel, such as viscosity, acidity, water and sulfur contents [237]. Some studies on biodiesel production from microalgae obtained relatively high sulfur contents (69 ppm) [146] compared to American Society for Testing and Materials (ASTM) standard (15 ppm) [214]. As a consequence, studies on biodiesel lubricity should be performed since, to our best knowledge, no study has measured lubricity of biodiesel produced from microalgae.

### Other properties

Concerning the density of biodiesel, that of microalgae biodiesel is higher than petrodiesel. For example, Miao and Wu [149] produced biodiesel from microalgae and measured a biodiesel density of 864  $\text{kg}/\text{m}^3$  by performing a direct transesterification of a heterotrophic microalgae (*Chlorella protothecoides*) using methanol and an acid catalyst ( $\text{H}_2\text{SO}_4$ ), while the standard for petrodiesel is 0.838  $\text{kg}/\text{m}^3$ .

### By-products to separate and valorize

Among the by-products, glycerol is obtained following the triglyceride transesterification at a mass ratio around 10 times lower than that of FAME [238,239]. Currently, an overproduction of glycerol (a direct consequence of biodiesel production that alone produces 1400 kton (worldwide) of crude glycerol per year) associated with a slight increase of world glycerol consumption from 600 to 870 kton/year from 2003 to 2013 [240,241] led, as a consequence, to a decrease of glycerol prices. From 2004 to 2011, the price of glycerol decreased from 110 to 7.5 US\$/ton [238,242]. In order to make biodiesel from microalgae cost effective, glycerol could be transformed into other products (added value products) by chemical [243,244], thermochemical [245] or biological [246] methods. A study suggests that if a microalga contains less than 40 wt% lipids, the anaerobic digestion of microalgae residues is essential to make microalgae biodiesel profitable [247].

Furthermore, microalgae contain non-transesterifiable components (chlorophyll, carotenoids) [58] that can be found in biodiesel after transesterification, which require further purification steps [202].

### Cost calculation

Several studies have attempted to determine if a process of biodiesel production from microalgae could be profitable. Some studies calculated the cost of microalgae biomass ranging from 10 to 32 US\$/kg biomass based on the initial investment [248,249]. On the other hand, some authors estimated that microalgae would be produced at a lower cost from 3.0 to 3.8 US\$/kg biomass for photobioreactors and for 'raceway' pond types, respectively [17]. Nevertheless, in more recent studies (2009–2013), some authors calculated lower biomass production costs for microalgae biomass at 2.7 \$US/kg biomass for raceway ponds [250,251], while the cost of microalgae cultivated in photobioreactors would be 7.4 \$US/kg biomass. According to the same sources, considering a microalga with a lipid content of 50 wt%, the cost of microalgae cultivated in fermenters would be 1.6\$US/kg biomass. It should be noted that 20 to 30% of the costs of biomass are associated with microalgae harvesting [252].

In order to calculate the energy balance from the microalgae biodiesel production process, Lardon *et al.* [148] estimated that with a lipid extraction without previous microalgae biomass drying using a solvent lipid extraction would produce a positive energy balance at 105 MJ/kg biodiesel, while microalgae drying would cause a negative energy balance at –2.6 MJ/kg biodiesel. However, only a process that would use low amounts of nitrogen (concentration non explicit) and an extraction of wet microalgae would consume less energy than biodiesel, which usually contains 37.8 MJ/kg biodiesel [148].

According to Davis *et al.* [253] microalgae biodiesel cost calculation modeling is at an early stage because of missing process data and further research has to be performed because several projects of industrial microalgae culture for biodiesel production process were not successful [254].

### Canadian biodiesel production

In 2012, in Canada, petrodiesel consumption represented more than 28% of the total refined oil products, 28 million m<sup>3</sup> [255]. Consequently, in order to meet a demand in biodiesel that would correspond to a blend of biodiesel into petrodiesel of 10 vol% [14], Canada should produce about 2.8 million of m<sup>3</sup> per year of biodiesel. As the density of biodiesel is around 864 kg/m<sup>3</sup> (0.864 kg/L), the biodiesel production should reach 2.4 million tons/year. Considering this production would come from microalgae, a biomass productivity of 0.035 kg/m<sup>2</sup>/day (culture in raceway pond) [17], a biodiesel yield of about 37% (g biodiesel/g dry biomass) [187] and a production of 6 months per year (180 days), the surface required for such production would be 1027 million m<sup>2</sup> (1027 km<sup>2</sup>).

### Biofuels: other applications

Despite the fact that biodiesel is one of the most studied biofuels, microalgae could be used to produce other biofuels such as bioethanol and biomethane [256]. Microalgae could be a raw material for other processes of valorization such as gasification, liquefaction, pyrolysis and hydrogenation [82]. However, these processes are performed at high temperatures and pressures ranging from 300 to 900°C or 7 to 55 MPa, respectively. As an example, researchers have developed patents to transform microalgae biomass by liquefaction using temperatures between 320 and 500°C at pressures between 20 and 55 MPa [257]. Direct combustion could be worked on to produce energy (for example to produce electricity) despite the fact that humidity would be an important factor for the cost efficiency of this process.

### Conclusion

The objective of this literature review was to study the possibility of using microalgae as a source of biodiesel: from microalgae culture through biodiesel synthesis to ASTM biodiesel production standards. There are a lot of valuable products contained in microalgae that could be exploited (such as lipid, proteins, sugar, etc.) which could make biodiesel production from microalgae cost effective. Otherwise, the fluctuation of oil prices encourages the government to reduce its oil dependency and reduce CO<sub>2</sub> emissions, which stimulates biofuel production from renewable biomass.

Among the different sources of biomass, microalgae has a high lipid content (up to 75 wt%), which could be used to produce biodiesel. The main advantage of microalgae is based on the fact that microalgae have high culture yields and do not use arable lands. On the other hand, the use of this biomass for industrial biodiesel production requires the improvement of the techniques of extraction, the transesterification of lipids and the purification of biodiesel.

### Acknowledgements

Michèle Heitz is grateful to Le Fond Québécois de la Recherche sur la Nature et les Technologies (FQRNT) for the grant for the research program in partnership contributing to the reduction of greenhouse gases. Marc Veillette wants also to express his gratitude to the National Sciences and Engineering Research Council of Canada (NSERC) (Alexander Graham Bell Canada Graduate Scholarship and Michael Smith Foreign Study Supplement) for the Rhône.

### Disclosure statement

No potential conflict of interest was reported by the authors.

## Funding

This work was supported by Le Fond Québécois de la Recherche sur la Nature et les Technologies (FQRNT) and the Natural Sciences and Engineering Research Council of Canada.

## References

- [1] International carbon dioxide emissions and carbon intensity [Internet]. Washington: Energy Information Administration; 2011 [cited 2013 Feb 13]. Available from: <http://www.eia.gov/cfapps/ipdbproject/IEDIn dex3.cfm?tid=90&pid=44&aid=8>.
- [2] Archived - Government of Canada announces 2030 emissions target [Internet]. Ottawa: Government of Canada; 2015 [cited 2015 Dec 25]. Available from: <http://news.gc.ca/web/article-en.do?nid=974959>
- [3] Environment Canada. Canada's emissions trends. Ottawa: Government of Canada; 2013. (Report number: En81-18/2013E-PDF).
- [4] Environment Canada. National Inventory Report 1990-2013: Greenhouse Gas Sources and Sinks in Canada. Ottawa: Government of Canada; 2015. (Report number: En81-4/1E-PDF).
- [5] Environment Canada. National Inventory Report 1990-2010: Greenhouse Gas Sources and Sinks in Canada. Ottawa: Government of Canada; 2012. (Report number: 1910-7064).
- [6] 2012 world oil production [Internet]. Washington: U.S. Energy Information Administration; 2013 [cited 2014 Jan 1]. Available from: <http://www.eia.gov/countries/index.cfm?view=production>
- [7] BP statistical review of world energy [Internet]. London: BP; 2013 [cited 2014 June 13]. Available from: <http://www.bp.com/>
- [8] World proved reserves of oil and natural gas, most recent estimates [Internet]. Washington: U.S. Energy Information Administration; 2012 [cited 2014 Jan 1]. Available from: <http://www.eia.doe.gov/international/reserves.html>
- [9] Shafiee S, Topal E. When will fossil fuel reserves be diminished? *Energy Policy*. 2009;37:181–189.
- [10] 2012 world oil consumption [internet]. Washington: U.S. Energy Information Administration; 2015 [cited 2016 Jan 1]. Available from: <http://www.eia.gov/countries/index.cfm?view=consumption>
- [11] McKendry P. Energy production from biomass (part 1): overview of biomass. *Bioresour Technol*. 2002;83:37–46.
- [12] World energy council. World energy resources [Internet]. London: World energy council; 2013 (ISBN: 978 0 946121 29 8).
- [13] Chapman L. Transport and climate change: a review. *J Transp Geogr*. 2007;15:354–367.
- [14] Bindraban PS, Bulte EH, Conijn SG. Can large-scale bio-fuels production be sustainable by 2020? *Agric Syst*. 2009;101:197–199.
- [15] Les législateurs européens a réduire leur consommation de biocarburant a base d'aliments [Internet]. Tulsa (OK): Krista D. Young; 2013 [cited 2014 Feb 1]. Available from <http://www.nouvellesdumonde.net>
- [16] Biodiesel Statistics [Internet]. Washington (DC): U.S. Department of Energy; 2010 [cited 2010 Mar 15]. Available from [http://www.afdc.energy.gov/afdc/fuels/bio diesel\\_statistics.html](http://www.afdc.energy.gov/afdc/fuels/bio diesel_statistics.html)
- [17] Chisti Y. Biodiesel from microalgae. *Biotechnol Adv*. 2007;25(3):294–306.
- [18] BP Statistical Review of World Energy [Internet]. London: BP; 2011 [cited 2011 July 13]. Available from <http://www.bp.com/>
- [19] Brent oil trading at \$121, Iran's President forecasts \$150 soon [Internet]. London: Live oil prices; 2011 [cited 2011 July 13]. Available from <http://www.liveoil prices.co.uk/>
- [20] Egan M. Oil sinks back below \$40. *CNN* [Internet]. Novembre 18 2015 [cited 2015 Jan 1]. Available from <http://money.cnn.com/2015/11/18/investing/oil-price-below-40/>
- [21] Ibañez E, Cifuentes A. Benefits of using algae as natural sources of functional ingredients. *J Sci Food Agric*. 2013;93(4):703–709.
- [22] Ambati RR, Phang S, Ravi S et al., Astaxanthin: Sources, extraction, stability, biological activities and its commercial applications—A review. *Mar Drugs*. 2014;12 (1):128–152.
- [23] Chen W, Lin B, Huang M et al., Thermochemical conversion of microalgal biomass into biofuels: a review. *Bioresour Technol*. 2015;184:314–327.
- [24] Gill S, Tsolakis A, Herreros J et al., Diesel emissions improvements through the use of biodiesel or oxygenated blending components. *Fuel*. 2012;95:578–586.
- [25] United States Environmental Protection Agency. A comprehensive analysis of biodiesel impacts on exhaust emissions. 2002;EPA420-P-02-001, 1–118.
- [26] Chauhan BS, Singh RK, Cho H et al., Practice of diesel fuel blends using alternative fuels: A review. *Ren Sus Energ Rev*. 2016;59:1358–1368.
- [27] Machado Corrêa S, Arbilla G. Carbonyl emissions in diesel and biodiesel exhaust. *Atmos Env*. 2008;42:769–775.
- [28] Demirbas T, Demirbas AH. Bioenergy, Green Energy. Biomass and Biofuels. *Energy Sources Part A*. 2010;32:1067–1075.
- [29] Singh J, Gu S. Commercialization potential of microalgae for biofuels production. *Renew Sust Energ Rev*. 2010;14(9):2596–2610.
- [30] FAOSTAT: Food Price Index [Internet]. Rome, Italy: FAO; 2011 [cited 2011 May 23]. Available from <http://www.fao.org/worldfoodsituation/wfs-home/foodpricesindex/en/>
- [31] A note on Rising Food Crisis [Internet]. D. Mitchell; 2008 [cited 2010 Apr 8]. Available from <http://www.wds.worldbank.org/>
- [32] Goldemberg J, Guardabassi P. Are biofuels a feasible option? *Energy Policy*. 2009;37(1):10–14.
- [33] Yong C, Pang WA. Gender impact of large-scale deforestation and oil palm plantations among indigenous groups in Sarawak, Malaysia. In: Gender and land tenure in the context of disaster in Asia. Berlin: Springer; 2015. p. 15–31.
- [34] Veillette M, Chamoumi M, Nikiema J et al., Production of biodiesel from microalgae. In: Mawaz Z, Naveed S, Editors. Chemical engineering. Rijeka, Croatia: Intech; 2012. p. 1–24.
- [35] Yen H, Hu I, Chen C et al., Microalgae-based biorefinery—from biofuels to natural products. *Bioresour Technol*. 2013;135:166–174.
- [36] Mata TM, Martins AA, Caetano NS. Microalgae for biodiesel production and other applications: A review. *Renew Sust Energ Rev*. 2010;14(1):217–232.
- [37] Wang K, Brown RC. Catalytic pyrolysis of microalgae for production of aromatics and ammonia. *Green Chem*. 2013;15(3):675–681.
- [38] Kim KH, Choi IS, Kim HM et al., Bioethanol production from the nutrient stress-induced microalga *Chlorella vulgaris* by enzymatic hydrolysis and immobilized

- yeast fermentation. *Bioresour Technol.* **2014**;153:47–54.
- [39] Templeton DW, Quinn M, Van Wycken S et al., Separation and quantification of microalgal carbohydrates. *J Chromatography A.* **2012**;1270:225–234.
- [40] Doan QC, Moheimani NR, Mastrangelo AJ et al., Microalgal biomass for bioethanol fermentation: Implications for hypersaline systems with an industrial focus. *Biomass Bioenerg.* **2012**;46:79–88.
- [41] Angermayr SA, Hellingwerf KJ, Lindblad P et al., Energy biotechnology with cyanobacteria. *Curr Opin Biotechnol.* **2009**;20(3):257–263.
- [42] Williams PJ, Laurens LM. Microalgae as biodiesel and biomass feedstocks: Review and analysis of the biochemistry, energetics and economics. *Energy Environ Sci.* **2010**;3:554–590.
- [43] Walker TL, Purton S, Becker DK et al., Microalgae as bioreactors. *Plant Cell. Rep.* **2005**;24:629–641.
- [44] Uslu L, İçik O, Koç K et al., The effects of nitrogen deficiencies on the lipid and protein contents of *Spirulina platensis*. *Afri J Biotechnol.* **2013**;10(3):386–389.
- [45] Kates M. Technique of lipidology: isolation, analysis and identification. Amsterdam, 1000 AE, Netherlands: Elsevier; **1986**.
- [46] Yang T, Fang S, Zhang H et al., N-3 PUFAs have antiproliferative and apoptotic effects on human colorectal cancer stem-like cells in vitro. *J Nutr Biochem.* **2013**;24(5):744–753.
- [47] Cicero AF, Tartagni E. Diet, dietary supplements and age-related macular degeneration (AMD). *Nutrafoods.* **2014**;13(1):5–11.
- [48] Chisti Y. Biodiesel from microalgae beats bioethanol. *Trends Biotechnol.* **2008**;26(3):126–131.
- [49] Deng X, Li Y, Fei X. Microalgae: A promising feedstock for biodiesel. *Afr J Microbiol Res.* **2009**;3(13):1008–1014.
- [50] Liang M, Jiang J. Advancing oleaginous microorganisms to produce lipid via metabolic engineering technology. *Prog Lipid Res.* **2013**;52(4):395–408.
- [51] Liu B, Lee Y. Secondary carotenoids formation by the green alga *Chlorococcum* sp. *J Appl Phycol.* **2000**;12:301–307.
- [52] Pires J, Alvim-Ferraz M, Martins F et al., Carbon dioxide capture from flue gases using microalgae: engineering aspects and biorefinery concept. *Renew Sust Energy Rev.* **2012**;16(5):3043–3053.
- [53] Mishra A, Jha B. Isolation and characterization of extracellular polymeric substances from micro-algae *Dunaliella salina* under salt stress. *Bioresour Technol.* **2009**;100(13):3382–3386.
- [54] Wang M, Kuo-Dahab WC, Dolan S et al., Kinetics of nutrient removal and expression of extracellular polymeric substances of the microalgae, *Chlorella* sp. and *Micractinium* sp., in wastewater treatment. *Bioresour Technol.* **2014**;154:131–137.
- [55] Becker W. Microalgae in human and animal nutrition. In: Richmond A, Editor. Handbook of microalgal culture: biotechnology and applied phycology. Oxford: Blackwell Publishing Ltd; **2004**. p. 312–351.
- [56] Christaki E, Florou-Paneri P, Bonos E. Microalgae: a novel ingredient in nutrition. *Int J Food Sci Nut.* **2011**;62(8):794–799.
- [57] Hosikian A, Lim S, Halim R et al., Chlorophyll extraction from microalgae: A review on the process engineering aspects. *Intern J Chem Eng.* **2010**;2010:1–11.
- [58] Bai M, Cheng C, Wan H et al., Microalgal pigments potential as byproducts in lipid production. *J Taiwan Ins Chem Eng.* **2011**;42(5):783–786.
- [59] De Clerck O, Guiry MD, Leliaert F et al., Algal taxonomy: a road to nowhere? *J Phycol.* **2013**;49:215–225.
- [60] Gopinathan CP. Marine microalgae. Presented at: Proceedings of Ocean Life Food and Medicine Expo; **2004 Feb**; Chennai, India.
- [61] Jianga L, Luoa S, Fana X et al., Biomass and lipid production of marine microalgae using municipal wastewater and high concentration of CO<sub>2</sub>. *Appl Energy.* **2011**;88:3336–3341.
- [62] de Moraes MG, Costa JAV. Isolation and selection of microalgae from coal fired thermoelectric power plant for biofixation of carbon dioxide. *Energy Convers Manag.* **2007**;48(7):2169–2173.
- [63] Kim S, Park J, Cho Y et al., Growth rate, organic carbon and nutrient removal rates of *Chlorella sorokiniana* in autotrophic, heterotrophic and mixotrophic conditions. *Bioresour Technol.* **2013**;144:8–13.
- [64] Cantin I. La production de biodiesel à partir des microalgues ayant un métabolisme heterotrophe [master's thesis]. Sherbrooke (Qc): Université de Sherbrooke; **2010**.
- [65] Li P, Miao X, Li R et al., In situ biodiesel production from fast-growing and high oil content *Chlorella pyrenoidosa* in rice straw hydrolysate. *J Biomed Biotechnol.* **2011**;2011:1–8.
- [66] Wu Z, Shi X. Rheological properties of *Chlorella pyrenoidosa* culture grown heterotrophically in a fermentor. *J Appl Phycol.* **2008**;20(3):279–282.
- [67] Adesanya VO, Vadillo DC, Mackley MR. The rheological characterization of algae suspensions for the production of biofuels. *J Rheol.* **2012**;56:925–939.
- [68] Tanoi T, Kawachi M, Watanabe MM. Effects of carbon source on growth and morphology of *Botryococcus braunii*. *J Appl Phycol.* **2011**;23:25–33.
- [69] Cheng Y, Lu Y, Gao C et al., Alga-based biodiesel production and optimization using sugar cane as the feedstock. *Energy Fuels.* **2009**;23(8):4166–4173.
- [70] Cheng Y, Zhou W, Gao C et al., Biodiesel production from Jerusalem artichoke (*Helianthus Tuberosus* L.) tuber by heterotrophic microalgae *Chlorella protothecoides*. *J Chem Technol Biotechnol.* **2009**;84(5):777–781.
- [71] Zhao X, Wu S, Hu C et al., Lipid production from Jerusalem artichoke by *Rhodospiridium toruloides* Y4. *J Ind Microbiol Biotechnol.* **2010**;37(6):581–585.
- [72] Wei A, Zhang X, Wei D et al., Effects of cassava starch hydrolysate on cell growth and lipid accumulation of the heterotrophic microalgae *Chlorella protothecoides*. *J Ind Microbiol Biotechnol.* **2009**;36(11):1383–1389.
- [73] Xu H, Miao X, Wu Q. High quality biodiesel production from a microalga *Chlorella protothecoides* by heterotrophic growth in fermenters. *J Biotechnol.* **2006**;126:499–507.
- [74] Chen C, Yeh K, Aisyah R et al., Cultivation, photobioreactor design and harvesting of microalgae for biodiesel production: A critical review. *Bioresour Technol.* **2011**;102(1):71–81.
- [75] Liang Y, Sarkany N, Cui Y. Biomass and lipid productivities of *Chlorella vulgaris* under autotrophic, heterotrophic and mixotrophic growth conditions. *Biotechnol Lett.* **2009**;31:1043–1049.
- [76] Burley JS. Algal culture: from laboratory to pilot plant. Washington, DC, USA: Carnegie Institution of Washington; **1976**.
- [77] Zeiler KG, Heacox DA, Toon ST et al., the use of microalgae for assimilation and utilization of carbon dioxide



- from fossil fuel-fired power plant flue gas. *Energy Convers. Manag.* **1995**;36(6-9):707–712.
- [78] Kao C, Chen T, Chang Y et al., Utilization of carbon dioxide in industrial flue gases for the cultivation of microalga *Chlorella* sp. *Bioresour Technol.* **2014**;166:485–493.
- [79] Chinnasamy S, Ramakrishnan B, Bhatnagar A et al., Biomass production potential of a wastewater *alga chlor-ellavulgaris* ARC 1 under elevated levels of CO<sub>2</sub> and temperature. *Int J Mol Sci.* **2009**;10:518–532.
- [80] Cai T, Park SY, Li Y. Nutrient recovery from wastewater streams by microalgae: status and prospects. *Renew Sust Energ Rev.* **2013**;19:360–369.
- [81] Grobbelaar JU. Microalgal biomass production: Challenges and realities. *Photosynthesis Res.* **2010**;106(1-2):135–144.
- [82] Amin S. Review on biofuel oil and gas production processes from microalgae. *Energy Convers Manag.* **2009**;40:1834–1840.
- [83] Arbiba Z, Ruiza J, Álvarez-Díaza P et al., Long term outdoor operation of a tubular airlift pilot photobioreactor and a high rate algal pond as tertiary treatment of urban wastewater. *Eco Eng.* **2013**;52:143–153.
- [84] Suh IS, Lee C. Photobioreactor engineering: Design and performance. *Biotechnol Bioprocess Eng.* **2003**;8(6):313–321.
- [85] Vandamme D, Foubert I, Muylaert K. Flocculation as a low-cost method for harvesting microalgae for bulk biomass production. *Trends Biotechnol.* **2013**;31:233–239.
- [86] Doucha J, Lívanský K. Productivity, CO<sub>2</sub>/O<sub>2</sub> exchange and hydraulics in outdoor open high density microalgal (*Chlorella* sp.) photobioreactors operated in a Middle and Southern European climate. *J Appl Phycol.* **2006**;18:811–826.
- [87] Xu L, Weathers PJ, Xiong X et al., Microalgal bioreactors: Challenges and opportunities. *Eng Life Sci.* **2009**;9(3):178–189.
- [88] Pulz O, Gross W. Valuable products from biotechnology of microalgae. *Appl Microbiol Biotechnol.* **2004**;65:635–648.
- [89] Spolaore P, Joannis-Cassan C, Duran E et al., Commercial applications of microalgae. *J Biosci Bioeng.* **2006**;101(2):87–96.
- [90] Menetrez MY. An Overview of Algae Biofuel Production and Potential Environmental Impact. *Environ Sci Technol.* **2012**;46(13):7073–7085.
- [91] Brennan L, Owende P. Biofuels from microalgae—A review of technologies for production, processing, and extractions of biofuels and co-products. *Renew Sust Energ Rev.* **2010**;14(2):557–577.
- [92] Alga Technologies: Technology Overview [Internet]. Kibbutz Ketura (Israel): Algatechnologies Ltd.; **2004** [cited 2011 May 28]. Available from <http://www.alga-tech.com/techno.htm>
- [93] Singh RN, Sharma S. Development of suitable photobioreactor for algae production – A review. *Renew Sust Energ Rev.* **2012**;16(4):2347–2353.
- [94] Acien Fernández FG, Hall DO, Cañizares Guerrero E et al., Outdoor production of *Phaeodactylum tricornutum* biomass in a helical reactor. *J Biotechnol.* **2003**;103(2):137–152.
- [95] Hall DO, Acien Fernández FG, Guerrero EC et al., Outdoor helical tubular photobioreactors for microalgal production: Modeling of fluid-dynamics and mass transfer and assessment of biomass productivity. *Biotechnol Bioeng.* **2003**;82(1):62–73.
- [96] Acien Fernández FG, Fernández Sevilla JM, Sánchez Pérez JA et al., Airlift-driven external-loop tubular photobioreactors for outdoor production of microalgae: Assessment of design and performance. *Chem Eng Sci.* **2001**;56(8):2721–2732.
- [97] Kurano N, Miyachi S. Microalgal studies for the 21st Century. *Hydrobiologia.* **2004**;512:27–32.
- [98] Xue S, Su Z, Cong W. Growth of *Spirulina platensis* enhanced under intermittent illumination. *J Biotechnol.* **2011**;151(3):271–277.
- [99] Barbosa MJ, Janssen M, Ham N et al., Microalgae cultivation in air-lift reactors: Modeling biomass yield and growth rate as a function of mixing frequency. *Biotechnol Bioeng.* **2003**;82(2):170–179.
- [100] Xu Z, Baicheng Z, Yiping Z et al., A simple and low-cost airlift photobioreactor for microalgal mass culture. *Biotechnol Lett.* **2002**;24(21):1767–1771.
- [101] Ranjbar R, Inoue R, Shiraishi H et al., High efficiency production of astaxanthin by autotrophic cultivation of *Haematococcus pluvialis* in a bubble column photobioreactor. *Biochem Eng J.* **2008**;39(3):575–580.
- [102] Rusch KA, Christensen JM. The hydraulically integrated serial turbidostat algal reactor (HISTAR) for microalgal production. *Aquacult Eng.* **2003**;27(4):249–264.
- [103] Li J, Xu NS, Su WW. Online estimation of stirred-tank microalgal photobioreactor cultures based on dissolved oxygen measurement. *Biochem Eng J.* **2003**;14(1):51–65.
- [104] Sato T, Usui S, Tsuchiya Y et al., Invention of outdoor closed type photobioreactor for microalgae. *Ener Conv Manag.* **2006**;47(6):791–799.
- [105] Huntley WE, Redalje DC, inventors; Huntley WE, Redalje DC, assignee. Continuous-batch hybrid process for production of oil and other useful products from photosynthetic microbes. United States patent US 10/582,029. **2006 June 7**.
- [106] Huntley M, Redalje DG. CO<sub>2</sub> mitigation and renewable oil from photosynthetic microbes: a new appraisal. *Mitig. Adapt. Strat. Glob. Chang.* **2007**;12:573–608.
- [107] Wu Z, Shi X. Optimization for high-density cultivation of heterotrophic *Chlorella* based on a hybrid neural network model. *Lett Appl Microbiol.* **2007**;44(1):13–18.
- [108] Meeting the growing need for renew-able fuels [Internet]. San Francisco (CA): Solazyme Inc.; **2011** [cited 2011 July 8]. Available from <http://www.solazyme.com/fuels>
- [109] Zhang XL, Yan S, Tyagi RD et al., Biodiesel production from heterotrophic microalgae through transesterification and nanotechnology application in the production. *Renew Sust Energ Rev.* **2013**;26:216–223.
- [110] Kitaya Y, Azuma H, Kiyota M. Effects of temperature, CO<sub>2</sub>/O<sub>2</sub> concentrations and light intensity on cellular multiplication of microalgae, *Euglena gracilis*. *Adv Space Res.* **2005**;35:1584–1588.
- [111] Cheng J, Huang Y, Feng J et al., Mutate *Chlorella* sp. by nuclear irradiation to fix high concentrations of CO<sub>2</sub>. *Bioresour Technol.* **2013**;136:496–501.
- [112] Yue L, Chen W. Isolation and determination of cultural characteristics of a new highly CO<sub>2</sub> tolerant fresh water microalgae. *Ener Conv Manag.* **2005**;46(11-12):1868–1876.
- [113] Wu LF, Chen PC, Huang AP et al., The feasibility of biodiesel production by microalgae using industrial wastewater. *Bioresour Technol.* **2012**;113:14–18.
- [114] Mohsenpour SF, Richards B, Willoughby N. Spectral conversion of light for enhanced microalgae growth



- rates and photosynthetic pigment production. *Biore-sour Technol.* **2012**;125:75–81.
- [115] Wahidin S, Idris A, Shaleh SRM. The influence of light intensity and photoperiod on the growth and lipid content of microalgae *Nannochloropsis* sp. *Biore-sour Technol.* **2013**;129:7–11.
- [116] Blanco AM, Moreno J, Del Campo JA et al., Outdoor cultivation of lutein-rich cells of *Muriellopsis* sp. in open ponds. *Appl Microbiol Biotechnol.* **2007**;73(6):1259–1266.
- [117] Sousa C, de Wintera L, Janssen M et al., Growth of the microalgae *Neochloris oleoabundans* at high partial oxygen pressures and sub-saturating light intensity. *Biore-sour Technol.* **2012**;104:565–570.
- [118] Xu N, Zhang X, Fan X et al., Effects of nitrogen source and concentration on growth rate and fatty acid composition of *Ellipsoidion* sp. (*Eustigmatophyta*). *J Appl Phycol.* **2001**;13(6):463–469.
- [119] Xin L, Hong-ying H, Ke G et al., Effects of different nitrogen and phosphorus concentrations on the growth, nutrient uptake, and lipid accumulation of a freshwater microalga *Scenedesmus* sp. *Biore-sour Technol.* **2010**;101(14):5494–5500.
- [120] Sheehan J, Dunahay T, Benemann J et al., A look back 1900 at the U.S. Department of Energy's aquatic species program—biodiesel from algae. Golden (CO): U.S. Department of Energy's Office of Fuels Development; **1998**. (Report number NREL/TP-580 24190).
- [121] Gao Y, Yang M, Wang C. Nutrient deprivation enhances lipid content in marine microalgae. *Biore-sour Technol.* **2013**;127:484–491.
- [122] Giordano M, Pezzoni V, Hell R. Strategies for the allocation of resources under sulfur limitation in the green alga *Dunaliella salina*. *Plant Physiol.* **2000**;124(2):857–864.
- [123] Yang S, Wang J, Cong W et al., Effects of bisulfite and sulfite on the microalga *Botryococcus braunii*. *Enzyme Microb Technol.* **2004**;35(1):46–50.
- [124] Ruangsomboon S. Effect of light, nutrient, cultivation time and salinity on lipid production of newly isolated strain of the green microalga, *Botryococcus braunii* KMITL 2. *Biore-sour Technol.* **2012**;109:261–265.
- [125] Renaud SM, Parry DL. Microalgae for use in tropical aquaculture II: Effect of salinity on growth, gross chemical composition and fatty acid composition of three species of marine microalgae. *J Appl Phycol.* **1994**;6(3):347–356.
- [126] Wang L, Min M, Li Y et al., Cultivation of green algae *Chlorella* sp. in different wastewaters from municipal wastewater treatment plant. *Appl Biochem Biotechnol.* **2010**;162(4):1174–1186.
- [127] Chinnasamy S, Bhatnagar A, Hunt RW et al., Microalgae cultivation in a wastewater dominated by carpet mill effluents for biofuel applications. *Biore-sour Technol.* **2010**;101(9):3097–3105.
- [128] Lüring M, Eshetu F, Faassen EJ et al., Comparison of cyanobacterial and green algal growth rates at different temperatures. *Freshwater Biol.* **2013**;58(3):552–559.
- [129] Abu-Rezq TS, Al-Musallam L, Al-Shimmari J et al., Optimum production conditions for different high-quality marine algae. *Hydrobiologia.* **1999**;403:97–107.
- [130] Sierra E, Acien FG, Fernández JM et al., Characterization of a flat plate photobioreactor for the production of microalgae. *Chem Eng J.* **2008**;138(1–3):136–147.
- [131] Briassoulis D, Panagakis P, Chionidis M et al., An experimental helical-tubular photobioreactor for continuous production of *Nannochloropsis* sp. *Biore-sour Technol.* **2010**;101(17):6768–6777.
- [132] Bitaubé Pérez E, Caro Pina I, Pérez Rodríguez L. Kinetic model for growth of *Phaeodactylum tricornutum* in intensive culture photobioreactor. *Biochem Eng J.* **2008**;40(3):520–525.
- [133] Al-Qasbi M, Raut S, Talebi S et al., A review of effect of light on microalgae growth. Presented at: Proceedings of the World Congress on Engineering; **2012 July 4–6**; London, UK.
- [134] Barbosa MJ, Albrecht M, Wijffels RH. Hydrodynamic stress and lethal events in sparged microalgae cultures. *Biotechnol Bioeng.* **2003**;83(1):112–120.
- [135] Carlsson AS, van Beilen JB, Moller R et al., Micro- and macro-algae: utility for industrial applications. Newbury: CPL Press; **2007**. (Report ISSN 978-1-872691-29-9).
- [136] Rashid N, Rehman SU, Han JI. Rapid harvesting of freshwater microalgae using chitosan. *Proc Biochem.* **2013**;48(4):1107–1110.
- [137] Lee Y, Kim B, Farooq W et al., Harvesting of oleaginous *Chlorella* sp. by organoclays. *Biore-sour Technol.* **2013**;132:440–445.
- [138] The SLS Lab Model Specs [Internet]. Columbus (OH): Algaeventure Inc.; **2011** [cited 2011 Oct 19]. Available from <http://algaevs.com>
- [139] Lotta Watta: algal innovators pioneering new extraction techniques. [Internet]. Columbus (OH): Algaeventure Inc.; **2011** [cited 2011 Oct 19]. Available from <http://algaevs.com>
- [140] Technical specification Evodos SPT 225 & SPT 325 [Internet]. Raamsdonksveer (The Netherlands): Evodos; **2010** [cited 2011 Oct 19]. Available from [http://www.dataopslag.net/storage/cmsserver/upload/files/200037\\_12542.pdf](http://www.dataopslag.net/storage/cmsserver/upload/files/200037_12542.pdf)
- [141] Hu Q, Sommerfeld M, Chen Y et al., inventors; Arizona Board Of Regents, A Body Corporate Of The State Of Arizona Acting For An On Behalf Of Arizona State University, assignee. Method of separation of algal biomass from aqueous or marine culture. United States patent OCT/US2010/0311181.2010, **2010 Apr 15**.
- [142] Lee J, Yoo C, Jun S et al., Comparison of several methods for effective lipid extraction from microalgae. *Biore-sour Technol.* **2010**;101:575–577.
- [143] Cooney M, Young G, Nagle N. Extraction of bio-oils from microalgae. *Sep Purif Rev.* **2009**;38:291–325.
- [144] Umdu ES, Tuncer M, Seker E. Transesterification of *Nannochloropsis oculata* microalga's lipid to biodiesel on Al<sub>2</sub>O<sub>3</sub> supported CaO and MgO catalysts. *Biore-sour Technol.* **2009**;100(11):2828–2831.
- [145] Li X, Xu H, Wu Q. Large-scale biodiesel production from microalga *Chlorella protothecoides* through heterotrophic cultivation in bioreactors. *Biotechnol Bioeng.* **2007**;98(4):764–771.
- [146] Johnson MB, Wen Z. Production of biodiesel fuel from the microalga *Schizochytrium limacinum* by direct transesterification of algal biomass. *Energy Fuels.* **2009**;23:5179–5183.
- [147] Pandolfi G, Miglio R, Carnelli L, inventors; Eni S.P.A., assignee. Process of the drying of algal biomass. United States patent WO2010/140037.2010. **2010 Dec 9**.
- [148] Lardon L, Hélias A, Sialve B et al., Life-cycle assessment of biodiesel production from microalgae. *Environ Sci Technol.* **2009**;43(17):6475–6481.
- [149] Miao X, Wu Q. Biodiesel production from heterotrophic microalgal oil. *Biore-sour Technol.* **2006**;97(6):841–846.

- [150] Folch J, Lees M, Stanley GHS. A simple method for the isolation and purification of total lipides from animal from animal tissues. *J Biol Chem.* **1957**;226:497–509.
- [151] Bligh EG, Dyer WJ. A rapid method for total lipid extraction and purification. *Can J Biochem Physiol.* **1959**;37:911–917.
- [152] Iverson SJ, Lang SLC, Cooper MH. Comparison of the Bligh and Dyer and Folch Methods for Total Lipid Determination in a Broad Range of Marine Tissu. *Lipids.* **2001**;36(11):1283–1287.
- [153] Lewis T, Nichols PD, McMeekina TA. Evaluation of extraction methods for recovery of fatty acids from lipid-producing microheterotrophs. *J Microbiol Methods.* **2000**;42:107–116.
- [154] Veillette M, Giroir-Fendler A, Fauchoux N et al., Biodiesel production from microalgae: Influence of pretreatment on lipid extraction. *J Sust Devel Plan.* **2015**;10(3):382–394.
- [155] Integrated Risk Information System: Chloroform [Internet]. Washington (DC): U.S. Environmental Protection Agency; **2011** [cited 2011 July 10]. Available from <http://www.epa.gov/iris/subst/0025.htm>
- [156] Halim R, Gladman B, Danquah MK et al., Oil extraction from microalgae for biodiesel production. *Bioresour Technol.* **2010**;102(1):178–185.
- [157] Lee S, Yoon B, Oh H. Rapid method for the determination of lipid from the green alga *Botryococcus braunii*. *Biotechnol Tech.* **1998**;12(7):553–556.
- [158] Samorì C, Torri C, Samorì G et al., Extraction of hydrocarbons from microalga *Botryococcus braunii* with switchable solvents. *Bioresour Technol.* **2010**;101(9):3274–3279.
- [159] Fleischer D, Jukie M, Thompson ARG., Inventors; Fleischer D, Jukie M, Thompson ARG., assignee. Systems and Methods for Extracting Lipids from and Dehydrating Wet Algal Biomass. United States patent US 7,868,195 B2.2011. **2011 May 5**.
- [160] Andrich G, Nesti U, Venturi F et al., Supercritical fluid extraction of bioactive lipids from the microalga *Nannochloropsis* sp. *Eur J Lipid Sci Technol.* **2005**;107(6):381–386.
- [161] Sajilata MG, Singhal RS, Kamat MY. Supercritical CO<sub>2</sub> extraction of  $\gamma$ -linolenic acid (GLA) from *Spirulina platensis* ARM 740 using response surface methodology. *J Food Eng.* **2008**;84(2):321–326.
- [162] Ruen-ngam D, Shotipruk A, Pavasant P et al., Selective Extraction of Lutein from Alcohol Treated *Chlorella vulgaris* by Supercritical CO<sub>2</sub>. *Chem Eng Technol.* **2012**;35(2):255–260.
- [163] Aresta M, Dibenedetto A, Carone M et al., Production of biodiesel from macroalgae by supercritical CO<sub>2</sub> extraction and thermochemical liquefaction. *Environ Chem Lett.* **2005**;3:136–139.
- [164] Santana A, Jesus S, Larrayoz MA et al., Supercritical carbon dioxide extraction of algal lipids for the biodiesel production. *Procedia Eng.* **2012**;42:1755–1761.
- [165] Yongli M, inventor; China Co. Ltd. Green Oil, assignee. Biodiesel production method taking algae as raw material. United States patent no 101747924A.2010. **2010 Dec 19**.
- [166] Perrut M. Supercritical fluid applications: Industrial developments and economic issues. *Ind Eng Chem Res.* **2000**;39(12):4531–4535.
- [167] Tan KT, Lee KT. A review on supercritical fluids (SCF) technology in sustainable biodiesel production: Potential and challenges. *Renew Sust Energ Rev.* **2011**;15(5):2452–2456.
- [168] Gordon R, Gorodnitsky I, Grichko V., inventors; Cavitation Technologies, Inc., assignee. Method for processing algae medium containing algae microorganism to produce algal oil and by-products. United States patent US 2010/0151540 A1.2010. **2009 Dec 14**.
- [169] Larach M., inventor; Kai Bioenergy Corporation, assignee. Hydrodynamic extraction of oils from photosynthetic cultures. United States Patent WO/2010/045392.2010. **2009 Oct 14**.
- [170] Dillon HF, Elefant D, Day AG et al., inventors; Solazyme, Inc., assignee. Fractionation of oil-bearing microbial biomass. United States patent PCT/US2010/036238. **2013 Oct 16**.
- [171] Ranjan A, Patil C, Moholkar VS. Mechanistic assessment of microalgal lipid extraction. *Ind Eng Chem Res.* **2010**;49(6):2979–2985.
- [172] Kempkes MA, Roth I, Gaudreau MPJ., inventor; Diversified Technologies, Inc., assignee. Pulsed electric field (PEF) method for continuous enhanced extraction of oil. United States patent US 2011/0107655 A1.2011. **2010 May 12**.
- [173] Machacek MT, Smith TG., inventor; Machacek MT, Smith TG, assignee. Continuous algal biodiesel production facility. United States patent US 2009/0071064 A1.2009. **2009 July 28**.
- [174] Di Serio M, Tesser R, Dimiccoli M et al., Synthesis of biodiesel via homogeneous Lewis acid catalyst. *J Mol Cat A Chem.* **2005**;239(1–2):111–115.
- [175] Liu Y, Lotero E, Goodwin Jr. JG. Effect of water on sulfuric acid catalyzed esterification. *J Mol Cat A Chem.* **2006**;245(1–2):132–140.
- [176] Chai M, Tu Q, Lu M et al., Esterification pretreatment of free fatty acid in biodiesel production, from laboratory to industry. *Fuel Process Technol.* **2014**;125:106–113.
- [177] Wase C, Ridderstrale R, Hasselrot El. Ett kontinuerligt förfarande för isolering av oljor från alger eller mikroorganismer. SE. **2010**;534:278.
- [178] Kuehnle AR, Nolasco NAB., inventor; Kuehnle Agrosystems, Inc., assignee. Enrichment of process feedstock. United States patent US2011/0124034A1.2011. **2010 Nov 24**.
- [179] Di Salvo R, Reich A, Dykes HWH et al., inventor; Streamline Automation, LLC, assignee. Method and apparatus using an active ionic liquid for algae biofuel harvest and extraction. United States patent US2011/0076748.2011. **2010 Dec 16**.
- [180] Poenie M, Jones J, Beach J., inventor; Board Of Regents, The University Of Texas System, assignee. Immobilized resins for algal oil extraction. United States patent US2011/0083360A1.2010. **2010 Oct 13**.
- [181] Thomas G, Lennart L., inventor; Thomas G, Lennart L., assignee. Adiabatic compaction of algae biomass for extraction of biofuel. US2009/0298158A1.2009. **2009 May 16**.
- [182] Echevarría Parres AJJSJB. Method and apparatus for fragmenting algae and extracting the oil therefrom, by means of either a continuous process or batch process. Patent number WO/2010/090506.2010. **2009 Feb 12**.
- [183] Zeiler H, Bullis DT, Light D., inventors; Synthetic Genomics, Inc., assignee. Delivery into cells using ultra-short pulse lasers. Patent number WO2009/140701A2.2009. **2009 May 18**.
- [184] Fu C, Hung T, Chen J et al., Hydrolysis of microalgal cell walls for production of reducing sugar and lipid extraction. *Bioresour Technol.* **2010**;101(22):8750–8754.

- [185] El-Shafie MA., inventor; El-Shafie MA., assignee. Method and systems for processing a biomass for producing biofuels and other products. Patent number WO2011/006019A2.2011. [2014 June 19](#).
- [186] Weaver CA, Kobzeff JM, Behrens PW et al., inventor; Dsm Ip Assets B.V., assignee. Method for producing lipids by liberation from biomass. United States patent US2005/0170479.2005. [2004 Oct 22](#).
- [187] Koberg M, Cohen M, Ben-Amotz A et al., Bio-diesel production directly from the microalgae biomass of *Nannochloropsis* by microwave and ultrasound radiation. *Bioresour Technol.* **2011**;102:4265–4269.
- [188] Nagle N, Lemke P. Production of methyl-ester fuel from microalgae. *Appl Biochem Biotechnol.* **1990**;24(24-25):355–361.
- [189] Lin VSY, Trewyn BG, Chung PW et al., inventor; Iowa State University Research Foundation, Inc., assignee. Sequestration of compounds from microorganism. United States patent US 2010/0196971 A1.2010. [2010 Feb 2](#).
- [190] McNichol J, MacDougall KM, Melanson JE et al., Suitability of Soxhlet extraction to quantify microalgal fatty acids as determined by comparison with in situ transesterification. *Lipids.* **2012**;47:195–207.
- [191] Knothe G. Biodiesel and renewable diesel: A comparison. *Prog Energy Combust Sci.* **2010**;36(3):364–373.
- [192] Altin R, Çetinkaya S, Yücesu HS. Potential of using vegetable oil fuels as fuel for diesel engines. *Ener Conv Manag.* **2001**;42(5):529–538.
- [193] Kiriamiti H, Aoyi O, Wafula W et al., Comparative Study of Petrodiesel and Biodiesel in Domestic and Engine Use. *African J Technol.* **2014**;1(1):1–7.
- [194] Xiong W, Li X, Xiang J et al., High-density fermentation of microalga *Chlorella protothecoides* in bioreactor for microbio-diesel production. *Appl Microbiol Biotechnol.* **2008**;78:29–36.
- [195] Leung DY, Wu X, Leung MKH. A review on biodiesel production using catalyzed transesterification. *Appl Energ.* **2010**;87:1083–1095.
- [196] Hanh HD, Dong NT, Okitsu K et al., Biodiesel production through transesterification of triolein with various alcohols in an ultrasonic field. *Renew Energ.* **2009**;34(3):766–768.
- [197] Likozar B, Levec J. Transesterification of canola, palm, peanut, soybean and sunflower oil with methanol, ethanol, isopropanol, butanol and tert-butanol to biodiesel: modelling of chemical equilibrium, reaction kinetics and mass transfer based on fatty acid composition. *Appl Energy.* **2014**;123:108–120.
- [198] Ehimen EA, Sun ZF, Carrington CG. Variables affecting the in situ transesterification of microalgae lipids. *Fuel.* **2010**;89(3):677–684.
- [199] Gole VL, Gogate PR. A review on intensification of synthesis of biodiesel from sustainable feed stock using sonochemical reactors. *Chem Eng Process.* **2012**;53:1–9.
- [200] Marchetti JM, Miguel VU, Errazu AF. Possible methods for biodiesel production. *Renew Sust Energ Rev.* **2007**;11(6):1300–1311.
- [201] Chauhan BS, Kumar N, Cho HM. A study on the performance and emission of a diesel engine fueled with *Jatropha* biodiesel oil and its blends. *Energy.* **2012**;37(1):616–622.
- [202] Chen L, Liu T, Zhang W et al., Biodiesel production from algae oil high in free fatty acids by two-step catalytic conversion. *Bioresour Technol.* **2012**;111:208–214.
- [203] Couto RM, Simões PC, Reis A et al., Supercritical fluid extraction of lipids from the heterotrophic microalga *Cryptocodinium cohnii*. *Eng Life Sci.* **2010**;10(2):158–164.
- [204] Wang Y, Wang F, Song Q et al., Heterogeneous ceria catalyst with water-tolerant lewis acidic sites for one-pot synthesis of 1, 3-diols via prins condensation and hydrolysis reactions. *J Am Chem Soc.* **2013**;135(4):1506–1515.
- [205] Dias JM, Alvim-Ferraz MC, Almeida MF et al., Selection of heterogeneous catalysts for biodiesel production from animal fat. *Fuel.* **2012**;94:418–425.
- [206] Li Y, Lian S, Tong D et al., One-step production of biodiesel from *Nannochloropsis* sp. on solid base Mg–Zr catalyst. *Appl Energ.* **2011**;88(10):3313–3317.
- [207] Carrero A, Vicente G, Rodríguez R et al., Hierarchical zeolites as catalysts for biodiesel production from *Nannochloropsis* microalga oil. *Catal Today.* **2011**;167(1):148–153.
- [208] Helwani Z, Othman MR, Aziz N et al., Solid heterogeneous catalysts for transesterification of triglycerides with methanol: A review. *Appl Catal.* **2009**;363:1–10.
- [209] Chai F, Cao F, Zhai F et al., Transesterification of vegetable oil to biodiesel using a hetero-polyacid solid catalyst. *Adv Synth Catal.* **2007**;349(7):1057–1065.
- [210] Omota F, Dimian AC, Blik A. Fatty acid esterification by reactive distillation: Part 2—kinetics-based design for sulphated zirconia catalysts. *Chem Eng Sci.* **2003**;58(14):3175–3185.
- [211] Liu C, Huang C, Wang Y et al., Biodiesel production by enzymatic transesterification catalyzed by *Burkholderia* lipase immobilized on hydrophobic magnetic particles. *Appl Energ.* **2012**;100:41–46.
- [212] Ehimen EA, Sun Z, Carrington GC. Use of ultrasound and co-solvents to improve the in-situ transesterification of microalgae biomass. *Procedia Environ Sci.* **2012**;15:47–55.
- [213] Sathish A, Sims RC. Biodiesel from mixed culture algae via a wet lipid extraction procedure. *Bioresour Technol.* **2012**;118:643–647.
- [214] American Society for Testing and Materials (ASTM). Standard specification for biodiesel fuel blend stock (B100) for middle distillate fuels. West Conshohocken (PA): ASTM. **2010**. Standards No. D6751-10.
- [215] Knothe G. Analyzing biodiesel: standards and other methods. *J Am Oil Chem Soc.* **2006**;83(10):823–833.
- [216] Ciubota-Rosie C, Ruiz JR, Ramos MJ et al., Biodiesel from *Camelina sativa*: a comprehensive characterisation. *Fuel.* **2013**;105:572–577.
- [217] Hart Energy Consulting. Establishment of the Guidelines for the Development of Biodiesel Standards in 2220 the APPEC Region. Houston (TX): Hart Energy; **2007**. (Report number APEC#208-RE-01.5).
- [218] Gouvernement du Québec. Règlement sur les produits pétroliers: Loi sur les produits pétroliers (L.R.Q., c. P-30.01, a. 5 et 96). CHAPITRE II: Normes relatives aux produits pétroliers, Article 10. 2011(July 22). **2011**.
- [219] Knothe G, Gerpen JV, Krahl J. The biodiesel handbook. Campaign, Illinois, USA: AOCS Press; **2005**.
- [220] Giakoumis EG. A statistical investigation of biodiesel physical and chemical properties, and their correlation with the degree of unsaturation. *Renew Energ.* **2013**;50:858–878.
- [221] Taux de cétane élevé: Qu'est-ce que le cétane et pourquoi est-ce important ? [Internet]. London (UK): BP; 2011 [15 May 2011]. Available from <http://www.bp.com/sectiongenericarticle.do?categoryId=9017537&contentId=7033512>



- [222] Knothe G, Matheaus AC, Ryan TW. Cetane numbers of branched and straight-chain fatty esters determined in an ignition quality tester. fuel technical review. San Ramon (CA): Chevron Corporation; 2007.
- [223] National Renewable Energy Laboratory. Biodiesel handling and use guide. Golden (CO): National Renewable Energy Laboratory; 2009. (Report No NREL/TP-540-4367).
- [224] Stansell GR, Gray VM, Sym SD. Microalgal fatty acid composition: implications for biodiesel quality. *J Appl Phycol*. 2011;24(4):791–801.
- [225] Canoira L, Alcántara R, Torcal S et al., Nitration of biodiesel of waste oil: Nitrated biodiesel as a cetane number enhancer. *Fuel*. 2007;86(7-8):965–971.
- [226] Dunn RO. Cold weather properties and performance of biodiesel. In: Knothe G, Krah J, Gerpen JV, Editors. The biodiesel handbook. Campaign, Illinois, USA: AOCS Press; 2005. p. 83–121.
- [227] Basha SA, Gopal KR, Jebaraj S. A review on biodiesel production, combustion, emissions and performance. *Renew Sust Energy Rev*. 2009;13(6-7):1628–1634.
- [228] Özener O, Yüsek L, Ergenç AT et al., Effects of soybean biodiesel on a DI diesel engine performance, emission and combustion characteristics. *Fuel*. 2014;115:875–883.
- [229] Knothe G. Oxidative stability of biodiesel. In: Knothe G, Krah J, Gerpen JV, Editors. The biodiesel handbook. Campaign, Illinois, USA: AOCS Press; 2005. p. 122–126.
- [230] Shahabuddin M, Kalam M, Masjuki H et al., An experimental investigation into biodiesel stability by means of oxidation and property determination. *Energy*. 2012;44(1):616–622.
- [231] Focke WW, Van der Westhuizen I, Grobler AL et al., The effect of synthetic antioxidants on the oxidative stability of biodiesel. *Fuel*. 2012;94:227–233.
- [232] Jakeria MR, Fazal MA, Haseeb ASMA. Influence of different factors on the stability of biodiesel: A review. *Renew Sust Energy Rev*. 2014;30:154–163.
- [233] Chevron Corporation. Diesel Fuel technical review. San Ramon (CA): Chevron Corporation; 2007.
- [234] Shumacher L. Biodiesel Lubricity. In: Knothe G, Krah J, Gerpen JV, Editors. The biodiesel handbook. Campaign, Illinois, USA: AOCS Press; 2005. p. 127–133.
- [235] Kalita P, Malshe AP, Kumar SA et al., Study of specific energy and friction coefficient in minimum quantity lubrication grinding using oil-based nanolubricants. *J Manuf Process*. 2012;14(2):160–166.
- [236] Knothe G, Steidley KR. Lubricity of components of biodiesel and petrodiesel. The origin of biodiesel lubricity. *Energy Fuels*. 2005;19(3):1192–1200.
- [237] Lois E, Arkoudeas P. Lubricating aspects of automobile fuels. In: Carmo JP, Ribeiro JE, Editors. New advances in vehicular technology and automotive engineering. Rijeka, Croatia: Intech; 2012. p. 91–118.
- [238] Yazdani SS, Gonzalez R. Anaerobic fermentation of glycerol: a path to economic viability for the biofuels industry. *Curr Opin Biotechnol*. 2007;18(3):213–219.
- [239] Hirkude JB, Padalkar AS. Performance and emission analysis of a compression ignition: engine operated on waste fried oil methyl esters. *Appl Energy*. 2012;90(1):68–72.
- [240] Bondioli P. From oil seeds to industrial products: present and near future of oleochemistry. *Ital J Agron*. 2003;7(2):129–135.
- [241] Global glycerol market from biodiesel, fatty acids, fatty alcohols for personal care, alkyl resins, polyether polyols applications, downstream opportunities is expected to reach USD 2.52 billion By 2020 [Internet]. San Francisco: Gran View Research; 2014 [cited 2014 Apr 4]. Available from <http://www.grandviewresearch.com/press-release/global-glycerol-market>
- [242] Renewable fuels: The biodiesel bulletin [Internet]. Chicago (IL): The Jacobsen; 2011 [cited 2011 avril 7]. Available from <http://www.thejacobsen.com>
- [243] Zakaria ZY, Linnekoski J, Amin N. Catalyst screening for conversion of glycerol to light olefins. *Chem Eng J*. 2012;207:803–813.
- [244] Li L, Korányi TI, Sels BF et al., Highly-efficient conversion of glycerol to solketal over heterogeneous Lewis acid catalysts. *Green Chem*. 2012;14(6):1611–1619.
- [245] Luo X, Hu S, Zhang X et al., Thermochemical conversion of crude glycerol to biopolyols for the production of polyurethane foams. *Bioresour Technol*. 2013;139:323–329.
- [246] Dobson R, Gray V, Rumbold K. Microbial utilization of crude glycerol for the production of value-added products. *J Ind Microbiol Biotechnol*. 2012;39(2):217–226.
- [247] Sialve B, Bernet N, Bernard O. Anaerobic digestion of microalgae as a necessary step to make microalgal biodiesel sustainable. *Biotechnol Adv*. 2009;27:409–416.
- [248] Molina Grima E, Belarbi EH, Acien Fernández FG et al., Recovery of microalgal biomass and metabolites: process options and economics. *Biotechnol Adv*. 2003;20(7-8):491–515.
- [249] Benemann JR, Oswald WJ. System and economic analysis of microalgae ponds for conversion of CO<sub>2</sub> to biomass. Washington (DC): U.S. Department of Energy; 1996. (Report No U.S.DE-FG22-93PC93204).
- [250] Alabi AO, Tampier M, Bibeau E. Microalgae technologies & processes for biofuels/bioenergy production in british columbia: Current technology, suitability & barriers to implementation. Vancouver (BC): The British Columbia Innovation Council; 2009.
- [251] Slade R, Bauen A. Micro-algae cultivation for biofuels: Cost, energy balance, environmental impacts and future prospects. *Biomass Bioenerg*. 2013;53:29–38.
- [252] Ozkan A, Kinney K, Katz L et al., Reduction of water and energy requirement of algae cultivation using an algae biofilm photobioreactor. *Bioresour Technol*. 2012;114:542–548.
- [253] Davis RE, Fishman DB, Frank ED et al., Integrated evaluation of cost, emissions, and resource potential for algal biofuels at the national scale. *Environ Sci Technol*. 2014;48(10):6035–6042.
- [254] Benemann JR. Open ponds and closed photobioreactors - comparative economics. Presented at: 5<sup>th</sup> Annual World Congress on Industrial Biotechnology & Bioprocessing; 2008 April 30; Chicago, USA.
- [255] The supply and disposition of refined petroleum products in Canada [Internet]. Ottawa: Statistics Canada; 2013 [cited 2013 Mar 27] Available from <http://www.statcan.gc.ca/pub/45-004-x/2013004/t109-eng.htm>
- [256] Lam MK, Lee KT. Microalgae biofuels: a critical review of issues, problems and the way forward. *Biotechnol Adv*. 2012;30(3):673–690.
- [257] Catallo JW, Junk T., inventors; Board Of Supervisors Of Louisiana State University And Agricultural And Mechanical College, assignee. Transforming biomass to hydrocarbon mixtures in near-critical or supercritical water. United States patent 6,180,845 B1.1999 Oct 7.
- [258] Fajardo AR, Cerdán LE, Medina AR et al., Lipid extraction from the microalga *Phaeodactylum tricornutum*. *Eur J Lipid Sci Technol*. 2007;109(2):120–126.

- [259] Kita K, Okada S, Sekino H et al., Thermal pre-treatment of wet microalgae harvest for efficient hydrocarbon recovery. *Appl Energ.* **2010**;87(7):2420–2423.
- [260] Wiltshire KH, Boersma M, Möller A et al., Extraction of pigments and fatty acids from the green alga *Scenedesmus obliquus* (Chlorophyceae). *Aquat Ecol.* **2000**;34(2):119–126.
- [261] Young G, Nippgen F, Titterbrandt S et al., Lipid extraction from biomass using co-solvent mixtures of ionic liquids and polar covalent molecules. *Sep Purif Technol.* **2010**;72(1):118–121.
- [262] Tran H, Hong S, Lee C. Evaluation of extraction methods for recovery of fatty acids from *Botryococcus braunii* LB 572 and *Synechocystis* sp. PCC 6803. *Biotechnol Bioprocess Eng.* **2009**;14(2):187–192.
- [263] Barakos N, Pasias S, Papayannakos N. Transesterification of triglycerides in high and low quality oil feeds over an HT2 hydrotalcite catalyst. *Bioresour Technol.* **2008**;99(11):5037–5042.
- [264] Shu Q, Nawaz Z, Gao J et al., Synthesis of biodiesel from a model waste oil feedstock using a carbon-based solid acid catalyst: Reaction and separation. *Bioresour Technol.* **2010**;101(14):5374–5384.
- [265] Zanette AF, Barella RA, Pergher SBC et al., Screening, optimization and kinetics of *Jatropha curcas* oil transesterification with heterogeneous catalysts. *Renew Energ.* **2011**;36(2):726–731.
- [266] Gao L, Xu B, Xiao G et al., Transesterification of palm oil with methanol to biodiesel over a KF/hydrotalcite solid catalyst. *Energ Fuels.* **2008**;22(5):3531–3535.
- [267] Gao L, Teng G, Xiao G et al., Biodiesel from palm oil via loading KF/Ca-Al hydrotalcite catalyst. *Biomass Bioener.* **2010**;34(9):1283–1288.
- [268] Trakarnpruk W, Porntangjitlikit S. Palm oil biodiesel synthesized with potassium loaded calcined hydrotalcite and effect of biodiesel blend on elastomer properties. *Renew Energ.* **2008**;33(7):1558–1563.
- [269] Ngamcharussrivichai C, Totarat P, Bunyakiat K. Ca and Zn mixed oxide as a heterogeneous base catalyst for transesterification of palm kernel oil. *Appl Catal A.* **2008**;341(1-2):77–85.
- [270] Zeng H, Feng Z, Deng X et al., Activation of Mg-Al hydrotalcite catalysts for transesterification of rape oil. *Fuel.* **2008**;87(13-14):3071–3076.
- [271] Meng X, Xin Z. Preparation of biodiesel from soybean oil by transesterification on KF/CaO catalyst. *Petrochem Technol.* **2005**;34(3):282–286.
- [272] Di Serio M, Ledda M, Cozzolino M et al., Transesterification of soybean oil to biodiesel by using heterogeneous basic catalysts. *Ind Eng Chem Res.* **2006**;45(9):3009–3014.
- [273] Xie W, Huang X. Synthesis of biodiesel from soybean oil using heterogeneous KF/ZnO catalyst. *Catal Lett.* **2006**;107(1-2):53–59.
- [274] Xie W, Huang X, Li H. Soybean oil methyl esters preparation using NaX zeolites loaded with KOH as a heterogeneous catalyst. *Bioresour Technol.* **2007**;98(4):936–939.
- [275] Yang Z, Xie W. Soybean oil transesterification over zinc oxide modified with alkali earth metals. *Fuel Process Technol.* **2007**;88(6):631–638.
- [276] Xie W, Li H. Alumina-supported potassium iodide as a heterogeneous catalyst for biodiesel production from soybean oil. *J Mol Cat A Chem.* **2006**;255(1-2):1–9.
- [277] Xie W, Peng H, Chen L. Calcined Mg-Al hydrotalcites as solid base catalysts for methanolysis of soybean oil. *J Mol Cat A Chem.* **2006**;246(1-2):24–32.
- [278] Xie W, Peng H, Chen L. Transesterification of soybean oil catalyzed by potassium loaded on alumina as a solid-base catalyst. *Appl Catal A.* **2006**;300(1):67–74.
- [279] Fu C, Hung T, Su C et al., Immobilization of calcium oxide onto chitosan beads as a heterogeneous catalyst for biodiesel production. *Polym Int.* **2011**;60(6):957–962.
- [280] Chakraborty R, Bepari S, Banerjee A. Transesterification of soybean oil catalyzed by fly ash and egg shell derived solid catalysts. *Chem Eng J.* **2010**;165(3):798–805.
- [281] Antunes WM, Veloso CdO, Henriques CA. Transesterification of soybean oil with methanol catalyzed by basic solids. *Catal Today.* **2008**;133-135(1-4):548–554.
- [282] Samart C, Chaiya C, Reubroycharoen P. Biodiesel production by methanolysis of soybean oil using calcium supported on mesoporous silica catalyst. *Energ Conv Manag.* **2010**;51(7):1428–1431.
- [283] Samart C, Sreetongkittikul P, Sookman C. Heterogeneous catalysis of transesterification of soybean oil using K/mesoporous silica. *Fuel Process Technol.* **2009**;90(7-8):922–925.
- [284] Kim M, DiMaggio C, Yan S et al., The synergistic effect of alcohol mixtures on transesterification of soybean oil using homogeneous and heterogeneous catalysts. *Appl Catal A.* **2010**;378(2):134–143.
- [285] Teng G, Gao L, Xiao G et al., Transesterification of soybean oil to biodiesel over heterogeneous solid base catalyst. *Energ Fuels.* **2009**;23(9):4630–4634.
- [286] Silva CCCM, Ribeiro NFP, Souza MMVM et al., Biodiesel production from soybean oil and methanol using hydrotalcites as catalyst. *Fuel Process Technol.* **2010**;91(2):205–210.
- [287] Kouzu M, Kasuno T, Tajika M et al., Active phase of calcium oxide used as solid base catalyst for transesterification of soybean oil with refluxing methanol. *Appl Catal A.* **2008**;334(1-2):357–365.
- [288] Kim M, Dimaggio C, Yan S et al., The effect of support material on the transesterification activity of CaO-La<sub>2</sub>O<sub>3</sub> and CaO-CeO<sub>2</sub> supported catalysts. *Green Chem.* **2011**;13(2):334–339.
- [289] Yan S, Kim M, Salley SO et al., Oil transesterification over calcium oxides modified with lanthanum. *Appl Catal A.* **2009**;360(2):163–170.
- [290] Wang Y, Zhang F, Xu S et al., Preparation of macro-spherical magnesia-rich magnesium aluminate spinel catalysts for methanolysis of soybean oil. *Chem Eng Sci.* **2008**;63(17):4306–4312.
- [291] Shu Q, Yang B, Yuan H et al., Synthesis of biodiesel from soybean oil and methanol catalyzed by zeolite beta modified with La<sub>3</sub><sup>+</sup>. *Catal Commun.* **2007**;8(12):2159–2165.
- [292] Chen L, Yin P, Liu X et al., Biodiesel production over copper vanadium phosphate. *Energy.* **2011**;36(1):175–180.
- [293] Guo F, Peng Z, Dai J et al., Calcined sodium silicate as solid base catalyst for biodiesel production. *Fuel Process Technol.* **2010**;91(3):322–328.
- [294] Kim MJ, Kim M, Kwon OZ et al., Transesterification of vegetable oils over a phosphazene hydroxide catalyst incorporated onto silica. *Fuel Process Technol.* **2011**;92(1):126–131.
- [295] Ramos MJ, Casas A, Rodríguez L et al., Transesterification of sunflower oil over zeolites using different metal loading: A case of leaching and agglomeration studies. *Appl Catal A Gen.* **2008**;346(1-2):79–85.
- [296] Lukić I, Krstić J, Jovanović D et al., Alumina/silica supported K<sub>2</sub>CO<sub>3</sub> as a catalyst for biodiesel synthesis from



- sunflower oil. *Bioresour Technol.* **2009**;100(20):4690–4696.
- [297] Albuquerque MCG, Jiménez-Urbistondo I, Santamaría-González J et al., CaO supported on mesoporous silicas as basic catalysts for transesterification reactions. *Appl Catal A Gen.* **2008**;334(1-2):35–43.
- [298] Albuquerque MCG, Santamaría-González J, Mérida-Robles JM et al., MgM (M = Al and Ca) oxides as basic catalysts in transesterification processes. *Appl Catalysis A Gen.* **2008**;347(2):162–168.
- [299] Qian W, Han P, Lü X. KF/CaO as solid base catalyst for transesterification to biodiesel by ultrasound. *Nat Prod Res.* **2010**;38(1):52–56.
- [300] Ramachandran K, Sivakumar P, Suganya T et al., Production of biodiesel from mixed waste vegetable oil using an aluminium hydrogen sulphate as a heterogeneous acid catalyst. *Bioresour Technol.* **2011**;102(15):7289–7293.
- [301] Shu Q, Gao J, Nawaz Z et al., Synthesis of biodiesel from waste vegetable oil with large amounts of free fatty acids using a carbon-based solid acid catalyst. *Appl Energ.* **2010**;87(8):2589–2596.
- [302] Li E, Rudolph V. Transesterification of vegetable oil to biodiesel over MgO-functionalized mesoporous catalysts. *Energ Fuels.* **2008**;22(1):145–149.
- [303] Zhang J, Chen S, Yang R et al., Biodiesel production from vegetable oil using heterogeneous acid and alkali catalyst. *Fuel.* **2010**;89(10):2939–2944.