



University College London
Department of Genetics, Evolution and Environment

Traits influence responses to land-use and climate change in terrestrial vertebrates

Adrienne Etard

Supervision: Dr. Tim Newbold, Dr. Alex Pigot

May 20, 2022

Submitted in part fulfilment of the requirements
for the degree of Doctor of Philosophy in Ecology at University College London

Word count: ~XXXXXX

Declaration

I, Adrienne Etard, confirm that the work presented in this thesis is my own. Some parts have been conducted in collaboration with other researchers, and the contributions of co-authors to specific Chapters are described in the *Thesis outline of contents* on page 6. All else is appropriately referenced, and where information has been derived from other sources, I confirm that this has been indicated in the Thesis.

Acknowledgements

Abstract

Human activities have profoundly impacted global biodiversity. Currently, anthropogenic land-use and climate change figure among the major threats to the world's fauna. However, not all species respond similarly to these pressures. Interspecific variability in responses to human threats is notably underpinned by the fact that different species possess different ecological characteristics, some of them allowing species to cope with environmental changes, while others confer a disadvantage to species in modified environments. Understanding what renders species sensitive to anthropogenic pressures is vital to inform and prioritise conservation efforts. Yet, in terrestrial vertebrates, a group for which ecological data is the most abundant, it remains unclear which traits are associated with higher sensitivity to human pressures. The aims of my thesis are to investigate whether and which traits are associated with land-use responses and climate-change sensitivity in terrestrial vertebrates, and to highlight some of the consequences for ecosystem functioning. I first assess the global availability of ecological trait data for terrestrial vertebrates, identifying understudied groups and regions (e.g., Central-African reptiles). I then show that, at global scales, disturbed land uses negatively impact the functional diversity of vertebrate assemblages. Further, I find that in all classes, higher sensitivity to land-use and climate change is associated with narrower ranges, smaller habitat breadth and inability to use human-modified habitats. Both land-use responses and climate-change sensitivity are unevenly distributed among dietary groups, highlighting potential food-web disruptions in assemblages under pressure. Finally, I show that land-use responses are influenced by species' energetic requirements, so that energetic fluxes within vertebrate assemblages are likely modified under human-driven land-use change. Although the large-scale consequences of biodiversity changes for ecosystem functioning remain to be fully understood, my thesis highlights a compositional reshaping of vertebrate assemblages under human pressure and furthers our understanding of anthropogenic impacts on biodiversity.

Impact statement

As anthropogenic pressures on the world's biota keep increasing, it is vital to put into place conservation measures to prevent and reverse further species loss. Beyond ethical and moral considerations, there is an urgent need to protect biodiversity because it sustains a range of ecological processes essential to human well-being and planetary health. Effectively managing biodiversity and related ecosystem processes in a changing world requires an understanding of how different species respond to anthropogenic disturbances. My thesis integrates various data sources to investigate the influence of traits on species land-use responses and on species climate-change sensitivity – two of the most pressing threats to biodiversity – at global scales and comparatively across the four terrestrial vertebrate classes. By asking whether interspecific trait variation is associated with species land-use responses and with climate-change sensitivity, my work consolidates our understanding of what renders species sensitive to environmental change, which can help prioritise conservation efforts.

Chapter 2 presents a trait data collection for terrestrial vertebrates, targeting seven commonly-used traits. I highlight the global taxonomic, geographical, and phylogenetic biases in the trait data, revealing knowledge gaps which could guide future data collection efforts. Chapter 2 was published in *Global Ecology and Biogeography*. The compiled data were made available and have since been used by other ecologists (e.g., Capdevila et al. (2022a)) and downloaded 272 times as of May 2022. Chapter 3 uses the collected trait data and reveals profound effects of land-use change on vertebrate functional diversity, which contributes to documenting global human impacts on vertebrates and also underlines the possible threats posed by land-use change to ecosystem processes sustained by vertebrates. Chapter 3 was published in *Ecology Letters*. In Chapter 4, I ask whether traits are associated with species land-use responses and with species climate-change sensitivity, comparatively across the four vertebrate classes. Chapter 4 thus puts into perspective the usefulness of trait data for understanding how species respond to these anthropogenic changes, which is valuable for conservation planning and prioritisation. In Chapter 5, I ask whether species energetic requirements, estimated from metabolic rates, influence species persistence in disturbed land uses. Chapter 5 thus

integrates physiological data to further our fundamental understanding of how vertebrate species respond to land-use change and of the potential consequences for ecosystem functioning.

Beyond publishing two of my PhD Chapters, I have been able to disseminate my work at various international conferences (BES annual meetings in 2019, 2020 and 2021; BES Macroecology conference in 2019; IBS early-career conference in 2021). I will also present my PhD work at the IBS conference (June 2022, 10th Biennial meeting), and at the BES Macroecology conference (July 2022). I have contributed to the Living Planet Report 2020 (WWF, 2020) and to other published papers (Newbold et al., 2019, 2020b). Overall, my PhD work consolidates our knowledge of the role of vertebrate traits for understanding species responses to human pressures and highlights the value of trait data, and more widely, of ecological knowledge, for preserving vertebrate species in a changing world.

Thesis outline of contents, authorship and collaborations

Chapter 1: General introduction

Chapter 1 presents the background for this thesis, exposes the fundamental concepts, and highlights the research questions I investigated in the different Chapters.

Chapter 2: Global gaps and biases in trait data for terrestrial vertebrates

In Chapter 2, I present an analysis of the global gaps and biases in terrestrial vertebrate trait data. To this end, I collate data on seven traits commonly measured in terrestrial vertebrates. I then evaluate the availability of these trait data across the vertebrate classes, assessing whether there are taxonomic, phylogenetic and spatial biases. This chapter was published in *Global Ecology and Biogeography* in 2020 (DOI: 10.1111/geb.13184; Etard et al. (2020)). The paper was co-authored by Sophie Morrill who collated some of the data on reptile traits as part of an MRes project at UCL, and by Tim Newbold, who participated in the development of the research questions, provided detailed feedback on the analyses, and contributed to the writing of the paper.

Chapter 3: Intensive human land uses negatively affect vertebrate functional diversity

In this Chapter, I investigate how land-use change affects the functional composition and functional diversity of local vertebrate assemblages. This Chapter was published in *Ecology Letters* in 2022 (DOI: 10.1111/ele.13926; Etard et al. (2022)) and co-authored by Alex Pigot and Tim Newbold, who helped construct the hypotheses, provided detailed feedback on the work, and took part in the writing of the paper.

Chapter 4: Geographical range area, habitat breadth and specialisation on natural habitats are associated with land-use responses and climate-change sensitivity more consistently than life-history and dietary traits in terrestrial vertebrates

In this Chapter, I assess whether ecological traits as well as geographical range area are associated with species' land-use responses and species' estimated climate-change sensitivity, comparatively among terrestrial vertebrate classes. Rhiannon Osborne-Tonner contributed to this Chapter by collecting data on amphibian and reptile diet during her MSci project at UCL, which I used to complement my datasets. This Chapter was conducted in collaboration with Tim Newbold who helped develop the research questions and provided detailed feedback on the work and on the writing.

Chapter 5: Energetic constraints and trophic group explain species persistence in disturbed land uses

In Chapter 5, I evaluate the impacts of land-use change on community-level energetic requirements, and I assess whether species' energetic requirements influence species persistence in disturbed land uses. To this end, I use physiological data, compiling species resting metabolic rates (used as a proxy for energetic requirements) from the literature. Meghan Hayden and Laura Dee of the University of Colorado, Boulder, as well as Tim Newbold, contributed to the elaboration of the research questions for this Chapter. Meghan Hayden further contributed to this Chapter by retrieving information on net primary productivity for PREDICTS sites, using data from MODIS satellite imagery. All collaborators also provided feedback on the work and participated in writing the manuscript. This Chapter was submitted to a scientific journal and underwent a round of peer-review. I am preparing this Chapter for resubmission.

Chapter 6: General discussion

This final chapter summarises the main findings of my thesis and assesses their contributions to the field.

Data and code access

Code access

The main pieces of R code for Chapters 2, 3, 4 and 5 are available at: https://github.com/AdrienneEtard/PhD_thesis_code.

This document was compiled with L^AT_EX; the source code and files are available at: <https://github.com/AdrienneEtard/Thesis>

Data

The data used in this thesis are freely accessible and sources are referenced throughout. In particular:

- the PREDICTS database (Hudson et al., 2014, 2017) can be downloaded from: <https://data.nhm.ac.uk/dataset/the-2016-release-of-the-predicts-database>;
- the trait datasets I compiled in Chapter 2 are available at: https://figshare.com/articles/dataset/Global_gaps_in_terrestrial_vertebrate_trait_data/10075421.

Contents

List of Tables	14
List of Figures	17
1 General introduction	23
1.1 Major drivers of global biodiversity change	24
1.1.1 Land-use change	24
1.1.2 Climate change	25
1.1.3 The future of biodiversity in the Anthropocene	27
1.2 Ecological importance of terrestrial vertebrates and current threats	27
1.3 Using trait-based approaches to understand global biodiversity change	28
1.3.1 Traits as common currencies across species	28
1.3.2 Using ecological traits to assess which species are at most risk from human-driven changes	29
1.3.3 Using physiological traits to understand land-use impacts on biodiversity and ecosystem functioning	30
1.3.4 Thesis aims	31
1.4 Detailed aims, hypotheses and outline of the following Chapters	31
2 Global gaps and biases in trait data for terrestrial vertebrates	35
2.1 Introduction	36
2.2 Methods	39
2.2.1 Trait data collection	39
2.2.2 Investigating gaps and biases in trait data	45
2.3 Results	47
2.3.1 Taxonomic biases in trait information	47

2.3.2	Phylogenetic biases in trait completeness	48
2.3.3	Spatial biases in trait completeness	51
2.4	Discussion	53
3	Intensive human land uses negatively affect vertebrate functional diversity	58
3.1	Introduction	58
3.2	Methods	60
3.2.1	Vertebrate assemblages	60
3.2.2	Functional traits and diversity metrics	62
3.2.3	Effects of land use and land-use intensity on FRic and FDis (Hypothesis 1)	63
3.2.4	Investigating functional under-dispersion (Hypothesis 2)	65
3.2.5	Functional loss and functional gain (Hypothesis 3)	65
3.3	Results	66
3.3.1	Effects of land use on FRic and FDis	66
3.3.2	Changes in the probability of occurrence of functional under-dispersion	69
3.3.3	Functional loss and gain	70
3.4	Discussion	72
4	Geographical range area, habitat breadth and specialisation on natural habitats are associated with land-use responses and climate-change sensitivity more consistently than life-history and dietary traits in terrestrial vertebrates	76
4.1	Introduction	77
4.2	Methods	81
4.2.1	Ecological characteristics (Figure 4.1a)	81
4.2.2	Imputations of missing trait values	82
4.2.3	Characterizing the influence of traits on species' land-use responses (Figure 4.1b) . .	83
4.2.4	Characterizing the influence of traits on species' sensitivity to climate change (Figure 4.1c)	87
4.3	Results	89
4.3.1	Land-use responses	89
4.3.2	Climate-change sensitivity	94
4.4	Discussion	98

5 Energetic constraints and trophic group explain species persistence in disturbed land uses	101
5.1 Introduction	102
5.2 Methods	107
5.2.1 Vertebrate assemblage composition	107
5.2.2 Energy availability by land-use type and land-use intensity	107
5.2.3 Resting Metabolic Rates (RMR) & imputations of missing RMR values	108
5.2.4 Trophic group and body mass information	109
5.2.5 Effects of land use, land-use intensity and trophic group on assemblage-level total RMR (prediction 1; Figure 5.2a)	109
5.2.6 Effects of land use, land-use intensity, trophic group and residual RMR on species occurrence probability (prediction 2; Figure 5.2b)	110
5.3 Results	111
5.3.1 Effects of land use, land-use intensity and trophic group on assemblage-level total RMR	111
5.3.2 Effects of land use, land-use intensity, trophic group and residual RMR on species' occurrence probability	112
5.4 Discussion	115
6 General discussion	118
Bibliography	119
Appendix 1: Supporting information for Chapter 2	144
S2.1 Taxonomic corrections	144
S2.2 Additional information for trait compilation	147
S2.3 Cutting distribution maps by altitudinal limits	148
S2.4 Impact of taxonomic corrections on trait coverage	149
S2.5 Assemblage-level median, mean and standard deviation of trait completeness (maps)	150
S2.6 Phylogenetic patterns in trait completeness	155
S2.7 Poisson model summaries	160
S2.8 Spatial model summaries	161
S2.9 Trait coverage and taxonomic matching	162
Appendix 2: Supporting information for Chapter 3	163

S3.1	Land-use categories in PREDICTS and sample sizes	163
S3.2	Trait data & imputation of missing trait values	166
S3.2.1	Choice of imputation technique	168
S3.2.2	Phylogenetic signal in traits	168
S3.2.3	Implementation of ‘missforest’ imputations	169
S3.3	Degree of multicollinearity among traits	170
S3.4	Imputation performance	171
S3.5	Functional loss and functional gain	174
S3.6	Diagnostic plots	177
S3.7	Model robustness	186
S3.8	Model robustness – time since land-use conversion	193
A	Appendix 3: Supporting information for Chapter 4	196
S4.1	Compiling diet information	196
S4.2	Imputing missing trait values	200
S4.2.1	Trait data coverage	200
S4.2.2	Phylogenetic signal in traits	200
S4.2.3	Implementation of missing value imputations	201
S4.2.4	Imputation error	201
S4.3	Land-use types in PREDICTS and sample sizes (number of sampled sites across classes) .	203
S4.4	Land-use responses: multicollinearity checks among the models’ explanatory variables .	205
S4.5	Implementing Climate-niche Factor Analysis across terrestrial vertebrates	208
S4.5.1	Historical climate data: groups of intercorrelated variables	208
S4.5.2	CENFA estimation and resolution	208
S4.6	Climate-change sensitivity models: multicollinearity checks among models’ explanatory variables	211
S4.7	Land-use responses: occurrence probability predictions from the partial models for artificial habitat use and diel activity	213
S4.8	Land-use responses: diagnostic plots for the full models	215
S4.9	Climate-change sensitivity: model summaries and diagnostic plots	218
S4.9.1	Summaries & diagnostic plots for models fitted on species with range area >100 km ²)	218

S4.9.2 Summaries for the PGLS models fitted on all species (including those with range area $\leqslant 100 \text{ km}^2$)	222
S4.9.3 Effects for the PGLS models fitted on all species, against effects for the PGLS models fitted on species whose range area was $>100 \text{ km}^2$	226
S4.10 Validations on complete trait data subsets	228
Appendix 4: Supporting information for Chapter 5	233
Bibliography for the Appendices	241

List of Tables

Chapter 2	35
2.1 Data sources for each trait	41
2.2 Number of species for each analysis	45
Chapter 4	76
4.1 Summary of the effects of the ecological characteristics (except for primary diet) on species' responses to disturbed land uses ('within land-use type' effects) and on species' climate-change sensitivity, for each Class of terrestrial vertebrates.	90
4.2 ANOVA summaries for the PGLS models investigating the associations between the species-level ecological characteristics and species' estimated climate-change sensitivity.	97
Appendix 1	144
S2.1 Coefficients of the model investigating whether species range size explained the number of sampled traits	160
S2.2 Coefficients of the model investigating whether species range size explained the number of sampled traits, using range maps not cut by altitudinal limits	160
S2.3 Spatial model summary for amphibians	161
S2.4 Spatial model summary for reptiles	161
Appendix 2	163
S3.1 Land-use categories in the PREDICTS database	163
S3.2 Phylogenetic signal in continuous and categorical traits	169
S3.3 Variance inflation factors across considered (imputed) traits	170
S3.4 Sample sizes (number of pairs of sites) for the calculation of functional loss and functional gain across all species	175

S3.5	Sample sizes (number of pairs of sites) for the calculation of functional loss and functional gain within each class	176
S3.6	Summary of the model explaining FRic by land use and time since land-use conversion, fitted on the subset of data for which there are information on time since land-use conversion	193
S3.7	Summary of the model explaining FDis by land use and time since land-use conversion, fitted on the subset of data for which I have information on time since land-use conversion	195
Appendix 3		196
S4.1	Phylogenetic signal in continuous and categorical traits	200
S4.2	Land-use categories in the PREDICTS database	203
S4.3	Land-use responses: Generalised Variance Inflation Factors (amphibians, with diet)	205
S4.4	Land-use responses: Generalised Variance Inflation Factors (amphibians, without diet) . .	205
S4.5	Land-use responses: Generalised Variance Inflation Factors (birds)	206
S4.6	Land-use responses: Generalised Variance Inflation Factors (mammals)	206
S4.7	Land-use responses: Generalised Variance Inflation Factors (reptiles, with diet)	206
S4.8	Land-use responses: Generalised Variance Inflation Factors (reptiles, without diet)	207
S4.9	PGLS models: Generalised Variance Inflation Factors (amphibians, with diet breadth) . .	211
S4.10	PGLS models: Generalised Variance Inflation Factors (amphibians, without diet breadth) .	211
S4.11	PGLS models: Generalised Variance Inflation Factors (birds)	211
S4.12	PGLS models: Generalised Variance Inflation Factors (mammals)	212
S4.13	PGLS models: Generalised Variance Inflation Factors (reptiles)	212
S4.14	Summary for the PGLS model fitted on amphibians	218
S4.15	Summary for the PGLS model fitted on birds	219
S4.16	Summary for the PGLS model fitted on mammals	220
S4.17	Summary for the PGLS model fitted on reptiles	221
S4.18	Summary for the PGLS model fitted on amphibians, with all species (including species whose range area was $\leq 100 \text{ km}^2$)	222
S4.19	Summary for the PGLS model fitted on birds, with all species (including species whose range area was $\leq 100 \text{ km}^2$)	223
S4.20	Summary for the PGLS model fitted on mammals, with all species (including species whose range area was $\leq 100 \text{ km}^2$)	224

S4.21	Summary for the PGLS model fitted on reptiles, with all species (including species whose range area was $\leq 100 \text{ km}^2$)	225
S4.22	Summary of the effects of the ecological characteristics (except for diet) on (a) species' responses to disturbed land uses ('within land-use type' effects) and on (b) species climate-change sensitivity, for each class of terrestrial vertebrates, from validation models using empirical, non imputed trait values	229
S4.23	Summary for the PGLS model fitted on amphibians: validations using the empirical trait data subset	231
S4.24	Summary for the PGLS model fitted on birds: validations using the empirical trait data subset	231
S4.25	Summary for the PGLS model fitted on mammals: validations using the empirical trait data subset	231
S4.26	Summary for the PGLS model fitted on reptiles: validations using the empirical trait data subset	232
Appendix 4		233
S5.1	Number of species for which RMR data was collected in each class & data sources, initial RMR coverage for PREDICTS species and phylogenetic signal in RMR	233

List of Figures

1.1	Characteristics of the ‘Great Acceleration’	26
General introduction		34
Chapter 2		35
2.1	Procedure used to identify the accepted names of species	40
2.2	Number of natural and artificial habitats used by species against number of strictly natural habitats used by species	44
2.3	Trait coverage and completeness across species	48
2.4	Within-family median trait completeness in herptiles	50
2.5	Relationship between number of sampled traits and geographical range size	51
2.6	Spatial distribution of assemblage-level median trait completeness in herptiles	52
2.7	Spatial model trends for herptiles	53
Chapter 3		58
3.1	Overview of the study design and functional diversity metrics	61
3.2	Effects of land use, land-use intensity and region on FRic (a) and on FDis (b) across all PREDICTS vertebrates.	67
3.3	Effects of land use, land-use intensity, region and taxonomic class on FRic (a) and FDis (b).	68
3.4	Effects of land use, land-use intensity and region on the probability of occurrence of functional under-dispersion.	69
3.5	Effects of land use and land-use intensity on (a) functional loss and (b) functional gain across pairs of sites in temperate areas and tropical areas.	70
3.6	Effects of land use, land-use intensity, region and taxonomic class on functional loss and functional gain across pairs of sites.	71

Chapter 4	76
4.1 Framework of the study	80
4.2 Assessing the effects of ecological characteristics on species' land-use responses: methodology for (a) categorical characteristics and (b) continuous characteristics.	86
4.3 Predicted occurrence probability as a function of land use, land-use intensity, diet and their interactions, for each class of terrestrial vertebrates.	92
4.4 Proportion of the explained variance attributable to each of the main effects in the land-use and climate-change sensitivity models	93
4.5 Estimated effects of the categorical traits on climate-change sensitivity, from the PGLS models fitted in each class.	95
4.6 Effects of the continuous ecological characteristics on climate-change sensitivity, estimated from the PGLS models in each class.	96
Chapter 5	101
5.1 Map of PREDICTS sites, sample sizes and NPP by land use and land-use intensity	105
5.2 Framework for the predictions and models	106
5.3 Effects of land use, land-use intensity and trophic group on assemblage-level total RMR.	112
5.4 Slope estimates and predictions for the relationship between occurrence probability and residual RMR	113
5.5 Predicted occurrence probabilities in primary vegetation and for the most disturbed land uses where residual RMR was found to have a significant effect on occurrence probability	114
Appendix 1	144
S2.1 Difference in species number before and after taxonomic correction (A) and distribution of number of synonyms across datasets (B)	146
S2.2 (a) Body mass versus body length and (b) longevity versus age at sexual maturity in amphibians	147
S2.3 Generation length versus longevity data in mammals and birds	147
S2.4 Availability of altitudinal limits across species	148
S2.5 Range sizes before versus after cutting by altitudinal limits	148
S2.6 Trait coverage when no taxonomic correction is applied before matching sources by binomial names, versus when I applied the described procedure	149

S2.7	Spatial distribution of assemblage-level median trait completeness in each class	151
S2.8	Spatial distribution of assemblage-level mean trait completeness in each class	152
S2.9	Spatial distribution of assemblage-level standard deviation of trait completeness in each class	153
S2.10	Assemblage-level species richness against standard deviation in trait completeness in each class	154
S2.11	Within-family median trait completeness in mammals	156
S2.12	Within-family median trait completeness in birds	157
S2.13	Within-family standard deviation in trait completeness (herptiles)	158
S2.14	Within-family species richness against the within-family standard deviation of trait completeness	159
S2.15	Relationships between number of sampled traits and geographical range size using distribution maps not cut by altitudinal limits	160
S2.16	Comparison of trait coverage among datasets corrected for taxonomy in different ways . .	162

Appendix 2 **163**

S3.1	Number of sites in each land use and land-use intensity for which FRic and FDis were calculated, across all species	164
S3.2	Number of sites in each land use, land-use intensity and for which FRic and FDis were calculated, within each class	165
S3.3	Relationship between habitat breadth and geographical range size across species in each class	166
S3.4	Trait coverage for the vertebrate species sampled in the PREDICTS database	167
S3.5	Distribution of trait completeness across the vertebrate species sampled in the PREDICTS database	167
S3.6	Distribution of continuous traits considered in the calculation of the functional diversity metrics	171
S3.7	Frequency distribution for categorical traits considered in the calculation of the functional diversity metrics	172
S3.8	Out-of-bag imputation errors for the continuous traits (a) and categorical traits (b)	173
S3.9	Diagnostic plots for Model 1a	177
S3.10	Diagnostic plots for Model 1b	178
S3.11	Diagnostic plots for Model 2a	179
S3.12	Diagnostic plots for Model 2b	180

S3.13	Diagnostic plots for Model 3	181
S3.14	Diagnostic plots for Model 4a	182
S3.15	Diagnostic plots for Model 4b	183
S3.16	Diagnostic plots for Model 5a	184
S3.17	Diagnostic plots for Model 5b	185
S3.18	Effects of land use, land-use intensity and region on FRic (a) and FDis (b) across vertebrates, for the subset of species with complete trait data	186
S3.19	Effects of land use, land-use intensity and region on FRic (a) and FDis (b), for the subset of species with complete trait data, with geographical range size as an additional trait	187
S3.20	Effects of land use and land-use intensity on FRic (a) and FDis (b), obtained when calculating FRic and FDis with each set of imputed traits	188
S3.21	Effects of land use and land-use intensity on FRic (a) and FDis (b), obtained when resampling primary vegetation sites twenty independent times	189
S3.22	Effects of land use, land-use intensity and taxonomic class on FRic (a) and FDis (c), for the subset of species with complete trait data	190
S3.23	Effects of land use, region, land-use intensity and taxonomic class on FRic and FDis obtained with the imputed trait data (black points) or with the complete data subsets (red points)	191
S3.24	Effects of land use, region, land-use intensity and taxonomic class on FRic and FDis, obtained when calculating FRic and FDis with each set of imputed traits	192
S3.25	Effects of land use on FRic for the model that includes time since land-use conversion (blue points) versus the model that does not take time since land-use conversion into account (red points)	194

Appendix 3 196

S4.1	Trait coverage across vertebrate classes, including coverage for diet information	200
S4.2	Out-of-bag estimation of imputation errors for the traits included in the analyses	202
S4.3	Sample sizes (number of PREDICTS sites) for the different land-use types, in each class . .	204
S4.4	Groups of intercorrelated climatic variables using a cutoff of 0.65 for Pearson's correlation coefficient	208
S4.5	Possible impact of resolution on estimated geographical range area	209
S4.6	Estimated climate-change sensitivity estimations at three different resolutions	210

S4.7	Predicted occurrence probability as a function of land use, land-use intensity, artificial habitat use and their interactions in each class	213
S4.8	Predicted occurrence probability as a function of land use, land-use intensity, diel activity and their interactions in each class	214
S4.9	Land-use responses: diagnostic plots for the mixed-effects model fitted on amphibians . . .	215
S4.10	Land-use responses: diagnostic plots for the mixed-effects model fitted on birds	215
S4.11	Land-use responses: diagnostic plots for the mixed-effects model fitted on mammals . . .	216
S4.12	Land-use responses: diagnostic plots for the mixed-effects model fitted on reptiles	217
S4.13	Diagnostic plots for the PGLS model fitted on amphibians	218
S4.14	Diagnostic plots for the PGLS model fitted on birds	219
S4.15	Diagnostic plots for the PGLS model fitted on mammals	220
S4.16	Diagnostic plots for the PGLS model fitted on reptiles	221
S4.17	Effects for the PGLS models investigating associations between species-level ecological characteristics and climate-change sensitivity, either fitted on all species (y-axis), or fitted on the species whose range area was >100 km ² (x-axis)	227
S4.18	Predicted occurrence probability as a function of land use, land-use intensity, diet and their interactions in each class: from validation models using empirical, non imputed trait values	230

Appendix 4 233

S5.1	Formula for the abundance model, used to understand the role of shifts in the body mass of species on observed changes in tRMR	233
S5.2	Effects of land use, land-use intensity, trophic group, body mass and their interactions on assemblage-level total abundance	234
S5.3	Diagnostic plots for the linear mixed-effects model looking at the effects of land use, land-use intensity, trophic group and their interactions on assemblage-level total RMR	235
S5.4	Estimates for the models looking at the effects of land use, land-use intensity, trophic group and their interactions on assemblage-level total RMR	236
S5.5	Effects of land use, land-use intensity, trophic level and their interactions on assemblage-level total RMR: empirical versus imputed	237
S5.6	Diagnostic plots for the generalised mixed-effects model looking at the effects of land use, land-use intensity, trophic group, residual RMR and their interactions on species' probability of occurrence	238

S5.7	Model's coefficients from the occurrence model fitted using the 'lme4' package against coefficients from the model fitted using the 'MCMCglmm' package	239
S5.8	Slope estimates and predictions for the relationship between occurrence probability and residual RMR for the model fitted on empirical RMR values only	240

¹ 1 | General introduction

² Humans have been modifying Earth's ecosystems for thousands of years. Archaeological and palaeontological evidence suggest that human activities may have played a major role in the extinction of Australia's megafauna as early as fifty thousand years ago (Johnson et al., 2016; Miller et al., 2016; Van Der Kaars et al., 2017). The subsequent arrival of modern humans in other parts of the world has also been associated with extinctions, of the megafauna in particular (Broughton and Weitzel, 2018; Sandom et al., 2014). However, the global signature of human presence on Earth has never been as prominent as in recent decades. The past two hundred years have been characterised by a sharp increase in the rates of human-driven changes at the planetary scale, a phenomenon that has been termed 'the Great Acceleration' (Steffen et al., 2015).

¹⁰ To emphasize the recent impacts of human activities on the Earth's systems, Crutzen and Stoermer (2000) proposed that we have entered a new geological epoch, which they called 'the Anthropocene'. Although the formal acceptance of this epoch and the timing of its start are still debated within the stratigraphic community (Lewis and Maslin, 2015; Monastersky, 2015), the coined term reflects the profound effects of humans on planetary processes and on the biosphere, such that its use has largely surpassed the geological field (Malhi, 2017).

¹⁶ The Anthropocene can be characterised by Earth-system and socio-economic indicators (Biermann et al., 2016; Steffen et al., 2011). Two of the major signatures are the human-driven transformations of the land surface, and changes to atmospheric composition, which have led to the onset of anthropogenic climate change (Lewis and Maslin, 2015). Altogether, the development of human activities at unprecedented scales and magnitude has led to the alteration of many ecosystems. As a result of combined anthropogenic pressures, the world's biodiversity has been changing (Daru et al., 2021; Dirzo et al., 2014; Johnson et al., 2017; McGill et al., 2015). Decreases in a range of biodiversity indicators have been reported for many taxonomic groups (Butchart et al., 2010). Human-mediated invasions and translocations of species, coupled with local declines in native species, have promoted biotic homogenisation (Daru et al., 2021; Finderup Nielsen et al., 2019; Newbold et al., 2018). In addition, species have gone extinct at rates higher than expected from natural

26 background variability, with current extinction rates estimated to exceed those inferred from fossil records by
27 a hundred to a thousand times (Barnosky et al., 2011; De Vos et al., 2015). Biodiversity loss and ecosystem
28 change have become such major issues in the 21st century that the prevention of biodiversity erosion and
29 the protection of ecosystems have become priority goals on international agendas (Convention on Biological
30 Diversity, 2022; Hoban et al., 2020). Indeed, it is now well established that biodiversity is tightly linked
31 with ecosystem functioning and ecosystem services delivery (Hooper et al., 2005; Millennium Ecosystem
32 Assessment, 2005; Oliver et al., 2015), and thus ultimately with human well-being (Millennium Ecosystem
33 Assessment, 2005). However, the difficulty in achieving global conservation goals – such as the failure to
34 reach the Aichi targets (Buchanan et al., 2020) – highlights the need to strengthen global conservation efforts
35 if we are to protect biodiversity and related ecosystem services from global threats (Butchart et al., 2016).

36 **1.1 Major drivers of global biodiversity change**

37 The biggest anthropogenic threats to biodiversity have been well characterised (Maxwell et al., 2016). Cur-
38 rently, land-use change is the primary driver of global biodiversity loss, and is responsible for causing global
39 declines in species richness and abundance through habitat modification (Chaudhary et al., 2018; Newbold
40 et al., 2015; Nowakowski et al., 2018; Powers and Jetz, 2019). Although climate change is not currently
41 the main driver of biodiversity change, the negative effects of climate change on biodiversity could equate
42 those of land-use change in their magnitude by 2070 (Newbold, 2018). Other major drivers of biodiversity
43 loss include overexploitation, pollution and the spread of non-native species. In this thesis, my focus is on
44 land-use and climate change as global drivers of biodiversity change.

45 **1.1.1 Land-use change**

46 Land cover describes the physical aspect and composition of the land surface from dominant biotic and abi-
47 otic features, typically classifying the Earth's surface into determined sets of natural and artificial ensembles;
48 land cover can notably be characterised with satellite imagery (Wulder et al., 2018). Land use, however, de-
49 scribes the human intent behind a particular land cover (Lambin et al., 2001). Land-use change thus refers
50 to the process by which humans transform the landscape to achieve socio-economic needs. Land-use change
51 includes transitions from natural to anthropogenic landscapes, as exemplified by agricultural-driven defor-
52 estation in tropical areas (Jayathilake et al., 2021). It also describes transitions between different forms of
53 human-dominated land uses, such as the expansion of urban areas over agricultural lands (Ustaoglu and
54 Williams, 2017). Land-use change can also describe transitions from anthropogenic land uses to natural

55 habitats, for example with the restoration of human-degraded landscapes (Banks-Leite et al., 2020). Al-
56 though humans have been modifying terrestrial ecosystems for millennia – between 75% and 95% of the
57 total land surface could have been altered by human activities at some point in history (Ellis et al., 2013,
58 2021) – it is only during the past three centuries that the terrestrial surface made the transition from mostly
59 wild to mostly human-dominated (Ellis et al., 2010). The most important driver behind this transition has
60 been agricultural expansion, with major increases in cropland and grazing areas from the mid-18th century
61 onward (Figure 1.1a). In recent decades, the expansion of grazing areas and animal feed crops, fuelled by
62 the rising demand for animal products, has been identified as the most important driver of land-use change
63 (Alexander et al., 2015).

64 The effects of land-use change on biodiversity have been characterised at local, regional and global
65 scales – although global-scale studies represent a small proportion of the published research (Davison et
66 al., 2021). Overall, land-use change has a negative impact on species richness and abundance (Newbold
67 et al., 2015). For example, habitat loss and fragmentation due to agricultural expansion is a major driver
68 of biodiversity declines (Foley et al., 2005). Urban areas, which have been expanding at faster rates than
69 urban populations themselves (Seto et al., 2010), can have a considerable negative impact on biodiversity
70 and ecosystem services, despite currently representing a small proportion of the terrestrial surface (about
71 1%; Goldewijk et al. (2017)). In particular, the expansion of impervious surfaces, which characterizes urban
72 development, has been linked to a reduction in species richness (Souza et al., 2019; Yan et al., 2019) and to
73 increases in environmental risks (e.g., due to flooding, Hou et al. (2022); or to heat-island effects). Another
74 important aspect of land-use change for biodiversity outcomes and ecosystem service delivery is the level of
75 intensity at which the land is used to fulfil its purpose. For instance, management practices in agricultural
76 areas are a major determinant of local biodiversity and related ecosystem services such as pollination and
77 pest control (Foley et al., 2005; Kehoe et al., 2015; Millard et al., 2021). In urban areas, introducing and
78 managing green spaces can lead to positive biodiversity outcomes (Aronson et al., 2017; Ives et al., 2016),
79 and can also help mitigate flooding risks and heat islands (Livesley et al., 2016). Yet, land-use intensity
80 has not been explicitly considered by a majority of past studies investigating impacts of land-use change on
81 biodiversity (Davison et al., 2021; Dullinger et al., 2021).

82 1.1.2 Climate change

83 According to the World Meteorological Organization, climate change is defined as long-term changes (i.e.,
84 over at least several decades) to the mean state or to the variability of the climate, attributable to human
85 activity or to natural causes. There is a strong scientific consensus that current climate change (from ap-




Figure 1.1: Characteristics of the ‘Great Acceleration’. (a) Land surface (and % land-surface) used for agricultural purposes between year 0 and 2016. Data from the HYDE database (Goldewijk et al., 2017), downloaded from <https://ourworldindata.org/land-use> (24/01/2022). (b) Annual land-surface temperature anomaly between 1880 and 2021. Data retrieved from the National Oceanic and Atmospheric Administration – National Centers for Environmental Information, Climate at a Glance: Global Time Series, published April 2022, retrieved 06/05/2022 from <https://www.ncdc.noaa.gov/cag/>. The anomalies are calculated with reference to the global temperature average for the 20th century.

proximately 1850) is the result of human-driven changes to atmospheric composition (Crowley, 2000; Intergovernmental Panel on Climate Change, 2013; Maibach et al., 2014). Current manifestations of climate change include rising average temperatures (Valipour et al. (2021); Figure 1.1b), increases in the frequency of extreme events (Seneviratne et al., 2012), and changes in global rainfall patterns (Dore, 2005; Trenberth, 2011).

There is accumulating empirical evidence that climate change affects biodiversity globally, with documented changes in phenology (Inouye, 2022), in the geographical distributions of species (Chen et al., 2011; Lenoir and Svenning, 2015; Soroye et al., 2020), and in species physiology (Chown et al., 2010; Pörtner and Farrell, 2008). Climate-change impacts on individual species have consequences for whole communities, through disruptions of species interactions, which can in turn exacerbate impacts on individual species (Cahill et al., 2013; Kharouba et al., 2018).

97 **1.1.3 The future of biodiversity in the Anthropocene**

98 Projecting future land-use and climate-change impacts on biodiversity highlights the key role of human-
99 development scenarios for global biodiversity outcomes (Newbold, 2018; Powell and Lenton, 2013), for the
100 long-term viability of animal populations (Spooner et al., 2018), and for ecosystem processes and services
101 (Lawler et al., 2014). As the world's population continues to grow and as the demand for food, energy and
102 other commodities keeps rising, rates of global land-use and climate change are unlikely to slow without
103 the implementation of strong international regulations and consumption changes (Intergovernmental Panel
104 on Climate Change, 2022; Stehfest et al., 2019). Under current development scenarios ('business as usual'),
105 future projections show that biodiversity will likely be negatively impacted overall, with decreases in species
106 abundance, increases in extinction rates, and shifts in the distribution of species (Newbold, 2018; Pereira
107 et al., 2010; Schipper et al., 2020). In this context, evaluating the effects of land-use and climate change
108 on biodiversity and associated ecosystem services has become vital in order to put into place mitigation
109 measures. Understanding how species have responded to past and current pressures can help assess how they
110 are likely to respond to future pressures. In particular, understanding what makes species more sensitive to
111 land-use and climate change can help conservation efforts and mitigate global human impacts on biodiversity.

112 **1.2 Ecological importance of terrestrial vertebrates and current threats**

113 In this thesis, I focus on terrestrial vertebrates, a group of more than 30,000 species that has been particu-
114 larly well sampled and studied (Titley et al., 2017), and for which there is available ecological information
115 for many species (such as geographical distributions, traits, occurrence, etc.), allowing for large-scale bio-
116 diversity assessments (e.g., Jenkins et al. (2013)). Terrestrial vertebrates play significant roles in ecosystem
117 functioning and support a wide range of processes, most notably as pollinators (Ratto et al., 2018), seed
118 dispersers (Tiffney, 2004), regulators of lower trophic levels (Barber et al., 2010; Lin et al., 2018; Luck
119 et al., 2012; Salo et al., 2010; Zhang, J. et al., 2018), nutrient cyclers (Cunningham et al., 2018; Inger et al.,
120 2016; Wilson and Wolkovich, 2011) and ecosystem engineers (Severtsov, 2012). Vertebrates are also impor-
121 tant for human societies, both culturally and as sources of proteins (Albert et al., 2018; Alves et al., 2018;
122 H irons et al., 2016), and feature among the most charismatic species in the public's eye (Albert et al., 2018;
123 Courchamp et al., 2018).

124 Despite their cultural and ecological importance, terrestrial vertebrates are highly threatened by hu-
125 man activities. The latest Living Planet Report revealed that vertebrate populations have decreased by
126 70% on average since 1970 (WWF, 2020). According to the IUCN Red List of Threatened Species,

about 41% of assessed amphibian species, 26% of mammals, 21% of reptiles and 13% of birds are classified as threatened with extinction (IUCN 2022, <https://www.iucnredlist.org/resources/summary-statistics>). A recent assessment of vertebrates listed in the IUCN Red List of Threatened Species highlights habitat destruction as the predominant human threat (Cox et al., 2022), but direct exploitation also features among the major factors of decline (Monastersky, 2014). Although climate change is not the principal driver of current population declines (Caro et al., 2022), the first extinction of a mammal (the Bramble Cay melomys, *Melomys rubicola*) attributed to anthropogenic climate was reported in 2016 (Watson, 2016). Future projections highlight that between 10% and 30% of vertebrate species could be locally lost by 2070 depending on climate-change scenarios (Newbold, 2018), and that up to one in six species could face extinction under current climate change (Urban, 2015). Further, despite having been well sampled and studied compared to other groups, there still remain important gaps and biases in our ecological knowledge of terrestrial vertebrates and of their responses to human threats (Hevia et al., 2017; Meiri and Chapple, 2016; Meyer et al., 2015; Oliver et al., 2021).

1.3 Using trait-based approaches to understand global biodiversity change

1.3.1 Traits as common currencies across species

Despite the global average declines reported for vertebrate populations, not all species respond similarly to environmental changes (Dornelas et al., 2019; Leung et al., 2020): while some species are impacted negatively, others benefit from global environmental changes (Newbold et al., 2018; Thomas, 2013). One of the reasons why species differ in their ability to cope with disturbances is that species present different intrinsic characteristics, or traits. Although the formal definition of a trait can vary depending on studies, in this thesis I consider traits to be characteristics that are measurable at an organismal level, comparable across different species, and that likely influence organismal fitness and performance (this is also the definition adopted in McGill et al. (2006)). The idea that species traits mediate species' responses to environmental change was formalised in the 'response-effect' framework, developed in the field of plant ecology (Lavorel and Garnier, 2002), where traits that influence species responses to environmental change were termed 'response traits' (and those that underpin ecological processes were termed 'effect traits'). One of the appeals of trait-based approaches is that individual species are no longer the fundamental unit of biodiversity investigations. Rather, traits become the focus and act as 'common currencies' across species, which is of particular interest for conservation when long-term population data are lacking. If species' responses to human threats consistently relate to certain traits, it may be possible to generalise patterns, and estimate the responses of species

157 for which population data are not available (Verberk et al., 2013).

158 **1.3.2 Using ecological traits to assess which species are at most risk from human-driven
159 changes**

160 Vertebrate ecological traits (which I define here as traits relating to the life-history, diet, morphology, and
161 habitat use of species) have been used to explain species' responses to global changes. Past studies have
162 notably investigated whether species extinction risk is associated with species traits, in order to understand
163 which types of species are likely to be at more risk of extinction (Chichorro et al., 2019; Lebreton, 2011;
164 Ripple et al., 2017), which is of high interest for conservation, but often lacks a consideration of specific
165 threats (González-Suárez et al., 2013).

166 Other studies have focused on the influence of traits on species' responses to particular human threats.
167 For example, a number of correlative trait-based approaches have been used to understand whether traits
168 are associated with species' responses to climate change (with studies focusing on explaining interspecific
169 variation in past or projected range shifts with traits (Di Marco et al., 2021; Mccain and King, 2014; Pacifici
170 et al., 2017; Schloss et al., 2012); with range-filling approaches, that is, looking at whether traits explain
171 differences between the realised and potentially suitable climatic niche of species (Estrada et al., 2018);
172 or with demographic approaches (Capdevila et al., 2022b; Spooner et al., 2018)). Ecological traits have
173 also been used in a predictive fashion, notably with frameworks aiming to assess species' vulnerability to
174 climate change (Foden et al., 2013; Pacifici et al., 2015), assuming that given traits confer higher sensitivity
175 to climate change.

176 In addition, past studies have investigated whether traits are associated with species' responses to habitat
177 disturbance. For example, Newbold et al. (2013) investigated whether traits explained interspecific varia-
178 tion in land-use responses in pantropical birds; Quesnelle et al. (2014) assessed whether traits influenced
179 responses to habitat loss in wetland vertebrates; and Nowakowski et al. (2017) investigated effects of am-
180 phibian traits on sensitivity to habitat modification.

181 Further, trait-based approaches have also been employed to understand the signature of human impacts
182 on the diversity and variability of traits within ecological communities. To this end, a range of functional
183 diversity indices have been developed (Legras et al., 2018; Schleuter et al., 2010). Functional diversity
184 indices have been employed to evaluate the effects of land-use disturbance on the trait diversity of vertebrate
185 assemblages (Flynn et al., 2009; La Sorte et al., 2018; Tinoco et al., 2018), or to assess the projected effects
186 of climate change on the trait diversity of vertebrate species (Stewart et al., 2022).

187 Finally, although trait-based approaches using ecological traits have been widely employed to understand

188 the effects of human-driven changes on vertebrate species, past studies have mostly been conducted at local
189 to regional scales (Davison et al., 2021), and have tended to focus on given taxa among the vertebrate classes
190 (Hevia et al., 2017). Global comparative assessments of the relationships between ecological traits and
191 species' responses to human pressures across terrestrial vertebrates are lacking.

192 **1.3.3 Using physiological traits to understand land-use impacts on biodiversity and ecosys- 193 tem functioning**

194 Physiological traits are typically measured to capture aspects of species' metabolism, performance or bio-
195 chemistry. Recent decades have seen advances in large-scale studies linking physiological traits to macroe-
196 cological patterns of species existence across levels of organisation (Chown et al., 2004; Robert Burger
197 et al., 2021), as exemplified by the development of the 'metabolic theory of ecology' (Brown et al., 2004;
198 Gillooly et al., 2001). In particular, metabolic rates reflect the amount of energy used at the organismal
199 level (I define metabolic rates here as the rates at which an organism processes available energy, which is
200 often measured in the lab by estimating the amount of consumed O₂ by a whole organism over a period of
201 time (Sadowska2015; Auer et al., 2017)). As energy is a fundamental currency across all living organisms,
202 metabolic rates can be employed comparatively across species to investigate interspecific variation in ener-
203 getic expenditure. Thus, past studies have focused on understanding both intraspecific variation in metabolic
204 rates (Auer et al., 2017; Burton et al., 2011) as well as interspecific variation (such as variation in metabolic
205 rates with temperature, Clarke and Fraser (2004); or variation in metabolic rates with longevity – e.g., testing
206 the 'pace of life' theory, Stark et al. (2020)).

207 How species allocate their energy impacts almost all aspects of their persistence. Energetic expenditure
208 relates to food intake, which itself is constrained by the amount of available energy in the environment. Thus,
209 species' energetic requirements are ultimately constrained and influenced by trade-offs between energetic-
210 expenditure allocation and resource intake (Auer et al., 2020). As land-use change profoundly modifies the
211 amount and the types of resources available, it follows that land-use change should impact the total amount
212 of energy processed by vertebrate assemblages. Further, the amount of energy required by species could
213 also be an important predictor of species' ability to cope with a disturbed resource landscape. However,
214 to my knowledge, no study has yet investigated how land-use change affects the energetic requirements of
215 vertebrate assemblages, or whether species' energetic requirements influence how species respond to land-
216 use change.

217 **1.3.4 Thesis aims**

218 As discussed above, studies investigating relationships between ecological traits and environmental change
219 have mostly been conducted at local to regional scale (Davison et al., 2021; Hevia et al., 2017), and have
220 mostly focused on single vertebrate classes or sub-taxa within particular classes. Thus, although response
221 traits to land-use and climate change have been identified in various vertebrate taxa, whether the effects of
222 such traits can be generalised geographically and taxonomically remains largely uncertain, emphasising the
223 need for global comparative assessments of the relationships between traits and species' responses to human
224 threats.

225 In this thesis, I set out to fill in this gap by asking whether interspecific variation in ecological traits is as-
226 sociated with species' land-use responses and with estimated climate-change sensitivity, at global scales, and
227 comparatively across the four terrestrial vertebrate classes. Such an assessment helps to understand which
228 species are at most risk from global changes, and may be useful for the prioritisation of conservation efforts.
229 My thesis also aims to highlight some of the consequences of land-use change for ecosystem functioning,
230 by investigating relationships between species energetic requirements (estimated from metabolic rates) and
231 land-use differences.

232 Throughout my thesis, assemblage-level and species-level responses to land use and land-use intensity
233 are assessed using a 'space-for-time' approach (De Palma et al., 2018). To this end, I use one of the most
234 comprehensive databases recording species occurrence and abundance in different land uses (the PREDICTS
235 database, Hudson et al. (2014, 2017)). I estimate sensitivity to climate change from properties of species'
236 climatic niche space, and thus it is important to emphasize that this does not allow a consideration of species'
237 *responses* to climate change. Indeed, it is difficult to capture the responses of many species to climate change,
238 given that this requires disentangling the effects of climate change from that of other drivers of change over
239 the considered time period (which can be complex; MacLean and Beissinger (2017)), and also requires
240 gathering data on the occurrence or abundance of species over several decades (which may be particularly
241 challenging when working at large taxonomic scales).

242 **1.4 Detailed aims, hypotheses and outline of the following Chapters**

243 The overarching aims of my thesis are to investigate whether species traits are associated with species' land-
244 use responses and species' estimated climate-change sensitivity in terrestrial vertebrates, and to highlight
245 some of the consequences of global changes for ecosystem processes sustained by terrestrial vertebrates.
246 One of the obstacles that has hindered the application of trait-based approaches at large scales in animal

247 taxa is the lack of a centralised repository for readily available trait data, as emphasized by the recent calls
248 to compile and release trait data for animals (Junker et al., 2022; Kissling et al., 2018). Thus, collecting
249 trait data and investigating the current availability of the data for terrestrial vertebrates is an important and
250 necessary prerequisite to any analysis. In Chapter 2, I present a data collection of ecological traits for
251 terrestrial vertebrates. Because using similar traits in the different vertebrate classes is necessary to be able
252 to make comparisons among vertebrate classes, I target seven classes of traits that are commonly used across
253 taxonomic groups: body mass/size, lifespan, litter/clutch size, trophic level, diel activity, habitat breadth,
254 and habitat specialisation (characterising whether a species is able to use artificial habitats). I am not able
255 to consider intraspecific variation in the data compilation, since multiple measurements of trait values do
256 not exist for many vertebrate species. Chapter 2 also assesses the availability of the trait data across the
257 terrestrial vertebrate classes, and investigates whether the trait data present global taxonomic, phylogenetic
258 and spatial biases. On the basis of past work (Titley et al., 2017), I predict that amphibians and reptiles are
259 under-sampled compared to mammals and birds, with more important taxonomic, phylogenetic and spatial
260 biases in the trait sampling. Further, I hypothesize that trait data are less abundant for the narrower-ranging
261 species and in species-richer regions.

262 At the assemblage level, the diversity of species traits can be summarised with functional diversity in-
263 dices (Legras et al., 2018; Schleuter et al., 2010; Villéger et al., 2008). Past research has shown that land-use
264 disturbances affect the functional composition of vertebrate assemblages (Flynn et al., 2009; Tinoco et al.,
265 2018). However, to the best of my knowledge, a global assessment of how the functional diversity of terres-
266 trial vertebrate assemblages respond to land use and land-use intensity, within and across taxonomic classes,
267 has not yet been undertaken. Chapter 3 aims to fill in this gap, by investigating the effects of land use and
268 land-use intensity on the functional diversity of local vertebrate assemblages. To this end, I combine the trait
269 data collected in Chapter 2 with the PREDICTS database, after imputing missing trait values (as described
270 in Chapter 2). I hypothesize that the functional diversity of vertebrate assemblages in disturbed land uses is
271 lower than in undisturbed areas. I further predict that decreases in functional diversity in disturbed land uses
272 are driven by high levels of functional loss and that observed declines in functional diversity exceed those
273 expected from random species loss.

274 Chapter 3 highlights the effects of land-use change on the functional composition of vertebrate assem-
275 blages, but does not allow an assessment of the effects of particular traits on species' land-use responses,
276 as multidimensional interspecific trait variation is summarised into single indices of functional diversity.
277 Chapter 4 aims at assessing such effects, by investigating whether species traits are associated with species'
278 land-use responses and species' climate-change sensitivity. In addition to the traits considered in Chapter 3,

279 Chapter 4 includes dietary traits and species geographical range area. Although geographical range area is
280 not a trait *per se*, it has been shown to influence species' responses to land-use and climate change (Newbold
281 et al., 2018; Thuiller et al., 2005), and it is likely an important determinant of climate-change sensitivity.
282 Thus, in Chapter 4, I enhance the trait data compiled in Chapter 2 with dietary traits and geographical range
283 area, which together with the traits previously collected I term 'ecological characteristics'. I investigate
284 whether these ecological characteristics are associated with species' land-use responses on the one hand and
285 with species' estimated climate-change sensitivity on the other hand, comparatively among the terrestrial
286 vertebrate classes. To the best of my knowledge, Chapter 4 constitutes the first global comparative assess-
287 ment, across terrestrial vertebrate classes, of associations between species' ecological characteristics and
288 both land-use responses and estimated climate-change sensitivity.

289 Chapter 5 develops our understanding of the impacts of land-use change on ecosystem functioning by
290 focusing on species' energetic requirements, which is interesting for at least two reasons: first, because en-
291 ergetic requirements relate to resource intake, and as such reflect the amount of energy that is processed by
292 different trophic groups, which can inform about ecosystem functioning; second, because species persistence
293 is constrained by trade-offs in energy allocation among diverse processes (e.g., maintenance, growth, repro-
294 duction), such that energetic requirements are likely important determinants of species' ability to cope with
295 disturbances. Yet, there has been no study so far investigating relationships between energetic requirements
296 and land-use change in terrestrial vertebrates. In Chapter 5, I collect resting metabolic rate estimates for
297 vertebrate species, that is, the estimated minimum amount of energy necessary for organismal maintenance.
298 I use these as a proxy for minimum species-level energetic expenditure, and I combine these estimates with
299 the PREDICTS database. I then assess the effects of land use on the total energetic requirements of verte-
300 brate assemblages (also referred to as community metabolism). Second, I assess whether species' energetic
301 requirements influence species persistence in disturbed land uses, after removing the effects of body mass
302 on energetic expenditure. Assuming that there is less energy available in disturbed land uses, I hypothesize
303 that the assemblage-level energetic requirements of vertebrates are lower in disturbed land uses compared
304 to natural habitats, and that species with lower mass-independent energetic requirements are favoured over
305 species with higher mass-independent energetic requirements in disturbed land uses. Chapter 5 highlights
306 the impacts of land-use change on vertebrate community metabolism and develops our understanding of the
307 factors that shape how species respond to changes in land use.

308 Finally, in Chapter 6, I summarise the findings of my thesis, I highlight some of the limitations, and I ex-
309 amine the relevance of my findings for the field. By investigating whether traits are associated with species'
310 land-use responses and climate-change sensitivity across the terrestrial vertebrates, my thesis furthers our

311 understanding of what could render species more sensitive to human threats, underlines possible modifica-
312 tions to ecosystem functioning, and stresses the role and usefulness of vertebrate trait data and ecological
313 knowledge for understanding species- and community-level responses to human pressures.

314 **2 | Global gaps and biases in trait data for**
315 **terrestrial vertebrates**

316 **Keywords**

317 Terrestrial vertebrates; traits; coverage; completeness; taxonomic biases; spatial biases; phylogenetic biases.

318 **Abstract**

319 Trait data are increasingly used in studies investigating the impacts of global changes on the structure and
320 functioning of ecological communities. Despite a growing number of trait data collations for terrestrial
321 vertebrates, there is to date no global assessment of the gaps and biases the data present. Here, I assess
322 whether terrestrial vertebrate trait data are taxonomically, spatially and phylogenetically biased. I compile
323 seven ecological traits and quantify coverage as the proportion of species for which an estimate is available.
324 For a species, I define completeness as the proportion of non-missing values across traits. I assess whether
325 coverage and completeness differ across classes and examine phylogenetic biases in trait data. To investi-
326 gate spatial biases, I test whether wider-ranging species have more complete trait data than narrow-ranging
327 species. Additionally, I test whether species-rich regions, which are of most concern for conservation, are
328 less well-sampled than species-poor regions. My results show that mammals and birds are well-sampled
329 even in species-rich regions. For reptiles and amphibians (herptiles), only body size presents a high cov-
330 erage (>80%), as well as habitat related variables (for amphibians). Herptiles are poorly sampled for other
331 traits. The shortfalls are particularly acute in some species-rich regions and for certain clades. Across all
332 classes, geographically rarer species have less complete trait information. Hence, trait information is less
333 available on average in some of the most diverse areas and in geographically rarer species, both critical for
334 biodiversity conservation. Gaps in trait data may impede our ability to conduct large scale analyses, while

335 biases can impact the validity of extrapolations. A short-term solution to the problem is to estimate missing
336 trait data using imputation techniques, while a longer-term and more robust filling of existing gaps requires
337 continued data collection efforts.

338 **2.1 Introduction**

339 Species traits are fundamental to ecological and evolutionary research. Comparative studies regularly use
340 trait data across organisms to understand evolutionary processes and species coexistence (Escudero and
341 Valladares, 2016; Zamudio et al., 2016), to investigate global patterns of life forms and functions (Díaz et
342 al., 2016), or to assess species' vulnerability to environmental changes (Böhm et al., 2016; Pacifici et al.,
343 2015; Pearson et al., 2014). Because traits influence species' ability to cope with environmental changes
344 (Newbold et al., 2013) and underpin species' contributions to ecosystem processes (Lavorel and Garnier,
345 2002; Violle et al., 2007; Wong et al., 2018), they play an increasingly important role in functional and
346 conservation ecology.

347 Past and recent efforts to collate and release trait data in the public domain have facilitated the develop-
348 ment of trait-based research. For instance, a global trait database has been published for plants (Kattge et al.,
349 2011). As of May 2020, data from this database had been used in 297 publications since its release (Ac-
350 tivity report, 18/06/2020, <https://www.try-db.org/TryWeb/Home.php>). Such databases hence
351 constitute invaluable research tools and have the potential to greatly advance the field.

352 Vertebrates are one of the most studied taxa (Titley et al., 2017). There are now diverse sources of eco-
353 logical traits for vertebrate groups (primates: Galán-Acedo et al., 2019; mammals: 'PanTHERIA', Jones et
354 al., 2009; amniotes: Myhrvold et al., 2015; amphibians: 'AmphiBIO', Oliveira et al., 2017). These datasets
355 stem from important efforts to collate published estimates of trait data and make them readily available. Trait
356 data have also been made available on online platforms (for instance, the Global Assessment of Reptile Dis-
357 tribution initiative: <http://www.gardinitiative.org/>; the IUCN Red List of Threatened Species:
358 <https://www.iucnredlist.org/>; BirdLife data zone: <http://datazone.birdlife.org/>
359 home).

360 Nevertheless, despite the importance of vertebrate species in global research outputs, there is no single
361 source for vertebrate ecological traits. Consequently, researchers wishing to conduct comparative studies
362 across vertebrate groups may have to collate trait data from a range of sources (such as in Cooke et al.
363 (2019a,b) or in González-Suárez et al. (2018)), a time-consuming prerequisite which may be a limiting
364 step of the research process. Indeed, collating data from heterogeneously-formatted sources presents many

365 challenges (Schneider et al., 2019), particularly when working across a large number of species. For instance,
366 traits may be measured differently across datasets; units may be inconsistent; and taxonomic resolution and
367 nomenclature may vary.

368 The lack of a curated, readily available global database for vertebrate ecological traits impedes our ability
369 to conduct cross-taxon comparative studies at global scales. However, efforts to collate data into a single
370 database are limited by the availability of underlying data. Because there exist important gaps in biodiversity
371 knowledge (Hortal et al., 2015), trait datasets are often incomplete, with many species lacking estimates for
372 many traits. The incompleteness of ecological trait data at the species level has been termed the ‘Raunkiær
373 shortfall’ by Hortal et al. (2015). Furthermore, incomplete trait data are likely to be biased. Biases in trait data
374 may be the consequence of uneven taxonomic and spatial collection effort, with a set of charismatic or easily
375 detectable species being more completely sampled. For instance, González-Suárez et al. (2012) investigated
376 biases in global trait information in mammals. They notably found that the availability of mammalian trait
377 data was geographically and phylogenetically biased, with larger and more widely distributed species being
378 better sampled. In addition, data availability also differed across IUCN Red List extinction risk categories,
379 with threatened species (Critically Endangered, Endangered or Vulnerable) being less well sampled for traits
380 than non-threatened species (Least Concern or Near Threatened).

381 A major issue with incomplete, biased data is the introduction of bias in subsequent analyses. Assessing
382 the amount of missing data as well as the so-called ‘missingness mechanism’ – whether missing data are
383 missing at random, as opposed to there being systematic biases in the way missing values are distributed, see
384 Baraldi and Enders (2010) – is an important prerequisite. Indeed, there exist diverse techniques to deal with
385 data missingness. The simplest one consists of retaining complete cases only by filtering out missing values
386 (case deletion, see Nakagawa and Freckleton (2008)). Nevertheless, case deletion may lead to biased param-
387 eter estimates and erroneous conclusions when values are not missing at random (González-Suárez et al.,
388 2012). Therefore, it is critical to determine the most appropriate way to deal with data incompleteness. For
389 instance, previous studies using terrestrial vertebrate trait data have employed multiple imputation techniques
390 to fill in the gaps (Cooke et al., 2019a; González-Suárez et al., 2012). Yet, imputation techniques could be
391 sensitive to non-randomness in trait data. Phylogenetic biases (where some clades are under-sampled com-
392 pared to other clades) could notably impact the performance of several imputation approaches. It is thus
393 vital to characterise the gaps in trait data prior to any analysis. However, there has been no study to date
394 investigating global patterns in the availability of trait data across terrestrial vertebrates.

395 Here, I aim to assess the global state of trait data in terrestrial vertebrates. I focus on a set of traits that are
396 available across the four classes and that are commonly used by ecologists: body size; litter or clutch size;

397 longevity; trophic level; activity time; habitat breath; and a measure of habitat specialisation. I quantify and
398 compare the gaps in trait data across classes by calculating the coverage of each trait across species, and the
399 completeness of trait estimates for each species (Box 1). I investigate taxonomic, spatial and phylogenetic
400 biases in trait coverage and completeness.

401 Given that biodiversity research is globally biased towards birds and mammals (Titley et al., 2017),
402 I hypothesise that herptiles are less well sampled for traits than mammals and birds, having both lower
403 coverage and completeness.

404 Furthermore, building upon previous studies conducted on mammals (González-Suárez et al., 2012), I
405 hypothesize that species rarity influences completeness, focusing on species geographical range size as one
406 aspect of rarity. Widely distributed species could be better sampled than narrowly distributed species because
407 their ranges overlap with more study sites, regardless of their abundance. As such, I test whether species
408 geographical range size explains trait completeness.

409 It is well established that global research effort is distributed unequally (United Nations Educational
410 Scientific and Cultural and Organization, 2015), with patterns underpinned by various geographical and so-
411 cieconomic factors. For instance, countries with higher gross domestic product tend to host a larger number
412 of research institutions (Martin et al., 2012). The proximity of research infrastructures and the accessibility
413 of survey sites play an important part in explaining the global distribution of knowledge (Hortal et al., 2015).
414 As a result of these factors, biodiversity data gaps tend to be greater in tropical areas (Collen et al., 2008).
415 Tropical areas have the greatest species richness, and so these data biases are of great concern for biodiversity
416 conservation. Whether species-rich regions are systematically under-sampled for traits compared to species-
417 poor regions is thus important to assess, given the significance of species-rich areas for global conservation.
418 Here, I investigate spatial biases in trait completeness, hypothesizing that species-rich areas are on average
419 less well sampled than species-poor areas.

420 Finally, I investigate phylogenetic biases in the trait data. I assess whether particular clades have received
421 more attention than others by looking for patterns in the distribution of trait completeness across the terminal
422 branches of phylogenetic trees in each class.

Box 1. Definitions

Trait: Sensu stricto, a characteristic measurable at the level of an individual and that influences organismal fitness or performance (Violle et al., 2007). In this thesis, I broaden this definition to include ‘ecological’ traits (e.g., the number of habitats used by a species), where the relationship of a species to the surrounding environment needs to be considered. Ecological traits may be estimated by aggregating data across multiple individuals.

Trait completeness: For a given species, the proportion of traits for which an estimate is available.

Trait coverage: For a given trait, the proportion of species for which an estimate is available.

423 2.2 Methods

424 I produced class-specific trait datasets that were made available on figshare (https://figshare.com/articles/dataset/Global_gaps_in_terrestrial_vertebrate_trait_data/10075421/)
425 2). Data compilation and all analyses were conducted with R version 3.5.1 (R Core Team, 2018). Distribu-
426 tion maps were processed using both R and the ‘ArcPy’ package available in ArcGIS v.10.6 (ESRI, 2011)
427 (implemented in Python 2.7; van Rossum, G. (1995)).

429 2.2.1 Trait data collection

430 Sources and taxonomic matching

431 I used freely accessible secondary sources in my compilation (Table 2.1), selected for their broad taxonomic
432 coverage and/or for their frequent use in macroecological studies. Across sources, similar species could
433 appear under synonymous names. This was a potential problem for matching sources by binomial names.
434 Indeed, synonymy can artefactually decrease trait coverage, when trait information is not available across
435 all synonyms. Notably, difficulties arise when species have been divided into several subspecies or when
436 different subspecies are clumped together. Systematic manual checks could not be applied considering the
437 scale of the data collection (there were >39,000 unique binomial names across sources). I developed a
438 procedure aiming at identifying one accepted name for each of the binomial names found across sources.
439 When I could not find an accepted name, I used the original name. Figure 2.1 summarizes the main steps;
440 similar solutions have been used in other large-scale studies (Cooke et al., 2019a).

441 Briefly, the procedure consisted of extracting synonyms from the IUCN (IUCN, 2020) or from the In-
442 tegrated Taxonomic Information System (ITIS; <https://www.itis.gov/>), using the ‘rredlist’ (Cham-
443 berlain, 2018) and ‘taxize’ (Chamberlain and Szöcs, 2013) R packages. One accepted name was assigned to

444 each synonym. I produced a "Synonym" dataset that I have also made available. I then normalized taxonomy
445 across sources by replacing binomial names with their identified accepted name where applicable.

446 Given that different taxonomic backbones could be used to correct for taxonomy, I make two versions
447 of the trait compilations available (corrected and not corrected for taxonomy), meaning that users are free to
448 apply their own corrections; for example, taxonomy could be aligned to that of class-specific sources, such
449 as The Reptile Database, the American Museum of Natural History's Amphibian Species of the World, the
450 Mammal Diversity Database or the International Ornithological Congress World Bird List. Datasets cor-
451 rected for taxonomy contain 11,634 species of birds, 5,381 mammals, 10,612 reptiles and 6,990 amphibians.
452 Where no taxonomic correction was applied when matching sources, the compiled datasets contain 13,501
453 birds, 5,791 mammals, 11,012 reptiles and 8,583 amphibians. For more information, see Appendix 1 (S2.1:
454 'Taxonomic corrections'; Figure S2.1).




Figure 2.1: Procedure used to identify the accepted names of species. I extracted, where possible, the accepted names of species from either the IUCN Red List or the Integrated Taxonomic Information System (ITIS). Figure reproduced from Etard et al. (2020).

455 **Compilation methods**

456 For continuous traits, I took the median value within species when multiple estimates were available from
457 different sources, after removal of any repeated values, which were assumed to represent estimates duplicated
458 across secondary compilations and derived from the same underlying primary sources. Although intraspe-
459 cific variation is increasingly being recognized to have important effects on ecological systems (Bolnick
460 et al., 2011; Des Roches et al., 2018; González-Suárez and Revilla, 2012; Siefert et al., 2015), it was not

Table 2.1: Data sources for each trait. Abbreviations: A = amphibians; B = birds; BL = body length; BM = body mass; DA = diel activity time; GL = generation length; H = habitat data; LCS = litter or clutch size; L/ML = longevity or maximum longevity; M = mammals; MA = age at sexual maturity; R = reptiles; RS = range size; TL = trophic level. Note. Data sources may contain more traits than shown here. Tick marks in parentheses indicate that the trait was present in the data source but that another closely related trait with a better coverage was used instead. The tilde character (~) before a tick mark indicates that I derived trophic levels from species diet. ¹ <http://datazone.birdlife.org/home>; ² <https://www.iucnredlist.org/resources/spatial-data-download>; ³ <http://apiv3.iucnredlist.org/api/v3/docs##general>. Table reproduced from Etard et al. (2020).

Sources	Taxa	Traits									
		BM	BL	L/ML	MA	GL	LCS	TL	DA	RS	H
Oliveira et al. (2017)	Amphibians	(✓)	✓	(✓)	✓		✓	~✓	✓		
Cooper et al. (2008)							✓				
Sodhi et al. (2008)			✓								
Wilman et al. (2014)	Birds	✓						~✓	✓		
BirdLife ¹		✓				✓					✓
Jones et al. (2009)	Mammals	✓	(✓)	(✓)	(✓)		✓		✓		
Kissling et al. (2014)								✓			
Gainsbury et al. (2018)								✓			
Wilman et al. (2014)		✓							✓		
Pacifici et al. (2015)		✓				✓					
Scharf et al. (2015)	Reptiles	✓		✓	(✓)		✓	✓	✓		
Vidan et al. (2017)										✓	
Stark et al. (2018)		✓		✓			✓		✓		
Schwarz and Meiri (2017)								✓			
Novosolov et al. (2017)		✓						✓			
Novosolov et al. (2013)							✓				
Slavenko et al. (2016)		✓									
Feldman et al. (2016)		✓									
Meiri (2018)					✓		✓	✓	✓		
Meiri et al. (2015)								✓	✓		
Roll et al. (2017)										✓	
Myhrvold et al. (2015)	B, M, R	✓	✓	✓	(✓)		✓				
IUCN (2020) ²	A, B, M										✓
IUCN (2020) ³	All										✓

461 feasible to obtain measures of intraspecific variability from all sources; therefore, estimates were provided
 462 as a single measure for each species. For some species and some traits, measures were provided separately
 463 for females and males. In such cases, I first obtained the mean of these two measures.

464 Across sources, there were multiple traits related to each of body size and lifespan. For instance, body
 465 mass and/or body length information could be provided. Different proxies were also available for lifespan,

466 such as the age at sexual maturity or generation length. In such cases, I focused on the trait presenting the
467 highest coverage.

468 • **Body size**

469 Adult body mass estimates were compiled for mammals, birds and reptiles. Body length information
470 was compiled for amphibians, because the coverage for body length was higher than that for body
471 mass. Body mass and body length are known to scale allometrically, although the allometric rela-
472 tionship differs across amphibian clades (Santini et al., 2018). In the amphibian dataset, Pearson’s
473 correlation coefficient between log(Body mass) and log(Body length) was 0.71 (data points shown in
474 Appendix 1, S2.2:‘Additional information for trait compilation’, Figure S2.2).

475 • **Longevity**

476 I defined longevity as the lifespan of an individual and maximum longevity as the longest lifespan
477 reported. I used closely related traits when longevity/maximum longevity was not available or when
478 longevity/maximum longevity had a poorer coverage than a related trait. I selected the age at sexual
479 maturity for amphibians; Pearson’s correlation coefficient between log(Age at sexual maturity) and
480 log(Maximum longevity) was 0.55 (Appendix 1, Figure S2.2). I compiled the generation length for
481 mammals and birds. The correlation between log(Generation length) and log(Longevity) was 0.74 for
482 mammals and 0.70 for birds (data points shown in Appendix 1, S2.2, Figure S2.3). Finally, I used
483 maximum longevity directly for reptiles.

484 • **Litter or clutch size**

485 The number of offspring (litter size) or eggs (clutch size) was compiled directly from the sources and
486 treated as equivalent across classes. I reported measures of central tendencies provided by the sources
487 where applicable; otherwise, I calculated range midpoints (mean of smallest and largest reported lit-
488 ter/clutch sizes).

489 • **Trophic level**

490 In all classes, species were described as omnivores, carnivores or herbivores. For reptiles and mam-
491 mals, this information was compiled directly from the sources. For amphibians and birds, trophic
492 levels were not provided. For these two classes, I inferred trophic levels from dietary information
493 (Table 2.1). For birds, I used the primary diet (based on food items recorded as composing $\geq 50\%$
494 of the diet of a species). Diet for amphibians was described without respect to the percentage use of
495 food items; simply as a binary record of whether or not food items were used. In both cases, species

496 recorded to only consume plant-based resources (seeds, nectar, fruit or other plant material) were
497 classified as herbivores. Species consuming only animal resources (invertebrates or vertebrates) were
498 classified as carnivores. Species consuming a mixture of plant and animal resources were classified as
499 omnivores.

500 • **Activity time**

501 Species were described as being either nocturnal or non-nocturnal. Despite a higher resolution of ac-
502 tivity time information in some of the sources (e.g., species being described as cathemeral, crepuscular
503 or diurnal), I adopted the classification of the source with the lowest resolution (EltonTraits: Wilman
504 et al. (2014), for birds), in order to have consistent information across classes. As such, all species
505 defined as diurnal, cathemeral or crepuscular were classified as non-nocturnal, as opposed to species
506 classified as strictly nocturnal.

507 • **Habitat breadth**

508 I used IUCN habitat data (IUCN, 2020), which describe species habitat preferences and the suitabil-
509 ity and importance of each habitat. I defined habitat breadth as the number of habitats a species was
510 known to use, using level 2 of the IUCN Habitat Classification Scheme for description of habitat
511 types (divided into: Forest, Savanna, Shrubland, Grassland, Wetland, Rocky areas, Caves and subter-
512 ranean, Desert, Marine, Marine intertidal or coastal/supratidal, Artificial, Introduced vegetation, and
513 Other/Unknown.) Note that the total number of habitats, determined by including those that qualify as
514 artificial, correlates positively with the number of natural habitats used (Figure 2.2).

515 • **Use of artificial habitats**

516 For a species, I recorded whether any artificial habitat was reported to be suitable in the IUCN habitat
517 data.

518 Finally, the compiled datasets contain an additional column, ‘Note’, where I reported species found to
519 be extinct or extinct in the wild (EW). I used species Red List status and information from Meiri (2018) to
520 flag such species. I reported 75 extinct/EW species for mammals, 160 for birds, 34 for amphibians and 53
521 for reptiles. It is likely that the datasets contain extinct species that I could not flag, because they were not
522 recorded as extinct in the sources I used.




Figure 2.2: Number of natural and artificial habitats used by species against number of strictly natural habitats used by species. Pearson's correlation coefficients show a high positive correlation between these two metrics of habitat breadth in all terrestrial vertebrate classes: 0.92 for amphibians (a), 0.95 for birds (b), 0.94 for mammals (c), and 0.90 for reptiles (d).

523 Phylogenies

524 I used class-specific phylogenetic trees downloaded on 13 April 2020. For mammals, I used ‘complete’
 525 trees from Faurby et al. (2018, 2020), downloaded from <https://zenodo.org/record/3690867#.Xyc5wyhKhPZ>. For amphibians, birds and squamates, I obtained trees from <https://data.vertlife.org/>. The original sources were as follows: Jetz et al. (2012) for birds; Jetz and Pyron (2018) for amphibians; and Tonini et al. (2016) for squamates. For each class, a distribution of 1,000 trees was available.
 526
 527 For plotting purposes, I obtained consensus trees using the TreeAnnotator program of the BEAST software
 528 (Bouckaert et al., 2019).
 529
 530

531 Species distributions

532 I obtained extent-of-occurrence distribution maps for reptiles from Roll et al. (2017), available at: <https://datadryad.org/stash/dataset/doi:10.5061/dryad.83s7k> (downloaded 13 April 2020).
 533
 534 For mammals and amphibians, species distribution maps were obtained from the IUCN Red List (IUCN

535 (2020), downloaded 13 April 2020); for birds, they were obtained from BirdLife International (<http://datazone.birdlife.org/species/requestdis>, downloaded 17 April 2020).

537 For amphibians, mammals and birds, I selected areas of extant or probably extant presence only. Addi-
538 tionally, I selected areas where species were resident or present during the breeding season, and I excluded
539 areas occupied during the non-breeding season or where species were considered vagrant.

540 In addition, for all classes, I excluded occupied areas that fell outside the known elevational limits of
541 species, where such data were available. Lower and upper elevational limits were retrieved from the IUCN
542 Red List (queried using the ‘rredlist’ package) and were available for approximately half of the species
543 (Appendix 1, S2.3: ‘Cutting distribution maps by altitudinal limits’, Figure S2.4). Decreases in range sizes
544 were observed after cutting distribution maps by the known elevational limits (Appendix 1, Figure S2.5).

545 **2.2.2 Investigating gaps and biases in trait data**

546 I used trait coverage and completeness to investigate taxonomic, phylogenetic and spatial biases in the trait
547 data. Table 2.2 summarizes the sample sizes (number of species) in each of the following analyses. Note
548 that species for which completeness was 0% were included in all analyses (for more details, see Figure 2.3).
549 Also note that I did not filter out species identified as extinct or extinct in the wild, because they represented
550 a small proportion of the datasets (0.48% for amphibians, 1.4% for both birds and mammals, and 0.50% for
551 reptiles) and also because I could not exclude such species systematically, because it is likely that I did not
552 flag them all.

Table 2.2: Number of species for each analysis. All species represented in the trait datasets were included in (1). All species from the class-specific phylogenetic trees or from the distribution maps that matched with species in the trait datasets were included in (2) and (3). *Table reproduced from Etard et al. (2020).*

	(1) Taxonomic biases	(2) Phylogenetic biases	(3) Spatial biases
Amphibians	6,990	6,170	5,650
Birds	11,634	8,315	10,802
Mammals	5,381	5,171	5,046
Reptiles	10,612	9,404	9,382

553 **Taxonomic biases**

554 I tested whether completeness varied across taxonomic class using pairwise Wilcoxon rank sum tests. I
555 tested for the extent and performance of the taxonomic corrections by looking at trait coverage when taxo-
556 nomic corrections are applied and when no correction is applied (Appendix 1, S2.4: ‘Impact of taxonomic
557 corrections on trait coverage’, Figure S2.6).

558 **Phylogenetic biases**

559 Initially, to assess whether more closely related species were more likely to be similar in trait completeness,
560 I estimated the phylogenetic signal in completeness with Pagel’s λ (Pagel, 1999) in each class. I used a boot-
561 strapping approach, calculating λ for each of 50 trees randomly sampled in each class (using the ‘phylosig’
562 function of the ‘phytools’ R package; Revell, 2012). I then estimated the mean and 95% confidence intervals
563 (95% CIs) of λ . Sample sizes for computing λ (number of species represented in both the phylogenies and
564 trait datasets) are shown in Table 2.2.

565 I then plotted within-family median completeness in phylogenetic trees built at the family level, using
566 the consensus trees. Within-family median completeness was calculated using taxonomic information in the
567 trait datasets (sample sizes shown in Table 2.2).

568 **Spatial biases**

569 I first investigated whether wider-ranging species were more likely to be better sampled than narrow-ranging
570 species. I tested for a relationship between species range size and trait completeness. I fitted a generalized
571 linear model with a Poisson error distribution (directly using the number of sampled traits, ‘N_{traits}’, rather
572 than the proportion (completeness)). Class was added as a predictor interacting with range size; thus the
573 model was:

574 $N_{traits} \sim \log(\text{Range size}) * \text{Class.}$

575 Second, I mapped assemblage-level median completeness. Assemblages were characterized at the pixel level
576 at 50 km² resolution. I determined pixel-level composition and richness by stacking species geographical
577 distributions. I then calculated median completeness across species in each pixel. I show the resulting maps
578 for herptiles in the main text, and for mammals and birds in Appendix 1 (Figure S2.7; median completeness
579 was very high across most pixels for mammals and birds). In addition, I provide maps of assemblage-
580 level mean completeness and standard deviation for all classes in Appendix 1 (Figures S2.8 and S2.9 show
581 corresponding maps; Figure S2.10 shows standard deviation against species richness).

I then tested for a spatial correlation between species richness and median completeness. Given that median completeness was very high across most pixels for mammals and birds, I fitted such models for herptiles only. I fitted spatial autoregressive lag models to explain assemblage-level median completeness as a function of species richness (using the function ‘lagsarlm’ of the ‘spatialreg’ package (Bivand and Piras, 2015; Bivand et al., 2013a; Bivand et al., 2013b)). Given that responses could vary geographically, I included the biogeographical realm as an interacting factor (using the World Wide Fund for Nature (WWF) ecoregion shapefile to characterise realms, obtained from <https://www.worldwildlife.org/publications/terrestrial-ecoregions-of-the-world>); the considered realms were Afrotropics, Australasia, Indo-Malayan, Nearctic, Neotropics and Palaearctic. To improve normality, I arc-sin square-root transformed completeness values and log-transformed species richness. The ‘lagsarlm’ function allows for a consideration of spatial autocorrelation in the dependent variable by estimating the autoregressive lag coefficient, ρ , associated with an n-by-n matrix of spatial weights, W . The final model was:

$$\text{arcsin}(\sqrt{\text{Completeness}}) \sim \log(\text{Species richness}) * \text{realm} + \rho \cdot W \cdot \text{arcsin}(\sqrt{\text{Completeness}}).$$

The value of W was estimated using the functions ‘tri2nb’ and ‘nb2listw’ of the ‘spdep’ package (Bivand and Wong, 2018; Bivand et al., 2013b). Fitting the model using all grid cells was computationally intractable; therefore, I randomly sampled cells for this analysis (using 30% of the grid cells in each realm). I selected grid cells where species richness was higher than three to avoid sampling issues. I fitted separate models for amphibians and reptiles, because when adding class as an interacting predictor, the same cells (with the same coordinates) might be sampled for multiple classes, whereas the ‘tri2nb’ function does not tolerate duplicated coordinates.

2.3 Results

2.3.1 Taxonomic biases in trait information

Trait coverage for mammals and birds was overall high (Figure 2.3a; mean and median coverage across traits: 89% and 95% for mammals; 84% and 85% for birds). In both cases, litter/clutch size was the trait with the poorest coverage (61% for mammals and 59% for birds). Coverage exceeded 80% for all other traits (except trophic level for birds, at 75% coverage).

Conversely, trait coverage was more variable for herptiles, and poorer overall (Figure 2.3a; mean and median trait coverage: 47% and 32% for amphibians, 46% and 38% for reptiles). Coverage exceeded 80% only for body size in both reptiles and amphibians and for habitat related traits in amphibians only. In all

other cases coverage was <55%, with very little information available for longevity-related traits. Trait completeness (proportion of non-missing trait values for a species) reflected similar biases (Figure 2.3b). The distribution of trait completeness varied significantly among classes (pairwise Wilcoxon rank sum test: p-value<0.0001 in all cases). Distributions were highly left skewed in mammals and birds (skewness: -2 and -1.6). 84% of all mammalian species and 80% of avian species fell in the 80–100% completeness range. Moreover, the completeness distribution was moderately right skewed for reptiles (skewness: 0.4), and slightly right skewed for amphibians (skewness: 0.02). 56% of all reptiles and 57% of amphibians fell in the 0-50% completeness range.




Figure 2.3: Trait coverage and completeness across species. (a) I defined coverage as the proportion of species for which an estimate is available for a given trait. The dashed line represents 50% coverage. (b) Trait completeness is the proportion of estimated traits for a species. Here, I show the distribution of completeness. Continuous lines represent the mean trait completeness for each class, whereas dashed lines represent the median trait completeness. Note that there were species with 0% completeness (230 species for amphibians – 3.3% of amphibian species in the trait dataset; 9 for birds – 0.077% of species; 7 for mammals – 0.13% of species; and 161 for reptiles – 1.5% of species). Species with 0% completeness were retained in the datasets when there was information for traits I did not select in the analyses, but no known value for the traits I did select. For instance, the body mass of the amphibian species *Rhinella centralis* was known, but other trait values (including body length) were missing, meaning that *Rhinella centralis* had 0% completeness for the set of traits I considered. *Figure reproduced from Etard et al. (2020).*

2.3.2 Phylogenetic biases in trait completeness

As expected from the distribution of trait completeness in mammals and birds (Figure 2.3), within-family median trait completeness was high across most tips of the phylogenetic trees (Appendix 1, Figures S2.11 and S2.12; I present the avian and mammalian phylogenies in Appendix 1 because there was little variation in completeness across tips). For birds, λ was 0.71 (± 0.0053). For mammals, λ was 0.78 (± 0.0035). This indicated that, despite completeness generally being high across tips, the sampling was not evenly distributed across the phylogeny.

626 In herptiles, clusters of families with similar median trait completeness appeared (Figure 2.4). In am-
627 phibians, groups of families belonging to the order *Anura* (frogs) showed both the best and worst median
628 completeness (Figure 2.4a). The best-sampled families included the tailed frogs of the family *Ascaphidae*
629 (two species) and species of the family *Leiopelmatidae* (four species endemic to New Zealand). The family
630 *Ceratobatrachidae* (containing c. 90 species occurring in Southeast Asia and in some Pacific islands), the
631 family *Ranidae* (true frogs, 450 species considered here) and the family *Rhacophoridae* (shrub frogs, 382
632 species considered here) figured among the worst-sampled families. For amphibians, λ was $0.63 (\pm 0.0039)$.
633 In reptiles, most snakes were poorly sampled, whereas families in other suborders appeared to be sampled
634 better overall (Figure 2.4b). Within snakes, the pythons, boas, the three species of the family *Acrochordidae*
635 and the python-like species of the family *Loxocemidae* were better sampled than other snake families. In
636 reptiles, λ was $0.69 (\pm 0.0032)$. The sampling in herptiles was thus also uneven with regard to the phylogeny.

637 It is important to underline that Figure 2.4 shows within-family median completeness, masking the con-
638 siderable variation in species richness across families, hence masking potential important variation in com-
639 pleteness across species within families. For example, in the amphibian family *Allophrynididae* (three recog-
640 nized species), the within-family median completeness was 50%; but the dataset comprised two species of
641 completeness 14% and 86%, respectively. I present similar plots to those in Figure 2.4 showing the within-
642 family standard deviation in completeness in Appendix 1 (Figure S2.13). Within-family standard deviation
643 tended to increase with within-family species richness (Appendix 1, Figure S2.14).




Figure 2.4: Within-family median trait completeness in herptiles. The number next to each family name represents the number of species included in the calculation of the median. *Figure reproduced from Etard et al. (2020).*

644 2.3.3 Spatial biases in trait completeness

645 Range size was significantly correlated with the number of sampled traits. Larger range sizes were associated
646 with a higher number of sampled traits (i.e., with higher completeness; Figure 2.5; Appendix 1, Table S2.1).
647 Similar results were obtained when using distribution maps not cut by elevational limits (Appendix 1, Table
648 S2.2; Figure S2.15). The rate of increase was steepest for reptiles, then for amphibians, then for birds and
649 mammals (slope estimates for birds and mammals were not significantly different from each other; Appendix
650 1, Table S2.1).





Figure 2.5: Relationship between number of sampled traits and geographical range size. The model was fitted using a Poisson error distribution. Class was added as a predictor interacting with range size. Rates of increase in number of sampled traits with range size were not significantly different for mammals and birds but differed for reptiles and amphibians, with the steepest rates of increase for reptiles. *Figure reproduced from Etard et al. (2020).*

651 There were marked spatial variations in median trait completeness in herptiles (Figure 2.6). North Amer-
652 ica and Europe were well sampled for both amphibians and reptiles. However, Southeast Asia and the Congo
653 basin were on average less well sampled. In other regions, contrasting patterns emerged between amphibians
654 and reptiles. For instance, median completeness was poorer for amphibians than for reptiles in Australia,
655 but opposite patterns were observed in South America. As in the phylogenetic analyses, assemblage-level
656 median completeness could mask potential important variation in completeness within species of a given
657 assemblage. Assemblage-level mean and standard deviation maps are shown in Appendix 1 (Figures S2.8
658 and S2.9). There was a trend for increasing standard deviation with increasing species richness, with a larger
659 spread in standard deviation at lower species richness (Appendix 1, Figure S2.10).

(a) Amphibians



Median completeness:

NA
0 – 10%
10 – 20%
20 – 30%
30 – 40%
40 – 50%
50 – 60%
60 – 70%
70 – 80%
80 – 90%
90 – 100%

(b) Reptiles




Figure 2.6: Spatial distribution of assemblage-level median trait completeness in herptiles. Similar maps for birds and mammals are shown in Appendix 1 (Figure S2.7). *Figure reproduced from Etard et al. (2020).*

660 Spatial models showed that species richness explained median trait completeness in herptiles in most
661 realms (Figure 2.7; Appendix 1, Tables S2.3 and S2.4); including spatial lags improved the models (am-
662 phibians: $\rho = 0.92$, p-value < 0.0001; reptiles: $\rho = 0.91$, p-value < 0.0001). For reptiles, completeness was
663 negatively correlated with species richness in the most species-rich realms (Afrotropics, Indo-Malayan and
664 Neotropics) and in the Palaearctic; the relationship was steepest in the Afrotropics and shallowest in the
665 Palaearctic. In the Australasian and Nearctic realms, completeness tended to increase with species richness.
666 For amphibians, negative relationships were observed in the Indo-Malay and Nearctic realms, whereas pos-
667 itive trends were observed in the Neotropics and the Palaearctic. The opposite trends between reptiles and
668 amphibians observed in the Australasian and Neotropical realms reflected patterns observed on the maps.
669 The Indo-Malayan was the only realm where median completeness tended to decrease with species richness
670 for both reptiles and amphibians.




Figure 2.7: Spatial model trends for herptiles. The lines represent in-sample predictions (\pm standard error) for the trend components of the spatial models (trends after accounting for spatial autocorrelation). *Figure reproduced from Etard et al. (2020).*

671 2.4 Discussion

672 The results of this Chapter illustrate the taxonomic, spatial and phylogenetic dimensions of the knowledge
 673 gaps in trait data, termed the Raunkiær shortfall by Hortal et al. (2015). To the best of my knowledge,
 674 this work constitutes the first comparative assessment of global gaps for terrestrial vertebrate trait data,
 675 despite their use in numerous studies. I showed that the trait data present important taxonomic, spatial and
 676 phylogenetic biases, with contrasts in the availability of trait information between, on the one hand, herptiles
 677 and, on the other hand, birds and mammals.

678 Birds and mammals are globally well sampled for the set of traits I considered, even in the most species-
 679 rich assemblages. However, the availability of trait information for herptiles is lower overall and phyloge-
 680 netically and geographically biased. Several factors could interplay to shape these patterns. For instance,
 681 species that are more easily detectable (for example, wider ranging) and more charismatic are likely to be
 682 better sampled. Diverse socio-economic predictors could also contribute to geographical biases in trait data
 683 sampling; global biases in primary data collection are likely to be one of the most important contributors to
 684 the patterns I highlighted. Nevertheless, biases in the data could have been introduced at later stages, notably
 685 with the selection of sources and traits. The global compilation I obtained in this Chapter reflects, in part, the

686 interest and focus of the secondary data sources I used. It is possible that the addition of new sources from
687 regional journals or other authorities could diminish spatial biases in the data by increasing coverage for
688 certain areas. Nevertheless, I argue that by focusing on widely used traits, these results are likely to reflect
689 the “true” availability of the data in primary sources and that the shortfalls for other, less used traits would
690 be more pronounced.

691 I believe that the results presented here are robust to taxonomic uncertainty, although taxonomic match-
692 ing might potentially be improved further using class-specific sources, such as the Reptile Database or Am-
693 phibiaWeb, for identification of synonyms (but see Appendix 1, Figure S2.16). I have made two versions of
694 the data compilations available, one in which my own corrections were applied and one using the original
695 binomial names of the sources, meaning that users are free to use their own taxonomic backbones and iden-
696 tify synonyms within the compilations. I believe that taxonomic matching is a recurring issue when working
697 across thousands of species. Taxonomic synonymy artefactually inflates the numbers of identified species,
698 potentially lowering trait coverage (whereas clumping subspecies together can have the opposite effect).
699 Tackling this problem is difficult (Isaac et al., 2004; Jones et al., 2012), notably because there is no global
700 curated database recording the status of species names, and also because of the nature of taxonomy and the
701 debates around the species concept (May, 2011). Nevertheless, taxonomic uncertainty can have important
702 consequences. For instance, Cardoso et al. (2017) showed that inaccuracies and errors in species checklists
703 contributed to the overestimation of plant diversity in the Amazon (but see Freeman and Pennell (2021): the
704 relative underdescription of species in tropical areas compared to temperate areas –‘taxonomic debt’, also
705 referred to as ‘latitudinal taxonomic gradient’ by the authors– may lead to the underestimation of species
706 richness at low latitudes).

707 Biases in trait data have important implications for conservation planning. Past studies have shown that
708 narrow-ranging species, for which fewer trait data are available on average, have higher extinction risks
709 (Collen et al., 2016; Purvis et al., 2000; Ripple et al., 2017) and are more negatively impacted by anthro-
710 pogenic pressures than wider-ranging species (Newbold et al., 2018). Trait information is also less available
711 for herptiles in tropical regions such as the Congo basin, Southeast Asia and South America, which are some
712 of the most diverse areas of crucial importance for worldwide conservation (Barlow et al., 2018). Conse-
713 quently, trait information is on average less available where potentially more crucial to conservation plan-
714 ning. Indeed, trait information can be incorporated into vulnerability assessments and, as such, can help to
715 prioritize conservation efforts. Species traits have been found to mediate species responses to environmental
716 changes across diverse taxonomic groups, and thus can inform on the sensitivity of species to anthropogenic
717 pressures (Flynn et al., 2009; Newbold et al., 2013; Nowakowski et al., 2017). Traits are now commonly

718 used to estimate species vulnerability or extinction risks (Pacifici et al., 2015; Ramírez-Bautista et al., 2020).
719 As opposed to trend-based approaches, which rely on historical population trends (changes in abundance
720 or shifts in distributions) to predict species' vulnerability and extinction risks, trait-based approaches rely
721 on species' intrinsic sensitivity to particular threats. The appeal of trait-based approaches to extinction risk
722 estimation is that, by providing mechanistic insights, they diminish the amount of population information
723 needed. If the responses of species to a threat consistently relate to certain traits, it is possible to generalize
724 patterns across species for which population data are less available (Verberk et al., 2013). Integrating traits
725 into vulnerability assessments is hence of particular interest when field monitoring of species population
726 sizes or distributions is difficult to achieve, but biases in the data could mean that such information is lacking
727 for some of the most vulnerable species.

728 Traits that influence species responses to environmental changes have been termed 'response traits' (or
729 'response-mediating traits'; Luck et al. (2012)), as opposed to 'effect traits' that underpin ecosystem func-
730 tioning (Lavorel and Garnier, 2002). For instance, relative brain size and longevity have been characterized
731 as response traits in birds (Newbold et al., 2013; Sayol et al., 2020), whereas dietary characteristics (e.g.,
732 trophic levels or guilds) are both response and effect traits. Hortal et al. (2015) highlighted that, for plants,
733 both response and effect traits have been investigated, whereas for vertebrates the research has been more
734 focused on understanding species responses. This could be because the way vertebrate traits interact to shape
735 some ecosystem processes has not yet been characterized well.

736 Ecosystem processes sustained by animals might be harder to quantify and might be influenced by a com-
737 bination of traits. The traits compiled in this work are likely to have a role in diverse processes. Nevertheless,
738 there was one important omission, in that I did not compile species diet in this Chapter, potentially the most
739 straightforward trait to link with diverse processes, such as grazing, pollination, scavenging and seed dis-
740 persal. From a practical perspective, I chose traits that had been estimated at least for some of the species
741 in each class, and that were readily available. Diet was excluded because although estimates were available
742 for amphibians, birds and mammals, there was no readily available database for reptilian diet. Movement or
743 dispersal abilities were also excluded because information was not readily available for any class. Although
744 I expect that species diet and dispersal abilities would present similar sampling biases to the ones presented
745 in this work, the addition of such traits to the compilation would represent a valuable contribution and would
746 notably facilitate studies looking at the functional roles of reptiles.

747 For practical reasons, I did not consider intraspecific trait variation. Intraspecific variation has been
748 shown to have important effects on ecological systems, and a growing body of literature encourages trait-
749 based research to include intraspecific variability (Guralnick et al., 2016). There have been several calls to

750 produce open-access, global trait datasets (Weiss and Ray, 2019), including a representation of intraspecific
751 trait variation (Kissling et al., 2018). Notably, Schneider et al. (2019) designed a framework to store and
752 share inter- and intraspecific trait data, accompanied by an R package to standardize the data in a proposed
753 format. Such a proposition could constitute an important step towards the unification of individual datasets
754 into a single, comprehensive database for ecological trait data.

755 The current spatial and taxonomic gaps in trait data might limit our ability to scale studies up, whereas
756 biases in the data can affect the validity of extrapolations to groups or areas that are undersampled. More gen-
757 erally, biases and gaps in biodiversity data can have important implications for ecological studies. Data gaps
758 can hinder our ability to draw conclusions on observed macroecological patterns. For example, Chaudhary
759 et al. (2016) proposed that marine species richness follows a bimodal distribution, peaking at mid-latitudinal
760 locations, and argued that these patterns were not underpinned by knowledge gaps in species distributions.
761 However, Menegotto and Rangel (2018) attributed the tropical dip in marine species richness to a lack of
762 species distribution data, explained by lower sampling efforts in tropical areas ('Wallacean' shortfall; Hor-
763 tal et al. (2015)). Biases and gaps in trait data could also affect studies in closely related fields, such as
764 functional ecology – for instance, past studies have shown that functional diversity indices are sensitive to
765 missing data (Májeková et al., 2016; Pakeman, 2014) – or community assembly (Perronne et al., 2017).

766 Ecologists should, therefore, take particular care when designing trait-based studies, because both data
767 quality and data gaps are likely to influence the results and the generality of the conclusions. There exist
768 diverse methods to deal with missing trait values, should data missingness be problematic. Complete removal
769 of missing values ('case deletion') is commonly used but presents several issues, because it reduces sample
770 size and statistical power and introduces potential bias in data subsamples (Nakagawa and Freckleton, 2008).
771 For example, retaining complete cases only from the trait datasets would generate trait data disproportionately
772 representative of mammals and birds, which would be problematic for conducting cross-taxon analyses in
773 terrestrial vertebrates. As such, it is recommended that case deletion be applied only when data are missing
774 completely at random, which is rarely the case (Peugh and Enders, 2004).

775 Alternatives to case deletion consist of filling in the gaps. In recent years, the development of impu-
776 tation techniques has provided robust methods to handle missing data. Such imputation techniques have
777 been used to complete trait datasets in recent studies (Cooke et al., 2019a). Penone et al. (2014) used a
778 simulation approach to evaluate the performance of four of these techniques, namely PhyloPars (Bruggeman
779 et al., 2009), random forest algorithms as implemented in R with missForest (Stekhoven, 2016; Stekhoven
780 and Bühlmann, 2012), multivariate imputation by chained equations (MICE; Buuren and Oudshoorn (2007))
781 and k-nearest neighbour (kNN; Troyanskaya et al. (2001)). Penone et al. (2014) introduced missing values

782 (10%–80%) in a complete trait dataset of carnivorans and measured imputation performance in different
783 scenarios. Given that phylogenetic non-randomness in missing trait values can impact imputation accuracy,
784 Penone et al. (2014) removed values in three different ways (completely at random; with a phylogenetic
785 bias; and with a body mass bias). Out of the four techniques, missForest and PhyloPars performed best when
786 species phylogenetic position was included as a predictor of missing trait values. Such imputations appeared
787 to be robust even when trait coverage was as low as 40%, which might be relevant for many reptilian and
788 amphibian traits. The performance was not significantly affected by phylogenetic non-randomness of the
789 data. Hence, missForest and PhyloPars appear to be well suited when traits are phylogenetically conserved,
790 because they allow species phylogenetic position to be included as a predictor of missing trait values. The
791 study by Penone et al. (2014) highlights that there are robust imputation techniques allowing to deal with
792 incomplete trait data where biases might otherwise be problematic. Nevertheless, it is important to highlight
793 that some imputation techniques, such as single or mean imputation, can be problematic because they do
794 not allow an estimation of uncertainty and suffer from a lack of accuracy (Nakagawa and Freckleton, 2008);
795 indeed, imputation techniques sometimes perform no better than case deletion. More work should be con-
796 ducted to assess imputation performance in various contexts (e.g., Johnson et al. (2021)), and the datasets
797 compiled in this Chapter might provide an opportunity for such studies.

798 Although robust imputation techniques can be useful for filling gaps in trait datasets, they are no substi-
799 tute for continued data collection efforts. The results of this Chapter show that data are particularly lacking
800 in herptiles, notably in the Afrotropics, the Neotropics and the Indo-Malayan realms. For these areas, incor-
801 porating regional databases into existing datasets could contribute to the reduction of global gaps. I believe
802 that both primary research and subsequent efforts to integrate new data and existing databases are required
803 if we are to collectively strive towards the unification of trait databases.

804 To conclude, this work constitutes, to my knowledge, the first assessment of the global gaps and biases
805 in terrestrial vertebrate trait information. I show that herptiles are undersampled compared with mammals
806 and birds, with important spatial and phylogenetic variability in the availability of trait information. Impu-
807 tation techniques are one possible solution to these problems. Nevertheless, I believe that primary research,
808 combined with efforts to complete existing datasets, is the only way to fill the current data gaps genuinely
809 and robustly. I hope that the compiled trait dataset and these findings can prove useful for guiding further
810 data collection efforts and for conducting macroecological analyses.

811 **3 | Intensive human land uses negatively af-**

812 **fect vertebrate functional diversity**

813 **Keywords**

814 Land use; land-use intensity; terrestrial vertebrates; functional diversity; traits.

815 **Abstract**

816 Land-use change is the leading driver of global biodiversity loss, thus characterising its impacts on the func-
817 tional structure of ecological communities is an urgent challenge. Using a database describing vertebrate
818 assemblages in different land uses, I assess how the type and intensity of land use affect the functional di-
819 versity of vertebrates globally. I find that human land uses alter local functional structure by driving declines
820 in functional diversity, with the strongest effects in the most disturbed land uses (intensely used urban sites,
821 cropland and pastures), and among amphibians and birds. Both tropical and temperate areas experience im-
822 portant functional losses, which are only partially offset by functional gains. Tropical assemblages are more
823 likely to show decreases in functional diversity that exceed those expected from species loss alone. These re-
824 sults indicate that land-use change non-randomly reshapes the functional structure of vertebrate assemblages,
825 raising concerns about the continuation of ecological processes sustained by vertebrates.

826 **3.1 Introduction**

827 Anthropogenic activities are profoundly transforming global biodiversity. Although multiple pressures act
828 in combination, land-use change currently poses the greatest threat to biodiversity (Maxwell et al., 2016;
829 Newbold et al., 2015). However, not all species respond similarly to land-use change. Traits have been found
830 to explain species' sensitivity to land-use change in diverse groups (Newbold et al., 2013; Nowakowski et

al., 2017; Quesnelle et al., 2014; Todd et al., 2017). Previous work has also shown that land-use change leads to non-random modification of assemblage trait composition (or functional diversity) (Chapman et al., 2018; Colin et al., 2018; Flynn et al., 2009; La Sorte et al., 2018; Newbold et al., 2013; Tinoco et al., 2018). Since it is widely acknowledged that biodiversity, and in particular trait diversity, may promote ecosystem functioning and stability, modification to the trait composition of assemblages could have far-reaching and adverse impacts on ecological processes (Hooper et al., 2012; Magioli et al., 2021; Oliver et al., 2015; Tilman and Downing, 1994).

Terrestrial vertebrates support many processes, ranging from pollination (Ratto et al., 2018), to seed dispersal to the regulation of lower trophic levels (Barber et al., 2010; Letnic et al., 2012; Salo et al., 2010; Zhang, Y. et al., 2018). However, we lack a global understanding of how the functional diversity of entire vertebrate assemblages responds to changes in land use. Most previous studies have been conducted at regional or local scales (Davison et al., 2021), but these may not be representative of global patterns. Indeed, recent global syntheses have highlighted how biodiversity responses can differ substantially between regions and across latitudes, with higher sensitivity reported for the tropics (Matuoka et al., 2020; Millard et al., 2021; Newbold et al., 2020b). Another key issue is the taxonomic coverage of past work. Few studies investigating effects of land use on functional diversity have considered several vertebrate classes together, and comparative studies remain rare. Thus, how land-use change affects the functional diversity of local vertebrate assemblages at global scales, and the potential geographical and taxonomic variation in the effects, still largely remains to be explored.

Here, I aim to assess how human land use and land-use intensity affect the functional diversity of vertebrate assemblages, across and within taxonomic classes. Building on recent work (Matuoka et al., 2020; Millard et al., 2021; Newbold et al., 2020b), I investigate differences in response between tropical and temperate regions. I use multiple response metrics to quantify functional diversity. First, functional richness measures the breadth and variety of trait combinations represented in an assemblage (Legras et al., 2018). Second, functional dispersion quantifies how similar species in a given assemblage are in terms of their traits (Laliberté and Legendre, 2010). These metrics can mask important alterations of assemblage composition if functional losses are compensated for by functional gains. To address this, I consider pairwise measures between assemblages, to explore levels of functional loss and functional gain across land uses (Figure 3.1).

To this end, I combine (1) the trait data across terrestrial vertebrates collected in Chapter 2 with (2) global records of species occurrence in eight land-use types of differing intensity of use (the PREDICTS database: Hudson et al. (2014, 2017); Figure 3.1; Appendix 2, Table S3.1). The PREDICTS database is currently the most comprehensive database of sampled species occurrence, and for most records also

863 abundance, across multiple land uses of different land-use intensity. Using the PREDICTS database allows
864 me to contrast biodiversity metrics among intact land uses (primary-vegetation sites, considered to be the
865 undisturbed reference condition), and all other human land-use types. Specifically, I test the following
866 hypotheses, both across and within taxonomic classes:

- 867 1. I expect decreases in functional diversity in human land uses compared to primary vegetation, caused
868 by contractions of occupied trait space. I expect such effects to be more pronounced where land is
869 used more intensively by humans. This hypothesis builds upon evidence that species with certain
870 traits are more sensitive to land-use disturbance (Newbold et al., 2013), meaning that disturbed land
871 uses will retain only disturbance-tolerant species, more functionally similar to one another. Given the
872 reported higher sensitivity of tropical assemblages to land-use disturbance, I predict that such effects
873 are stronger in the tropics.
- 874 2. I hypothesise that decreases in functional diversity in disturbed land uses exceed decreases expected
875 by chance, given local species loss. Thus, I expect disturbed land uses to promote functional under-
876 dispersion. Functional under-dispersion occurs when species within an assemblage are more similar,
877 in term of their traits, than expected by chance (Cadotte and Tucker, 2017; Wong et al., 2018) – or,
878 in other words, when functional dispersion is lower than expected given local species richness. I
879 predict that under-dispersion is more likely to occur in the highly disturbed sites, in both temperate
880 and tropical areas. This hypothesis is based on the idea that species are being removed non-randomly
881 from sensitive areas of the trait space, and increasingly so with higher disturbance level.
- 882 3. Finally, I expect decreases in functional diversity in human land uses to be driven by high functional
883 loss, whereby species are being removed from previously occupied areas of the trait space; I expect
884 no functional gain. This hypothesis is based on the idea that the functional trait space in undisturbed
885 land uses represents all of the possible regional trait combinations and that species with functional
886 attributes rendering them unable to persist in altered conditions will be filtered out (Cornwell et al.,
887 2006).

888 3.2 Methods

889 3.2.1 Vertebrate assemblages

890 I used vertebrate occurrence data from the PREDICTS database (Hudson et al., 2014, 2017), a collection of
891 studies that recorded species occurrence across multiple land uses and land-use intensities. In PREDICTS,




Figure 3.1: Overview of the study design and functional diversity metrics. I used occurrence data for vertebrate species from the PREDICTS database ((Hudson et al., 2014, 2017); 180 studies; 431,170 records; 4,339 species; 6,758 sampled sites). (a) shows the spatial distribution of sites I consider. I combine occurrence data with trait data compiled in Chapter 2 to calculate functional diversity metrics. (b) is a representation of the trait data in two dimensions, plotted across PREDICTS vertebrates. Traits that contributed most to dimension 1 were lifespan (29%) and litter/clutch size (22%), while traits that contributed most to dimension 2 were habitat breadth (47%) and use of artificial habitats (35%). (c) and (d) present the conceptual framework for the calculation of the functional diversity metrics: local measures (c) and pairwise metrics (d). (c) Given a trait space, functional richness is calculated as the hypervolume occupied by the minimum convex hull encompassing all species (Villéger et al., 2008). Functional dispersion is calculated as the mean distance of the species to the centroid, g (Laliberté and Legendre, 2010). (d) I compute functional loss as the proportion of hypervolume lost from the reference assemblage, and I define functional gain as the proportion of hypervolume of the disturbed assemblage that was gained (proportion of novel trait space in the disturbed assemblage). *Figure reproduced from Etard et al. (2021).*

each study contains several sites, which may be clustered into spatial blocks. Assemblage and land-use data are available at the site level: one site is characterised by a unique land use of given land-use intensity and provides occurrence data for a set of sampled taxa (and the same set of taxa is sought at all other sites within a study). Sites located between 23.5°N and 23.5°S of latitude were considered tropical, and otherwise temperate (Figure 3.1).

Land uses in PREDICTS were assigned to the following categories, based on the descriptions of the habitat given by the original collectors of the data: primary vegetation (considered to be the undisturbed reference); secondary vegetation; plantation forest; pasture; cropland; urban (considered human, or disturbed; Appendix 2, Table S3.1; Hudson et al., 2014, 2017). Secondary vegetation is further divided into three cat-

901 egories: mature, intermediate and young, depending on the stage of recovery of the vegetation. Land-use
902 intensity is reported as minimal, light or intense, according to criteria that depended on the land-use type in
903 question (e.g., crop diversity, degree of mechanisation and chemical inputs in cropland, or bushmeat harvest-
904 ing and selective logging in primary vegetation; Hudson et al., 2014). I excluded sites for which the land use
905 could not be characterised or for which the stage of recovery of secondary vegetation was unclear. As the
906 PREDICTS database is a collection of independent studies, the design of this study was not balanced: the
907 sample size varied across land uses (Appendix 2, Figures S3.1 & S3.2), and across taxonomic groups (3,103
908 species of birds; 531 mammals; 379 amphibians; 326 reptiles).

909 **3.2.2 Functional traits and diversity metrics**

910 Trait choice is a critical step when calculating functional diversity metrics, which are highly sensitive to
911 trait selection (Mouillot et al., 2021). However, trait selection trades off with data availability. Here, a
912 constraint was to use similar traits across the different classes. Thus, I used the seven traits compiled in
913 Chapter 2 across terrestrial vertebrates. Most of these traits were available for at least 50% of the species
914 in each class (except trophic level in amphibians and lifespan in herptiles; Appendix 2, Figure S3.4). In
915 addition, I chose traits that were ecologically relevant, broadening the biological definition of traits (i.e.,
916 a characteristic measurable at the level of an individual) to include measures of habitat breadth and habitat
917 specialisation (still theoretically measurable at the level of an individual). The final set constituted seven traits
918 that influence species' responses to environmental change: body mass, trophic level, lifespan, litter/clutch
919 size, diel activity, habitat breadth and use of artificial habitats. These traits related to life-history, habitat
920 specialisation and use of geographical space (e.g., habitat breadth significantly explains geographical range
921 size in all classes; Appendix 2, Figure S3.3). Here, I did not consider estimations of dispersal abilities
922 or home range size as these were available for a small fraction of the species (<3%, Alex Smith and M.
923 Green, 2005; Paradis et al., 1998; Sutherland et al., 2000; Whitmee and Orme, 2013), neither did I include
924 geographical range size which is measured across many individuals, and hence cannot be considered a trait.
925 As in Chapter 2, I did not consider intraspecific trait variation, thus assuming no effect of the environment
926 on trait values.

927 Trait coverage was variable among classes and traits, with important gaps for reptiles and amphibians
928 (Appendix 2, Figures S3.4 & S3.5; Chapter 2). I imputed missing trait values using random forest algorithms
929 ('missForest' package: Stekhoven and Bühlmann (2012), Stekhoven (2016)), including traits, taxonomic
930 order and phylogenetic eigenvectors as predictors (Debastiani et al., 2021; Penone et al., 2014). To further
931 assess the sensitivity of the results to imputation (see next section), I imputed missing trait values eight times,

932 thereby obtaining eight sets of imputed traits. I randomly selected one imputed trait set for the calculation of
933 functional metrics. Imputations of missing trait values & imputation performance are detailed in Appendix
934 2, S3.2: ‘Trait data & imputation of missing trait values’ and S3.4: ‘Imputation performance’ (and Figures
935 S3.6-S3.8). Post-imputation, continuous traits were \log_{10} -transformed (except habitat breadth which was
936 square-rooted) and z-scored (standardised to unit variance and zero mean). In addition, I assessed whether
937 the results were robust to imputation error using a subset of the PREDICTS data considering only species
938 for which I had complete trait information (see next section).

939 Correlation among traits can be a safeguard against high sensitivity of functional metrics to trait omis-
940 sion, notably where omitted traits correlate strongly with traits that are already included in the calculation
941 (Mouillot et al., 2021). Nevertheless, high multicollinearity among traits has been reported as potentially
942 problematic for the calculation of functional diversity (Cadotte et al., 2011). Thus, I verified that the degree
943 of multicollinearity among traits was not problematically high (with a threshold of 5 for variance inflation
944 factors, Appendix 2, Table S3.3). Furthermore, I tested the sensitivity of the results to trait omission, by
945 investigating whether adding geographical range size in the calculation of functional metrics was likely to
946 affect the results.

947 3.2.3 Effects of land use and land-use intensity on FRic and FDis (Hypothesis 1)

948 For each assemblage, I measured functional richness using ‘FRic’ (Villéger et al., 2008), and functional
949 dispersion using ‘FDis’ (Laliberté and Legendre (2010); Figure 3.1), from the ‘FD’ package (Laliberté and
950 Legendre (2010); Laliberté et al. (2015)). I assessed the effects of land use, land-use intensity, and region
951 (temperate versus tropical) on FRic and FDis across and within taxonomic classes using linear mixed-effects
952 models (‘lme4’ package, Bates et al. (2015)). Land use and land-use intensity were not ranked in the models.
953 A random intercept of study identity accounted for variation in experimental design across studies, while a
954 random intercept representing spatial blocks of sampled sites, nested within study, accounted for spatial
955 structuring within studies. To improve normality and bound predictions between 0 and 1, I transformed FRic
956 and FDis using an arcsin-square-root transformation. The best-fitting model was sought using backwards
957 stepwise model selection, starting with the most complex model that included all two-way interactions among
958 the specified main effects. Model fits were compared using likelihood-ratio tests at each iteration of the
959 selection procedure.

960 Across vertebrates, the starting models included the effects of land use, land-use intensity and region
961 (temperate versus tropical). The best-fitting model for FRic was:

962 $\text{arcsin}(\sqrt{\text{FRic}}) \sim \text{Land use} + \text{Land-use intensity} + \text{Region} + \text{Land use : Land-use intensity} + \text{Land use : Region}$.

964 (Model 1a)

965 For FDis, the best-fitting model did not include interactions between land use and region, but the main
966 effect of region was retained:

967 $\text{arcsin}(\sqrt{\text{FDis}}) \sim \text{Land use} + \text{Land-use intensity} + \text{Region} + \text{Land use : Land-use intensity}$.

968 (Model 1b)

969 To investigate differences in responses across classes, I pooled some of the land uses together, because
970 otherwise, sample sizes would have been too low. Mature, intermediate and young secondary vegetation were
971 grouped together as ‘Secondary vegetation’, and cropland and pasture were grouped together as ‘Agricultural
972 land uses’. The starting models included the effects of land use, land-use intensity, region and taxonomic
973 class. For FRic, the best model was:

974 $\text{arcsin}(\sqrt{\text{FRic}}) \sim \text{Land use} + \text{Land-use intensity} + \text{Region} + \text{Class} + \text{Land use : Land-use intensity} + \text{Land use : Class} + \text{Land-use intensity : Region} + \text{Class : Region}$.

976 (Model 2a)

977 For FDis, regional effects were dropped:

978 $\text{arcsin}(\sqrt{\text{FDis}}) \sim \text{Land use} + \text{Land-use intensity} + \text{Class} + \text{Land use : Land-use intensity} + \text{Land use : Class} + \text{Land-use intensity : Class}$.

980 (Model 2b)

981 To assess whether the results were robust to imputation error, I used a subset of the PREDICTS data
982 considering only species for which there were complete trait information (6,212 sites; 442 mammals; 1,975
983 birds; 78 reptiles; 9 amphibians), and I fitted models again to this data subset. I did not have enough complete
984 trait data among amphibians to be able to consider this class separately, so I first considered amphibians and
985 reptiles together (herptiles), and reptiles only. In addition, I complemented this validation with a sensitivity
986 analysis to variation in imputed values. I calculated FDis and FRic using each of the eight imputed trait
987 datasets and fitted the previous models to each set. I then qualitatively evaluated the congruence of the
988 estimates from the different models. Finally, because there tended to be more sites sampled in primary
989 vegetation than in other land uses (Appendix 2, Figures S3.1 & S3.1), I ran additional sensitivity tests to
990 assess whether the results were robust to resampling primary vegetation sites to a number equal to 50 (a
991 sample size close to the median number of sites sampled in land uses other than primary vegetation in both
992 regions (median = 37 for the temperate subset and 57 for the tropical subset, Appendix 2, Figure S3.1)).

993 **3.2.4 Investigating functional under-dispersion (Hypothesis 2)**

994 To assess whether effects of land use and land-use intensity on FDis differed from what would be expected
995 by chance given changes in local species richness, I generated null expectations of FDis at each site. I
996 randomised assemblage composition 500 times, drawing species from the corresponding study's species
997 pool while maintaining local species richness. For each site, I thus obtained a null distribution for FDis.
998 Then, I tested whether FDis differed from null expectations using Wilcoxon signed-rank tests. I created
999 a binary variable which was assigned 1 if FDis was significantly lower than null expectations at a given
1000 site (significant under-dispersion), and 0 otherwise. I investigated how land use, land-use intensity, region
1001 and taxonomic class affected the probability of occurrence of under-dispersion using a generalised linear
1002 mixed-effects model with a binomial distribution of errors. The best-fitting model did not retain any effect
1003 of taxonomic class:

1004 $P_{\text{under-dispersion}} \sim \text{Land use} + \text{Land-use intensity} + \text{Region} + \text{Land use : Land-use intensity} + \text{Land use :}$
1005 Region.

1006 (Model 3)

1007 **3.2.5 Functional loss and functional gain (Hypothesis 3)**

1008 I calculated the proportion of trait space that was lost in disturbed land uses compared to reference land uses
1009 (functional loss) and the proportion of trait space that was gained in disturbed land uses (functional gain)
1010 (Figure 3.1c), across and within taxonomic classes. I selected studies where at least one site was sampled in
1011 primary vegetation. I then made within study pairwise comparisons between reference assemblages, sampled
1012 in primary vegetation, and disturbed assemblages. In addition, I considered all comparisons between pairs
1013 of primary-vegetation sites, to create reference pairs. I then investigated how land use, land-use intensity
1014 and region affected functional loss and gain across and within taxonomic classes using linear mixed-effects
1015 models, controlling for study identity in the random effects. Across vertebrates, the best-fitting model for
1016 functional loss was:

1017 $\text{arcsin}(\sqrt{\text{loss}}) \sim \text{Land use} + \text{Land-use intensity} + \text{Region} + \text{Land use : Land-use intensity} + \text{Land use :}$
1018 Region.

1019 (Model 4a)

1020 For functional gain, one interaction term (land use with region) was dropped:

1021 $\arcsin(\sqrt{\text{gain}}) \sim \text{Land use} + \text{Land-use intensity} + \text{Region} + \text{Land use : Land-use intensity}$.

1022 (Model 4b)

When considering the effects of taxonomic class, the best-fitting model for functional loss was:

1024 $\text{arcsin}(\sqrt{\text{loss}}) \sim \text{Land use} + \text{Land-use intensity} + \text{Class} + \text{Region} + \text{Land use :}$

1025 Land-use intensity + Land use : Class + Land use : Region + Land-use intensity : Class.

1026 (Model 5a)

For functional gain (Model 5b), the fitted effects were the same as those of Model 2b. More details about the calculation of functional loss and gain can be found in Appendix S3.5.

1029 All data analyses were conducted using R version 3.5.1 (R Core Team, 2018). I made the code available
1030 on figshare (DOIs: <https://doi.org/10.6084/m9.figshare.14161883> and <https://doi.org/10.6084/m9.figshare.15163926>), as well as the main result datasets (<https://doi.org/10.6084/m9.figshare.15163971>).

1033 3.3 Results

1034 3.3.1 Effects of land use on FRic and FDis

1035 Across all vertebrates, land use and land-use intensity significantly affected FRic and FDis (Figure 3.2).
1036 FRic tended to decrease with increasing disturbance level and higher intensity of land use. For FRic, relative
1037 effects differed between regions (Figure 3.2a). Although declines were overall more important for disturbed
1038 tropical assemblages, significant declines were observed for the temperate assemblages (e.g., a 37% average
1039 decline in intensely used urban areas; a 49% decline in pastoral areas of high land-use intensity). Never-
1040 theless, tropical assemblages typically showed more important reductions in FRic. For instance, declines
1041 averaged 63% for intensely used tropical cropland and 76% for urban areas. For FDis, relative effects were
1042 similar in both regions (Figure 3.2b). The most important average declines were observed for urban assem-
1043 blages of intense use (20% decline), and for lightly- and intensely used cropland (by 15% and 26%). Note
1044 that confidence intervals around the estimated average declines were large in some cases, highlighting some
1045 heterogeneity in the responses.

1046 Fitting the same models to the subset of species with complete trait data, I detected important declines in
1047 functional diversity in a number of land uses, showing that the conclusions are robust to trait imputation un-
1048 certainty (for example, FRic declined on average by 75% in intensely used temperate pastoral assemblages;
1049 by 48% for intensely used tropical cropland; and FDis declined by an average 37% in intensely used tropical




Figure 3.2: Effects of land use, land-use intensity and region on FRic (a) and on FDis (b) across all PREDICTS vertebrates. Effects were rescaled with reference to primary vegetation (PV) and are plotted as a % difference relative to PV within each land-use intensity. For FRic, the best-fitting model included interactions between land use and region, while these interactions were dropped for FDis, explaining the similar relative effects in both regions. Error bars represent 95% confidence intervals. MSV: mature secondary vegetation; ISV, intermediate secondary vegetation; YSV, young secondary vegetation; PF, plantation forest; PA, pasture; CR, cropland; UR, urban. Effects for intense use in MSV could not be estimated as there were not enough sampled sites. *Figure reproduced from Etard et al. (2021).*

1050 urban assemblages; Appendix 2, Figure S3.18). Furthermore, using the subset of species with complete trait
 1051 data, I found that the results were not sensitive to the inclusion of geographical range size as an additional
 1052 trait (Appendix 2, Figure S3.19). Finally, the results were not sensitive to variation across imputed trait val-
 1053 ues (Appendix 2, Figure S3.20) and were also robust to resampling in primary-vegetation sites (Appendix 2,
 1054 Figure S3.21).

1055 Responses of FRic and FDis to land use and land-use intensity differed among taxonomic classes (Figure
 1056 3.3). Within-class effects for FDis were similar between regions. The most notable decreases were observed
 1057 in lightly- and intensely used agricultural land uses in amphibians, birds and reptiles; and in intensely used
 1058 urban land uses for birds and mammals. For FRic, the effects in tropical and temperate regions were qual-
 1059 itatively similar in three out of four classes (birds, mammals and reptiles), although effect sizes tended to be
 1060 bigger for tropical assemblages. Birds and reptiles showed reductions in disturbed land uses in both tropical

and temperate regions, whereas I detected few significant effects for mammals. For birds, the most important average decline, of 50%, was observed in intensely used tropical urban land uses, while for reptiles I detected significant decreases in lightly- and intensely used agricultural sites (but I could not estimate effects for urban land uses due to the small sample size). Finally, the effects differed between tropical and temperate regions for amphibians, with no significant effects detected across temperate assemblages, but important reductions across tropical agricultural and urban assemblages.




Figure 3.3: Effects of land use, land-use intensity, region and taxonomic class on FRic (a) and FDis (b). Effects were rescaled with reference to primary vegetation (PV) and are plotted as a % difference relative to PV within each land-use intensity. Error bars represent 95% confidence intervals. Effects for FRic were estimated from Model 2a, and from Model 2b for FDis. SV, secondary vegetation; PF, plantation forest; AGR, agricultural land uses (pasture and cropland); UR, urban. Effects for reptiles in urban land uses could not be estimated as there were not enough sampled sites. *Figure reproduced from Etard et al. (2021).*

Fitting similar models only for species with complete trait data showed that these patterns are unlikely to be affected by imputation uncertainty for birds; for mammals and reptiles, the main results could even be conservative (Appendix 2, Figures S3.22, S3.23). Indeed, although confidence intervals around the estimates

were large, I typically observed larger decreases in functional diversity when using the complete data subset, including an 86% decline in FRic for mammals in intensely used tropical agricultural areas. The results were also unaffected by variation across replicate sets of imputed trait values (Appendix 2, Figure S3.24).

3.3.2 Changes in the probability of occurrence of functional under-dispersion

Land use, land-use intensity and region significantly affected the probability of occurrence of functional under-dispersion across vertebrates. Functional under-dispersion was more likely to occur in tropical crop-land of all land-use intensities (Figure 3.4b), as well as in some of the lightly-used land uses (notably urban and plantation forest). Contrary to my expectations, and with the exception of tropical cropland, functional under-dispersion was not more likely to occur in intensely-used land uses. For minimally-used sites, changes in FDis were mostly consistent with changes expected given local species richness.




Figure 3.4: Effects of land use, land-use intensity and region on the probability of occurrence of functional under-dispersion. Error bars represent 95% confidence intervals. PV: primary vegetation; MSV, mature secondary vegetation; ISV, intermediate secondary vegetation; YSV, young secondary vegetation; PF, plantation forest; PA, pasture; CR, cropland; UR, urban. Effects are rescaled and represent the average difference in the probability of occurrence of functional under-dispersion between the reference (PV, probability of functional under-dispersion set at 0 within each land-use intensity) and the disturbed land uses. *Figure reproduced from Etard et al. (2021).*

1080 3.3.3 Functional loss and gain

1081 Across and within vertebrate classes, I detected high levels of functional loss, exceeding the natural turnover
 1082 between primary-vegetation sites, both in temperate and tropical regions. Across vertebrates (Figure 3.5a),
 1083 functional loss was notably high in temperate pastures (+27% above reference for minimal use; +73% for
 1084 intense use), temperate urban sites (+27% for light use; +50% for intense use; effects for tropical urban sites
 1085 could not be estimated), temperate and tropical cropland (+44% and +56% respectively for light use; effects
 1086 for intense use could not be estimated). Important levels of functional loss were also observed in tropical
 1087 plantation forest of light use intensity (+51%; effects for the intense use could not be estimated). High
 1088 levels of functional loss were also observed within each class (Figure 3.6a) (although not all effects could
 1089 be estimated because of limited sample sizes, Appendix 2, Table S3.5). The highest losses were observed
 1090 in agricultural areas for amphibians and reptiles, with important losses also observed in temperate urban
 1091 areas for both birds and amphibians (+35% for minimal use; effects for tropical urban areas could not be
 1092 estimated).




Figure 3.5: Effects of land use and land-use intensity on (a) functional loss and (b) functional gain across pairs of sites in temperate areas and tropical areas. PV: primary vegetation; SV, secondary vegetation; PF, plantation forest; PA, pasture; CR, cropland; UR, urban. Error bars represent 95% confidence intervals. Effects are rescaled and represent the average difference in functional loss and gain between the reference pairs (PV/PV, with loss and gain set at 0% within each land-use intensity), and pairs of primary vegetation with disturbed land uses. Negative effects mean that, on average, levels of functional loss (or gain) were lower than functional loss (or gain) observed between pairs of primary-vegetation sites, and not that there were absolute ‘negative losses’ or ‘negative gains’. *Figure reproduced from Etard et al. (2021).*

1093 Across vertebrates, average functional gain (average proportion of novel trait space in the disturbed
 1094 assemblage) was moderate and on average did not exceed 20% in any disturbed land uses (Figure 3.5b).
 1095 Patterns of functional gain were similar in both regions. The highest functional gains were observed for
 1096 minimally-used urban sites and plantation forest (range: +16% to +20%). On the other hand, important
 1097 levels of functional gain were observed in some classes (Figure 3.6b), with the highest functional gain for
 1098 mammals (+80% in intensely used urban sites).

1099 Diagnostic plots (qq-plots and residual distributions) for the models are shown in Appendix 2, Figures
 1100 S3.9–S3.17. Overall, the model residuals were appropriately distributed (but with some leptokurtic residual
 1101 distributions, to which mixed-effect models are generally robust (Schielzeth et al., 2020)).




Figure 3.6: Effects of land use, land-use intensity, region and taxonomic class on functional loss (a) and functional gain (b) across pairs of sites. PV: primary vegetation; SV: secondary vegetation; PF: plantation forest; AGR: agricultural land uses (pasture and cropland); UR: urban. Error bars represent 95% confidence intervals. Effects are rescaled and represent the average difference in functional loss and gain between the reference pairs (PV/PV, with loss and gain set at 0% within each land-use intensity), and pairs of primary vegetation with disturbed land uses. Negative effects mean that, on average, levels of functional loss (or gain) were lower than functional loss (or gain) observed between pairs of primary-vegetation sites, and not that there were absolute ‘negative losses’ or ‘negative gains’. Figure reproduced from Etard et al. (2021).

1102 **3.4 Discussion**

1103 Here, I showed that the functional diversity of vertebrate assemblages is negatively impacted in human land
1104 uses, particularly in the most intensely used land types. The results of this Chapter extend previous studies
1105 that have been more taxonomically or geographically restricted (Flynn et al., 2009; Matuoka et al., 2020).
1106 Matuoka et al. (2020) found that the functional diversity of tropical bird assemblages was negatively affected
1107 by human disturbance, a pattern that did not appear in temperate assemblages. Yet, I found that functional
1108 diversity was negatively affected in both tropical and temperate areas, with important functional losses in all
1109 four vertebrate classes.

1110 Using multiple metrics allowed me to explore different facets of functional diversity. For instance, func-
1111 tional gain could locally offset functional loss in some disturbed land uses. This could indicate that despite
1112 no apparent negative effect on FRic, some disturbed land uses (e.g. lightly-used temperate cropland) could
1113 experience important functional loss, and highlights the importance of using a variety of indicators. This
1114 mechanism could be at play in mammalian assemblages, for which important levels of functional gain were
1115 observed in agricultural and urban sites. Further, functional gain in disturbed land uses could indicate that
1116 disturbances facilitate the introduction of functionally novel species, falling into previously unoccupied parts
1117 of the trait space. This may be because non-native species are more likely to become established in disturbed
1118 assemblages. Previous work has shown that land-use disturbance facilitates biological invasions in island
1119 ecosystems (Jesse et al., 2018; Sanchez-Ortiz et al., 2019), but to my knowledge, this has not been tested
1120 specifically across continental areas for invasive vertebrates (but see Pyšek et al. (2010)). It is also pos-
1121 sible that disturbed areas harbour synanthropic species that do not occur in primary vegetation, leading to
1122 substantial functional gain.

1123 Overall, the negative effects of land use on functional richness tended to be more pronounced in the
1124 tropics. This is congruent with past studies that have found tropical biodiversity to be disproportionately
1125 sensitive to human pressures (Martins and Pereira, 2017; Newbold et al., 2020b). There are a number of
1126 potential explanations for this. First, it could be that a long history of intense land-use disturbance at large
1127 scales in many temperate regions (e.g. Western Europe; Stephens et al. (2019)) means that biodiversity is
1128 now less sensitive to new disturbances, because the most sensitive species have been filtered out (Balmford,
1129 1996; Krauss et al., 2010; Le Provost et al., 2020; Munteanu et al., 2020). Species unable to cope with such
1130 disturbances may have gone extinct in the past, while the remaining species would be more disturbance-
1131 tolerant (Betts et al., 2019). Tropical regions, historically less disturbed at large scales, would then contain
1132 a higher proportion of disturbance-sensitive species than temperate regions. Consequently, the functional

richness in undisturbed tropical sites could be less resilient to new disturbances. This also highlights that time since land-use conversion could have important impacts on local functional diversity. Although I did not consider the effects of time since land-use conversion in this work (notably because PREDICTS contained data on time since land-use conversion only for about 22% of the sites), I expect that time since land-use conversion may affect assemblage composition, and thus, functional diversity, with potentially land-use-specific relationships between time since conversion and functional diversity (e.g., a positive relationship for recovering secondary vegetation or a negative relationship for urban areas; but I did not detect such effects when using the data subset for which there were information on time since land-use conversion [see Appendix 2, S3.8: ‘Model robustness – time since land-use conversion’]).

Second, it could be that tropical species are intrinsically more sensitive to disturbances than temperate species because of their evolutionary history. Natural climatic variability experienced by species as well as species history of exposure to disturbances have been proposed to influence sensitivity to disturbance. For instance, tropical species are, on average, nearer to their climatic limits than temperate species (Deutsch et al., 2008; Sunday et al., 2014). Tropical species could therefore experience more deleterious effects from interacting drivers of change, with land-use change bringing about novel climatic conditions pushing them beyond their tolerance limits (Frishkoff et al., 2016; Williams and Newbold, 2020).

In addition to filtering out sensitive species, land-use change is also expected to modify interactions among species, thereby influencing species persistence (Tylianakis et al., 2008; Valiente-Banuet et al., 2015). Although I detected a signal of functional under-dispersion (particularly in tropical cropland), which indicates that assemblages may be locally structured by environment filtering (Bregman et al., 2015), it is likely that several assembly rules underpin assemblage composition (Fournier et al., 2016). For instance, land-use changes could enhance competition among species, promoting over-dispersion by removing species that share similar resources. Such opposite signatures of environmental filtering and enhanced competition on functional dispersion could explain why I did not detect stronger effects of land use on functional under-dispersion occurrence.

Studies looking at impacts of global land use on functional diversity computed with species from all four terrestrial vertebrate classes remain rare. Lack of availability of standardised trait data across terrestrial vertebrates may have hindered such studies from being conducted in the past. To overcome this problem, I based the analyses on a large-scale collation of trait data (Chapter 2; Etard et al. (2020)), and I imputed missing trait values to obtain complete trait datasets in each class. I used random forest algorithms, currently thought to be one of the most robust technique for missing value imputations in trait datasets (Debastiani et al., 2021; Johnson et al., 2021; Penone et al., 2014). Replicating the analyses on complete trait data

1165 subsets showed that imputation uncertainty did not affect the main conclusions of this work and that the
1166 negative effects of human land uses were in some cases even stronger when using the complete data subsets.
1167 Furthermore, the results were highly consistent across imputed datasets and so insensitive to variation across
1168 imputed values. Although missing value imputation can offer a robust filling of missing entries, this study
1169 highlights the existing taxonomic biases both in trait data availability and in PREDICTS studies, and thus
1170 stresses the need to pursue data compilation efforts, particularly for the least-sampled classes (reptiles and
1171 amphibians).

1172 Another implication of trait data availability for vertebrates is that the choice of traits was constrained.
1173 Mouillot et al. (2021) showed that functional diversity metrics are sensitive to trait omission and that the
1174 sensitivity to trait omission decreases with increasing levels of correlation among traits. Here, I chose seven
1175 traits that were available across all classes at least for a subset of the species and that have been implicated in
1176 shaping species responses to environmental change. A notable omission was any metric of dispersal ability,
1177 which is likely to influence species' ability to respond to land-use change but is difficult to obtain for most
1178 species. In fact, past studies have shown that dispersal abilities can be predicted from ecological correlates,
1179 such as body mass, diet or geographical range size (Schloss et al., 2012; Sutherland et al., 2000). Since the
1180 results were robust to the omission of geographical range size, I am confident that the omission of dispersal
1181 abilities also does not affect the conclusions of this work.

1182 Functional diversity metrics are often used as a proxy for ecosystem functioning because of the concep-
1183 tual and mechanistic link between functional 'effect' traits and ecosystem processes (Lavorel and Garnier,
1184 2002; Violle et al., 2007). In many studies focused on vertebrates, however, functional diversity metrics do
1185 not correlate with a given ecosystem function (Hatfield et al., 2018). Here, I did not explicitly target given
1186 ecosystem functions, but I argue that evidence of functional loss of vertebrate assemblages indicates that
1187 processes sustained by vertebrates are put at risk by land-use change. My results further show that some
1188 disturbed land uses are more likely to experience functional under-dispersion, particularly tropical cropland
1189 and tropical urban areas, which again indicates a potential imperilment of ecological processes. Indeed, in
1190 such cases, decreases in functional dispersion exceed changes expected from the chance removal of species;
1191 such non-random modifications indicate that certain areas of the functional trait space are more sensitive to
1192 land-use disturbance. Future work could investigate the impacts of land-use change on particular ecosystem
1193 functions. The integration of trophic information (beyond the trophic levels I used here) to the species-trait
1194 dataset could be an interesting step in that direction, as dietary traits relate to resource use and are, as such,
1195 probably the most straightforward traits to link with ecosystem functions. Furthermore, my results suggest
1196 that the functional loss experienced within a class is unlikely to be compensated for by the persistence of

1197 functionally similar species in other classes. Indeed, I detected negative effects of human land use on func-
1198 tional richness in at least three out of four vertebrate classes (amphibians, birds, and reptiles), in accordance
1199 with past studies focusing on each of these groups (Gallmetzer and Schulze, 2015; Marcacci et al., 2021;
1200 Riemann et al., 2017; Sol et al., 2020). Although overall mammalian functional richness was less affected,
1201 high levels of functional gain suggest that the functional composition of mammalian assemblages is heavily
1202 modified in disturbed land uses.

1203 To conclude, the results of this Chapter highlight the negative impacts of human land uses on multiple
1204 dimensions of functional diversity, within and across terrestrial vertebrate classes, at a global scale. In
1205 many disturbed sites, decreases in functional diversity exceed changes expected from species loss alone,
1206 showing that human activities non-randomly reshape ecological assemblages. By intensifying functional
1207 loss and promoting functional under-dispersion, land-use change could have deleterious effects on ecosystem
1208 functioning, highlighting the necessity of putting into place effective conservation measures in the face of
1209 anthropogenic change.

1210 **4 | Geographical range area, habitat breadth**

1211 **and specialisation on natural habitats**

1212 **are associated with land-use responses**

1213 **and climate-change sensitivity more con-**

1214 **sistently than life-history and dietary traits**

1215 **in terrestrial vertebrates**

1216 **Keywords**

1217 Land use; land-use intensity; climate change; sensitivity; CENFA; dietary traits ; life-history traits; speciali-
1218 sation; geographical range area; terrestrial vertebrates.

1219 **Abstract**

1220 Land-use and climate change are two of the most important pressures on terrestrial biodiversity, however the
1221 factors that explain interspecific variation in responses to these pressures remain unclear. Although it is well
1222 established that extinction risk and some species' responses to human pressures relate to species traits, we
1223 lack large-scale comparative assessments across multiple clades linking traits to multiple human pressures.
1224 Here, I investigated whether a set of ecological characteristics that are commonly measured across terrestrial
1225 vertebrates (that is, ecological traits and geographical range area) are associated with (1) species' responses
1226 to different land-use types and (2) species' sensitivity to climate change. My aim was to test whether general-

1227 isable patterns in species' response to these pressures arise with regards to species ecological characteristics,
1228 which helps assess the global signature of human pressures on vertebrate biodiversity and is also of interest
1229 for the prioritisation of conservation efforts. Among the sets of characteristics I considered, I found that
1230 only three were consistently associated with both land-use responses and climate-change sensitivity across
1231 terrestrial vertebrate classes: geographical range area, habitat breadth and specialisation on natural habitats.
1232 The association of other traits with species' land-use responses and with climate-change sensitivity often de-
1233 pended on class and land-use type. My work highlights that narrow-ranged species with small habitat breadth
1234 and natural habitat specialism are typically more sensitive to human pressures. Further, in all classes, I found
1235 that invertebrate eaters and fruit/nectar eaters tended to be negatively affected in disturbed land uses, and that
1236 invertebrate- and plant/seed- eating birds had higher climate-change sensitivity, raising concerns about the
1237 continuation of ecological processes sustained by these species under global changes. My work stresses the
1238 need for putting into place conservation and mitigation measures to protect biodiversity and related services
1239 from human impacts.

1240 4.1 Introduction

1241 Land-use change is currently the most important driver of global biodiversity loss (Newbold et al., 2015)
1242 and is likely to continue to cause species loss in the coming decades (Li et al., 2022; Powers and Jetz, 2019;
1243 Stehfest et al., 2019). However, biodiversity faces multiple pressures acting in combination (Maxwell et al.,
1244 2016). In particular, the impacts of climate change on biodiversity are projected to equate or even surpass
1245 those of land-use change in their magnitude by 2070 (Newbold, 2018). Thus, it has become more vital than
1246 ever to put into place mitigation and conservation measures to protect biodiversity from human pressures.

1247 It is now well established that species differ in their ability to cope with environmental changes (Fer-
1248reira et al., 2022; Matich and Schalk, 2019; Newbold et al., 2013). As such, global average declines in
1249 biodiversity indices mask substantial interspecific variation in responses to disturbances (Leung et al., 2020).
1250 Such interspecific variation has important consequences for the prioritisation of conservation efforts and the
1251 definition of protected areas (Morelli et al., 2021). Mitigating land-use and climate change impacts on the
1252 world's biota requires to understand which species are put at most risk by these pressures, in other words to
1253 understand the factors that are associated with species' sensitivity to land-use and climate change.

1254 By capturing key aspects of species' morphology, life-history, ecological strategies or demography, traits
1255 can inform on species' use of resources and space, as well as on some community and population-level
1256 processes (Capdevila et al., 2022b). As such, traits can help understand what drives species' responses

1257 to environmental change. Thus, to explain interspecific differences in responses to human disturbance, a
1258 number of studies have investigated whether species traits influence species' responses to human pressures,
1259 in particular to land-use change (Newbold et al., 2013; Nowakowski et al., 2017; Quesnelle et al., 2014;
1260 Tinoco et al., 2018) and climate change (Angert et al., 2011; Di Marco et al., 2021; Estrada et al., 2018;
1261 Mccain and King, 2014; Pacifici et al., 2017; Pearson et al., 2014; Schloss et al., 2012).

1262 From these past studies, several traits have been identified as important correlates of species' responses
1263 to land-use and climate change within vertebrate taxa (for example, body mass and generation length were
1264 found to influence bird responses to land-use change in Newbold et al. (2013); and body mass and activity
1265 time were found to be associated with mammal responses to climate change in Mccain and King (2014)).
1266 However, past work has mostly been conducted at local to regional scales (Davison et al., 2021; Hevia et
1267 al., 2017), such that it remains unclear whether the effects of traits on species' responses to environmental
1268 change can be generalised across vertebrate taxa and regions. Yet, at least two metanalyses have investigated
1269 whether traits explained responses to human pressures across diverse taxa, one focused on climate-change
1270 responses (MacLean and Beissinger, 2017), and one on species extinction risk (Chichorro et al., 2019).
1271 MacLean and Beissinger (2017) found that habitat breadth and historic range limit were consistently as-
1272 sociated with variation in species range shifts under contemporary climate change across a range of taxa
1273 (including plants, birds and butterflies), but they did not detect any effect of life-history traits, such as body
1274 size or fecundity. Similarly, Chichorro et al. (2019) highlighted the effects of geographical range area and
1275 habitat breadth on species extinction risks in different taxa (including terrestrial vertebrates), with other
1276 traits having inconsistent effects. However, as underlined by Chichorro et al. (2019), the studies included in
1277 the metanalysis often considered extinction risk without an explicit consideration of the pressures to which
1278 the species were exposed. Yet, a given trait could be associated with opposite responses depending on the
1279 pressure in consideration (González-Suárez et al., 2013).

1280 Further, previous studies have often been restricted in their taxonomic coverage, with very few studies
1281 considering several vertebrate classes together, so that comparative investigations among vertebrate classes
1282 remain rare. In addition, there has not yet been a global assessment of the association between vertebrate
1283 traits and both land-use responses and climate-change sensitivity. Here, I test whether general patterns in
1284 species' land-use responses and climate-change sensitivity arise with regards to species traits. I include
1285 species geographical range area in the analysis, as it is one aspect of rarity that has been shown to influence
1286 species' responses to land use and to climate change (Newbold, 2018; Thuiller et al., 2005). Since geo-
1287 graphical range area does not meet the strict definition of a trait, I henceforth refer to all traits and range
1288 area as 'ecological characteristics'. Thus, I examine associations with a set of ecological characteristics that

1289 are commonly measured across terrestrial vertebrates, at global scales (Figure 4.1a). Considering ecological
1290 characteristics that are available at least for a subset of the species in each class allows for a cross-taxon
1291 comparative assessment. Further, it also allows me to ask whether such commonly measured ecological
1292 characteristics show consistent associations with species' land-use responses and climate-change sensitivity.
1293 I ask two questions: (1) are any ecological characteristics associated with interspecific variation in responses
1294 to land use and with climate-change sensitivity? (2) If so, are these ecological characteristics similar across
1295 classes; are they similar between land-use responses and climate-change sensitivity; and are associations in
1296 the same direction, such that I can identify a set of characteristics that are associated with a high sensitivity
1297 of species to human pressures? Conversely, are such associations both taxon- and pressure- dependent?

1298 Given the different nature of the threats I consider, I use two independent approaches, one for land-use
1299 change and one for climate-change sensitivity. Thus, I do not consider interactive effects between these
1300 pressures. To infer species' responses to land-use change, I use a space-for-time substitution approach,
1301 modelling occurrence probability across different land-use types (Figure 1b). I estimate species' expected
1302 sensitivity to future climate change from properties of species' climatic niches (Figure 1c); species niche
1303 properties have been shown to be strong indicators of species' climate-change sensitivity (Thuiller et al.
1304 2005), and are also straightforward to use at large scales given the availability of species distribution data,
1305 from which climatic niche space can be constructed. I then bring these two approaches together to look for
1306 any emerging pattern in species' responses to land use or in their climate-change sensitivity, with regards to
1307 species' ecological characteristics.

1308 Among the characteristics I consider (Figure 4.1a), some may directly influence species survival by me-
1309 diating resource acquisition and use. These characteristics are body mass, diet, and diet breadth. Other
1310 characteristics (e.g., lifespan and litter/clutch size) may indirectly affect species persistence over time by
1311 influencing species reproductive output and demographic processes (Capdevila et al., 2022b). Finally, re-
1312 sponds to human pressures are known to be dependent on species' degree of specialisation, which I capture
1313 with characteristics reflecting specialisation in time (i.e., diel activity) and reflecting use of space (e.g., habi-
1314 tation breadth and geographical range area).




Figure 4.1: Framework of the study. (a) I collected ecological trait data and geographical range areas across terrestrial vertebrates (termed ‘ecological characteristics’). I then used two independent approaches to assess the influence of these characteristics on species’ responses to land-use and on species’ climate-change sensitivity. (b) To assess the influence of traits on responses to land use and land-use intensity in each vertebrate class, I combined the ecological characteristics with the PREDICTS database. (c) To estimate species sensitivity to climate change, I used the CENFA framework (Rinnan and Lawler, 2019), which relies on the combination of species’ distributions with climatic variables to estimate sensitivity from properties of the species’ climatic niche space. I then built class-specific models to assess whether the ecological characteristics were associated with species sensitivity to climate change.

1315 **4.2 Methods**

1316 **4.2.1 Ecological characteristics (Figure 4.1a)**

1317 **Traits**

1318 I obtained the six following traits from Chapter 2 (in which I presented a trait data compilation across
1319 terrestrial vertebrates): body size (body mass and/or length, depending on the class); a proxy for species
1320 lifespan (generation length for mammals and birds; age at sexual maturity for amphibians; and maximum
1321 longevity for reptiles); litter or clutch size; diel activity; habitat breadth; and use of artificial habitats. I
1322 chose these traits because 1) they were available across all vertebrate classes, at least for a subset of species,
1323 allowing for a comparative assessment; and 2) they relate to species life-history, ecology, and resource use,
1324 such that they might influence species' land-use responses and climatic niche properties (and thus expected
1325 climate-change sensitivity). I couldn't capture intraspecific variation in trait values, and instead I used single
1326 mean values for all traits.

1327 I enhanced the trait data from Chapter 2 with species-level estimates of diet, lacking in the published
1328 database but likely important for understanding species sensitivity to human pressures. For birds and mam-
1329 mals, I collected estimates of species primary diet (i.e., the diet inferred from the combination of food items
1330 totalling more than 50% of species' consumption), from the EltonTraits database (Wilman et al., 2014). For
1331 amphibians and reptiles, obtaining species *primary* diet was not possible, as there were no data available
1332 on the relative consumption of different food items. For amphibians, the AmphiBIO database (Oliveira
1333 et al., 2017) provided information on species consumption of different food items (just in terms of pres-
1334 ence/absence in the diet, but without estimation of their percent use), so I inferred diet on the basis of these
1335 reported food items (however the coverage was low, with more than 75% of the species missing diet infor-
1336 mation; Appendix 3, Figure S4.1). For reptiles, there was no available data collection describing diet. For
1337 both reptiles and amphibians, I supplemented the existing datasets by collecting data on species consumption
1338 from published sources (recording the presence/absence of different food items in species consumption), for
1339 an additional 108 amphibians and for 239 reptiles (see Appendix 3, S4.1: 'Compiling diet information').

1340 I standardised the diet data across the vertebrate classes, by grouping species in five different diet cate-
1341 gories: vertebrate eaters; invertebrate eaters; plant/seed eaters; fruit/nectar eaters; and omnivores (for mam-
1342 mals and birds, species were classified as omnivores when all food items had a percent use $\leq 50\%$; and for
1343 amphibians and reptiles, when species where known to consume both plant and animal matter). I also cal-
1344 culated species diet breadth – the total number of recorded food items (in terms of presence/absence) known

1345 to be consumed by a species. More information on the compilation of dietary information can be found in
1346 Appendix 3, S4.1: ‘Compiling diet information’.

1347 **Geographical range area**

1348 I used extent-of-occurrence maps from BirdLife International for birds (<http://datazone.birdlife.org/species/requestdis>), from the IUCN Red List for mammals and amphibians (IUCN, 2020), and
1349 from Roll et al. (2017) for reptiles (all downloaded in April 2020). I excluded areas occupied during non-
1350 breeding seasons and areas falling outside species known elevational limits (following Chapter 2). The range
1351 maps were then converted to the raster format (‘raster’ package, version 3.5.15 Hijmans (2022)), and I esti-
1352 mated species geographical range areas using a resolution of 1 km² with Behrmann’s equal-area projection.
1353 Although range area cannot be considered a trait (which is a property measurable at the level of individual
1354 organisms), I included range area in the analyses because past work has shown that range area is an important
1355 correlate of species’ responses to land use (Newbold et al., 2018) and climate change (Thuiller et al., 2005).
1356 In addition, range area may correlate with other aspects of species’ ecology that I could not include directly
1357 in the analysis because of limited data availability, such as dispersal ability (Capurucho et al., 2020).

1359 **Phylogenies**

1360 I used information on species’ phylogenetic position in the imputations of missing trait values (see next sec-
1361 tion), and also to control for phylogenetic relationships in the models investigating effects of ecological char-
1362 acteristics on species’ estimated climate-change sensitivity. Class-specific phylogenetic trees were down-
1363 loaded April 2020 from <https://zenodo.org/record/3690867#.Xyc5wyhKhPZ> for mammals
1364 (Phylacine 1.2; Faurby et al. (2018, 2020)); and from <https://data.vertlife.org/> for amphibians
1365 (Jetz and Pyron, 2018), birds (Jetz et al., 2012) and squamates (Tonini et al., 2016). For each class, I used a
1366 consensus tree obtained with the TreeAnnotator programme of the BEAST software (Bouckaert et al. 2014),
1367 from an available distribution of 1000 trees.

1368 **4.2.2 Imputations of missing trait values**

1369 For some of the traits and classes, there was a substantial proportion of missing trait values (Figure S4.1).
1370 To fill these gaps, I imputed missing trait values using random forests, implemented with the ‘missforest’
1371 function of the ‘missForest’ package in R (version 1.4, Stekhoven (2016) and Stekhoven and Bühlmann
1372 (2012)). ‘missforest’ is one of the best methods for missing-value imputations when working with continuous
1373 and categorical variables, and when including species phylogenetic position as a predictor (Debastiani et al.,

1374 2021; Penone et al., 2014). After showing that several traits were strongly phylogenetically conserved
1375 (Table S4.1), I included ten phylogenetic eigenvectors in the imputations (Penone et al., 2014), as well as
1376 taxonomic orders as a categorical variable (included to account for the taxonomic positions of species that
1377 were not represented in the phylogenies). Full details are given in Appendix 3 (S4.2: ‘Imputing missing
1378 trait values’). After imputation, continuous traits were \log_{10} -transformed to improve normality (except for
1379 habitat and diet breadth, which I square-root transformed; this transformation was more appropriate here
1380 because the distributions of habitat breadth and diet breadth tended to be less right-skewed than that of the
1381 other traits, and the range of values was smaller).

1382 **4.2.3 Characterizing the influence of traits on species’ land-use responses (Figure 4.1b)**

1383 **Vertebrate assemblage composition**

1384 To compare vertebrate assemblages in different land-use types, I used the PREDICTS database (Hudson
1385 et al., 2014, 2017). PREDICTS is a collection of independent studies that have sampled biodiversity in sites
1386 of varying land use and land-use intensity. Samples are mostly of species abundance, sometimes species
1387 occurrence, and rarely just overall species richness. It is one of the most comprehensive such databases
1388 to date, with 4,107 vertebrate species sampled across 7,689 sites considered in this work (Figure 4.1b). In
1389 PREDICTS, sites are assigned to one of the following land-use categories: primary vegetation (native vege-
1390 tation); secondary vegetation, plantation forest, pasture, cropland, and urban (disturbed land uses; see Table
1391 S4.2 and Hudson et al. (2014, 2017) for more details). Each site is also characterised in terms of land-use
1392 intensity based on land-use-specific criteria (such as mechanisation degree, crop diversity and agricultural
1393 inputs for cropland; Hudson et al. (2014)). Land-use intensity is divided into three categories to reflect the
1394 degree of human transformation and impacts on the land: minimal, light or intense. Here, I considered
1395 minimally-used primary vegetation to be the least-disturbed reference land use against which I compare all
1396 other land-use types. I grouped pasture and cropland together into a category I termed ‘agricultural’. As the
1397 design of the PREDICTS database is not balanced, sample sizes varied among classes and land-use types
1398 (Figure S4.3).

1399 **Full models (all-predictor models)**

1400 Within each vertebrate class, I investigated whether interactions among the ecological characteristics, land
1401 use and land-use intensity explained species occurrence probability. I fitted four binomial mixed-effects
1402 models (one for each class), using the ‘lme4’ package (version 1.1-23; Bates et al. (2015)), with random

1403 effects accounting for study, site and species identity to account for the nested design of the database, taxo-
1404 nomic non-independence, and repeated observations among species. I did not consider interactions among
1405 the ecological characteristics, but I included interactions between land use and ecological characteristics, and
1406 between land-use intensity and ecological characteristics. Before fitting the models, I checked the degree of
1407 multicollinearity among explanatory variables using generalised variance inflation factors (GVIF; Fox and
1408 Monette (1992)), with a threshold of 5 for the detection of multicollinearity (Tables S4.3-S4.8). For amphibi-
1409 ans and reptiles, including both diet and diet breadth was problematic, so I excluded diet from the set of
1410 predictors for these classes on the basis of the GVIF scores . Models investigating the effects of diet were
1411 built separately (see next section, ‘Partial models’).

1412 I did not use phylogenetic random effects directly in the models because of the computational load re-
1413 quired by such models when working with several hundred or thousands of species. However, I checked the
1414 phylogenetic signal in the models’ residuals using Pagel’s λ (Pagel, 1999). Thus, in each class, the model
1415 fitted was:

1416
1417 $P_{\text{occurrence}} \sim \text{land use} + \text{land-use intensity} + \text{species-level ecological characteristics} +$
1418 $\text{land use : species-level ecological characteristics} +$
1419 $\text{land-use intensity : species-level ecological characteristics} +$
1420 $(1|\text{study identity}) + (1|\text{site identity}) + (1|\text{species identity}).$

1421
1422 To verify that the models’ estimates were robust to any violation of distributional assumptions, I fitted
1423 the models again using a Bayesian framework (using the ‘MCMCglmm’ package version 2.32, Hadfield
1424 (2010)).

1425 **Partial models (single-predictor models)**

1426 In addition to the full models, I fitted partial models for each class. These were fitted to visualise occurrence
1427 patterns for each trait independently of other traits. The structure of the models was similar to that of the full
1428 models, except that I included a single species-level characteristic at a time in each model.

1429 **Effects of categorical ecological characteristics on species’ occurrence probability (Figure 4.2a)**

1430 The influence of categorical traits on species’ responses to land use and land-use intensity can be visualised
1431 in two ways: either by comparing occurrence probability in different land-use types relative to species with
1432 similar traits (I term such effects ‘among land-use type effects’, Figure 4.2a); or by comparing occurrence


1433 probability in a given land-use type relative to species with different traits (I term such effects ‘within land-
1434 use type effects’, Figure 4.2a).

- 1435 • Within land-use type effects (Figure 4.2a): from the full, all-predictor models fitted for each class, I
1436 focused on the interactive effects between land use and ecological characteristics (and between land-
1437 use intensity and ecological characteristics). These interactions indicated whether, in a given land-use
1438 type, there were any significant differences in occurrence probability between species with different
1439 traits. In other words, I looked at whether any trait level lowered or increased occurrence probability
1440 in each land-use type, compared to a reference trait level. I used this approach for all the categorical
1441 predictors, except diet (interpreting within land-use type effects for primary diet being complicated by
1442 the fact that there were more than two levels for this trait).
- 1443 • Among land-use type effects (Figure 4.2a): from a partial model, I predicted occurrence probability
1444 in the different land uses for all different levels of the trait. The partial models allowed to visualise
1445 occurrence patterns across land-use types for single explanatory variables, without having to account
1446 for the values of other variables. I used this approach to evaluate the influence of diet on species’
1447 land-use responses.

1448 **Effects of continuous ecological characteristics on species’ occurrence probability (Figure 4.2b)**

1449 For a given continuous ecological characteristic, any effect of land use or land-use intensity can be as-
1450 sessed through changes in the slope of the relationship between the ecological characteristic and occurrence
1451 probability (Figure 4.2b). When an ecological characteristic negatively impacts occurrence probability in a
1452 disturbed land use, I expect the slope of the relationship to be more negative than the slope for the reference
1453 land use (minimally-used primary vegetation). Focussing on slopes does not allow to infer absolute changes
1454 in occurrence probability across land-use types (e.g., a positive slope in a disturbed land use does not mean
1455 that there are absolute increases in occurrence probability in that land use, but only that higher values of the
1456 ecological characteristic are associated with relatively higher occurrence probability in that land-use type).
1457 This is because I do not assess changes in the mean occurrence probability here (which would require to
1458 consider the intercept of the relationship between the ecological characteristic and occurrence probability in
1459 different land-use types). Thus, I only capture ‘within land-use type’ effects for continuous predictors.

(a) Assessing the effects of categorical ecological characteristics on land-use responses




'Among land-use type' effects: main effect of given characteristic on occurrence probability, among land-use types

'Within land-use type' effects: interactive effect between land-use type and characteristic: difference in occurrence probability for two different values of the characteristic, within land-use types

Categorical characteristic:

- level 1
- level 2

(b) Assessing the effects of continuous ecological characteristics on land-use responses



Slope of the relationship between continuous characteristic & occurrence probability:

1: higher values of ecological characteristics associated with relatively **lower probability of occurrence** in disturbed relative to reference land-use type compared to lower values of ecological characteristics

2: lower values of ecological characteristics associated with relatively **lower probability of occurrence** in disturbed relative to reference land-use type compared to higher values of ecological characteristics

(Does not inform on absolute changes in occurrence probability among land-use types)

Figure 4.2: Assessing the effects of ecological characteristics on species' land-use responses: methodology for (a) categorical characteristics and (b) continuous characteristics. (a) For all categorical characteristics, except diet, I look at 'within land-use type' effects, asking whether there are significant differences in occurrence probability among species with different ecological characteristics in a given land-use type. For diet, I look at 'among land-use type' effects, comparing species occurrence probability in disturbed land uses versus that in primary vegetation (I chose this approach here because visualising 'within land-use type' effects for diet is complicated by the fact that there were more than two levels for this categorical trait). (b) For continuous characteristics, I focus on the relationship with occurrence probability, and I investigate how the slope of this relationship is affected by land-use type, i.e. a 'within land-use type' effect.

1460 **Validation on complete trait data subset (no imputed trait values)**

1461 To assess whether the results were robust to trait imputation uncertainty, I fitted the models again for the
 1462 subset of species for which I had complete, non-imputed data for all ecological characteristics. The models'
 1463 structure was unchanged for birds and mammals. For amphibians, I excluded both diet and litter/clutch size
 1464 because of multicollinearity issues, and I also excluded lifespan proxy and body mass (as there were too
 1465 many missing values in the dataset, 85% and 59% respectively). For reptiles, I excluded both diet and body
 1466 mass because of multicollinearity issues.

1467 **4.2.4 Characterizing the influence of traits on species' sensitivity to climate change (Figure**
1468 **4.1c)**

1469 I estimated climate-change sensitivity across vertebrate species using the ‘Climate-niche Factor Analysis’
1470 (CENFA) approach developed by Rinnan and Lawler (2019), implemented with the ‘CENFA’ R package
1471 version 1.1.1 (Rinnan, 2021). CENFA is a spatial approach for estimating species’ climate-change sensitivity,
1472 exposure, and vulnerability. CENFA combines distribution data with climatic variables to estimate sensitivity
1473 and vulnerability from properties of species’ climatic niches (see Rinnan and Lawler (2019) for details).
1474 CENFA has been used in previous studies focused on a small number of species or on a few taxonomic
1475 groups, but to my knowledge has not yet been applied across all terrestrial vertebrates.

1476 **Historical climate data**

1477 I used global climate data from WorldClim version 2.1 (Fick and Hijmans, 2017). I downloaded 19 climatic
1478 variables at a resolution of 2.5 arcminutes ($\sim 4.6 \text{ km}^2$ at the Equator). I removed variables that were strongly
1479 collinear with any other climatic variables (using a threshold of 0.65 for Spearman correlation coefficients).
1480 I obtained six groups of intercorrelated variables (using the ‘removeCollinearity’ function from the ‘virtu-
1481 alspecies’ R package version 1.5.1 (Leroy et al., 2015); Figure S4.4), and randomly selected one climatic
1482 variable in each group. The final set comprised six climatic variables: annual mean temperature (bio1), mean
1483 diurnal temperature range (bio2), maximum temperature of the warmest month (bio5), annual precipitation
1484 (bio12), precipitation seasonality (bio15), and precipitation of the coldest quarter (bio19).

1485 **Estimating climate-change sensitivity from CENFA**

1486 All climatic variables and distribution files were re-projected to a resolution of 5 km^2 in the Behrmann
1487 equal-area projection. I picked this resolution because the coarser the resolution, the more climate-change
1488 sensitivity tended to be underestimated for narrowly distributed species (Figures S4.5 & S4.6). However,
1489 finer resolutions demand a large amount of memory space when working at global scales across all terrestrial
1490 vertebrates. I found the 5-km^2 resolution to be an acceptable trade-off between computational load and
1491 accuracy of the sensitivity estimations. However, when working at 5-km^2 resolution, there were still some
1492 narrowly distributed species for which sensitivity was likely underestimated (Figure S4.6). Thus, I chose to
1493 exclude species with a range area $\leq 100 \text{ km}^2$ from further analyses (i.e., excluding narrow-ranging species
1494 whose distributions could intersect up to 4 grid cells). In doing so, the sample size was reduced by 660
1495 species for amphibians, by 142 species for birds, by 129 species for mammals, and by 615 species for

1496 reptiles (the final sample sizes were: n=4,537 for amphibians; n=10,198 for birds; n=4,721 for mammals;
1497 n=7,330 for reptiles). My results were overall robust to the exclusion of these species (see Results section).

1498 I then combined the climate data with the species' distributions to estimate sensitivity to climate change,
1499 applying the CENFA framework across terrestrial vertebrates (Figure 4.1c). Further details of the imple-
1500 mentation of the CENFA framework are given in Appendix 3 (S4.5: 'Implementing Climate-niche Factor
1501 Analysis across terrestrial vertebrates').

1502 **Climate-change sensitivity models**

1503 I used phylogenetic least-square (PGLS) regressions, implemented in the 'caper' R package version 1.0.1
1504 (Orme et al., 2018), to assess the effects of ecological characteristics on species estimated sensitivity to
1505 climate change, while controlling for phylogenetic relationships among species. I combined the ecological
1506 characteristics and the phylogenies using the 'comparative.data' function from the 'caper' package, and then
1507 built class-specific models to explain climate-change sensitivity with the ecological characteristics (Figure
1508 4.1c). Before fitting the models, I checked for multicollinearity among the predictors using GVIF scores.
1509 Across all classes, the models included all the main effects of the ecological characteristics, except for am-
1510 phibians, for which I dropped diet breadth (which was strongly collinear with diet; Tables S4.9-S4.13). For
1511 the continuous predictors, I fitted third-order polynomials to allow for non-linearity of the responses (I in-
1512 cluded third order polynomials for the climate-change sensitivity models but not for the land-use models
1513 because the PGLS model had a simpler structure than the land-use models, were less computationally inten-
1514 sive, and also because the number of estimated parameters was already high for the land-use models without
1515 allowing for third-order polynomials). As such, the general form of the PGLS models was:
1516 $\log_{10}(\text{climate-change sensitivity}) \sim \text{poly}(\log_{10}(\text{continuous ecological characteristics}), 3) +$
1517 categorical ecological characteristics + phylogenetic random effects.

1518 **Models' robustness**

1519 To check whether the results were robust to the exclusion of species whose range area was $\leq 100 \text{ km}^2$, I
1520 repeated the models on all species (including those with range area $\leq 100 \text{ km}^2$: n=5,208 for amphibians;
1521 n=10,340 for birds; n=4,844 for mammals; n=7,951 for reptiles).

1522 Finally, to assess the degree to which the results were robust to trait imputation uncertainty, I fitted the
1523 models again for the subset of species for which I had empirical (i.e., non-imputed) trait estimates. Diet
1524 was excluded for amphibians and reptiles on the basis of high collinearity (GVIF>5). I fitted first-order

1525 polynomials here because of the substantially reduced sample size compared to the main models.

1526 4.3 Results

1527 4.3.1 Land-use responses

1528 ‘Within land-use type’ effects (Table 4.1a)

1529 Land-use, land-use intensity, species’ ecological characteristics and their interactions had significant effects
1530 on species occurrence probability. Significant interactive effects between land use and ecological charac-
1531 teristics (and between land-use intensity and ecological characteristics) reflected differences in the ability
1532 of species with different ecological characteristics to cope within the disturbed land-use types (Table 4.1a).
1533 Across all classes, species with narrower geographical range areas, smaller habitat breadth and inability to
1534 exploit artificial habitats showed greater decreases in occurrence probability within disturbed land uses, than
1535 species with larger range areas, larger habitat breadth and ability to exploit artificial habitats (the only ex-
1536 ceptions were opposite effects found for mammals and reptiles for habitat breadth in two of the land-use
1537 types). The effects of the other ecological characteristics differed in direction depending on class and land
1538 use, impeding any generalisation (Table 4.1a). For instance, I found that being smaller and longer-lived was
1539 associated with decreases in occurrence probability for birds found in agricultural areas, but with increases
1540 in occurrence probability for urban birds; and that longer-lived species tended to be more negatively affected
1541 for mammals and reptiles, whereas I found evidence of opposite trends for amphibians.

1542 I would like to highlight that the ‘within land-use type’ effects summarised in Table 4.1a do not nec-
1543 essarily reflect occurrence patterns among land-use types. For example, in all classes, ‘among land-use
1544 type’ effects derived from partial models showed that occurrence probability in disturbed land uses was
1545 strongly negatively affected for natural habitat specialists, compared with primary vegetation levels Figure
1546 S4.7. On the other hand, in most classes and disturbed land uses, artificial habitat users either increased or
1547 showed no significant difference in occurrence probability. One exception was for reptiles, where the effect
1548 of habitat specialisation was mostly non-significant within land-use types (Table 4.1a), with both natural
1549 habitat specialists and artificial habitat users showing important declines in some disturbed land uses (e.g.,
1550 intensely-used agricultural areas, Figure S4.7d). Similarly, the occurrence probability of both nocturnal and
1551 non-nocturnal species was negatively impacted in disturbed land uses compared with primary vegetation
1552 (Figure S4.8), such that land-use responses were not distinguishable between nocturnal and non-nocturnal
1553 species for all classes and land-use types.

Table 4.1: Summary of the effects of the ecological characteristics (except for primary diet) on (a) species' responses to disturbed land uses ('within land-use type' effects) and on (b) species' climate-change sensitivity, for each class of terrestrial vertebrates. The symbol - indicates where a characteristic has a significant negative effect on occurrence probability within a disturbed land-use type (within any of the land-use intensities), or where the characteristic renders species significantly more sensitive to climate change. A + indicates a significantly positive effect of a characteristic on occurrence probability in a land-use type (within any of the land-use intensities), or significantly lower sensitivity to climate change. For the land-use effects, I report 'within land-use type effects' here, that is, within a disturbed land use whether there were significant differences in occurrence probability among species with different trait values (see Figure 4.2). These effects were derived from the interactive terms of the full, all-predictor models.

(a) "Within land-use type" effects of the species-level characteristics on occurrence probability - <=> more sensitive; + <=> less sensitive															Predictors				(b) Climate-change sensitivity: - <=> more sensitive + <=> less sensitive			
Secondary vegetation			Plantation forest				Agricultural				Urban											
-	-	-	-	(-)	-	0	-	0	-	-	-	0	-	0	NA	More narrowly-distributed	Habitat affinity	-	-	-	-	
0	-	0	+	-	-	-	0	-	-	0	-	0	-	+	NA	Smaller habitat breadth		-	-	-	-	
-	-	(-)	-	-	-	0	0	-	-	-	0	0	-	-	NA	Natural habitat specialist	Specialisation	(-)	-	-	-	
0	+	+	-	-	+	0	+	-	+	+	-	0	0	0	NA	Non-nocturnal		-	0	0	(+)	
-	0	+	(-)	-	0	+	+	-	0	0	+	0	-	0	NA	Narrower diet breadth	Resource use	NA	0	-	0	
0	0	+	0	(+)	+	+	0	0	-	0	-	0	+	0	NA	Smaller body mass		-	+	-	-	
+	0	0	0	0	-	0	0	+	-	0	0	0	0	-	NA	Smaller litter/clutch size	Life-history	-	+	+	0	
(-)	0	(+)	+	0	0	0	+	-	+	(+)	+	0	-	0	NA	Shorter-lived		(+)	-	0	-	

1554 **Effects of diet on species' occurrence probability (Figure 4.3)**

1555 In all classes, diet had significant effects on occurrence probability in disturbed land uses (Figure 4.3).
1556 Changes in occurrence probability in disturbed land uses differed among classes and dietary groups. Overall,
1557 invertebrate eaters tended to be negatively affected in disturbed land uses (e.g., -66% average declines in
1558 occurrence probability for amphibians in intensely used agricultural areas, compared with minimally-used
1559 primary vegetation). Omnivores were both negatively and positively impacted, showing both important
1560 decreases (e.g., -81% for reptiles in intensely used plantation forest) as well as strong increases (e.g., +43%
1561 for lightly used urban areas in birds). Overall, fruit/nectar eaters showed important declines in occurrence
1562 probability for mammals and birds, as opposed to plants/seeds eaters, whose occurrence probability tended
1563 to be strongly positively affected for birds, and dependent on land-use intensity for mammals (with increases
1564 in minimally-used land-types, but not in more intensely-used land-types). Finally, I also detected significant
1565 changes in occurrence probability for vertebrate eaters, with some declines for mammals in agricultural areas
1566 (-75% on average in intense uses), but also some increases (e.g., +43% on average for birds in lightly used
1567 agricultural areas).

1568 **Explanatory power for the full models & variance explained by each characteristic (Figure 4.4)**

1569 Overall, land use, land-use intensity and the ecological characteristics explained a small amount of the total
1570 variation in species' occurrence probability (marginal R²: 0.15 for amphibians; 0.054 for birds; 0.15 for
1571 mammals; 0.13 for reptiles), in part because the random effects explained a substantial proportion (condi-
1572 tional R²: 0.59 for amphibians; 0.61 for birds; 0.72 for mammals; 0.57 for reptiles). The relative importance
1573 of traits explaining the most variation differed among classes, with interactions between land use and habitat
1574 breadth explaining the most variation in amphibians and birds, but interactions between land use and body
1575 mass explaining the most variation for mammals, and interactions between land use and lifespan explaining
1576 the most variation for reptiles (Figure 4.4a).

1577 Finally, the models' diagnostics showed evidence of deviations from distributional assumptions (diag-
1578 nistic plots for the full models are shown in Figures S4.9-S4.12). However, when estimated from a Bayesian
1579 framework, the models' estimates were mostly congruent (results not shown), so the frequentist approach I
1580 used with 'lme4' was robust despite the deviations from distributional assumptions. The phylogenetic sig-
1581 nals in the models' residuals were low and not significant (Pagel's $\lambda < 0.01$ for amphibians and reptiles, $p \approx$
1582 1; $\lambda = 0.13$ for mammals, $p = 0.09$; $\lambda = 0.01$ for birds, $p = 0.56$), despite not having included phylogenetic
1583 random effects.




Figure 4.3: Predicted occurrence probability as a function of land use, land-use intensity, diet and their interactions, for each class of terrestrial vertebrates (median \pm 95% confidence interval; the predictions are rescaled with reference to minimally-used primary vegetation). The predictions were obtained from the partial models fitted for each class, considering only diet among the ecological characteristics. Empty plots are drawn where there were no data for a diet category for a given class. Effects could not be estimated for urban reptiles, as well as for urban vertebrate, fruit/nectar and plant/seed eaters for mammals because there were no sampled species. Primary: primary vegetation; Secondary: secondary vegetation; Plantation: plantation forest; Agricultural: cropland and pasture.




Figure 4.4: Proportion of the explained variance attributable to each of the main effects for (a) the mixed-effects models fitting the effects of land use, land-use intensity, and ecological characteristics on species occurrence probability (after factoring out residual variation); (b) for the phylogenetic least-squares regressions investigating whether the ecological characteristics explained climate-change sensitivity (after factoring out residual variation); and (c) for the phylogenetic least-squares regressions investigating whether the ecological characteristics explained climate-change sensitivity (after factoring out the variance explained by geographical range area and the residual variation). The dashed vertical lines mark 10% explained variance (for visualisation purposes). I individually show all the effects that explain more than 5% of the overall variation. Effects that individually explain less than 5% of the overall variation are grouped together as ‘Other effects’. LU: land use; LUI: land-use intensity.

1584 **4.3.2 Climate-change sensitivity**

1585 The ecological characteristics were significantly associated with climate-change sensitivity in all classes
1586 (Tables 4.1b & 4.2, Figures 4.5 & 4.6); models' coefficients shown in Tables ??-??). Overall, climate-
1587 change sensitivity was highest for amphibians (median \log_{10} -sensitivity: 1.1; 95% interpercentile range:
1588 0.40-2.2), then reptiles (median \log_{10} -sensitivity: 1.0; 95% interpercentile range: 0.32-2.1), then mammals
1589 (median \log_{10} -sensitivity: 0.76; 95% interpercentile range: 0.22-2.0) and birds (median \log_{10} -sensitivity:
1590 0.62; 95% interpercentile range: 0.21-1.77). In all classes, narrower geographical range area, smaller habitat
1591 breadth and being specialised on natural habitats were consistently associated with higher climate-change
1592 sensitivity (Table 4.1b). However, other characteristics did not have consistent associations with climate-
1593 change sensitivity across classes, in different cases varying in both significance and direction. For instance,
1594 I found opposite associations between body mass and climate-change sensitivity for mammals, amphibians
1595 and reptiles on the one hand, and birds on the other hand.

1596 The PGLS models explained an important proportion of the overall variation in estimated climate-change
1597 sensitivity (adjusted $R^2 = 0.64$ for amphibians; 0.62 for birds; 0.63 for mammals and reptiles). Geographical
1598 range area explained the majority of this variation in climate-change sensitivity (about 60% in all classes;
1599 Figure 4.4b), which largely reflects the design of the CENFA approach. When factoring out residual variation
1600 and variation explained by geographical range area, the relative importance of the traits as correlates of
1601 climate-change sensitivity varied among classes (Figure 4.4c), with body mass explaining the most variation
1602 for mammals and reptiles, and litter/clutch size explaining the most variation for amphibians and birds.




Figure 4.5: Estimated effects of the categorical traits on climate-change sensitivity, from the PGLS models fitted in each class (mean effect \pm 95% confidence interval). For each categorical trait, I show the effect size for all levels referring to the reference level (vertical dashed line). (a) For artificial habitat use, the reference level is ‘Artificial habitat user’; (b) for diel activity, the reference level is ‘Nocturnal’; (c) for diet, the reference level for mammals and birds is ‘Fruit/nectar’ eaters, but it is ‘Invertebrate’ eaters for amphibians and reptiles.




Figure 4.6: Effects of the continuous ecological characteristics on climate-change sensitivity, estimated from the PGLS models in each class. The lines represent the estimated relationships between climate-change sensitivity and ecological characteristics; the shaded areas are 95% confidence intervals. I plotted the estimated relationships only when they were found to be significant.

1603 Robustness of the PGLS models

1604 The PGLS models were robust to distributional assumptions (Figures S4.13-S4.16). When fitting the models
 1605 on all species (including those with range area $\leq 100\text{km}^2$), I found that the relationship between climate-
 1606 change sensitivity and geographical range area was reversed in all classes (with smaller-ranging species
 1607 estimated to be less sensitive). This result is likely an artefact caused by the underestimation of climate-
 1608 change sensitivity for the most narrow-ranging species, which would support the exclusion of such species
 1609 from the analysis. Other results were generally not sensitive to the exclusion of species whose range area
 1610 was $\leq 100 \text{ km}^2$ (Figure ??; models' summaries shown in Tables ??-??).

1611 Running the models again using data subsets for which I had empirical, non-imputed values only for the
 1612 ecological characteristics showed that the conclusions of this work are likely robust to imputation uncertainty.
 1613 Overall, across all classes, the associations of geographical range area, habitat breadth and use of artificial
 1614 habitats with sensitivity to climate change and land use were consistent with the main models (Tables ??-??;
 1615 Figure ??).

Table 4.2: ANOVA summaries for the PGLS models investigating the associations between the species-level ecological characteristics and species' estimated climate-change sensitivity.

Class	Model terms	Df	Sum Square	Mean Square	F-value	Pr(>F)
Amphibians	poly(log ₁₀ (Body mass, 3))	3	9.85E-02	3.28E-02	3.41E+01	9.03E-22 ***
	poly(log ₁₀ (Lifespan proxy), 3)	3	5.62E-02	1.87E-02	1.94E+01	1.60E-12 ***
	poly(log ₁₀ (Litter/clutch size), 3)	3	1.22E-01	4.07E-02	4.22E+01	6.68E-27 ***
	poly(log ₁₀ (Range area), 3)	3	7.30E+00	2.43E+00	2.52E+03	0.00E+00 ***
	poly(sqrt(Habitat breadth), 3)	3	5.86E-02	1.95E-02	2.03E+01	4.69E-13 ***
	Specialisation	1	3.17E-03	3.17E-03	3.29E+00	6.96E-02 .
	Diel activity	1	1.03E-02	1.03E-02	1.07E+01	1.07E-03 **
Birds	Diet	3	8.61E-04	2.87E-04	2.98E-01	8.27E-01
	poly(log ₁₀ (Body mass, 3))	3	5.78E-02	1.93E-02	2.87E+01	1.87E-18 ***
	poly(log ₁₀ (Lifespan proxy), 3)	3	1.55E-02	5.18E-03	7.70E+00	3.87E-05 ***
	poly(log ₁₀ (Litter/clutch size), 3)	3	1.15E-01	3.83E-02	5.70E+01	1.63E-36 ***
	poly(log ₁₀ (Range area), 3)	3	1.10E+01	3.66E+00	5.45E+03	0.00E+00 ***
	poly(sqrt(Habitat breadth), 3)	3	1.29E-02	4.30E-03	6.39E+00	2.53E-04 ***
	poly(sqrt(Diet breadth, 3))	3	1.01E-02	3.38E-03	5.02E+00	1.77E-03 **
Mammals	Specialisation	1	7.10E-02	7.10E-02	1.06E+02	1.15E-24 ***
	Diel activity	1	1.58E-03	1.58E-03	2.34E+00	1.26E-01
	Primary diet	4	3.15E-02	7.88E-03	1.17E+01	1.70E-09 ***
	poly(log ₁₀ (Body mass, 3))	3	9.00E-02	3.00E-02	3.44E+01	5.41E-22 ***
	poly(log ₁₀ (Lifespan proxy), 3)	3	7.32E-03	2.44E-03	2.80E+00	3.85E-02 *
	poly(log ₁₀ (Litter/clutch size), 3)	3	3.46E-02	1.15E-02	1.32E+01	1.33E-08 ***
	poly(log ₁₀ (Range area), 3)	3	6.83E+00	2.28E+00	2.61E+03	0.00E+00 ***
Reptiles	poly(sqrt(Habitat breadth), 3)	3	2.73E-02	9.11E-03	1.05E+01	7.50E-07 ***
	poly(sqrt(Diet breadth, 3))	3	8.64E-03	2.88E-03	3.31E+00	1.94E-02 *
	Specialisation	1	8.99E-03	8.99E-03	1.03E+01	1.33E-03 **
	Diel activity	1	6.08E-06	6.08E-06	6.98E-03	9.33E-01
	Primary diet	4	9.32E-03	2.33E-03	2.67E+00	3.04E-02 *
	poly(log ₁₀ (Body mass), 3)	3	8.53E-01	2.84E-01	3.18E+02	3.24E-194 ***
	poly(log ₁₀ (Lifespan proxy), 3)	3	6.40E-02	2.13E-02	2.39E+01	2.25E-15 ***

1616 **4.4 Discussion**

1617 Here, I investigated whether species' ecological characteristics were associated with species' sensitivity to
1618 two human pressures (climate change and land-use change), across terrestrial vertebrate classes. Overall, I
1619 found that geographical range area, habitat breadth and specialisation on natural habitats were **the only char-**
1620 **acteristics that showed consistent effects across both pressures and across the terrestrial vertebrate classes;**
1621 narrower ranges, narrower habitat breadth and inability to exploit artificial habitats were associated with
1622 more negative land-use responses and with higher climate-change sensitivity. My results are in line with pre-
1623 vious **metanalyses** that have found species' extinction risks to be associated with habitat breadth and range
1624 area (Chichorro et al., 2019), range shifts under contemporary climate change to be associated with species'
1625 historical range limits and habitat breadth (MacLean and Beissinger, 2017), and with many other studies on
1626 land-use responses or extinction risks (Newbold et al., 2018; Nowakowski et al., 2017; Ripple et al., 2017).
1627 However, to the best of my knowledge, this work constitutes the first global study to compare patterns among
1628 vertebrate classes and between the two major human pressures of climate change and land use. The results
1629 of this Chapter have important implications for conservation, as they mean that land-use and climate change
1630 are non-randomly affecting all terrestrial vertebrates, with a consistently higher risk for geographically rarer
1631 species and habitat specialists. Geographical rarity has been employed by the IUCN for many years in vul-
1632 nerability assessments (Rodrigues et al., 2006), and this work provides further support for its integration in
1633 large-scale assessments. My results also lend support to the idea that habitat specialisation is a likely indica-
1634 tor of species' sensitivity to both land-use and climate change across all vertebrates, thus indices reflecting
1635 habitat specialisation should be highly relevant to consider in large-scale vulnerability assessments, **such as**
1636 **in Foden et al. (2013).**

1637 This work further highlights the class-specific associations between most traits and likely responses to
1638 human-driven environmental changes, but again highlighting a non-random reshaping of vertebrate biodi-
1639 versity under global changes. In the case of land use, I find that the directionality of the responses not only
1640 often depends on taxonomic class but also on land-use type, further complicating the patterns. In line with
1641 past work highlighting the low explanatory power of traits when used to explain species' responses to human
1642 pressures (Angert et al., 2011; Cannistra and Buckley, 2021; Verberk et al., 2013), I found that most traits
1643 explained a small proportion of the interspecific variation in land-use responses and in climate-change sen-
1644 sitivity. The only exception was range area, which explained a large proportion of the interspecific variation
1645 in climate-change sensitivity. Given that the estimates of climate-change sensitivity were based on proper-
1646 ties of species' climatic niches (Rinnan and Lawler, 2019), it was not surprising that range area explained

1647 such an important part of the interspecific variation in climate-change sensitivity, as it is likely that broader
1648 range areas are associated with broader climatic tolerances, and thus with lower estimated climate-change
1649 sensitivity on average.

1650 Despite their generally low explanatory power, traits have been used to assess species vulnerability to
1651 human threats, in particular to climate change (Böhm et al., 2016; Foden et al., 2013). One of the conceptual
1652 appeals behind the use of traits is that if clear patterns in responses to environmental change can be identified
1653 across taxa, then it could be possible to generalize their effects in space and time (Hamilton et al., 2020;
1654 Verberk et al., 2013), which is of interest for conservation -- notably for data deficient species and/or those
1655 lacking estimates of abundance or population sizes. This work does not provide support for the generalisation
1656 at large scales of the effects of life-history and dietary traits on sensitivity to either land-use or climate change.
1657 The class-specific influence of traits on climate-change sensitivity, coupled with their low explanatory power,
1658 could be one of the reasons why trait-based approaches have been shown to perform less well and show less
1659 congruence than trend-based approaches (which rely on the use of long-term population data) for climate-
1660 change vulnerability assessments (Wheatley et al., 2017). I would like to emphasize that this does not mean
1661 that life-history and dietary traits are unimportant for understanding species' responses, but it means that their
1662 effects and relative importance could depend on interactions between the considered taxa and the pressures
1663 affecting them.

1664 Further, it is possible that general underlying patterns remain undetected or are being masked by inter-
1665 actions and trade-offs among traits, which I did not consider here. For instance, larger species tend to have
1666 larger dispersal distances and movement abilities (Jenkins et al., 2007), which could be beneficial to resource
1667 acquisition in disturbed areas (Hillaert et al., 2018); however, such species also tend to have higher energetic
1668 requirements (White, 2011) and lower reproductive output, which could be detrimental to their persistence in
1669 the face of environmental change. Interactions and trade-offs among traits have been shown to be important
1670 for understanding which species are likely to persist in disturbed environments (Sayol et al., 2020), but little
1671 is known about their potential effects at large scales and for different types of pressure. Thus, developments
1672 of my work could focus on the effects of trait interactions on species' sensitivity to climate change and land-
1673 use responses. I was also unable to consider intraspecific variation in this work. Intraspecific variation and
1674 potential for acclimation and evolutionary adaptation are likely important determinants of species' ability to
1675 cope with human disturbance (Carlson et al., 2014; Rohr et al., 2018), but considering such effects at large
1676 scales is challenging because of the lack of available data.

1677 Moreover, I investigated climate-change sensitivity and land-use responses separately, meaning that I
1678 did not consider the combined effects of these pressures. However, human pressures act in combination

1679 (Capdevila et al., 2022a; Harfoot et al., 2021; Segan et al., 2016), and a number of confounding factors and
1680 threats that I could not account for could possibly influence species' sensitivity. For example, larger species
1681 might be more sensitive to warming than smaller species (Hantak et al., 2021; Merckx et al., 2018), but they
1682 could also be better able to persist in fragmented landscapes, such that habitat fragmentation and climate
1683 warming may have opposite effects on the responses of such species. Further, larger species might also
1684 be disproportionately exposed to other threats such as overexploitation and human-wildlife conflicts (Ripple
1685 et al., 2014, 2017). Interactions among traits, among types of pressure, and among traits and pressures
1686 should ideally be considered together to understand species' responses to human disturbances (Hantak et al.,
1687 2021). However, considering all these effects simultaneously may be challenging because of data-limitation
1688 issues, model complexity, and difficulty in assessing and disentangling individual and interactive effects.
1689 In addition, I would like to emphasize that my results, based on correlative assessments of the associations
1690 between ecological characteristics and species' sensitivity to climate change and species' land-use responses,
1691 do not allow to infer any causal links between traits and species' responses to global changes. Reinforcing
1692 our mechanistic understanding of how traits influence species' ability to cope with different disturbances
1693 may help understand opposite signatures of human impacts on vertebrate trait diversity. Further work could
1694 help elucidate the mechanistic links between species traits and responses to environmental change, perhaps
1695 supported by long-term population data and demographic models (Hernández-Yáñez et al., 2022).

1696 To conclude, the results of this Chapter indicate that land-use and climate change are likely to impact
1697 terrestrial vertebrates non-randomly with respect to their ecological characteristics, which could have im-
1698 portant consequences for ecosystem functioning (Duffy, 2003; Luck et al., 2012). For instance, I detected
1699 substantial declines in occurrence probability of certain dietary groups in disturbed land-use types, most
1700 notably invertebrate eaters and fruit/nectars eaters in all classes. I also found higher climate-change sensitiv-
1701 ity for invertebrate- and plant/seed-eating birds. My findings thus highlight the potential risks from global
1702 changes for ecosystem processes and services sustained by those species, such as pest control, seed dispersal
1703 or pollination (Civantos et al., 2012; Fricke et al., 2022; González-Varo et al., 2013), highlighting the need
1704 to put into place mitigation and conservation measures in the face of global changes.

5 | Energetic constraints and trophic group explain species persistence in disturbed land uses

Keywords

Land use; land-use intensity; metabolic rates; energetic constraints; energetic requirements; terrestrial vertebrates; trophic group; occurrence.

Abstract

Land-use change is the primary driver of global biodiversity loss. In terrestrial vertebrates, previous work has shown that sensitivity to land-use change depends on species traits, but the extent to which energetic constraints explain species responses to disturbed land uses remains largely unexplored. Here, I investigate relationships between the energetic requirements of terrestrial vertebrates (estimated from resting metabolic rates) and land-use change, at two levels of organisation. First, at the assemblage level I hypothesize that total energetic requirements in disturbed land uses are lower than in undisturbed land uses, assuming that there is less energy available in these areas overall. Second, after controlling for the effects of body mass and taxonomy on metabolic rates, I predict that species with relatively lower energetic expenditure are favoured over species with relatively higher energetic expenditure in disturbed land uses, as resource efficiency will be beneficial in these resource-poor environments. Because trophic group influences species ability to assimilate various types of food, I investigate whether my predictions are consistent among trophic groups (here, omnivores, carnivores or herbivores). The results challenged both hypotheses. I found that total assemblage-level energetic requirements did not systematically decrease in disturbed land uses. For instance, I detected

1725 significant increases for urban areas in all trophic groups, highlighting that disturbed areas may not be as
1726 energy-poor as I initially assumed. Second, I found a positive effect of metabolic rates (after controlling
1727 for body mass and taxonomy) on species probability of occurrence across all trophic groups for at least one
1728 of the most disturbed land uses I considered (pasture, cropland and urban). Species for which there are
1729 exploitable resources in disturbed environments may benefit from having larger energetic expenditure: they
1730 may display a set of characteristics rendering them more able to cope with disturbances and more able to
1731 acquire available resources, such as higher activity levels or bigger brain sizes. The findings of this Chapter
1732 highlight that land-use change has substantial impacts on vertebrate community metabolism.

1733 5.1 Introduction

1734 Land-use change is currently the strongest driver of global biodiversity declines (Maxwell et al., 2016;
1735 Newbold et al., 2015), with major and long-lasting impacts on the structure and functioning of ecological
1736 communities (Bregman et al., 2016; Fukasawa and Akasaka, 2019; Magioli et al., 2021; Marcacci et al.,
1737 2021). With land-use change likely to continue to intensify (Stehfest et al., 2019), it is vital to put into
1738 place conservation and mitigation measures to minimise future losses of biodiversity and negative impacts
1739 on ecosystem functioning. To this end, pressing questions remain as to what renders species able or unable
1740 to cope with human disturbance, and how losses of sensitive species might influence ecosystem functioning
1741 (Dirzo et al., 2014; Young et al., 2016).

1742 Land-use change acts as an environmental filter affecting species persistence (Edwards et al., 2021;
1743 Evans et al., 2018). Past studies have shown that sensitivity to land-use change is distributed unevenly across
1744 the tree of life (Nowakowski et al., 2018), and across behavioural (Lowry et al., 2013; Samia et al., 2015)
1745 and ecological strategies (De Palma et al., 2015; Møller, 2009; Newbold et al., 2013). For instance, long-
1746 lived and large forest specialist birds respond more negatively to land-use change than generalist species
1747 (Newbold et al., 2013). In addition, human land uses impose energetic constraints on species, by modifying
1748 the amount and type of available resources (Inger et al., 2016; Zuñiga-Palacios et al., 2021). Consequently,
1749 land-use change brings about modifications in dietary and foraging strategies (Møller, 2009; Ramesh et al.,
1750 2017; Sévèque et al., 2020), which can have cascading effects on local trophic structure (e.g., mesopredator
1751 release; Crooks and Soulé (2010)). The impacts of land-use change on biodiversity also depend on land-
1752 use intensity (Davison et al., 2021; Newbold et al., 2015), that is, the magnitude of human disturbance in
1753 a given land-use type, which can vary importantly according to local practices and management (Dullinger
1754 et al., 2021). In particular, land-use intensity affects the types and availability of resources (by homogenising

1755 resources for example), which in turn may affect biodiversity patterns (Weiner et al., 2011).

1756 Energetic requirements are likely to be important predictors of species' ability to cope with a disturbed
1757 environment where resource availability and resource types are strongly impacted (Auer et al., 2020). Past
1758 research has shown that metabolic rates (the amount of energy required for organismal maintenance and
1759 for other processes such as growth and reproduction) are heritable and subject to selection, for instance
1760 with species in energy-rich environments having evolved faster metabolic rates than species in less energy-
1761 rich environments (Mueller and Diamond, 2001). However, metabolic rates have received less attention
1762 than other life-history traits in studies aiming to understand different species' responses to land-use change.
1763 Body mass, known to explain much of the variation in metabolic rates (Bushuev et al., 2018; Hudson et
1764 al., 2013; White and Seymour, 2003), has been considered in past studies examining effects of land-use
1765 disturbance on assemblage composition (Hevia et al., 2017). Since metabolic rates correlate positively with
1766 body mass, the effects of land-use change on local energetic requirements have been indirectly assessed
1767 through investigations of changes in body mass composition along land-use gradients (Newbold et al., 2020a;
1768 Tinoco et al., 2018). However, to the best of our knowledge, no study has yet directly reported the effects of
1769 land-use change on assemblage-level energetic requirements.

1770 Thus, the first objective of this Chapter is to investigate how land-use change impacts assemblage-level
1771 energetic requirements in terrestrial vertebrates. To this end, I use metabolic rates as a proxy for maintenance
1772 energetic requirements at the species level. I predict that, at the assemblage level, total energetic requirements
1773 in disturbed land uses are lower than in undisturbed land uses. This prediction relies on the assumption that
1774 there is less energy available overall in disturbed land uses than in undisturbed land uses (Figure 5.1c), with
1775 disturbed land uses being characterised by human appropriation of net primary productivity (Krausmann et
1776 al., 2013). Because species' ability to exploit particular resources and meet energetic demands also depends
1777 on food types and diet (McNab, 1986; Mendoza and Araújo, 2019), I test whether this prediction holds
1778 true within trophic groups (i.e., within herbivores, omnivores, and carnivores). Here, I consider energetic
1779 requirements at the assemblage level, not controlling for species body mass, to obtain estimates of total
1780 energy expenditure in different land-use types (Figure 5.2a). Thus, any changes in assemblage-level energetic
1781 requirements could be ultimately driven by shifts in the size-spectrum of local assemblages.

1782 Body mass explains much of the variation in metabolic rates, but species with similar body masses still
1783 display important variation in metabolic rates (Mueller and Diamond, 2001). Some of this variation can
1784 be explained by taxonomic position (which also accounts for differences in thermoregulatory strategy be-
1785 tween endotherms and ectotherms). However, there remains substantial interspecific variation in metabolic
1786 rates even after accounting for body mass and taxonomic affiliation (Mueller and Diamond, 2001; White

1787 and Kearney, 2013). Food availability and food type might explain this residual variation in metabolic rates
1788 among species (Mueller and Diamond, 2001). Species that evolved in areas with abundant resources assim-
1789 ilate and burn calories at high rates, regardless of their body mass – while species that evolved with limiting
1790 resources could process resources at lower rates, irrespectively of their body mass (Mueller and Diamond,
1791 2001). Building upon this idea and the ‘allocation’ principle (i.e., the trade-offs in resource allocation; Auer
1792 et al. (2020)), species with lower energetic expenditure (than expected from body mass and taxonomy) may
1793 be able to cope with food scarcity better than those with larger energy expenditure (than expected from body
1794 mass and taxonomy). Indeed, regardless of their body mass, such species should need fewer resources to
1795 meet energetic demands (Clarke and Fraser, 2004). Thus, as land-use disturbance modifies the types and
1796 abundance of resources, we expect species energetic requirements –after removing the effects of body mass
1797 and taxonomic position– to influence responses to land-use change. Yet, whether energetic constraints ex-
1798 plain species responses to land-use change in terrestrial vertebrates has not been investigated before (Hevia et
1799 al., 2017). Thus, the second objective of this Chapter is to characterise the influence of energetic constraints
1800 on species occurrence probability in disturbed land uses, compared to undisturbed land uses. I approximate
1801 energetic constraints at the species level from residual variation in metabolic rates, that is, the variation not
1802 explained by body mass or taxonomy. I predict that species with negative deviations (lower rates than ex-
1803 pected from body mass and taxonomy) are more likely to persist in disturbed land uses than species with
1804 positive deviations (higher rates than expected from body mass and taxonomy; Figure 5.2b). Given that
1805 trophic group can influence resource acquisition and species ability to assimilate various types of food, I
1806 investigate whether this prediction is consistent among trophic groups.

1807 To test both predictions, I use a space-for-time substitution approach, obtaining vertebrate assemblage
1808 composition data from the PREDICTS database (Hudson et al., 2014, 2017). I combine this database with
1809 trophic-group information (characterising species as either omnivores, herbivores, or carnivores), and with
1810 species-level estimates of resting metabolic rates and body mass.





Figure 5.1: (a) Spatial distribution of the sampled sites from the PREDICTS database for terrestrial vertebrates (6,484 sites); (b) Number of sites in each land-use and land-use-intensity category; (c) Net primary productivity by land use and land-use intensity (derived from MODIS satellite imagery), with model predictions plotted relative to minimally used primary vegetation (and rescaled with reference to that land-use type). Primary: primary vegetation; secondary: secondary vegetation; plantation: plantation forest.


(a) Assemblage-level: there is less energy available to utilize by vertebrate assemblages in disturbed land uses than in undisturbed land uses.



Prediction 1: within all trophic groups, total assemblage-level **mass-dependent** metabolic rates decrease in disturbed land uses compared to undisturbed land uses.


$$\text{Assemblage level:} \\ t\text{RMR} = \sum_i (\text{RMR}_i * a_i)$$

$$\text{Across assemblages - model 1:} \\ \log(t\text{RMR}) \sim \text{LU} + \text{LUI} + \text{TG} + \\ \text{LU:LUI} + \text{LU:TG} + \text{LUI:TG} + \\ \text{LU:LUI:TG} + RE$$



(b) Species-level: species with lower energetic expenditure than expected from body mass and taxonomy (negative deviations) do better in disturbed land uses than species with higher energetic expenditure than expected from body mass and taxonomy (positive deviations).

residuals of
 $\log(\text{RMR}) \sim \log(\text{BM}) + (1|\text{Class/Order/Family})$



Prediction 2: within all trophic groups, species occurrence probability in disturbed land uses is negatively affected by **residual RMR**. The slope of the relationship between occurrence probability and residual RMR is significantly lower than the slope estimate for the undisturbed land uses.

Across species - model 2:

$$\text{P}_{\text{occurrence}} \sim \text{LU} + \text{LUI} + \text{resRMR} + \text{TG} + \\ \text{LU:LUI} + \text{LU:TG} + \text{LU:resRMR} + \\ \text{LUI:TG} + \text{LUI:resRMR} + \text{TG:resRMR} + \\ \text{LU:TG:resRMR} + \text{LUI:TG:resRMR} + RE$$




Figure 5.2: Framework for the predictions and models. I use resting metabolic rates (RMR) as a proxy for the amount of energy processed by vertebrate species. (a) I hypothesize that there is less energy overall that can be utilized by vertebrate species in disturbed land uses compared to undisturbed land uses. As such, I expect decreases in assemblage-level total metabolic rates ($t\text{RMR}$). $t\text{RMR}$ is calculated as the sum, over all species occurring at a site, of species RMR (RMR_i) weighted by species abundance (a_i). (b) After removing the effects of body mass and taxonomic position, I expect species with lower energy expenditure than expected to do better than species with higher energy expenditure than expected. I predict that the slope of the relationship between residual RMR and occurrence probability is more negative in disturbed land uses than in undisturbed land uses. For both (a) and (b), I test whether the predictions are consistent among trophic groups. LU: Land use; LUI: land-use intensity; TG: trophic group; RE: random effects. In model 1, random effects include the identity of the PREDICTS studies from which the assemblages are derived. In model 2, random effects include study, site, and species identity.

1811 **5.2 Methods**

1812 **5.2.1 Vertebrate assemblage composition**

1813 I obtained vertebrate assemblage composition in different land uses from the PREDICTS database (Hudson
1814 et al., 2014, 2017). The PREDICTS database is a large collection of published studies that measure bio-
1815 diversity across different land uses and is one of the most comprehensive global databases of its type. In
1816 each PREDICTS study, species occurrence and often abundance were recorded across different sites. Each
1817 site was assigned to one of the following land-use types: primary vegetation (natural, undisturbed vegeta-
1818 tion), secondary vegetation (recovering after complete destruction of primary vegetation), plantation forest
1819 (woody crops), pasture (areas grazed by livestock), cropland (herbaceous crops) and urban (built-up areas).
1820 The land-use categories were assigned based on habitat descriptions from the original studies (Hudson et al.,
1821 2014), sometimes in consultation with the original study authors. Each site was also classified in terms of
1822 land-use intensity as either minimal, light or intense. The land-use-intensity assignment was also made on
1823 the basis of the habitat description in the original studies, and depended on criteria specific to each land use
1824 (such as degree of mechanisation, yield or chemical inputs for cropland; or the amount of green space in
1825 urban areas; Hudson et al. (2014)).

1826 I subset the PREDICTS database for studies that sampled terrestrial vertebrates, and for which both land
1827 use and land-use intensity had been characterised. I thus obtained 181 studies for 4,238 species sampled
1828 across 6,484 sites (Figure 5.1a). Sample sizes varied across land uses and land-use intensities (Figure 5.1b).

1829 **5.2.2 Energy availability by land-use type and land-use intensity**

1830 The predictions of this Chapter rely on the assumption that resource types and abundance are modified in dis-
1831 turbed environments, with less energy available in disturbed compared to undisturbed land uses overall. To
1832 test this assumption, I used terrestrial net primary productivity (NPP) across land uses as a proxy for available
1833 energy. NPP quantifies the amount of atmospheric carbon fixed by plants and accumulated as biomass. NPP
1834 estimates were derived using imagery from the Moderate Resolution Imaging Spectroradiometer (MODIS)
1835 on board NASA's Terra satellite. NPP estimates were based on a yearly composite of measures made at
1836 8-day intervals, captured at 500-m spatial resolution (Running and Zhao, 2015). I obtained NPP for 4,062
1837 of the PREDICTS sites used in the analysis (matching the sites to the NPP data using the sampling year
1838 available in PREDICTS). I fit a linear mixed-effects model ('lme4' package, version 1.1-23, Bates et al.
1839 (2015)) explaining site-level NPP by land use and land-use intensity, with a random intercept accounting for

1840 study identity, to control for differences in experimental design across studies. Model predictions showed
1841 that NPP decreased significantly in several land uses (e.g., pasture and cropland) compared with the primary
1842 vegetation reference level, although the strength and in some cases direction of the difference varied among
1843 land-use and land-use intensity combinations (e.g., increases in urban land uses; Figure 5.1c).

1844 **5.2.3 Resting Metabolic Rates (RMR) & imputations of missing RMR values**

1845 As a proxy for species-level energetic expenditure, I used estimates of the minimum amount of energy re-
1846 quired for organismal maintenance, i.e., basal metabolic rates (BMR) for endotherms, and resting metabolic
1847 rates (RMR) for ectotherms. From the literature, I obtained estimates of BMR for 719 species of birds and
1848 685 mammals, and estimates of RMR for 126 amphibians and 173 reptiles (Appendix 4, Table S5.1). For
1849 endotherms, BMR are measured when species are in their thermoneutral zone, that is, when there is little to
1850 no energy expenditure allocated to thermoregulation. Thus, BMR estimates were derived from lab studies
1851 that mostly measured oxygen consumption of the organisms at rest under controlled conditions and in the
1852 thermoneutral zone of the species. For an ectotherm, there is no ‘basal’ metabolic rate, as body tempera-
1853 ture mainly depends on environmental temperature. Their metabolic rates follow a hump-shaped relationship
1854 with environmental temperature, highest at an optimal temperature which corresponds to a performance peak.
1855 To be able to compare endotherms’ BMR with ectotherms’ RMR, Stark et al. (2020) used the metabolic rates
1856 that correspond to a performance peak for both groups (i.e., BMR in the thermoneutral zone for endotherms,
1857 and metabolic rates at optimal temperature for ectotherms). Thus, I used the data compiled in Stark et al.
1858 (2020) for ectotherms, and from the sources specified in Table S5.1 for endotherms. The units for BMR
1859 and RMR were standardized to mL of dioxygen consumed per hour (mLO₂/h). As in Stark et al. (2020), I
1860 henceforth refer to both basal and resting metabolic rates as RMR.

1861 For the species occurring in PREDICTS, initial data coverage for RMR was poor (Table S5.1), necessi-
1862 tating imputation of missing values. To do so, I first measured the phylogenetic signal in BMR and RMR
1863 (\log_e -transformed), using Pagel’s λ (Pagel, 1999), to assess whether metabolic rates were sufficiently phylo-
1864 genetically conserved to be estimated from species phylogenetic position. I obtained class-specific phyloge-
1865 netic trees from Jetz et al. (2012) for birds, from Faurby et al. (2018, 2020) for mammals, from (Tonini et al.,
1866 2016) for reptiles (squamates), and from Jetz and Pyron (2018) for amphibians (all downloaded in April
1867 2020). For each class, I randomly sampled 100 trees. To account for phylogenetic uncertainty, I calculated
1868 Pagel’s λ for each sampled tree and reported the median value, as well as the 2.5th and 97.5th percentiles
1869 (Table S5.1).

1870 In addition to being highly phylogenetically conserved (Table S5.1), RMR correlate strongly with body

mass (Figure 5.2a). Thus, I imputed missing values using body-mass information (see next section), phylogenetic relationships and taxonomic orders as predictors (Penone et al., 2014). For each class, I used a consensus phylogenetic tree from which I summarised phylogenetic relationships in the form of five phylogenetic eigenvectors. Including more eigenvectors had little impact on the imputed values (results not shown). Consensus trees were obtained with the TreeAnnotator programme of the BEAST software (Bouckaert et al., 2014). Missing RMR values were imputed using random forests algorithms implemented in R using the ‘missForest’ package (version 1.4; Stekhoven (2016) and Stekhoven and Bühlmann (2012)).

5.2.4 Trophic group and body mass information

I used body mass and trophic group information for terrestrial vertebrates compiled in Chapter 2. Body mass was compiled as a single measure at the species level, meaning I was unable to consider intraspecific variation. Trophic group described species as either carnivores, omnivores, or herbivores. Because there were gaps in the availability of the data, more so for trophic group than for body mass (see Chapter 2), I imputed the missing trait values (independently of RMR imputations), then used both imputed and empirical body mass values for imputations of missing RMR values. To impute missing body mass and trophic groups, I used random forests algorithms (again, using the ‘missForest’ R package), including as additional predictors phylogenetic information, added in the form of 10 phylogenetic eigenvectors (Diniz-Filho et al., 2012) following Penone et al. (2014), and also taxonomic order. I considered a wider set of life-history traits in the missing values imputations: lifespan, litter/clutch size, habitat breadth and use of artificial habitats (compiled in Chapter 2). Phylogenetic eigenvectors were extracted from the class-specific phylogenies using the ‘PVR’ package (Santos, 2018).

5.2.5 Effects of land use, land-use intensity and trophic group on assemblage-level total RMR (prediction 1; Figure 5.2a)

Assemblage-level total RMR (tRMR) was obtained by summing abundance-weighted RMR for the species occurring in each site; abundance data were available for 125 of the 181 PREDICTS studies I considered (sampling 3,487 species across 4,644 sites). I fitted a linear mixed-effects model to explain \log_e -tRMR as a function of land use, land-use intensity and trophic level, with a random intercept accounting for study identity to control for differences in experimental design across studies. I started with a model allowing all two-way interactions among the predictors. I then tested whether adding the three-way interaction among land use, land-use intensity and trophic level improved the fit of the model, using a likelihood-ratio test. The model that included the three-way interaction was retained ($P \ll 0.01$; model 1, Figure 5.2). In addi-

1901 tion, because it is well established that resting metabolic rates are influenced by temperature (Clarke and
1902 Fraser, 2004), I checked whether including annual mean temperature in the model affected the conclusions.
1903 Annual mean temperature at each PREDICTS site was estimated from WorldClim version 2.1 (Fick and
1904 Hijmans, 2017), using a 2.5 arc-minute resolution. Adding annual mean temperature did not improve model
1905 fit (likelihood-ratio test: $P=0.113$), thus I did not consider its effects any further.

1906 **Model validation.**

1907 To ensure that imputation uncertainty did not affect the conclusions, I refitted model 1 using the subset of
1908 species ($n = 426$) from PREDICTS for which there were empirical RMR information (i.e., excluding imputed
1909 RMR values).

1910 **Disentangling the effects of body mass and abundance on tRMR.**

1911 Since RMR correlates strongly with body mass, changes in tRMR are likely to be driven in part by changes
1912 in the size-spectrum of ecological assemblages. I fitted an additional model to explain changes in species'
1913 abundance (given presence) by land use, land-use intensity, trophic group, body mass and their interactions,
1914 to understand the role of shifts in the body mass of species on observed changes in tRMR (Appendix 4,
1915 Figure S5.1).

1916 **5.2.6 Effects of land use, land-use intensity, trophic group and residual RMR on species
1917 occurrence probability (prediction 2; Figure 5.2b)**

1918 To control for the effects of body mass and taxonomy on RMR, I used the residual variation in RMR after
1919 accounting for these variables, from a linear mixed-effects model fitting \log_e -RMR as a function of \log_e -
1920 body mass with nested random taxonomic effects (1|Class/Order/Family; Figure 5.2). Hence, I used a metric
1921 that describes how much more energy (positive deviations) or less energy (negative deviations) than expected
1922 from body mass and taxonomic position a species spends for organismal maintenance. Similar approaches
1923 have been used in previous papers (Furness and Speakman, 2008; Naya et al., 2013). As detailed earlier, I
1924 expect species with lower residual RMR to do better in disturbed land uses than species with higher residual
1925 RMR (prediction 2; Figure 5.2b) because, given any body mass, investing less energy in maintenance could
1926 contribute to persistence in a context of resource scarcity.

1927 To test the second prediction, I fitted a binomial mixed-effects model explaining species occurrence
1928 with land use, land-use intensity, trophic group and residual RMR. I started with a complete model that
1929 included all two-way interactions among the main effects. Because I wanted to test whether the second

1930 prediction was valid for each trophic group, I needed to account for potential differences in the slope of the
1931 relationships between occurrence probability and residual RMR among trophic groups. Thus, I performed
1932 a forward stepwise selection procedure to test whether adding three-way interactions among (1) land use,
1933 trophic group and residual RMR and (2) among land-use intensity, trophic group and residual RMR improved
1934 model fit, using likelihood-ratio tests. The final model included both three-way interactions (Figure 5.2b;
1935 model 2). I fitted random effects that accounted for species identity, as well as for study and site identity
1936 within PREDICTS.

1937 **Model validation.**

1938 I checked the phylogenetic signal in the model residuals using Pagel's λ (Pagel, 1999). Non-significant
1939 phylogenetic signal in the residuals would indicate that fitting species identity in the model's random effects
1940 was sufficient to account for residual phylogenetic variation in RMR. Further, to assess the potential effects
1941 of imputation uncertainty on the results, I again fitted model 2 on the data subset for the 489 species with
1942 collected empirical RMR values, across 5,948 sites in 151 studies (i.e., excluding imputed values).

1943 **5.3 Results**

1944 **5.3.1 Effects of land use, land-use intensity and trophic group on assemblage-level total
1945 RMR**

1946 Land use, land-use intensity, trophic group and their interactions had significant effects on assemblage-level
1947 total RMR (Figure 5.3). Overall, and contrary to our expectations, assemblage-level total RMR did not
1948 show systematic decreases in disturbed land uses. In fact, urban land uses were associated with strong
1949 significant increases in tRMR in all trophic groups (e.g., a 200% average increase in tRMR in lightly-used
1950 urban areas for carnivores, compared with primary vegetation levels; +207% on average in lightly-used urban
1951 areas for herbivores; +107% for minimally-used urban areas for omnivores). In other land uses, responses
1952 depended on trophic group and land-use intensity. Whilst for herbivores, disturbed land uses were typically
1953 associated with increases in tRMR, we detected decreases in tRMR for omnivores and carnivores in several
1954 land uses, most notably in intensely-used pasture for carnivores (-84%). Such effects could reflect changes
1955 in the size-spectrum of local assemblages (Appendix 4, Figure S5.2). For instance, in minimally-used urban
1956 areas, larger herbivores tended to occur at higher abundances compared to primary vegetation level; and in
1957 intensely-used pastures, carnivores tended to occur at lower abundances overall (Figure S5.2).

1958 The model residuals were appropriately distributed (see diagnostic plots in Appendix 4, Figure S5.3).
 1959 Investigating the sensitivity of our results to imputation uncertainty showed that our results and conclusions
 1960 were robust to the removal of all imputed estimates of RMR (the correlation coefficient was 0.72 between the
 1961 two sets of model coefficients; Figure S5.4). Comparing model predictions showed that effects were mostly
 1962 congruent, although there were some differences (Figure S5.5). In particular, for herbivores, effect sizes
 1963 tended to be bigger for the model fitted on empirical data compared with the model that included imputed
 1964 data. Thus, our main results appear to be conservative if anything. The model fitted on empirical data had
 1965 larger standard errors, likely due to the reduction in sample size.





Figure 5.3: Effects of land use, land-use intensity and trophic group on assemblage-level total RMR. Model predictions are rescaled with reference to minimally-used primary vegetation, considered to be the undisturbed baseline. Primary: primary vegetation; secondary: secondary vegetation; plantation: plantation forest.

1966 5.3.2 Effects of land use, land-use intensity, trophic group and residual RMR on species' 1967 occurrence probability

1968 Species' occurrence probability was significantly affected by land use, land-use intensity, trophic level, resid-
 1969 ual RMR and their interactions (Figures 5.4, 5.5). Contrary to our expectations, species with higher residual
 1970 RMR (relative to their body mass and taxonomic position) tended to do better than species with lower resid-
 1971 ual RMR in a number of disturbed land uses. Overall, land-use type was more important for determining the
 1972 relationship between occurrence probability and residual RMR than land-use intensity (Figure 5.4a).

(a) Slope of the relationships between occurrence probability (log-odds) and residual RMR



(b) Predicted effects of residual RMR on occurrence probability (minimal use intensity)




Figure 5.4: (a) Slope estimates for the relationship between residual RMR and occurrence probability in each land-use type and for the three levels of land-use intensity. The black horizontal line indicates the slope for the reference level (primary vegetation) for minimal land-use intensity. The grey dashed line marks 0. Error bars are 95% confidence intervals. (b) Effect of residual RMR on species probability of occurrence within each trophic level and for each land use-type. I plotted the predictions for minimal land-use intensity only. Solid lines represent significant relationships. Primary: primary vegetation; secondary: secondary vegetation; plantation: plantation forest.

1973 For minimally-used primary vegetation (reference), the model predicted negative effects of residual RMR
 1974 on species occurrence probability in all trophic levels (but with a significant slope for herbivores only; Figure
 1975 5.4a). However, the directionality of this relationship was reversed in some disturbed land uses in all trophic
 1976 groups (secondary vegetation, cropland and urban for omnivores; cropland for herbivores; urban for carni-
 1977 vores), with significant positive slopes, also significantly higher than those observed for primary vegetation
 1978 (Figure 5.4a). The only exception was the opposite pattern for urban herbivores (Figure 5.4b), where residual
 1979 RMR had a more negative effect on occurrence probability than in minimally-used primary vegetation.

1980 I would like to emphasize that positive effects of residual RMR on occurrence probability in some of the
 1981 most disturbed land uses (e.g., urban for carnivores) do not mean that there were absolute increases in species
 1982 occurrence probability in disturbed land uses compared to primary vegetation (and vice-versa). I illustrate
 1983 this point in Figure 5.5. For carnivores with a median value for residual RMR, occurrence probability was
 1984 reduced by an average 24% in urban land uses compared to primary vegetation.




Figure 5.5: Predicted occurrence probabilities (+/- 95% confidence interval) in primary vegetation (primary) and for the most disturbed land uses where residual RMR was found to have a significant effect on occurrence probability. For visualisation purposes, I discretised residual RMR in three levels. The predicted probabilities of occurrence were rescaled with reference to primary vegetation for the lowest value of residual RMR (25% quantile). Here, the predictions are plotted for minimal land-use intensity (effects would be similar for light and intense land-use intensities). Black points and error bars are plotted where the relationship between occurrence probability and residual RMR was significant (and dark grey points and error bars represent non-significant trends).

1985 Finally, the model showed some degree of deviation from distributional assumptions (diagnostic plots,
 1986 Appendix 4, Figure S5.6). Nevertheless, the model's coefficients were similar when estimated with a
 1987 Bayesian framework, suggesting that the estimates were robust (Figure S5.7). The phylogenetic signal in
 1988 the model residuals was weak and non-significant ($\lambda=0.004$, $P=0.24$). Re-fitting the model using the com-
 1989 plete data subset (i.e., excluding imputed RMR estimates) showed that our conclusions are likely robust to
 1990 imputation uncertainty (Figure S5.8), with congruent results overall, although there were a few differences
 1991 in the predictions between the two models – notably, for herbivores in urban land uses (Figure S5.8).

1992 5.4 Discussion

1993 The results of this Chapter provide insights into the relationship between land-use change and energetic
1994 requirements at two levels of organisation (at the assemblage level and the species level), and contradict
1995 both my initial predictions. I found that total energetic requirements did not show systematic decreases in
1996 disturbed land uses. On the contrary, there were strong increases in all trophic groups, most notably in urban
1997 land uses. Further, in all trophic groups, species with higher energetic expenditure (than expected from body
1998 mass and taxonomy) tended to do better in some of the most disturbed land uses (that is, cropland and urban)
1999 than species with lower energetic expenditure (than expected from body mass and taxonomy).

2000 At the assemblage level, the results first highlight the effects of land-use change on vertebrate com-
2001 munity metabolism. Contrary to my expectations, I found differing effects of land use on total energetic
2002 requirements among trophic groups, reflecting changes in the size-spectrum of ecological assemblages (i.e.,
2003 changes in the distribution of abundance along the body-mass gradient). On the one hand, decreases in total
2004 energetic requirements, such as observed for carnivores in intensely used pastures, are likely due to overall
2005 reductions in local abundance (with carnivorous species in pastures perceived as a threat to livestock; Eeden
2006 et al., 2018; or because of increased competition for fewer resources). On the other hand, increases in total
2007 energetic requirements could reflect higher levels of resource intake in some disturbed land-use types. On the
2008 basis of net primary productivity, I hypothesized that there would be fewer resources available in disturbed
2009 areas. However, it could be that the carrying capacity of some disturbed land-use types actually increases
2010 for some trophic groups compared to that of primary vegetation, because of the presence of novel or more
2011 abundant food sources. For instance, in urban areas, both wildlife feeding and the presence of anthropogenic
2012 food sources, such as human refusal or pet food, could lead to an increase in resource availability all year-
2013 round for species with a non-specialised diet that are able to exploit such food sources (Fischer et al., 2012).
2014 Past research on urban carnivores has shown that some species have adapted to urban environments by ex-
2015 ploiting anthropogenic food sources, and also benefit from physical protection (with some human structures
2016 providing shelter) and from reduced occurrence of natural enemies (Bateman and Fleming, 2012). Herbi-
2017 vores and omnivores could also benefit from increased resources in urban areas, notably where the urban
2018 matrix includes semi-natural habitats, which is congruent with observed increases in occurrence probability
2019 in minimally used urban areas for these trophic groups. Further, anthropogenic ‘bottom-up food forcing’ has
2020 been shown to affect food-web dynamics, with decreases in predation rates, loss of anti-predator abilities
2021 (Geffroy et al., 2020) and mesopredator release (Fischer et al., 2012). Thus, in urban areas, increases in
2022 anthropogenic food sources could provide support for a higher abundance of medium-sized species released

2023 from predation, increasing total energetic requirements.

2024 I hypothesized that resource availability would be the main driver behind changes in energetic require-
2025 ments. However, resource availability likely interacts with other factors to explain the patterns. In particular,
2026 past work has shown that brain size is an important predictor of species ability to cope with disturbance,
2027 notably in urban settings (Sayol et al., 2020). Since brain size scales allometrically with body mass, it is
2028 possible that I observe increases in total energetic requirements where having larger brains is advantageous
2029 for resource acquisition, and in turn where resources are both exploitable and abundant enough to sustain
2030 the larger energetic expenditures associated with bigger brain sizes. In addition, body size, which explains
2031 an important proportion of the interspecific variation in mobility, likely interacts with characteristics of the
2032 landscape (such as fragmentation) to predict species survival in disturbed land uses (Merckx et al., 2018).
2033 Hence, the patterns I observe in this Chapter could emerge from interactions and trade-offs among resource
2034 availability, diet, body size and human management of the landscape.

2035 At the species level, the results underline the role of energetic constraints on species responses to land-
2036 use change. After controlling for the effects of body mass and taxonomy, I found that residual energetic
2037 expenditure was a significant predictor of species occurrence probability in disturbed land uses. Contrary
2038 to my expectations, in several disturbed land uses and in all trophic groups, species with higher residual ener-
2039 getic expenditure tended to have a higher occurrence probability than species with lower residual ener-
2040 getic expenditure. It could be that species with lower residual energetic requirements are less well equipped
2041 than species with higher residual energetic requirements at making use of the available food sources in dis-
2042 turbed land uses. Species in disturbed land uses may need to display higher levels of feeding innovation
2043 (Coogan et al., 2018), have larger brains (Sayol et al., 2020), or be bolder and more active to make use of
2044 the available resources. In mammals, past research has shown that larger brains are associated with larger
2045 residual energetic expenditure (Isler and Van Schaik, 2006). At the individual level, past research suggests
2046 that metabolic rates are linked to differences in behaviour, with bolder and more active individuals exhibiting
2047 higher metabolic rates than less active individuals, with consequences for food acquisition (Biro and Stamps,
2048 2010). Although I am not aware of similar evidence at the species level, I propose that residual metabolic
2049 rates interact with species' ecological traits, behavioural traits and foraging strategy in influencing species'
2050 responses to land use. Among species with an adaptable diet, able to make use of the resources available in
2051 a disturbed landscape, those with higher residual metabolic rates could present a set of behavioural charac-
2052 teristics that render them better at acquiring the available resources, hence more able to cope with land-use
2053 disturbance. This could be the case for urban carnivores and omnivores, as well as herbivores in cropland.
2054 Thus, it is possible that disturbed land uses favour species with higher residual resting metabolic rates be-

cause such species are overall better competitors when faced with disturbance. In addition, release from predation in some disturbed land uses – most notably in urban areas – could favour bolder species, as they are less exposed to predation risks. Conversely, and in accordance with my initial hypothesis, the results suggest that when there are no or few exploitable resources in disturbed areas, spending less energy than expected from body mass and taxonomy might be beneficial for persistence (this could be the case for urban herbivores, which include species specialised on fruit, nectar, and other plant materials that likely become less abundant and less substitutable in disturbed areas).

To summarise, I propose that, for species with adaptable diets, increases in exploitable resources coupled with decreases in predation rates leads to increases in overall energetic expenditure, associated with shifts towards assemblages composed of a higher abundance of medium-sized, bolder, more active and larger-brained species. On the other hand, species with specialised diets might be better able to persist and cope with resource scarcity when their energetic expenditure are lower than expected from body mass. These ideas could be further tested by considering more refined dietary groups (as those compiled in Chapter 4) rather than trophic groups, as diet should be easier to link with resource availability in a given land-use type. This could constitute a development for this work.

Finally, I would like to emphasize that energetic budget allocation is a fundamental aspect of organismal fitness (Robert Burger et al., 2021). Past studies suggest that individuals exposed to food scarcity can lower their intrinsic energy expenditure to increase their chances of survival. For instance, Zhang^{2018a} showed that, subjected to food restrictions, the Chinese bulbul (*Pycnonotus sinensis*) was able to lower its basal metabolic rate not only passively through body and organ mass reductions, but also by modifying enzymatic activity and metabolism in the muscles and liver. Other studies have also found that bird nestlings down-regulate energetic costs allocated to maintenance when exposed to food shortage (Brzék and Konarzewski, 2001; Moe et al., 2004), or to disturbances such as urban noise (Brischoux et al., 2017). Intraspecific variation in energetic expenditure could affect individual responses to land-use change, which in turn could affect species' ability to adapt to human pressures. However, the lack of spatially explicit estimates of metabolic rates precluded its consideration in this Chapter.

To conclude, the findings of this Chapter suggest important effects of land-use change on energetic fluxes, and thus potentially on ecosystem functioning. Further research is warranted as to the effects of these energetic patterns for ecosystem processes. Indeed, assemblage-level energy fluxes may serve as important indicators of change for ecosystem processes, such as decomposition (Barnes et al., 2014). Understanding these impacts is particularly important given increasing land-use change and biodiversity loss across the globe.

2087 **6 | General discussion**

2088 **Bibliography**

- 2089 Albert, C., Luque, G. M., and Courchamp, F. (2018). The twenty most charismatic species. *PLoS ONE*. DOI: 10.1371/journal.pone.0199149.
- 2090
- 2091 Alex Smith, M. and M. Green, D. (2005). Dispersal and the metapopulation paradigm in amphibian ecology and conservation: are all amphibian populations metapopulations? *Ecography*, 28.1, 110–128.
- 2092
- 2093 Alexander, P., Rounsevell, M. D., Dislich, C., Dodson, J. R., Engström, K., and Moran, D. (2015). Drivers for global agricultural land use change: The nexus of diet, population, yield and bioenergy. *Global Environmental Change*, 35, 138–147. DOI: 10.1016/j.gloenvcha.2015.08.011.
- 2094
- 2095
- 2096 Alves, R. R. N., Souto, W. M. S., Fernandes-Ferreira, H., Bezerra, D. M. M., Barboza, R. R. D., and Vieira, W. L. S. (2018). *The Importance of Hunting in Human Societies*. Elsevier Inc., 95–118. DOI: 10.1016/B978-0-12-809913-1.00007-7.
- 2097
- 2098
- 2099 Angert, A. L., Crozier, L. G., Rissler, L. J., Gilman, S. E., Tewksbury, J. J., and Chunco, A. J. (2011). Do species' traits predict recent shifts at expanding range edges? DOI: 10.1111/j.1461-0248.2011.01620.x.
- 2100
- 2101
- 2102 Aronson, M. F., Lepczyk, C. A., Evans, K. L., Goddard, M. A., Lerman, S. B., MacIvor, J. S., Nilon, C. H., and Vargo, T. (2017). Biodiversity in the city: key challenges for urban green space management. *Frontiers in Ecology and the Environment*, 15.4, 189–196. DOI: 10.1002/fee.1480.
- 2103
- 2104
- 2105 Auer, S. K., Killen, S. S., and Rezende, E. L. (2017). Resting vs. active: a meta-analysis of the intra- and inter-specific associations between minimum, sustained, and maximum metabolic rates in vertebrates. *Functional Ecology*, 31.9, 1728–1738. DOI: 10.1111/1365-2435.12879.
- 2106
- 2107
- 2108 Auer, S. K., Solowey, J. R., Rajesh, S., and Rezende, E. L. (2020). Energetic mechanisms for coping with changes in resource availability. *Biology Letters*, 16.11, 1–8. DOI: 10.1098/rsbl.2020.0580.
- 2109
- 2110 Balmford, A. (1996). Extinction filters and current resilience: The significance of past selection pressures for conservation biology. *Trends in Ecology and Evolution*. DOI: 10.1016/0169-5347(96)10026-4.
- 2111
- 2112 Banks-Leite, C., Ewers, R. M., Folkard-Tapp, H., and Fraser, A. (2020). Counteracting the effects of habitat loss, fragmentation, and degradation through habitat restoration. *One Earth*, 3.6, 672–676. DOI: 10.1016/j.oneear.2020.11.016.
- 2113
- 2114
- 2115 Baraldi, A. N. and Enders, C. K. (2010). An introduction to modern missing data analyses. *Journal of School Psychology*. DOI: 10.1016/j.jsp.2009.10.001.
- 2116
- 2117 Barber, N. A., Mooney, K. A., Greenberg, R., Philpott, S. M., Van Bael, S. A., and Gruner, D. S. (2010). Interactions among predators and the cascading effects of vertebrate insectivores on arthropod communities and plants. *Proceedings of the National Academy of Sciences*. DOI: 10.1073/pnas.1001934107.
- 2118
- 2119

- 2120 Barlow, J., França, F., Gardner, T. A., Hicks, C. C., Lennox, G. D., Berenguer, E., Castello, L., Economo,
2121 E. P., Ferreira, J., Guénard, B., Gontijo Leal, C., Isaac, V., Lees, A. C., Parr, C. L., Wilson, S. K., Young,
2122 P. J., and Graham, N. A. (2018). The future of hyperdiverse tropical ecosystems. *Nature*. DOI: 10 .
2123 1038/s41586-018-0301-1.
- 2124 Barnes, A. D., Jochum, M., Mumme, S., Haneda, N. F., Farajallah, A., Widarto, T. H., and Brose, U. (2014).
2125 Consequences of tropical land use for multitrophic biodiversity and ecosystem functioning. *Nature Com-
2126 munications*, 5, 1–7. DOI: 10.1038/ncomms6351.
- 2127 Barnosky, A. D., Matzke, N., Tomaia, S., Wogan, G. O., Swartz, B., Quental, T. B., Marshall, C., McGuire,
2128 J. L., Lindsey, E. L., Maguire, K. C., Mersey, B., and Ferrer, E. A. (2011). Has the Earth’s sixth mass
2129 extinction already arrived? *Nature*. DOI: 10.1038/nature09678.
- 2130 Bateman, P. W. and Fleming, P. A. (2012). Big city life: Carnivores in urban environments. *Journal of Zool-
2131 ogy*, 287.1, 1–23. DOI: 10.1111/j.1469-7998.2011.00887.x.
- 2132 Bates, D., Mächler, M., Bolker, B. M., and Walker, S. C. (2015). Fitting linear mixed-effects models using
2133 lme4. *Journal of Statistical Software*. DOI: 10.18637/jss.v067.i01.
- 2134 Betts, M. G. et al. (2019). Extinction filters mediate the global effects of habitat fragmentation on animals.
2135 *Science*. DOI: 10.1126/science.aax9387.
- 2136 Biermann, F., Bai, X., Bondre, N., Broadgate, W., Arthur Chen, C. T., Dube, O. P., Erisman, J. W., Glaser, M.,
2137 Hel, S. van der, Lemos, M. C., Seitzinger, S., and Seto, K. C. (2016). Down to Earth: Contextualizing the
2138 Anthropocene. *Global Environmental Change*, 39, 341–350. DOI: 10.1016/j.gloenvcha.2015.
2139 11.004.
- 2140 Biro, P. A. and Stamps, J. A. (2010). Do consistent individual differences in metabolic rate promote con-
2141 sistent individual differences in behavior? *Trends in Ecology and Evolution*, 25.11, 653–659. DOI: 10 .
2142 1016/j.tree.2010.08.003.
- 2143 Bivand, R. and Piras, G. (2015). Comparing Implementations of Estimation Methods for Spatial Economet-
2144 rics. *Journal of Statistical Software*, 63.18, 1–36.
- 2145 Bivand, R. and Wong, D. W. S. (2018). Comparing implementations of global and local indicators of spatial
2146 association. *TEST*, 27.3, 716–748.
- 2147 Bivand, R., Hauke, J., and Kossowski, T. (2013a). Computing the Jacobian in Gaussian spatial autoregressive
2148 models: An illustrated comparison of available methods. *Geographical Analysis*, 45.2, 150–179.
- 2149 Bivand, R. S., Pebesma, E., and Gomez-Rubio, V. (2013b). *Applied spatial data analysis with R, Second
2150 edition*. Springer, NY.
- 2151 Böhm, M., Cook, D., Ma, H., Davidson, A. D., García, A., Tapley, B., Pearce-Kelly, P., and Carr, J. (2016).
2152 Hot and bothered: Using trait-based approaches to assess climate change vulnerability in reptiles. *Bio-
2153 logical Conservation*. DOI: 10.1016/j.biocon.2016.06.002.
- 2154 Bolnick, D. I., Amarasekare, P., Araújo, M. S., Bürger, R., Levine, J. M., Novak, M., Rudolf, V. H., Schreiber,
2155 S. J., Urban, M. C., and Vasseur, D. A. (2011). Why intraspecific trait variation matters in community
2156 ecology. *Trends in Ecology and Evolution*. DOI: 10.1016/j.tree.2011.01.009.
- 2157 Bouckaert, R., Heled, J., Kühnert, D., Vaughan, T., Wu, C. H., Xie, D., Suchard, M. A., Rambaut, A., and
2158 Drummond, A. J. (2014). BEAST 2: A Software Platform for Bayesian Evolutionary Analysis. *PLoS
2159 Computational Biology*. DOI: 10.1371/journal.pcbi.1003537.

- 2160 Bouckaert, R. et al. (2019). BEAST 2.5: An advanced software platform for Bayesian evolutionary analysis.
2161 *PLoS Computational Biology*. DOI: 10.1371/journal.pcbi.1006650.
- 2162 Bregman, T. P., Lees, A. C., Seddon, N., Macgregor, H. E., Darski, B., Aleixo, A., Bonsall, M. B., and
2163 Tobias, J. A. (2015). Species interactions regulate the collapse of biodiversity and ecosystem function in
2164 tropical forest fragments. *Ecology*, 96.10, 2692–2704. DOI: 10.1890/14-1731.1.
- 2165 Bregman, T. P., Lees, A. C., MacGregor, H. E., Darski, B., Moura, N. G. de, Aleixo, A., Barlow, J., and
2166 Tobias, J. A. (2016). Using avian functional traits to assess the impact of land-cover change on ecosystem
2167 processes linked to resilience in tropical forests. *Proceedings of the Royal Society B: Biological Sciences*,
2168 283.1844. DOI: 10.1098/rspb.2016.1289.
- 2169 Brischoux, F., Meillère, A., Dupoué, A., Lourdais, O., and Angelier, F. (2017). Traffic noise decreases
2170 nestlings' metabolic rates in an urban exploiter. *Journal of Avian Biology*, 48.7, 905–909. DOI: 10 .
2171 1111/j.avb.01139.
- 2172 Broughton, J. M. and Weitzel, E. M. (2018). Population reconstructions for humans and megafauna suggest
2173 mixed causes for North American Pleistocene extinctions. *Nature Communications*, 9.1, 1–12. DOI: 10 .
2174 1038/s41467-018-07897-1.
- 2175 Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M., and West, G. B. (2004). Toward a metabolic theory
2176 of ecology. *Ecology*, 85.7, 1771–1789. DOI: 10.2106/JBJS.J.01311.
- 2177 Bruggeman, J., Heringa, J., and Brandt, B. W. (2009). PhyloPars: Estimation of missing parameter values
2178 using phylogeny. *Nucleic Acids Research*. DOI: 10.1093/nar/gkp370.
- 2179 Brzék, P. and Konarzewski, M. (2001). Effect of food shortage on the physiology and competitive abilities
2180 of sand martin (*Riparia riparia*) nestlings. *Journal of Experimental Biology*, 204.17, 3065–3074. DOI:
2181 10.1242/jeb.204.17.3065.
- 2182 Buchanan, G. M., Butchart, S. H., Chandler, G., and Gregory, R. D. (2020). Assessment of national-level
2183 progress towards elements of the Aichi Biodiversity Targets. *Ecological Indicators*, 116.May, 106497.
2184 DOI: 10.1016/j.ecolind.2020.106497.
- 2185 Burton, T., Killen, S. S., Armstrong, J. D., and Metcalfe, N. B. (2011). What causes intraspecific variation
2186 in resting metabolic rate and what are its ecological consequences? *Proceedings of the Royal Society B:*
2187 *Biological Sciences*, 278.1724, 3465–3473. DOI: 10.1098/rspb.2011.1778.
- 2188 Bushuev, A., Tolstenkov, O., Zubkova, E., Solovyeva, E., and Kerimov, A. (2018). Basal metabolic rate
2189 in free-living tropical birds: The influence of phylogenetic, behavioral, and ecological factors. *Current
2190 Zoology*, 64.1, 33–43. DOI: 10.1093/cz/zox018.
- 2191 Butchart, S. H. M. et al. (2010). Global Biodiversity: Indicators of Recent Declines. *Science*, 328.5982,
2192 1164–1168. DOI: 10.1126/science.1187512.
- 2193 Butchart, S. H., Di Marco, M., and Watson, J. E. (2016). Formulating Smart Commitments on Biodiversity:
2194 Lessons from the Aichi Targets. *Conservation Letters*, 9.6, 457–468. DOI: 10.1111/conl.12278.
- 2195 Buuren, S van and Oudshoorn, C. G. M. (2007). MICE: multivariate imputation by chained equations. *R
2196 package version*.
- 2197 Cadotte, M. W. and Tucker, C. M. (2017). Should Environmental Filtering be Abandoned? *Trends in Ecology
2198 and Evolution*. DOI: 10.1016/j.tree.2017.03.004.

- 2199 Cadotte, M. W., Carscadden, K., and Mirochnick, N. (2011). Beyond species: Functional diversity and the
2200 maintenance of ecological processes and services. *Journal of Applied Ecology*. DOI: 10.1111/j.
2201 1365-2664.2011.02048.x.
- 2202 Cahill, A. E., Aiello-Lammens, M. E., Caitlin Fisher-Reid, M., Hua, X., Karanewsky, C. J., Ryu, H. Y.,
2203 Sbeglia, G. C., Spagnolo, F., Waldron, J. B., Warsi, O., and Wiens, J. J. (2013). How does climate change
2204 cause extinction? *Proceedings of the Royal Society B: Biological Sciences*, 280.1750. DOI: 10.1098/
2205 rspb.2012.1890.
- 2206 Cannistra, A. F. and Buckley, L. B. (2021). Improving range shift predictions: enhancing the power of traits.
2207 *bioRxiv*, 2021.02.15.431292. DOI: 10.1101/2021.02.15.431292.
- 2208 Capdevila, P., Noviello, N., McRae, L., Freeman, R., and Clements, C. (2022a). Body mass and latitude
2209 as global predictors of vertebrate populations exposure to multiple threats. *bioRxiv*. DOI: 10.1101/
2210 2022.02.04.479091.
- 2211 Capdevila, P., Stott, I., Cant, J., Beger, M., Rowlands, G., Grace, M., and Salguero-Gómez, R. (2022b). Life
2212 history mediates the trade-offs among different components of demographic resilience. *Ecology Letters*,
2213 July 2021, 1–14. DOI: 10.1111/ele.14004.
- 2214 Capuracho, J. M., Ashley, M. V., Tsuru, B. R., Cooper, J. C., and Bates, J. M. (2020). Dispersal ability
2215 correlates with range size in Amazonian habitat-restricted birds. *Proceedings of the Royal Society B:
2216 Biological Sciences*, 287.1939. DOI: 10.1098/rspb.2020.1450 rspb20201450.
- 2217 Cardoso, D. et al. (2017). Amazon plant diversity revealed by a taxonomically verified species list. *Proceed-
2218 ings of the National Academy of Sciences of the United States of America*, 114.40, 10695–10700. DOI:
2219 10.1073/pnas.1706756114.
- 2220 Carlson, S. M., Cunningham, C. J., and Westley, P. A. (2014). Evolutionary rescue in a changing world.
2221 *Trends in Ecology and Evolution*, 29.9, 521–530. DOI: 10.1016/j.tree.2014.06.005.
- 2222 Caro, T., Rowe, Z., Berger, J., Wholey, P., and Dobson, A. (2022). An inconvenient misconception: Climate
2223 change is not the principal driver of biodiversity loss. *Conservation Letters*, October 2021, 1–6. DOI:
2224 10.1111/conl.12868.
- 2225 Chamberlain, S. (2018). *rredlist: 'IUCN' Red List Client*. R package version 0.5.0.
- 2226 Chamberlain, S. A. and Szöcs, E. (2013). taxize : taxonomic search and retrieval in R [version 2; referees: 3
2227 approved]. *F1000Research*. DOI: 10.12688/f1000research.2-191.v2.
- 2228 Chapman, P. M., Tobias, J. A., Edwards, D. P., and Davies, R. G. (2018). Contrasting impacts of land-use
2229 change on phylogenetic and functional diversity of tropical forest birds. *Journal of Applied Ecology*.
2230 DOI: 10.1111/1365-2664.13073.
- 2231 Chaudhary, A., Pourfaraj, V., and Mooers, A. O. (2018). Projecting global land use-driven evolutionary
2232 history loss. *Diversity and Distributions*, 24.2, 158–167. DOI: 10.1111/ddi.12677.
- 2233 Chaudhary, C., Saeedi, H., and Costello, M. J. (2016). Bimodality of Latitudinal Gradients in Marine Species
2234 Richness. *Trends in Ecology & Evolution*, 31.9, 670–676. DOI: <https://doi.org/10.1016/j.tree.2016.06.001>.
- 2236 Chen, I. C., Hill, J. K., Ohlemüller, R., Roy, D. B., and Thomas, C. D. (2011). Rapid range shifts of
2237 species associated with high levels of climate warming. *Science*, 333.6045, 1024–1026. DOI: 10.1126/
2238 science.1206432.

- 2239 Chichorro, F., Juslén, A., and Cardoso, P. (2019). A review of the relation between species traits and extinc-
2240 tion risk. *Biological Conservation*, 237.June, 220–229. DOI: 10.1016/j.biocon.2019.07.001.
- 2241 Chown, S. L., Gaston, K. J., and Robinson, D (2004). Macrophysiology: large-scale patterns in physiological.
2242 *Functional Ecology*, 18, 159–167.
- 2243 Chown, S. L., Hoffmann, A. A., Kristensen, T. N., Angilletta, M. J., Stenseth, N. C., and Pertoldi, C. (2010).
2244 Adapting to climate change: A perspective from evolutionary physiology. *Climate Research*, 43.1-2, 3–
2245 15. DOI: 10.3354/cr00879.
- 2246 Civantos, E., Thuiller, W., Maiorano, L., Guisan, A., and Arajo, M. B. (2012). Potential impacts of climate
2247 change on ecosystem services in Europe: The case of pest control by vertebrates. *BioScience*, 62.7, 658–
2248 666. DOI: 10.1525/bio.2012.62.7.8.
- 2249 Clarke, A. and Fraser, K. P. (2004). Why does metabolism scale with temperature? *Functional Ecology*, 18.2,
2250 243–251. DOI: 10.1111/j.0269-8463.2004.00841.x.
- 2251 Collin, N., Villéger, S., Wilkes, M., Sostoa, A. de, and Maceda-Veiga, A. (2018). Functional diversity mea-
2252 sures revealed impacts of non-native species and habitat degradation on species-poor freshwater fish
2253 assemblages. *Science of the Total Environment*. DOI: 10.1016/j.scitotenv.2017.12.316.
- 2254 Collen, B., Ram, M., Zamin, T., and McRae, L. (2008). The Tropical Biodiversity Data Gap: Address-
2255 ing Disparity in Global Monitoring. *Tropical Conservation Science*, 1.2, 75–88. DOI: 10.1177/
2256 194008290800100202.
- 2257 Collen, B., Dulvy, N. K., Gaston, K. J., Gärdenfors, U., Keith, D. A., Punt, A. E., Regan, H. M., Böhm,
2258 M., Hedges, S., Seddon, M., Butchart, S. H., Hilton-Taylor, C., Hoffmann, M., Bachman, S. P., and
2259 Akçakaya, H. R. (2016). Clarifying misconceptions of extinction risk assessment with the IUCN Red
2260 List. *Biology Letters*. DOI: 10.1098/rsbl.2015.0843.
- 2261 Convention on Biological Diversity (2022). *Report of the open-ended working group on the Post-2020 Global
2262 Biodiversity Framework on its third meeting (part II)*. Tech. rep. United Nations Convention on Biologi-
2263 cal Diversity.
- 2264 Coogan, S. C., Raubenheimer, D., Zantis, S. P., and Machovsky-Capuska, G. E. (2018). Multidimensional
2265 nutritional ecology and urban birds. *Ecosphere*, 9.4. DOI: 10.1002/ecs2.2177.
- 2266 Cooke, R. S., Bates, A. E., and Eigenbrod, F. (2019a). Global trade-offs of functional redundancy and func-
2267 tional dispersion for birds and mammals. *Global Ecology and Biogeography*, October 2018, 1–12. DOI:
2268 10.1111/geb.12869.
- 2269 Cooke, R. S., Eigenbrod, F., and Bates, A. E. (2019b). Projected losses of global mammal and bird ecological
2270 strategies. *Nature Communications*. DOI: 10.1038/s41467-019-10284-z.
- 2271 Cooper, N., Bielby, J., Thomas, G. H., and Purvis, A. (2008). Macroecology and extinction risk correlates of
2272 frogs. *Global Ecology and Biogeography*. DOI: 10.1111/j.1466-8238.2007.00355.x.
- 2273 Cornwell, W. K., Schwilk, D. W., and Ackerly, D. D. (2006). A trait-based test for habitat filtering: Convex
2274 hull volume. *Ecology*. DOI: 10.1890/0012-9658 (2006) 87 [1465:ATTFHF] 2.0.CO;2.
- 2275 Courchamp, F., Jaric, I., Albert, C., Meinard, Y., Ripple, W. J., and Chapron, G. (2018). Loved and ignored
2276 to death: the paradoxical extinction of the most charismatic animals. *PLoS Biology*, 16.4, e2003997.
- 2277 Cox, N. et al. (2022). A global reptile assessment highlights shared conservation needs of tetrapods. *Nature*,
2278 April 2021. DOI: 10.1038/s41586-022-04664-7.

- 2279 Crooks, K. and Soulé, M. (2010). Mesopredator release and avifaunal extinctions in a fragmented system.
2280 *Nature*, 563–566.
- 2281 Crowley, T. J. (2000). Causes of climate change over the past 1000 years. *Science*, 289.5477, 270–277. DOI:
2282 10.1126/science.289.5477.270.
- 2283 Crutzen, P. and Stoermer, E. (2000). Sustaining Earth's Life Support Systems – the Challenge for the Next
2284 Decade and Beyond. *IGBP newsletter*, 41, 17–18.
- 2285 Cunningham, C. X., Johnson, C. N., Barmuta, L. A., Hollings, T., Woehler, E. J., and Jones, M. E. (2018).
2286 Top carnivore decline has cascading effects on scavengers and carrion persistence. *Proceedings of the
2287 Royal Society B: Biological Sciences*. DOI: 10.1098/rspb.2018.1582.
- 2288 Daru, B. H., Davies, T. J., Willis, C. G., Meineke, E. K., Ronk, A., Zobel, M., Pärtel, M., Antonelli, A.,
2289 and Davis, C. C. (2021). Widespread homogenization of plant communities in the Anthropocene. *Nature
2290 Communications*, 12.1, 1–10. DOI: 10.1038/s41467-021-27186-8.
- 2291 Davison, C. W., Rahbek, C., and Morueta-Holme, N. (2021). Land-use change and biodiversity: Challenges
2292 for assembling evidence on the greatest threat to nature. *Global Change Biology*, 27.21, 5414–5429. DOI:
2293 10.1111/gcb.15846.
- 2294 De Palma, A., Kuhlmann, M., Roberts, S. P., Potts, S. G., Börger, L., Hudson, L. N., Lysenko, I., Newbold,
2295 T., and Purvis, A. (2015). Ecological traits affect the sensitivity of bees to land-use pressures in European
2296 agricultural landscapes. *Journal of Applied Ecology*, 52.6, 1567–1577. DOI: 10.1111/1365-2664.
2297 12524.
- 2298 De Palma, A., Sanchez-Ortiz, K., Martin, P. A., Chadwick, A., Gilbert, G., Bates, A. E., Börger, L., Contu,
2299 S., Hill, S. L., and Purvis, A. (2018). Challenges With Inferring How Land-Use Affects Terrestrial Bio-
2300 diversity: Study Design, Time, Space and Synthesis. *Advances in Ecological Research*. DOI: 10.1016/
2301 bs.aecr.2017.12.004.
- 2302 De Vos, J. M., Joppa, L. N., Gittleman, J. L., Stephens, P. R., and Pimm, S. L. (2015). Estimating the normal
2303 background rate of species extinction. *Conservation Biology*. DOI: 10.1111/cobi.12380.
- 2304 Debastiani, V. J., Bastazini, V. A. G., and Pillar, D (2021). Ecological Informatics Using phylogenetic infor-
2305 mation to impute missing functional trait values in ecological databases. 63.April. DOI: 10.1016/j.
2306 ecoinf.2021.101315.
- 2307 Des Roches, S., Post, D. M., Turley, N. E., Bailey, J. K., Hendry, A. P., Kinnison, M. T., Schweitzer, J. A.,
2308 and Palkovacs, E. P. (2018). The ecological importance of intraspecific variation. *Nature Ecology and
2309 Evolution*. DOI: 10.1038/s41559-017-0402-5.
- 2310 Deutsch, C. A., Tewksbury, J. J., Huey, R. B., Sheldon, K. S., Ghilambor, C. K., Haak, D. C., and Martin,
2311 P. R. (2008). Impacts of climate warming on terrestrial ectotherms across latitude. *Proceedings of the
2312 National Academy of Sciences of the United States of America*, 105.18, 6668–6672. DOI: 10.1073/
2313 pnas.0709472105.
- 2314 Di Marco, M., Pacifici, M., Maiorano, L., and Rondinini, C. (2021). Drivers of change in the realised climatic
2315 niche of terrestrial mammals. *Ecography*, 44.8, 1180–1190. DOI: 10.1111/ecog.05414.
- 2316 Díaz, S. et al. (2016). The global spectrum of plant form and function. *Nature*. DOI: 10.1038/nature16489.
- 2317 Diniz-Filho, J. A. F., Bini, L. M., Rangel, T. F., Morales-Castilla, I., Olalla-Tárraga, M. Á., Rodríguez,
2318 M. Á., and Hawkins, B. A. (2012). On the selection of phylogenetic eigenvectors for ecological analyses.
2319 *Ecography*. DOI: 10.1111/j.1600-0587.2011.06949.x.

- 2320 Dirzo, R., Young, H. S., Galetti, M., Ceballos, G., Isaac, N. J., and Collen, B. (2014). Defaunation in the
2321 Anthropocene. *Science*, 345.6195, 401–406. DOI: 10.1126/science.1251817.
- 2322 Dore, M. H. (2005). Climate change and changes in global precipitation patterns: What do we know? *Envi-
2323 ronment International*, 31.8, 1167–1181. DOI: 10.1016/j.envint.2005.03.004.
- 2324 Dornelas, M., Gotelli, N. J., Shimadzu, H., Moyes, F., Magurran, A. E., and McGill, B. J. (2019). A balance of
2325 winners and losers in the Anthropocene. *Ecology Letters*, 22.5, 847–854. DOI: 10.1111/ele.13242.
- 2326 Duffy, J. E. (2003). Biodiversity loss, trophic skew and ecosystem functioning. *Ecology Letters*, 6.8, 680–
2327 687. DOI: 10.1046/j.1461-0248.2003.00494.x.
- 2328 Dullinger, I., Essl, F., Moser, D., Erb, K., Haberl, H., and Dullinger, S. (2021). Biodiversity models need to
2329 represent land-use intensity more comprehensively. *Global Ecology and Biogeography*, 30.5, 924–932.
2330 DOI: 10.1111/geb.13289.
- 2331 Edwards, F. A., Edwards, D. P., Hamer, K. C., and Fayle, T. M. (2021). Tropical land-use change alters
2332 trait-based community assembly rules for dung beetles and birds. *Oecologia*, 195.3, 705–717. DOI: 10.
2333 1007/s00442-020-04829-z.
- 2334 Eeden, L. M. van et al. (2018). Carnivore conservation needs evidence-based livestock protection. *PLoS
2335 Biology*, 16.9, 1–8. DOI: 10.1371/journal.pbio.2005577.
- 2336 Ellis, E. C., Goldewijk, K. K., Siebert, S., Lightman, D., and Ramankutty, N. (2010). Anthropogenic trans-
2337 formation of the biomes, 1700 to 2000. *Global Ecology and Biogeography*, 19.5, 589–606. DOI: 10.
2338 1111/j.1466-8238.2010.00540.x.
- 2339 Ellis, E. C., Kaplan, J. O., Fuller, D. Q., Vavrus, S., Goldewijk, K. K., and Verburg, P. H. (2013). Used planet:
2340 A global history. *Proceedings of the National Academy of Sciences of the United States of America*,
2341 110.20, 7978–7985. DOI: 10.1073/pnas.1217241110.
- 2342 Ellis, E. C., Gauthier, N., Goldewijk, K. K., Bird, R. B., Boivin, N., Díaz, S., Fuller, D. Q., Gill, J. L., Kaplan,
2343 J. O., Kingston, N., Locke, H., McMichael, C. N., Ranco, D., Rick, T. C., Rebecca Shaw, M., Stephens,
2344 L., Svenning, J. C., and Watson, J. E. (2021). People have shaped most of terrestrial nature for at least
2345 12,000 years. *Proceedings of the National Academy of Sciences of the United States of America*, 118.17,
2346 1–8. DOI: 10.1073/pnas.2023483118.
- 2347 Escudero, A. and Valladares, F. (2016). Trait-based plant ecology: moving towards a unifying species coex-
2348 istence theory: Features of the Special Section. *Oecologia*. DOI: 10.1007/s00442-016-3578-5.
- 2349 ESRI (2011). ArcGIS desktop: Release 10. *Environmental Systems Research Institute*.
- 2350 Estrada, A., Morales-Castilla, I., Meireles, C., Caplat, P., and Early, R. (2018). Equipped to cope with climate
2351 change: traits associated with range filling across European taxa. *Ecography*. DOI: 10.1111/ecog.
2352 02968.
- 2353 Etard, A., Morrill, S., and Newbold, T. (2020). Global gaps in trait data for terrestrial vertebrates. *Global
2354 Ecology and Biogeography*, November 2019, 1–16. DOI: 10.1111/geb.13184.
- 2355 Etard, A., Pigot, A. L., and Newbold, T. (2021). Intensive human land uses negatively affect vertebrate
2356 functional diversity. *Ecology Letters*. DOI: <https://doi.org/10.1111/ele.13926>.
- 2357 — (2022). Intensive human land uses negatively affect vertebrate functional diversity. *Ecology Letters*, 25.2,
2358 330–343. DOI: <https://doi.org/10.1111/ele.13926>.

- 2359 Evans, B. S., Reitsma, R., Hurlbert, A. H., and Marra, P. P. (2018). Environmental filtering of avian com-
2360 munities along a rural-to-urban gradient in Greater Washington, D.C., USA. *Ecosphere*, 9.11. DOI: 10 .
2361 1002/ecs2.2402.
- 2362 Faurby, S., Davis, M., Pedersen, R., Schowanek, S. D., Antonelli1, A., and Svenning, J.-C. (2018). PHY-
2363 LACINE 1.2: The Phylogenetic Atlas of Mammal Macroecology. *Ecology*, 99.11, 2626–2626. DOI:
2364 10 .1002/ecy.2443.
- 2365 Faurby, S., Pedersen, R., Davis, M., Schowanek, S. D., Jarvie, S., Antonelli, A., and Svenning, J.-C. (2020).
2366 PHYLACINE 1.2.1: An update to the Phylogenetic Atlas of Mammal Macroecology. DOI: 10 .5281/
2367 zenodo.3690867.
- 2368 Feldman, A., Sabath, N., Pyron, R. A., Mayrose, I., and Meiri, S. (2016). Body sizes and diversification rates
2369 of lizards, snakes, amphisbaenians and the tuatara. *Global Ecology and Biogeography*, 25.2, 187–197.
2370 DOI: 10 .1111/geb .12398.
- 2371 Ferreira, D. F., Gibb, R., López-Baucells, A., Nunes, N. J., Jones, K. E., and Rocha, R. (2022). Species-
2372 specific responses to land-use change in island insectivorous bats. *Journal for Nature Conservation*,
2373 126177. DOI: 10 .1016/j.jnc .2022.126177.
- 2374 Fick, S. E. and Hijmans, R. J. (2017). WorldClim 2: new 1-km spatial resolution climate surfaces for global
2375 land areas. *International Journal of Climatology*, 37.12, 4302–4315. DOI: 10 .1002/joc .5086.
- 2376 Finderup Nielsen, T., Sand-Jensen, K., Dornelas, M., and Bruun, H. H. (2019). More is less: net gain in
2377 species richness, but biotic homogenization over 140 years. *Ecology Letters*, 22.10, 1650–1657. DOI:
2378 10 .1111/ele .13361.
- 2379 Fischer, J. D., Cleeton, S. H., Lyons, T. P., and Miller, J. R. (2012). Urbanization and the predation paradox:
2380 The role of trophic dynamics in structuring vertebrate communities. *BioScience*, 62.9, 809–818. DOI:
2381 10 .1525/bio .2012 .62 .9 .6.
- 2382 Flynn, D. F., Gogol-Prokurat, M., Nogeire, T., Molinari, N., Richers, B. T., Lin, B. B., Simpson, N., May-
2383 field, M. M., and DeClerck, F. (2009). Loss of functional diversity under land use intensification across
2384 multiple taxa. *Ecology Letters*, 12.1, 22–33. DOI: 10 .1111/j .1461-0248 .2008 .01255 .x.
- 2385 Foden, W. B., Butchart, S. H. M., Stuart, S. N., Vié, J. C., Akçakaya, H. R., Angulo, A., DeVantier, L. M.,
2386 Gutsche, A., Turak, E., Cao, L., Donner, S. D., Katariya, V., Bernard, R., Holland, R. A., Hughes, A. F.,
2387 O'Hanlon, S. E., Garnett, S. T., Şekercioğlu, Ç. H., and Mace, G. M. (2013). Identifying the World's Most
2388 Climate Change Vulnerable Species: A Systematic Trait-Based Assessment of all Birds, Amphibians and
2389 Corals. *PLoS ONE*. DOI: 10 .1371/journal .pone .0065427.
- 2390 Foley, J. A., DeFries, R., Asner, G. P., Barford, C., Bonan, G., Carpenter, S. R., Chapin, F. S., Coe, M. T.,
2391 Daily, G. C., Gibbs, H. K., Helkowski, J. H., Holloway, T., Howard, E. A., Kucharik, C. J., Monfreda, C.,
2392 Patz, J. A., Prentice, I. C., Ramankutty, N., and Snyder, P. K. (2005). Global consequences of land use.
2393 *Science*, 309.5734, 570–574. DOI: 10 .1126/science .1111772.
- 2394 Fournier, B., Mouly, A., and Gillet, F. (2016). Multiple assembly rules drive the co-occurrence of orthopteran
2395 and plant species in grasslands: Combining network, functional and phylogenetic approaches. *Frontiers
2396 in Plant Science*, 7.AUG2016, 1–12. DOI: 10 .3389/fpls .2016.01224.
- 2397 Fox, J. and Monette, G. (1992). Generalized collinearity diagnostics. *Journal of the American Statistical
2398 Association*. DOI: 10 .1080/01621459 .1992 .10475190.

- 2399 Freeman, B. G. and Pennell, M. W. (2021). The latitudinal taxonomy gradient. *Trends in Ecology and Evolution*, 36.9, 778–786. DOI: 10.1016/j.tree.2021.05.003.
- 2400
- 2401 Fricke, E. C., Ordóñez, A., Rogers, H. S., and Svenning, J.-C. (2022). The effects of defaunation on plants' capacity to track climate change. *Science*, 375.6577, 210–214. DOI: 10.1126/science.abk3510.
- 2402
- 2403 Frishkoff, L. O., Karp, D. S., Flanders, J. R., Zook, J., Hadly, E. A., Daily, G. C., and M'Gonigle, L. K. (2016). Climate change and habitat conversion favour the same species. *Ecology letters*. DOI: 10.1111/ele.12645.
- 2404
- 2405
- 2406 Fukasawa, K. and Akasaka, T. (2019). Long-lasting effects of historical land use on the current distribution of mammals revealed by ecological and archaeological patterns. *Scientific Reports*, 9.1, 1–11. DOI: 10.1038/s41598-019-46809-1.
- 2407
- 2408
- 2409 Furness, L. J. and Speakman, J. R. (2008). Energetics and longevity in birds. *Age*, 30.2-3, 75–87. DOI: 10.1007/s11357-008-9054-3.
- 2410
- 2411 Gainsbury, A. M., Tallowin, O. J., and Meiri, S. (2018). An updated global data set for diet preferences in terrestrial mammals: testing the validity of extrapolation. *Mammal Review*. DOI: 10.1111/mam.12119.
- 2412
- 2413
- 2414 Galán-Acedo, C., Arroyo-Rodríguez, V., Andresen, E., and Arasa-Gisbert, R. (2019). Ecological traits of the world's primates. *Scientific data*. DOI: 10.1038/s41597-019-0059-9.
- 2415
- 2416 Gallmetzer, N. and Schulze, C. H. (2015). Impact of oil palm agriculture on understory amphibians and reptiles: A Mesoamerican perspective. *Global Ecology and Conservation*, 4, 95–109. DOI: 10.1016/j.gecco.2015.05.008.
- 2417
- 2418
- 2419 Geffroy, B., Sadoul, B., Putman, B. J., Berger-Tal, O., Garamszegi, L. Z., Møller, A. P., and Blumstein, D. T. (2020). Evolutionary dynamics in the anthropocene: Life history and intensity of human contact shape antipredator responses. *PLoS Biology*, 18.9, 1–17. DOI: 10.1371/journal.pbio.3000818.
- 2420
- 2421
- 2422 Gillooly, J. F., Brown, J. H., West, G. B., Savage, V. M., and Charnov, E. L. (2001). Effects of Size and Temperature on Metabolic Rate. *Science*, 293.5538, 2248–2251. DOI: 10.1126/science.1061967.
- 2423
- 2424 Goldewijk, K. K., Beusen, A., Doelman, J., and Stehfest, E. (2017). Anthropogenic land use estimates for the Holocene - HYDE 3.2. *Earth System Science Data*, 9.2, 927–953. DOI: 10.5194/essd-9-927-2017.
- 2425
- 2426
- 2427 González-Suárez, M. and Revilla, E. (2012). Variability in life-history and ecological traits is a buffer against extinction in mammals. *Ecology Letters*, 16.2, 242–251. DOI: 10.1111/ele.12035.
- 2428
- 2429 González-Suárez, M., Lucas, P. M., and Revilla, E. (2012). Biases in comparative analyses of extinction risk: Mind the gap. *Journal of Animal Ecology*. DOI: 10.1111/j.1365-2656.2012.01999.x.
- 2430
- 2431 González-Suárez, M., Gómez, A., and Revilla, E. (2013). Which intrinsic traits predict vulnerability to extinction depends on the actual threatening processes. *Ecosphere*. DOI: 10.1890/ES12-00380.1.
- 2432
- 2433 González-Suárez, M., Zanchetta Ferreira, F., and Grilo, C. (2018). Spatial and species-level predictions of road mortality risk using trait data. *Global Ecology and Biogeography*, 27.9, 1093–1105. DOI: 10.1111/geb.12769.
- 2434
- 2435
- 2436 González-Varo, J. P., Biesmeijer, J. C., Bommarco, R., Potts, S. G., Schweiger, O., Smith, H. G., Steffan-Dewenter, I., Szentgyörgyi, H., Woyciechowski, M., and Vilà, M. (2013). Combined effects of global
- 2437

- 2438 change pressures on animal-mediated pollination. *Trends in Ecology and Evolution*, 28.9, 524–530. DOI:
2439 10.1016/j.tree.2013.05.008.
- 2440 Guralnick, R. P., Zermoglio, P. F., Wieczorek, J., LaFrance, R., Bloom, D., and Russell, L. (2016). The
2441 importance of digitized biocollections as a source of trait data and a new VertNet resource. *Database*.
2442 DOI: 10.1093/database/baw158.
- 2443 Hadfield, J. D. (2010). MCMC Methods for Multi-Response Generalized Linear Mixed Models: The MCM-
2444 Cglmm R Package. *Journal of Statistical Software*, 33.2, 1–22.
- 2445 Hamilton, A. T., Schäfer, R. B., Pyne, M. I., Chessman, B., Kakouei, K., Boersma, K. S., Verdonschot,
2446 P. F., Verdonschot, R. C., Mims, M., Khamis, K., Bierwagen, B., and Stamp, J. (2020). Limitations of
2447 trait-based approaches for stressor assessment: The case of freshwater invertebrates and climate drivers.
2448 *Global Change Biology*, 26.2, 364–379. DOI: 10.1111/gcb.14846.
- 2449 Hantak, M. M., McLean, B. S., Li, D., and Guralnick, R. P. (2021). Mammalian body size is determined by
2450 interactions between climate, urbanization, and ecological traits. *Communications Biology*, 4.1, 1–10.
2451 DOI: 10.1038/s42003-021-02505-3.
- 2452 Harfoot, M. B., Johnston, A., Balmford, A., Burgess, N. D., Butchart, S. H., Dias, M. P., Hazin, C., Hilton-
2453 Taylor, C., Hoffmann, M., Isaac, N. J., Iversen, L. L., Outhwaite, C. L., Visconti, P., and Geldmann, J.
2454 (2021). Using the IUCN Red List to map threats to terrestrial vertebrates at global scale. *Nature Ecology
and Evolution*, 5.11, 1510–1519. DOI: 10.1038/s41559-021-01542-9.
- 2455 Hatfield, J. H., Harrison, M. L. K., and Banks-Leite, C. (2018). Functional Diversity Metrics: How They Are
2456 Affected by Landscape Change and How They Represent Ecosystem Functioning in the Tropics. *Current
2457 Landscape Ecology Reports*. DOI: 10.1007/s40823-018-0032-x.
- 2458 Hernández-Yáñez, H., Kim, S. Y., and Che-Castaldo, J. P. (2022). Demographic and life history traits ex-
2459 plain patterns in species vulnerability to extinction. *PLoS ONE*, 17.2 February, 1–15. DOI: 10.1371/journal.pone.0263504.
- 2460 Hevia, V., Martín-López, B., Palomo, S., García-Llorente, M., Bello, F. de, and González, J. A. (2017). Trait-
2461 based approaches to analyze links between the drivers of change and ecosystem services: Synthesizing
2462 existing evidence and future challenges. *Ecology and Evolution*, 7.3, 831–844. DOI: <https://doi.org/10.1002/ece3.2692>.
- 2463 Hijmans, R. J. (2022). *raster: Geographic Data Analysis and Modeling*. R package version 3.5-15.
- 2464 Hillaert, J., Hovestadt, T., Vandegehuchte, M. L., and Bonte, D. (2018). Size-dependent movement explains
2465 why bigger is better in fragmented landscapes. *Ecology and Evolution*, 8.22, 10754–10767. DOI: 10.1002/ece3.4524.
- 2466 Hirons, M., Comberti, C., and Dunford, R. (2016). Valuing Cultural Ecosystem Services. *Annual Review of
2467 Environment and Resources*, 41, 545–574. DOI: 10.1146/annurev-environ-110615-085831.
- 2468 Hoban, S. et al. (2020). Genetic diversity targets and indicators in the CBD post-2020 Global Biodiver-
2469 sity Framework must be improved. *Biological Conservation*, 248.July, 108654. DOI: 10.1016/j.biocon.2020.108654.
- 2470 Hooper, D. U., Chapin, F. S., Ewel, J. J., Hector, A., Inchausti, P., Lavorel, S., Lawton, J. H., Lodge, D. M.,
2471 Loreau, M., Naeem, S., Schmid, B., Setälä, H., Symstad, A. J., Vandermeer, J., and Wardle, D. A. (2005).
2472 Effects of biodiversity on ecosystem functioning: A consensus of current knowledge. *Ecological Mono-
2473 graphs*. DOI: 10.1890/04-0922.

- 2479 Hooper, D. U., Adair, E. C., Cardinale, B. J., Byrnes, J. E., Hungate, B. A., Matulich, K. L., Gonzalez, A.,
2480 Duffy, J. E., Gamfeldt, L., and Connor, M. I. (2012). A global synthesis reveals biodiversity loss as a
2481 major driver of ecosystem change. *Nature*. DOI: 10.1038/nature11118.
- 2482 Hortal, J., Bello, F. de, Diniz-Filho, J. A. F., Lewinsohn, T. M., Lobo, J. M., and Ladle, R. J. (2015). Seven
2483 Shortfalls that Beset Large-Scale Knowledge of Biodiversity. *Annual Review of Ecology, Evolution, and
2484 Systematics*. DOI: 10.1146/annurev-ecolsys-112414-054400.
- 2485 Hou, Y., Ding, W., Liu, C., Li, K., Cui, H., Liu, B., and Chen, W. (2022). Influences of impervious sur-
2486 faces on ecological risks and controlling strategies in rapidly urbanizing regions. *Science of The Total
2487 Environment*, 153823. DOI: 10.1016/j.scitotenv.2022.153823.
- 2488 Hudson, L. N., Isaac, N. J., and Reuman, D. C. (2013). The relationship between body mass and field
2489 metabolic rate among individual birds and mammals. *Journal of Animal Ecology*, 82.5, 1009–1020. DOI:
2490 10.1111/1365-2656.12086.
- 2491 Hudson, L. N. et al. (2014). The PREDICTS database: A global database of how local terrestrial biodiversity
2492 responds to human impacts. *Ecology and Evolution*. DOI: 10.1002/ece3.1303.
- 2493 — (2017). The database of the PREDICTS (Projecting Responses of Ecological Diversity In Changing
2494 Terrestrial Systems) project. *Ecology and Evolution*. DOI: 10.1002/ece3.2579.
- 2495 Inger, R., Cox, D. T., Per, E., Norton, B. A., and Gaston, K. J. (2016). Ecological role of vertebrate scavengers
2496 in urban ecosystems in the UK. *Ecology and Evolution*. DOI: 10.1002/ece3.2414.
- 2497 Inouye, D. W. (2022). Climate change and phenology. *Wiley Interdisciplinary Reviews: Climate Change*,
2498 November 2021, 1–17. DOI: 10.1002/wcc.764.
- 2499 Intergovernmental Panel on Climate Change (2013). *Climate Change 2013 – The Physical Science Basis*.
2500 Tech. rep. WMO, UNEP, 104.
- 2501 — (2022). *Climate Change 2022 Mitigation of Climate Change – Summary for Policymakers*. Tech. rep.
2502 WMO, UNEP.
- 2503 Isaac, N. J., Mallet, J., and Mace, G. M. (2004). Taxonomic inflation: Its influence on macroecology and
2504 conservation. *Trends in Ecology and Evolution*. DOI: 10.1016/j.tree.2004.06.004.
- 2505 Isler, K. and Van Schaik, C. P. (2006). Metabolic costs of brain size evolution. *Biology Letters*, 2.4, 557–560.
2506 DOI: 10.1098/rsbl.2006.0538.
- 2507 IUCN (2020). The IUCN Red List of Threatened Species. <http://www.iucnredlist.org>. Version 2020.1.
- 2508 Ives, C. D., Lentini, P. E., Threlfall, C. G., Ikin, K., Shanahan, D. F., Garrard, G. E., Bekessy, S. A., Fuller,
2509 R. A., Mumaw, L., Rayner, L., Rowe, R., Valentine, L. E., and Kendal, D. (2016). Cities are hotspots for
2510 threatened species. *Global Ecology and Biogeography*, 25.1, 117–126. DOI: 10.1111/geb.12404.
- 2511 Jayathilake, H. M., Prescott, G. W., Carrasco, L. R., Rao, M., and Symes, W. S. (2021). Drivers of deforesta-
2512 tion and degradation for 28 tropical conservation landscapes. *Ambio*, 50.1, 215–228. DOI: 10.1007/
2513 s13280-020-01325-9.
- 2514 Jenkins, C. N., Pimm, S. L., and Joppa, L. N. (2013). Global patterns of terrestrial vertebrate diversity and
2515 conservation. *Proceedings of the National Academy of Sciences of the United States of America*, 110.28,
2516 E2603–E2610. DOI: 10.1073/pnas.1302251110.
- 2517 Jenkins, D. G., Brescacin, C. R., Duxbury, C. V., Elliott, J. A., Evans, J. A., Grablow, K. R., Hillegass, M.,
2518 Lyon, B. N., Metzger, G. A., Olandese, M. L., Pepe, D., Silvers, G. A., Suresch, H. N., Thompson,

- 2519 T. N., Trexler, C. M., Williams, G. E., Williams, N. C., and Williams, S. E. (2007). Does size matter
2520 for dispersal distance? *Global Ecology and Biogeography*, 16.4, 415–425. DOI: 10.1111/j.1466-
2521 8238.2007.00312.x.
- 2522 Jesse, W. A., Behm, J. E., Helmus, M. R., and Ellers, J. (2018). Human land use promotes the abundance
2523 and diversity of exotic species on Caribbean islands. *Global Change Biology*, 24.10, 4784–4796. DOI:
2524 10.1111/gcb.14334.
- 2525 Jetz, W., Thomas, G. H., Joy, J. B., Hartmann, K., and Mooers, A. O. (2012). The global diversity of birds in
2526 space and time. *Nature*. DOI: 10.1038/nature11631.
- 2527 Jetz, W. and Pyron, R. A. (2018). The interplay of past diversification and evolutionary isolation with
2528 present imperilment across the amphibian tree of life. *Nature Ecology and Evolution*. DOI: 10.1038/
2529 s41559-018-0515-5.
- 2530 Johnson, C. N., Alroy, J., Beeton, N. J., Bird, M. I., Brook, B. W., Cooper, A., Gillespie, R., Herrando-Pérez,
2531 S., Jacobs, Z., Miller, G. H., Prideaux, G. J., Roberts, R. G., Rodríguez-Rey, M., Saltré, F., Turney, C. S.,
2532 and Bradshaw, C. J. (2016). What caused extinction of the pleistocene megafauna of Sahul? *Proceedings
2533 of the Royal Society B: Biological Sciences*, 283.1824. DOI: 10.1098/rspb.2015.2399.
- 2534 Johnson, C. N., Balmford, A., Brook, B. W., Buettel, J. C., Galetti, M., Guangchun, L., and Wilmshurst,
2535 J. M. (2017). *Biodiversity losses and conservation responses in the Anthropocene*. DOI: 10.1126/
2536 science.aam9317.
- 2537 Johnson, T. F., Isaac, N. J., Paviolo, A., and González-Suárez, M. (2021). Handling missing values in trait
2538 data. *Global Ecology and Biogeography*. DOI: 10.1111/geb.13185.
- 2539 Jones, K. E. et al. (2009). PanTHERIA: a species-level database of life history, ecology, and geography of
2540 extant and recently extinct mammals. *Ecology*. DOI: 10.1890/08-1494.1.
- 2541 Jones, O. R., Purvis, A., and Quicke, D. L. (2012). Latitudinal gradients in taxonomic overdescription rate
2542 affect macroecological inferences using species list data. *Ecography*. DOI: 10.1111/j.1600-0587.
2543 2011.06956.x.
- 2544 Junker, R. R., Albrecht, J., Becker, M., Keuth, R., Farwig, N., and Schleuning, M. (2022). Towards an animal
2545 economics spectrum for ecosystem research. *Functional Ecology*, March, 1–16. DOI: 10.1111/1365-
2546 2435.14051.
- 2547 Kattge, J. et al. (2011). TRY - a global database of plant traits. *Global Change Biology*. DOI: 10.1111/j.
2548 1365-2486.2011.02451.x.
- 2549 Kehoe, L., Kuemmerle, T., Meyer, C., Levers, C., Václavík, T., and Kreft, H. (2015). Global patterns of
2550 agricultural land-use intensity and vertebrate diversity. *Diversity and Distributions*, 21.11, 1308–1318.
2551 DOI: 10.1111/ddi.12359.
- 2552 Kharouba, H. M., Ehrlén, J., Gelman, A., Bolmgren, K., Allen, J. M., Travers, S. E., and Wolkovich, E. M.
2553 (2018). Global shifts in the phenological synchrony of species interactions over recent decades. *Pro-
2554 ceedings of the National Academy of Sciences of the United States of America*, 115.20, 5211–5216. DOI:
2555 10.1073/pnas.1714511115.
- 2556 Kissling, W. D. et al. (2018). Towards global data products of Essential Biodiversity Variables on species
2557 traits. *Nature Ecology and Evolution*. DOI: 10.1038/s41559-018-0667-3.
- 2558 Kissling, W. D., Dalby, L., Fløjgaard, C., Lenoir, J., Sandel, B., Sandom, C., Trøjelsgaard, K., and Svenning,
2559 J. C. (2014). Establishing macroecological trait datasets: Digitalization, extrapolation, and validation of

- 2560 diet preferences in terrestrial mammals worldwide. *Ecology and Evolution*. DOI: 10.1002/ece3.
 2561 1136.
- 2562 Krausmann, F., Erb, K. H., Gingrich, S., Haberl, H., Bondeau, A., Gaube, V., Lauk, C., Plutzar, C., and
 2563 Searchinger, T. D. (2013). Global human appropriation of net primary production doubled in the 20th
 2564 century. *Proceedings of the National Academy of Sciences of the United States of America*, 110.25,
 2565 10324–10329. DOI: 10.1073/pnas.1211349110.
- 2566 Krauss, J., Bommarco, R., Guardiola, M., Heikkinen, R. K., Helm, A., Kuussaari, M., Lindborg, R., Öckinger,
 2567 E., Pärtel, M., Pino, J., Pöyry, J., Raatikainen, K. M., Sang, A., Stefanescu, C., Teder, T., Zobel, M., and
 2568 Steffan-Dewenter, I. (2010). Habitat fragmentation causes immediate and time-delayed biodiversity loss
 2569 at different trophic levels. *Ecology Letters*. DOI: 10.1111/j.1461-0248.2010.01457.x.
- 2570 La Sorte, F. A., Lepczyk, C. A., Aronson, M. F., Goddard, M. A., Hedblom, M., Katti, M., MacGregor-Fors,
 2571 I., Mörtberg, U., Nilon, C. H., Warren, P. S., Williams, N. S., and Yang, J. (2018). The phylogenetic and
 2572 functional diversity of regional breeding bird assemblages is reduced and constricted through urbaniza-
 2573 tion. *Diversity and Distributions*. DOI: 10.1111/ddi.12738.
- 2574 Laliberté, E. and Legendre, P. (2010). A distance-based framework for measuring functional diversity from
 2575 multiple traits. *Ecology*, 91.1, 299–305. DOI: 10.1890/08-2244.1.
- 2576 Laliberté, E., Legendre, P., and Shipley, B. (2015). FD: measuring functional diversity from multiple traits,
 2577 and other tools for functional ecology. *R Package*.
- 2578 Lambin, E. F. et al. (2001). The causes of land-use and land-cover change : Moving beyond the myths.
 2579 *Global Environmental Change*, 11.December, 261–269.
- 2580 Lavorel, S. and Garnier, E. (2002). Predicting changes in community composition and ecosystem functioning
 2581 from plant traits: Revisiting the Holy Grail. *Functional Ecology*. DOI: 10.1046/j.1365-2435.
 2582 2002.00664.x.
- 2583 Lawler, J. J., Lewis, D. J., Nelson, E., Plantinga, A. J., Polasky, S., Withey, J. C., Helmers, D. P., Martin-
 2584 uzzi, S., Penningtonh, D., and Radeloff, V. C. (2014). Projected land-use change impacts on ecosystem
 2585 services in the United States. *Proceedings of the National Academy of Sciences of the United States of
 2586 America*, 111.20, 7492–7497. DOI: 10.1073/pnas.1405557111.
- 2587 Le Provost, G., Badenauer, I., Le Bagousse-Pinguet, Y., Clough, Y., Henckel, L., Violle, C., Bretagnolle,
 2588 V., Roncoroni, M., Manning, P., and Gross, N. (2020). Land-use history impacts functional diversity
 2589 across multiple trophic groups. *Proceedings of the National Academy of Sciences of the United States of
 2590 America*. DOI: 10.1073/pnas.1910023117.
- 2591 Lebreton, J. D. (2011). The impact of global change on terrestrial vertebrates. *Comptes Rendus - Biologies*,
 2592 334.5-6, 360–369. DOI: 10.1016/j.crvi.2011.01.005.
- 2593 Legras, G., Loiseau, N., and Gaertner, J. C. (2018). Functional richness: Overview of indices and underlying
 2594 concepts. *Acta Oecologica*. DOI: 10.1016/j.actao.2018.02.007.
- 2595 Lenoir, J. and Svenning, J. C. (2015). Climate-related range shifts - a global multidimensional synthesis and
 2596 new research directions. *Ecography*, 38.1, 15–28. DOI: 10.1111/ecog.00967.
- 2597 Leroy, B., Meynard, C. N., Bellard, C., and Courchamp, F. (2015). virtualspecies, an R package to generate
 2598 virtual species distributions. *Ecography*. DOI: 10.1111/ecog.01388.

- 2599 Letnic, M., Ritchie, E. G., and Dickman, C. R. (2012). Top predators as biodiversity regulators: The dingo
2600 Canis lupus dingo as a case study. *Biological Reviews*. DOI: 10.1111/j.1469-185X.2011.
2601 00203.x.
- 2602 Leung, B., Hargreaves, A. L., Greenberg, D. A., McGill, B., Dornelas, M., and Freeman, R. (2020). Clustered
2603 versus catastrophic global vertebrate declines. *Nature*, 588.7837, 267–271. DOI: 10.1038/s41586-
2604 020-2920-6.
- 2605 Lewis, S. L. and Maslin, M. A. (2015). Defining the Anthropocene. *Nature*, 519.7542, 171–180. DOI: 10.
2606 1038/nature14258.
- 2607 Li, G., Fang, C., Li, Y., Wang, Z., Sun, S., He, S., Qi, W., Bao, C., Ma, H., Fan, Y., Feng, Y., and Liu, X.
2608 (2022). Global impacts of future urban expansion on terrestrial vertebrate diversity. *Nature Communications*,
2609 13.1, 1–12. DOI: 10.1038/s41467-022-29324-2.
- 2610 Lin, F., Jia, S., Luskin, M. S., Ye, J., Hao, Z., Wang, X., and Yuan, Z. (2018). Global signal of top-down
2611 control of terrestrial plant communities by herbivores. *Proceedings of the National Academy of Sciences*.
2612 DOI: 10.1073/pnas.1707984115.
- 2613 Livesley, S. J., McPherson, E. G., and Calfapietra, C. (2016). The Urban Forest and Ecosystem Services:
2614 Impacts on Urban Water, Heat, and Pollution Cycles at the Tree, Street, and City Scale. *Journal of
2615 Environmental Quality*, 45.1, 119–124. DOI: 10.2134/jeq2015.11.0567.
- 2616 Lowry, H., Lill, A., and Wong, B. B. (2013). Behavioural responses of wildlife to urban environments.
2617 *Biological Reviews*, 88.3, 537–549. DOI: 10.1111/brv.12012.
- 2618 Luck, G. W., Lavorel, S., McIntyre, S., and Lumb, K. (2012). Improving the application of vertebrate trait-
2619 based frameworks to the study of ecosystem services. *Journal of Animal Ecology*. DOI: 10.1111/j.
2620 1365-2656.2012.01974.x.
- 2621 MacLean, S. A. and Beissinger, S. R. (2017). Species' traits as predictors of range shifts under contemporary
2622 climate change: A review and meta-analysis. *Global Change Biology*, 23.10, 4094–4105. DOI: 10.
2623 1111/gcb.13736.
- 2624 Magioli, M., Ferraz, K. M. P. M. d. B., Chiarello, A. G., Galetti, M., Setz, E. Z. F., Paglia, A. P., Abrego,
2625 N., Ribeiro, M. C., and Ovaskainen, O. (2021). Land-use changes lead to functional loss of terrestrial
2626 mammals in a Neotropical rainforest. *Perspectives in Ecology and Conservation*, 19.2, 161–170. DOI:
2627 10.1016/j.pecon.2021.02.006.
- 2628 Maibach, E., Myers, T., and Leiserowitz, A. (2014). Climate scientists need to set the record straight: There
2629 is a scientific consensus that human-caused climate change is happening. *Earth's Future*, 2.5, 295–298.
2630 DOI: 10.1002/2013ef000226.
- 2631 Májeková, M., Paal, T., Plowman, N. S., Bryndová, M., Kasari, L., Norberg, A., Weiss, M., Bishop, T. R.,
2632 Luke, S. H., Sam, K., Le Bagousse-Pinguet, Y., Lepš, J., Götzenberger, L., and De Bello, F. (2016).
2633 Evaluating Functional diversity: Missing trait data and the importance of species abundance structure
2634 and data transformation. *PLoS ONE*. DOI: 10.1371/journal.pone.0149270.
- 2635 Malhi, Y. (2017). The Concept of the Anthropocene. *Annual Review of Environment and Resources*, 42, 77–
2636 104. DOI: 10.1146/annurev-environ-102016-060854.
- 2637 Marcacci, G., Westphal, C., Wenzel, A., Raj, V., Nölke, N., Tscharntke, T., and Grass, I. (2021). Taxonomic
2638 and functional homogenization of farmland birds along an urbanization gradient in a tropical megacity.
2639 *Global Change Biology*, June, 1–15. DOI: 10.1111/gcb.15755.

- 2640 Martin, L. J., Blossey, B., and Ellis, E. (2012). Mapping where ecologists work: biases in the global distri-
2641 bution of terrestrial ecological observations. *Frontiers in Ecology and the Environment*, 10.4, 195–201.
2642 DOI: 10.1890/110154.
- 2643 Martins, I. S. and Pereira, H. M. (2017). Improving extinction projections across scales and habitats using the
2644 countryside species-area relationship. *Scientific Reports*. DOI: 10.1038/s41598-017-13059-y.
- 2645 Matich, P. and Schalk, C. M. (2019). Move it or lose it: Interspecific variation in risk response of pond-
2646 breeding anurans. *PeerJ*, 7, 1–16. DOI: 10.7717/PEERJ.6956.
- 2647 Matuoka, M. A., Benchimol, M., Almeida-Rocha, J. M. de, and Morante-Filho, J. C. (2020). Effects of
2648 anthropogenic disturbances on bird functional diversity: A global meta-analysis. *Ecological Indicators*.
2649 DOI: 10.1016/j.ecolind.2020.106471.
- 2650 Maxwell, S. L., Fuller, R. A., Brooks, T. M., and Watson, J. E. M. (2016). Biodiversity: The ravages of guns,
2651 nets and bulldozers. *Nature*. DOI: 10.1038/536143a.
- 2652 May, R. M. (2011). Why worry about how many species and their loss? *PLoS Biology*. DOI: 10.1371/
2653 journal.pbio.1001130.
- 2654 McCain, C. M. and King, S. R. B. (2014). Body size and activity times mediate mammalian responses to
2655 climate change. *Global Change Biology*. DOI: 10.1111/gcb.12499.
- 2656 McGill, B. J., Enquist, B. J., Weiher, E., and Westoby, M. (2006). Rebuilding community ecology from
2657 functional traits. *Trends in Ecology and Evolution*. DOI: 10.1016/j.tree.2006.02.002.
- 2658 McGill, B. J., Dornelas, M., Gotelli, N. J., and Magurran, A. E. (2015). Fifteen forms of biodiversity trend in
2659 the anthropocene. *Trends in Ecology and Evolution*, 30.2, 104–113. DOI: 10.1016/j.tree.2014.
2660 11.006.
- 2661 McNab, B. K. (1986). The Influence of Food Habits on the Energetics of Eutherian Mammals. *Ecological
2662 Monographs*, 56.1, 1–19.
- 2663 Meiri, S., Feldman, A., and Kratochvíl, L. (2015). Squamate hatchling size and the evolutionary causes of
2664 negative offspring size allometry. *Journal of Evolutionary Biology*, 28.2, 438–446. DOI: 10.1111/jeb.12580.
- 2666 Meiri, S. (2018). Traits of lizards of the world: Variation around a successful evolutionary design. *Global
2667 Ecology and Biogeography*, 27.10, 1168–1172. DOI: 10.1111/geb.12773.
- 2668 Meiri, S. and Chapple, D. G. (2016). Biases in the current knowledge of threat status in lizards, and bridging
2669 the ‘assessment gap’. *Biological Conservation*, 204, 6–15. DOI: 10.1016/j.biocon.2016.03.
2670 009.
- 2671 Mendoza, M. and Araújo, M. B. (2019). Climate shapes mammal community trophic structures and humans
2672 simplify them. *Nature Communications*, 10.1. DOI: 10.1038/s41467-019-12995-9.
- 2673 Menegotto, A. and Rangel, T. F. (2018). Mapping knowledge gaps in marine diversity reveals a latitudi-
2674 nal gradient of missing species richness. *Nature Communications*. DOI: 10.1038/s41467-018-
2675 07217-7.
- 2676 Merckx, T. et al. (2018). Body-size shifts in aquatic and terrestrial urban communities. *Nature*. DOI: 10.
2677 1038/s41586-018-0140-0.
- 2678 Meyer, C., Kreft, H., Guralnick, R., and Jetz, W. (2015). Global priorities for an effective information basis
2679 of biodiversity distributions. *Nature Communications*, 6. DOI: 10.1038/ncomms9221.

- 2680 Millard, J., Outhwaite, C. L., Kinnersley, R., Freeman, R., Gregory, R. D., Adedoja, O., Gavini, S., Kioko, E.,
2681 Kuhlmann, M., Ollerton, J., Ren, Z.-x., and Newbold, T. (2021). Global effects of land-use intensity on
2682 local pollinator biodiversity. *Nature Communications*, 1–11. DOI: 10.1038/s41467-021-23228-
2683 3.
- 2684 Millennium Ecosystem Assessment (2005). *Millennium Ecosystem Assessment: Ecosystems and Human
2685 Well-being: Synthesis*. Tech. rep.
- 2686 Miller, G., Magee, J., Smith, M., Spooner, N., Baynes, A., Lehman, S., Fogel, M., Johnston, H., Williams, D.,
2687 Clark, P., Florian, C., Holst, R., and DeVogel, S. (2016). Human predation contributed to the extinction
2688 of the Australian megafaunal bird *Genyornis newtoni* ~ 47 ka. *Nature Communications*, 7, 1–7. DOI:
2689 10.1038/ncomms10496.
- 2690 Moe, B., Brunvoll, S., Mork, D., Brobak, T. E., and Bech, C. (2004). Developmental plasticity of physi-
2691 ology and morphology in diet-restricted European shag nestlings (*Phalacrocorax aristotelis*). *Journal of
2692 Experimental Biology*, 207.23, 4067–4076. DOI: 10.1242/jeb.01226.
- 2693 Möller, A. P. (2009). Successful city dwellers: A comparative study of the ecological characteristics of urban
2694 birds in the Western Palearctic. *Oecologia*, 159.4, 849–858. DOI: 10.1007/s00442-008-1259-8.
- 2695 Monastersky, R. (2014). Life - a status report. *Nature*, 516, 161.
2696 — (2015). The human age. *Nature*, 519, 144–147.
- 2697 Morelli, F., Benedetti, Y., Hanson, J. O., and Fuller, R. A. (2021). Global distribution and conservation of
2698 avian diet specialization. *Conservation Letters*, 14.4, 1–12. DOI: 10.1111/conl.12795.
- 2699 Mouillot, D., Loiseau, N., Grenié, M., Algar, A. C., Allegra, M., Cadotte, M. W., Casajus, N., Denelle, P.,
2700 Guéguen, M., Maire, A., Maitner, B., McGill, B. J., McLean, M., Mouquet, N., Munoz, F., Thuiller, W.,
2701 Villéger, S., Violle, C., and Auber, A. (2021). The dimensionality and structure of species trait spaces.
2702 *Ecology Letters*, March, 1–22. DOI: 10.1111/ele.13778.
- 2703 Mueller, P. and Diamond, J. (2001). Metabolic rate and environmental productivity: Well-provisioned ani-
2704 mals evolved to run and idle fast. *Proceedings of the National Academy of Sciences of the United States
2705 of America*, 98.22, 12550–12554. DOI: 10.1073/pnas.221456698.
- 2706 Munteanu, C., Kamp, J., Nita, M. D., Klein, N., Kraemer, B. M., Müller, D., Koshkina, A., Prishchepov,
2707 A. V., and Kuemmerle, T. (2020). Cold War spy satellite images reveal long-term declines of a philopatric
2708 keystone species in response to cropland expansion. *Proceedings of the Royal Society B: Biological
2709 Sciences*, 287.1927. DOI: 10.1098/rspb.2019.2897.
- 2710 Myhrvold, N. P., Baldridge, E., Chan, B., Sivam, D., Freeman, D. L., and Ernest, S. K. M. (2015). An amniote
2711 life-history database to perform comparative analyses with birds, mammals, and reptiles. *Ecology*. DOI:
2712 10.1890/15-0846R.1.
- 2713 Nakagawa, S. and Freckleton, R. P. (2008). Missing inaction: the dangers of ignoring missing data. *Trends
2714 in Ecology and Evolution*. DOI: 10.1016/j.tree.2008.06.014.
- 2715 Naya, D. E., Spangenberg, L., Naya, H., and Bozinovic, F. (2013). How does evolutionary variation in basal
2716 metabolic rates arise? A statistical assessment and a mechanistic model. *Evolution*, 67.5, 1463–1476.
2717 DOI: 10.1111/evo.12042.
- 2718 Newbold, T. (2018). Future effects of climate and land-use change on terrestrial vertebrate community diver-
2719 sity under different scenarios. *Proceedings of the Royal Society of London Series B, Biological Sciences*,
2720 20180792. DOI: 10.1098/rspb.2018.0792.

- 2721 Newbold, T., Scharlemann, J. P. W., Butchart, S. H. M., Sekercioğlu, C. H., Alkemade, R., Booth, H., and
2722 Purves, D. W. (2013). Ecological traits affect the response of tropical forest bird species to land-use
2723 intensity. *Proceedings. Biological sciences / The Royal Society*. DOI: 10.1098/rspb.2012.2131.
- 2724 Newbold, T. et al. (2015). Global effects of land use on local terrestrial biodiversity. *Nature*. DOI: 10.
2725 1038/nature14324.
- 2726 Newbold, T., Hudson, L. N., Contu, S., Hill, S. L., Beck, J., Liu, Y., Meyer, C., Phillips, H. R., Scharlemann,
2727 J. P., and Purvis, A. (2018). Widespread winners and narrow-ranged losers: Land use homogenizes biodi-
2728 versity in local assemblages worldwide. *PLoS Biology*. DOI: 10.1371/journal.pbio.2006841.
- 2729 Newbold, T., Adams, G. L., Robles, G. A., Boakes, E. H., Ferreira, G. B., Chapman, A. S., Etard, A., Gibb,
2730 R., Millard, J., Outhwaite, C. L., and Williams, J. J. (2019). Climate and land-use change homogenise
2731 terrestrial biodiversity, with consequences for ecosystem functioning and human well-being. *Emerging
2732 Topics in Life Sciences*, 3.2, 207–219. DOI: 10.1042/ETLS20180135.
- 2733 Newbold, T., Bentley, L. F., Hill, S. L., Edgar, M. J., Horton, M., Su, G., Şekercioğlu, Ç. H., Collen, B., and
2734 Purvis, A. (2020a). Global effects of land use on biodiversity differ among functional groups. *Functional
2735 Ecology*, 34.3, 684–693. DOI: 10.1111/1365-2435.13500.
- 2736 Newbold, T., Oppenheimer, P., Etard, A., and Williams, J. J. (2020b). Tropical and Mediterranean biodiver-
2737 sity is disproportionately sensitive to land-use and climate change. *Nature Ecology and Evolution*. DOI:
2738 10.1038/s41559-020-01303-0.
- 2739 Novosolov, M., Raia, P., and Meiri, S. (2013). The island syndrome in lizards. *Global Ecology and Biogeog-
2740 raphy*. DOI: 10.1111/j.1466-8238.2012.00791.x.
- 2741 Novosolov, M., Rodda, G. H., North, A. C., Butchart, S. H., Tallowin, O. J., Gainsbury, A. M., and Meiri,
2742 S. (2017). Population density–range size relationship revisited. *Global Ecology and Biogeography*. DOI:
2743 10.1111/geb.12617.
- 2744 Nowakowski, A. J., Thompson, M. E., Donnelly, M. A., and Todd, B. D. (2017). Amphibian sensitivity to
2745 habitat modification is associated with population trends and species traits. *Global Ecology and Biogeog-
2746 raphy*. DOI: 10.1111/geb.12571.
- 2747 Nowakowski, A. J., Frishkoff, L. O., Thompson, M. E., Smith, T. M., and Todd, B. D. (2018). Phylogenetic
2748 homogenization of amphibian assemblages in human-altered habitats across the globe. *Proceedings of
2749 the National Academy of Sciences of the United States of America*, 115.15, E3454–E3462. DOI: 10.
2750 1073/pnas.1714891115.
- 2751 Oliveira, B. F., São-Pedro, V. A., Santos-Barrera, G., Penone, C., and Costa, G. C. (2017). AmphiBIO, a
2752 global database for amphibian ecological traits. *Scientific Data*. DOI: 10.1038/sdata.2017.123.
- 2753 Oliver, R. Y., Meyer, C., Ranipeta, A., Winner, K., and Jetz, W. (2021). Global and national trends, gaps,
2754 and opportunities in documenting and monitoring species distributions. *PLoS Biology*, 19.8, 1–14. DOI:
2755 10.1371/journal.pbio.3001336.
- 2756 Oliver, T. H., Heard, M. S., Isaac, N. J., Roy, D. B., Procter, D., Eigenbrod, F., Freckleton, R., Hector, A.,
2757 Orme, C. D. L., Petchey, O. L., Proença, V., Raffaelli, D., Suttle, K. B., Mace, G. M., Martín-López,
2758 B., Woodcock, B. A., and Bullock, J. M. (2015). Biodiversity and Resilience of Ecosystem Functions.
2759 *Trends in Ecology and Evolution*. DOI: 10.1016/j.tree.2015.08.009.
- 2760 Orme, D., Freckleton, R., Thomas, G., Petzoldt, T., Fritz, S., Isaac, N., and Pearse, W. (2018). *caper: Com-
2761 parative Analyses of Phylogenetics and Evolution in R*. R package version 1.0.1.

- 2762 Pacifici, M. et al. (2015). Assessing species vulnerability to climate change. *Nature Climate Change*. DOI:
2763 10.1038/nclimate2448.
- 2764 Pacifici, M., Visconti, P., Butchart, S. H. M., Watson, J. E. M., Cassola, F. M., and Rondinini, C. (2017).
2765 Species' traits influenced their response to recent climate change. *Nature Climate Change*. DOI: 10 .
2766 1038/nclimate3223.
- 2767 Pagel, M. (1999). Inferring the historical patterns of biological evolution. *Nature*. DOI: 10.1038/44766.
- 2768 Pakeman, R. J. (2014). Functional trait metrics are sensitive to the completeness of the species' trait data?
2769 *Methods in Ecology and Evolution*. DOI: 10.1111/2041-210X.12136.
- 2770 Paradis, E., Baillie, S. R., Sutherland, W. J., and Gregory, R. D. (1998). Patterns of natal and breeding
2771 dispersal in birds. *Journal of Animal Ecology*, 67.4, 518–536. DOI: 10 . 1046 / j . 1365 – 2656 .
2772 1998.00215.x.
- 2773 Pearson, R. G., Stanton, J. C., Shoemaker, K. T., Aiello-Lammens, M. E., Ersts, P. J., Horning, N., Ford-
2774 ham, D. A., Raxworthy, C. J., Ryu, H. Y., Mcnees, J., and Akçakaya, H. R. (2014). Life history and
2775 spatial traits predict extinction risk due to climate change. *Nature Climate Change*. DOI: 10 . 1038 /
2776 nclimate2113.
- 2777 Penone, C., Davidson, A. D., Shoemaker, K. T., Di Marco, M., Rondinini, C., Brooks, T. M., Young, B. E.,
2778 Graham, C. H., and Costa, G. C. (2014). Imputation of missing data in life-history trait datasets: Which
2779 approach performs the best? *Methods in Ecology and Evolution*. DOI: 10.1111/2041-210X.12232.
- 2780 Pereira, H. M. et al. (2010). Scenarios for global biodiversity in the 21st century. *Science*, 330.6010, 1496–
2781 1501. DOI: 10.1126/science.1196624.
- 2782 Perronne, R., Munoz, F., Borgy, B., Reboud, X., and Gaba, S. (2017). How to design trait-based analyses
2783 of community assembly mechanisms: Insights and guidelines from a literature review. *Perspectives in
2784 Plant Ecology, Evolution and Systematics*. DOI: 10.1016/j.ppees.2017.01.004.
- 2785 Peugh, J. L. and Enders, C. K. (2004). Missing data in educational research: A review of reporting practices
2786 and suggestions for improvement. *Review of Educational Research*. DOI: 10 . 3102 / 00346543074004525.
- 2787 Pörtner, H. O. and Farrell, A. P. (2008). Physiology and Climate Change. *Science*, 322.5902, 690–692. DOI:
2788 10.1126/science.1163156.
- 2789 Powell, T. W. R. and Lenton, T. M. (2013). Scenarios for future biodiversity loss due to multiple drivers
2790 reveal conflict between mitigating climate change and preserving biodiversity. *Environmental Research
2791 Letters*, 8.2, 025024. DOI: 10.1088/1748-9326/8/2/025024.
- 2792 Powers, R. P. and Jetz, W. (2019). Global habitat loss and extinction risk of terrestrial vertebrates under
2793 future land-use-change scenarios. *Nature Climate Change*, 9.4, 323–329. DOI: 10 . 1038 / s41558 –
2794 019 -0406-z.
- 2795 Purvis, A., Gittleman, J. L., Cowlishaw, G., and Mace, G. M. (2000). Predicting extinction risk in declining
2796 species. *Proceedings of the Royal Society B: Biological Sciences*. DOI: 10.1098/rspb.2000.1234.
- 2797 Pyšek, P., Bacher, S., Chytrý, M., Jarošík, V., Wild, J., Celesti-Grapow, L., Gassó, N., Kenis, M., Lambdon,
2798 P. W., Nentwig, W., Pergl, J., Roques, A., Sádlo, J., Solarz, W., Vilà, M., and Hulme, P. E. (2010).
2799 Contrasting patterns in the invasions of European terrestrial and freshwater habitats by alien plants,
2800 insects and vertebrates. *Global Ecology and Biogeography*, 19.3, 317–331. DOI: 10.1111/j.1466 –
2801 8238.2009.00514.x.

- 2802 Quesnelle, P. E., Lindsay, K. E., and Fahrig, L. (2014). Low reproductive rate predicts species sensitivity to
2803 habitat loss: A meta-analysis of wetland vertebrates. *PLoS ONE*. DOI: 10.1371/journal.pone.0090926.
- 2804
- 2805 R Core Team (2018). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical
2806 Computing. Vienna, Austria.
- 2807 Ramesh, T., Kalle, R., and Downs, C. T. (2017). Space use in a South African agriculture landscape by the
2808 caracal (Caracal caracal). *European Journal of Wildlife Research*, 63.1. DOI: 10.1007/s10344-
2809 016-1072-3.
- 2810 Ramírez-Bautista, A., Thorne, J. H., Schwartz, M. W., and Williams, J. N. (2020). Trait-based climate vulner-
2811 ability of native rodents in southwestern Mexico. *Ecology and Evolution*. DOI: 10.1002/ece3.6323.
- 2812 Ratto, F., Simmons, B. I., Spake, R., Zamora-Gutierrez, V., MacDonald, M. A., Merriman, J. C., Tremlett,
2813 C. J., Poppy, G. M., Peh, K. S., and Dicks, L. V. (2018). Global importance of vertebrate pollinators for
2814 plant reproductive success: a meta-analysis. *Frontiers in Ecology and the Environment*. DOI: 10.1002/
2815 fee.1763.
- 2816 Revell, L. J. (2012). phytools: An R package for phylogenetic comparative biology (and other things). *Meth-
2817 ods in Ecology and Evolution*, 3, 217–223.
- 2818 Riemann, J. C., Ndriantsoa, S. H., Rödel, M. O., and Glos, J. (2017). Functional diversity in a fragmented
2819 landscape — Habitat alterations affect functional trait composition of frog assemblages in Madagascar.
2820 *Global Ecology and Conservation*, 10, 173–183. DOI: 10.1016/j.gecco.2017.03.005.
- 2821 Rinnan, D. S. (2021). *CENFA: Climate and Ecological Niche Factor Analysis*. R package version 1.1.1.
- 2822 Rinnan, D. S. and Lawler, J. (2019). Climate-niche factor analysis: a spatial approach to quantifying species
2823 vulnerability to climate change. *Ecography*, 42.9, 1494–1503. DOI: <https://doi.org/10.1111/ecog.03937>.
- 2825 Ripple, W. J., Estes, J. A., Beschta, R. L., Wilmers, C. C., Ritchie, E. G., Hebblewhite, M., Berger, J.,
2826 Elmhagen, B., Letnic, M., Nelson, M. P., Schmitz, O. J., Smith, D. W., Wallach, A. D., and Wirsing,
2827 A. J. (2014). Status and ecological effects of the world's largest carnivores. *Science*, 343.6167. DOI:
2828 10.1126/science.1241484.
- 2829 Ripple, W. J., Wolf, C., Newsome, T. M., Hoffmann, M., Wirsing, A. J., and McCauley, D. J. (2017). Ex-
2830 tinction risk is most acute for the world's largest and smallest vertebrates. *Proceedings of the National
2831 Academy of Sciences*. DOI: 10.1073/pnas.1702078114.
- 2832 Robert Burger, J., Hou, C., A. S. Hall, C., and Brown, J. H. (2021). Universal rules of life: metabolic rates,
2833 biological times and the equal fitness paradigm. *Ecology Letters*, 24.6, 1262–1281. DOI: 10.1111/
2834 ele.13715.
- 2835 Rodrigues, A. S., Pilgrim, J. D., Lamoreux, J. F., Hoffmann, M., and Brooks, T. M. (2006). The value of the
2836 IUCN Red List for conservation. *Trends in Ecology and Evolution*, 21.2, 71–76. DOI: 10.1016/j.
2837 tree.2005.10.010.
- 2838 Rohr, J. R., Civitello, D. J., Cohen, J. M., Roznik, E. A., Sinervo, B., and Dell, A. I. (2018). The complex
2839 drivers of thermal acclimation and breadth in ectotherms. *Ecology Letters*, 21.9, 1425–1439. DOI: 10.
2840 1111/ele.13107.
- 2841 Roll, U. et al. (2017). The global distribution of tetrapods reveals a need for targeted reptile conservation.
2842 *Nature Ecology and Evolution*. DOI: 10.1038/s41559-017-0332-2.

- 2843 Running, S. W. and Zhao, M. (2015). Daily GPP and Annual NPP (MOD17A2/A3) products NASA Earth
2844 Observing System MODIS Land Algorithm (User's guide V3). *User Guide*.
- 2845 Salo, P., Banks, P. B., Dickman, C. R., and Korpimäki, E. (2010). Predator manipulation experiments: Im-
2846 pacts on populations of terrestrial vertebrate prey. *Ecological Monographs*. DOI: 10.1890/09-1260 .
2847 1.
- 2848 Samia, D. S., Nakagawa, S., Nomura, F., Rangel, T. F., and Blumstein, D. T. (2015). Increased tolerance to
2849 humans among disturbed wildlife. *Nature Communications*, 6. DOI: 10.1038/ncomms9877.
- 2850 Sanchez-Ortiz, K., Taylor, K. J., Palma, A. de, König, C., Pyšek, P., Weigelt, P., Essl, F., and Purvis, A.
2851 (2019). Effects of land-use change and related pressures on alien and native subsets of island communi-
2852 ties. DOI: 10.1101/2019.12.16.878041.
- 2853 Sandom, C., Faurby, S., Sandel, B., and Svenning, J. C. (2014). Global late Quaternary megafauna extinc-
2854 tions linked to humans, not climate change. *Proceedings of the Royal Society B: Biological Sciences*,
2855 281.1787. DOI: 10.1098/rspb.2013.3254.
- 2856 Santini, L., Benítez-López, A., Ficetola, G. F., and Huijbregts, M. A. J. (2018). Length–mass allometries in
2857 amphibians. *Integrative Zoology*, 13.1, 36–45. DOI: 10.1111/1749-4877.12268.
- 2858 Santos, T. (2018). Package ‘PVR’. Phylogenetic Eigenvectors Regression and Phylogenetic Signal-Representation
2859 Curve.
- 2860 Sayol, F., Sol, D., and Pigot, A. L. (2020). Brain Size and Life History Interact to Predict Urban Tolerance
2861 in Birds. *Frontiers in Ecology and Evolution*, 8, 58. DOI: 10.3389/fevo.2020.00058.
- 2862 Scharf, I., Feldman, A., Novosolov, M., Pincheira-Donoso, D., Das, I., Böhm, M., Uetz, P., Torres-Carvajal,
2863 O., Bauer, A., Roll, U., and Meiri, S. (2015). Late bloomers and baby boomers: Ecological drivers of
2864 longevity in squamates and the tuatara. *Global Ecology and Biogeography*. DOI: 10.1111/geb .
2865 12244.
- 2866 Schielzeth, H., Dingemanse, N. J., Nakagawa, S., Westneat, D. F., Allegue, H., Teplitsky, C., Réale, D.,
2867 Dochtermann, N. A., Garamszegi, L. Z., and Araya-Ajoy, Y. G. (2020). Robustness of linear mixed-
2868 effects models to violations of distributional assumptions. *Methods in Ecology and Evolution*, 11.9,
2869 1141–1152. DOI: 10.1111/2041-210X.13434.
- 2870 Schipper, A. M., Hilbers, J. P., Meijer, J. R., Antão, L. H., Benítez-López, A., Jonge, M. M. de, Leemans,
2871 L. H., Scheper, E., Alkemade, R., Doelman, J. C., Mylius, S., Stehfest, E., Vuuren, D. P. van, Zeist, W. J.
2872 van, and Huijbregts, M. A. (2020). Projecting terrestrial biodiversity intactness with GLOBIO 4. *Global
2873 Change Biology*, 26.2, 760–771. DOI: 10.1111/gcb.14848.
- 2874 Schleuter, D., Daufresne, M., Massol, F., and Argillier, C. (2010). A user's guide to functional diversity
2875 indices. *Ecological Monographs*. DOI: 10.1890/08-2225.1.
- 2876 Schloss, C. A., Nunez, T. A., and Lawler, J. J. (2012). Dispersal will limit ability of mammals to track
2877 climate change in the Western Hemisphere. *Proceedings of the National Academy of Sciences*. DOI:
2878 10.1073/pnas.1116791109.
- 2879 Schneider, F. D., Fichtmueller, D., Gossner, M. M., Güntsch, A., Jochum, M., König-Ries, B., Le Provost,
2880 G., Manning, P., Ostrowski, A., Penone, C., and Simons, N. K. (2019). Towards an ecological trait-data
2881 standard. *Methods in Ecology and Evolution*. DOI: 10.1111/2041-210X.13288.

- 2882 Schwarz, R. and Meiri, S. (2017). The fast-slow life-history continuum in insular lizards: a comparison
2883 between species with invariant and variable clutch sizes. *Journal of Biogeography*. DOI: 10.1111/jbi.13067.
- 2885 Segan, D. B., Murray, K. A., and Watson, J. E. (2016). A global assessment of current and future biodiversity
2886 vulnerability to habitat loss-climate change interactions. *Global Ecology and Conservation*, 5, 12–21.
2887 DOI: 10.1016/j.gecco.2015.11.002.
- 2888 Seneviratne, S. I. et al. (2012). Changes in climate extremes and their impacts on the natural physical en-
2889 vironment. *Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adapta-*
2890 *tion: Special Report of the Intergovernmental Panel on Climate Change*, 9781107025, 109–230. DOI:
2891 10.1017/CBO9781139177245.006.
- 2892 Seto, K. C., Sánchez-Rodríguez, R., and Fragkias, M. (2010). The new geography of contemporary urbaniza-
2893 tion and the environment. *Annual Review of Environment and Resources*, 35, 167–194. DOI: 10.1146/annurev-environ-100809-125336.
- 2895 Sévêque, A., Gentle, L. K., López-Bao, J. V., Yarnell, R. W., and Uzal, A. (2020). Human disturbance has
2896 contrasting effects on niche partitioning within carnivore communities. *Biological Reviews*, 95.6, 1689–
2897 1705. DOI: 10.1111/brv.12635.
- 2898 Severtsov (2012). The Significance of Vertebrates in the Structure and Functioning of Ecosystems. *Biology*
2899 *Bulletin*, 40.7, 571–579.
- 2900 Siefert, A. et al. (2015). A global meta-analysis of the relative extent of intraspecific trait variation in plant
2901 communities. *Ecology Letters*, 18.12, 1406–1419. DOI: 10.1111/ele.12508.
- 2902 Slavenko, A., Tallowin, O. J., Itescu, Y., Raia, P., and Meiri, S. (2016). Late Quaternary reptile extinctions:
2903 size matters, insularity dominates. *Global Ecology and Biogeography*. DOI: 10.1111/geb.12491.
- 2904 Sodhi, N. S., Bickford, D., Diesmos, A. C., Lee, T. M., Koh, L. P., Brook, B. W., Sekercioglu, C. H., and
2905 Bradshaw, C. J. (2008). Measuring the meltdown: Drivers of global amphibian extinction and decline.
2906 *PLoS ONE*. DOI: 10.1371/journal.pone.0001636.
- 2907 Sol, D., Trisos, C., Múrria, C., Jeliazkov, A., González-Lagos, C., Pigot, A. L., Ricotta, C., Swan, C. M.,
2908 Tobias, J. A., and Pavoine, S. (2020). The worldwide impact of urbanisation on avian functional diversity.
2909 *Ecology Letters*, 23.6, 962–972. DOI: 10.1111/ele.13495.
- 2910 Soroye, P., Newbold, T., and Kerr, J. (2020). Climate change contributes to widespread declines among
2911 bumble bees across continents. *Science*, 367.6478, 685–688. DOI: 10.1126/science.aax8591.
- 2912 Souza, F. L., Valente-Neto, F., Severo-Neto, F., Bueno, B., Ochoa-Quintero, J. M., Laps, R. R., Bolzan, F.,
2913 and Roque, F. d. O. (2019). Impervious surface and heterogeneity are opposite drivers to maintain bird
2914 richness in a Cerrado city. *Landscape and Urban Planning*, 192.August, 103643. DOI: 10.1016/j.landurbplan.2019.103643.
- 2916 Spooner, F. E., Pearson, R. G., and Freeman, R. (2018). Rapid warming is associated with population decline
2917 among terrestrial birds and mammals globally. *Global Change Biology*. DOI: 10.1111/gcb.14361.
- 2918 Stark, G., Tamar, K., Itescu, Y., Feldman, A., and Meiri, S. (2018). Cold and isolated ectotherms: drivers
2919 of reptilian longevity. *Biological Journal of the Linnean Society*. DOI: 10.1093/biolinnean/bly153/5145102.
- 2921 Stark, G., Pincheira-Donoso, D., and Meiri, S. (2020). No evidence for the ‘rate-of-living’ theory across the
2922 tetrapod tree of life. *Global Ecology and Biogeography*, 29.5, 857–884. DOI: 10.1111/geb.13069.

- 2923 Steffen, W., Grinevald, J., Crutzen, P., and Mcneill, J. (2011). The anthropocene: Conceptual and historical
2924 perspectives. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 369.1938, 842–867. DOI: 10.1098/rsta.2010.0327.
- 2925
- 2926 Steffen, W., Broadgate, W., Deutsch, L., Gaffney, O., and Ludwig, C. (2015). *The trajectory of the Anthropocene: The Great Acceleration*. DOI: 10.1177/2053019614564785.
- 2927
- 2928 Stehfest, E., Zeist, W. J. van, Valin, H., Havlik, P., Popp, A., Kyle, P., Tabeau, A., Mason-D'Croz, D.,
2929 Hasegawa, T., Bodirsky, B. L., Calvin, K., Doelman, J. C., Fujimori, S., Humpenöder, F., Lotze-Campen,
2930 H., Meijl, H. van, and Wiebe, K. (2019). Key determinants of global land-use projections. *Nature Communications*, 10.1, 1–10. DOI: 10.1038/s41467-019-09945-w.
- 2931
- 2932 Stekhoven, D. J. (2016). Nonparametric Missing Value Imputation using Random Forest. *R Package version 1.4*. DOI: 10.1093/bioinformatics/btr597.
- 2933
- 2934 Stekhoven, D. J. and Bühlmann, P. (2012). Missforest-Non-parametric missing value imputation for mixed-type data. *Bioinformatics*. DOI: 10.1093/bioinformatics/btr597.
- 2935
- 2936 Stephens, L. et al. (2019). Archaeological assessment reveals Earth's early transformation through land use. *Science*, 365.6456, 897 LP –902. DOI: 10.1126/science.aax1192.
- 2937
- 2938 Stewart, P. S., Voskamp, A., Santini, L., Biber, M. F., Devenish, A. J., Hof, C., Willis, S. G., and Tobias, J. A.
2939 (2022). Global impacts of climate change on avian functional diversity. *Ecology Letters*, 25.3, 673–685.
2940 DOI: 10.1111/ele.13830.
- 2941 Sunday, J. M., Bates, A. E., Kearney, M. R., Colwell, R. K., Dulvy, N. K., Longino, J. T., and Huey,
2942 R. B. (2014). Thermal-safety margins and the necessity of thermoregulatory behavior across latitude
2943 and elevation. *Proceedings of the National Academy of Sciences of the United States of America*. DOI:
2944 10.1073/pnas.1316145111.
- 2945 Sutherland, G. D., Harestad, A. S., Price, K., and Lertzman, K. P. (2000). Scaling of natal dispersal distances
2946 in terrestrial birds and mammals. *Ecology and Society*, 4.1. DOI: 10.5751/es-00184-040116.
- 2947 Thomas, C. D. (2013). The Anthropocene could raise biological diversity. *Nature*, 502.7469, 7. DOI: 10.
2948 1038/502007a.
- 2949 Thuiller, W., Lavorel, S., and Araújo, M. B. (2005). Niche properties and geographical extent as predictors
2950 of species sensitivity to climate change. *Global Ecology and Biogeography*, 14.4, 347–357. DOI: 10.
2951 1111/j.1466-822X.2005.00162.x.
- 2952 Tiffney, B. H. (2004). Vertebrate dispersal of seed plants through time. *Annual Review of Ecology, Evolution,
2953 and Systematics*, 35, 1–29. DOI: 10.1146/annurev.ecolsys.34.011802.132535.
- 2954 Tilman, D. and Downing, J. A. (1994). Biodiversity and stability in grasslands. *Nature*. DOI: 10.1038/
2955 367363a0.
- 2956 Tinoco, B. A., Santillán, V. E., and Graham, C. H. (2018). Land use change has stronger effects on functional
2957 diversity than taxonomic diversity in tropical Andean hummingbirds. *Ecology and Evolution*. DOI: 10.
2958 1002/ece3.3813.
- 2959 Titley, M. A., Snaddon, J. L., and Turner, E. C. (2017). Scientific research on animal biodiversity is sys-
2960 tematically biased towards vertebrates and temperate regions. *PLoS ONE*. DOI: 10.1371/journal.
2961 pone.0189577.

- 2962 Todd, B. D., Nowakowski, A. J., Rose, J. P., and Price, S. J. (2017). Species traits explaining sensitivity
2963 of snakes to human land use estimated from citizen science data. *Biological Conservation*. DOI: 10 .
2964 1016/j.biocon.2016.12.013.
- 2965 Tonini, J. F. R., Beard, K. H., Ferreira, R. B., Jetz, W., and Pyron, R. A. (2016). Fully-sampled phylogenies
2966 of squamates reveal evolutionary patterns in threat status. *Biological Conservation*. DOI: 10 .1016/j.
2967 biocon.2016.03.039.
- 2968 Trenberth, K. E. (2011). Changes in precipitation with climate change. *Climate Research*, 47.1-2, 123–138.
2969 DOI: 10 .3354/cr00953.
- 2970 Troyanskaya, O., Cantor, M., Sherlock, G., Brown, P., Hastie, T., Tibshirani, R., Botstein, D., and Altman,
2971 R. B. (2001). Missing value estimation methods for DNA microarrays. *Bioinformatics*. DOI: 10 .1093/
2972 bioinformatics/17.6.520.
- 2973 Tylianakis, J. M., Didham, R. K., Bascompte, J., and Wardle, D. A. (2008). Global change and species
2974 interactions in terrestrial ecosystems. *Ecology Letters*, 11.12, 1351–1363. DOI: 10 .1111/j.1461-
2975 0248.2008.01250.x.
- 2976 United Nations Educational Scientific and Cultural and Organization (2015). *UNESCO Global Science Re-
2977 port: Towards 2030*. Tech. rep.
- 2978 Urban, M. C. (2015). Accelerating extinction risk from climate change. *Science*, 348.6234, 571–573.
- 2979 Ustaoglu, E. and Williams, B. (2017). Determinants of Urban Expansion and Agricultural Land Conversion
2980 in 25 EU Countries. *Environmental Management*, 60.4, 717–746. DOI: 10 .1007/s00267-017-
2981 0908-2.
- 2982 Valiente-Banuet, A., Aizen, M. A., Alcántara, J. M., Arroyo, J., Cocucci, A., Galetti, M., García, M. B.,
2983 García, D., Gómez, J. M., Jordano, P., Medel, R., Navarro, L., Obeso, J. R., Oviedo, R., Ramírez, N., Rey,
2984 P. J., Traveset, A., Verdú, M., and Zamora, R. (2015). Beyond species loss: The extinction of ecological
2985 interactions in a changing world. *Functional Ecology*, 29.3, 299–307. DOI: 10 .1111/1365-2435.
2986 12356.
- 2987 Valipour, M., Bateni, S. M., and Jun, C. (2021). Global surface temperature: A new insight. *Climate*, 9.5,
2988 1–4. DOI: 10 .3390/cli9050081.
- 2989 Van Der Kaars, S., Miller, G. H., Turney, C. S., Cook, E. J., Nürnberg, D., Schönfeld, J., Kershaw, A. P.,
2990 and Lehman, S. J. (2017). Humans rather than climate the primary cause of Pleistocene megafaunal
2991 extinction in Australia. *Nature Communications*, 8, 1–7. DOI: 10 .1038/ncomms14142.
- 2992 van Rossum, G. (1995). *Python tutorial*. Centrum voor Wiskunde en Informatica (CWI).
- 2993 Verberk, W. C. E. P., Noordwijk, C. G. E. van, and Hildrew, A. G. (2013). Delivering on a promise: integrating
2994 species traits to transform descriptive community ecology into a predictive science. *Freshwater Science*.
2995 DOI: 10 .1899/12-092.1.
- 2996 Vidan, E., Roll, U., Bauer, A., Grismer, L., Guo, P., Maza, E., Novosolov, M., Sindaco, R., Wagner, P.,
2997 Belmaker, J., and Meiri, S. (2017). The Eurasian hot nightlife: Environmental forces associated with
2998 nocturnality in lizards. *Global Ecology and Biogeography*. DOI: 10 .1111/geb.12643.
- 2999 Villéger, S., Mason, N. W. H., and Mouillot, D. (2008). New multidimensional functional diversity indices
3000 for a multifaceted framework in functional ecology. *Ecology*.

- 3001 Violette, C., Navas, M. L., Vile, D., Kazakou, E., Fortunel, C., Hummel, I., and Garnier, E. (2007). Let the
3002 concept of trait be functional! *Oikos*. DOI: 10.1111/j.0030-1299.2007.15559.x.
- 3003 Watson, J. (2016). Bring climate change back from the future. *Nature*, 534.437.
- 3004 Weiner, C. N., Werner, M., Linsenmair, K. E., and Blüthgen, N. (2011). Land use intensity in grasslands:
3005 Changes in biodiversity, species composition and specialisation in flower visitor networks. *Basic and*
3006 *Applied Ecology*, 12.4, 292–299. DOI: 10.1016/j.baae.2010.08.006.
- 3007 Weiss, K. C. and Ray, C. A. (2019). Unifying functional trait approaches to understand the assemblage of
3008 ecological communities: synthesizing taxonomic divides. *Ecography*. DOI: 10.1111/ecog.04387.
- 3009 Wheatley, C. J., Beale, C. M., Bradbury, R. B., Pearce-Higgins, J. W., Critchlow, R., and Thomas, C. D.
3010 (2017). Climate change vulnerability for species—Assessing the assessments. *Global Change Biology*.
3011 DOI: 10.1111/gcb.13759.
- 3012 White, C. R. (2011). Allometric estimation of metabolic rates in animals. *Comparative Biochemistry and*
3013 *Physiology - A Molecular and Integrative Physiology*, 158.3, 346–357. DOI: 10.1016/j.cbpa.3014
2010.10.004.
- 3015 White, C. R. and Kearney, M. R. (2013). Determinants of inter-specific variation in basal metabolic rate.
3016 *Journal of Comparative Physiology B: Biochemical, Systemic, and Environmental Physiology*, 183.1,
3017 1–26. DOI: 10.1007/s00360-012-0676-5.
- 3018 White, C. R. and Seymour, R. S. (2003). Mammalian basal metabolic rate is proportional to body mass^{2/3}.
3019 *Proceedings of the National Academy of Sciences of the United States of America*, 100.7, 4046–4049.
3020 DOI: 10.1073/pnas.0436428100.
- 3021 Whitmee, S. and Orme, C. D. L. (2013). Predicting dispersal distance in mammals: A trait-based approach.
3022 *Journal of Animal Ecology*, 82.1, 211–221. DOI: 10.1111/j.1365-2656.2012.02030.x.
- 3023 Williams, J. J. and Newbold, T. (2020). Local climatic changes affect biodiversity responses to land use: A
3024 review. *Diversity and Distributions*. DOI: 10.1111/ddi.12999.
- 3025 Wilman, H., Belmaker, J., Simpson, J., Rosa, C. de la, Rivadeneira, M. M., and Jetz, W. (2014). EltonTraits
3026 1.0: Species-level foraging attributes of the world's birds and mammals. *Ecology*. DOI: 10.1890/13-
3027 1917.1.
- 3028 Wilson, E. E. and Wolkovich, E. M. (2011). *Scavenging: How carnivores and carrion structure communities*.
3029 DOI: 10.1016/j.tree.2010.12.011.
- 3030 Wong, M. K., Guénard, B., and Lewis, O. T. (2018). Trait-based ecology of terrestrial arthropods. *Biological*
3031 *Reviews*. DOI: 10.1111;brv.12488.
- 3032 Wulder, M. A., Coops, N. C., Roy, D. P., White, J. C., and Hermosilla, T. (2018). Land cover 2.0. *International*
3033 *Journal of Remote Sensing*, 39.12, 4254–4284. DOI: 10.1080/01431161.2018.1452075.
- 3034 WWF (2020). *Living Planet Report 2020 - Bending the curve of biodiversity loss*. Tech. rep. Almond, R.E.A.,
3035 Grooten M. and Petersen, T. (Eds). WWF, Gland, Switzerland., 1–9.
- 3036 Yan, Z., Teng, M., He, W., Liu, A., Li, Y., and Wang, P. (2019). Impervious surface area is a key predictor
3037 for urban plant diversity in a city undergone rapid urbanization. *Science of the Total Environment*, 650,
3038 335–342. DOI: 10.1016/j.scitotenv.2018.09.025.

- 3039 Young, H. S., McCauley, D. J., Galetti, M., and Dirzo, R. (2016). Patterns, Causes, and Consequences of
3040 Anthropocene Defaunation. *Annual Review of Ecology, Evolution, and Systematics*, 47. December, 333–
3041 358. DOI: 10.1146/annurev-ecolsys-112414-054142.
- 3042 Zamudio, K. R., Bell, R. C., and Mason, N. A. (2016). Phenotypes in phylogeography: Species' traits, en-
3043 vironmental variation, and vertebrate diversification. *Proceedings of the National Academy of Sciences*,
3044 113.29, 8041–8048. DOI: 10.1073/pnas.1602237113.
- 3045 Zhang, J., Qian, H., Girardello, M., Pellissier, V., Nielsen, S. E., and Svenning, J. C. (2018). Trophic inter-
3046 actions among vertebrate guilds and plants shape global patterns in species diversity. *Proceedings of the*
3047 *Royal Society B: Biological Sciences*. DOI: 10.1098/rspb.2018.0949.
- 3048 Zhang, Y., Yang, K., Yang, P., Su, Y., Zheng, W., and Liu, J. (2018). Food restriction decreases BMR, body
3049 and organ mass, and cellular energetics, in the Chinese Bulbul (*Pycnonotus sinensis*). *Avian Research*,
3050 9.1, 1–11. DOI: 10.1186/s40657-018-0131-8.
- 3051 Zuñiga-Palacios, J., Zuria, I., Castellanos, I., Lara, C., and Sánchez-Rojas, G. (2021). What do we know (and
3052 need to know) about the role of urban habitats as ecological traps? Systematic review and meta-analysis.
3053 *Science of the Total Environment*, 780. DOI: 10.1016/j.scitotenv.2021.146559.

Appendix 1: Supporting information for Chapter 2

S2.1 Taxonomic corrections

Across the different sources of trait data, similar species could appear under different binomial names. This was a problem when matching datasets by species. Moreover, it is possible that within a source, a given species was appearing under two or more different, synonymous names. As such, taxonomic synonymy created duplicated rows for the same species, overall falsely increasing the total number of species and potentially inflating the number of missing trait values. Taxonomic synonymy was hence a major issue. Due to the large number of species across datasets, extensive manual checks could not be applied. The presence of typos in species names had the same effect as synonymy, erroneously duplicating species. I attempted to correct for taxonomy first by correcting for typos, and second by identifying species which were entered under non-accepted names and replacing these with the accepted name. To this end, I developed an automated procedure, complemented with a few manual entries where errors were opportunistically spotted. Such errors in taxonomy were notably spotted when attempting to retrieve trait data for subsets of species, for analyses not directly related to the work conducted here. Taxonomic synonymy was as such checked manually for 91 species (56 birds, 7 mammals and 28 reptiles); in that case, information was extracted from other diverse sources (such as the Reptile Database (<http://www.reptile-database.org/>); Avibase (<https://avibase.bsc-eoc.org/avibase.jsp?lang=EN&pg=home>); AmphibiaWeb (<https://amphibiaweb.org/>); and additional manual checks using the IUCN Red List for mammals). A column in the Synonym dataset mentions where manual checks were applied (in which case the Synonym dataset was manually corrected).

Automated procedure and outputs

Extracting names from the IUCN Red List and the Integrated Taxonomic Information System (ITIS)

The objectives of the automated procedure were to (1) extract species synonymous binomial names from the

IUCN Red List or from the ITIS, using the ‘rredlist’ (Chamberlain, 2018) and ‘taxize’ (Chamberlain and Szöcs, 2013) R packages; and (2) identify the status of each name (accepted or not accepted). I started by generating a list of all names featuring in any of the sources. These ‘original’ names were corrected for typos (using the ‘gnr_resolve’ function in the ‘taxize’ package). Then, the IUCN Red List was queried and any listed synonyms were stored, as well as the status of each synonym (accepted or not accepted). When species were not found in the IUCN Red List, synonyms were extracted from the ITIS. When species were not found in the ITIS either, corrected names (original names corrected for typos) were used. Family and order designations were extracted using the same procedure and some entries were retrieved from the Global Biodiversity Information Facility taxonomic backbone when not available in the IUCN Red List or in the ITIS (GBIF, <https://www.gbif.org/tools/species-lookup>).

NB: for species entered with the forms *Genus cf.*, *Genus aff.* or *Genus spp.*, the accepted binomial name was left empty.

Output. I generated a ‘Synonym’ dataset containing records of binomial names (14,124 recorded names for birds; 8,743 for amphibians; 6,090 for mammals; and 11,678 for reptiles), and for each I recorded their status and their potential synonyms.

Harmonising taxonomy in trait datasets. Taxonomy across datasets was finally homogenised by replacing synonyms with a uniquely identified accepted name. As a consequence, the total number of identified unique species decreased (Figure S2.1a). The species presenting the highest number of synonyms was the East African mole rat (*Tachyoryctes splendens*), for which I found 12 synonymous names (Figure S2.1b).

The automated procedure was not perfect, and taxonomic errors are likely to have persisted in the trait datasets. The IUCN Red List and the ITIS were not comprehensive taxonomic sources, and for clades with high degrees of synonymy in names, such as reptiles or amphibians, neither the IUCN Red List or the ITIS contained enough information. Taxonomy may be further improved by using class-specific sources in an automated procedure.




Figure S2.1: Difference in species number before and after taxonomic correction (A) and distribution of number of synonyms across datasets (B). (A) shows the number of species (binomial names) extracted from all sources (blue bars), and the number of uniquely identified accepted names (in red). Replacing non-accepted synonyms by one identified accepted name reduced the number of species in all classes, with the largest reduction for birds. (B) shows the distribution of the number of synonymous names. In all four classes, more than 5,000 species were known under one name only. Nevertheless, a large number of species had two identified synonyms (range: 400 species for mammals - 2086 for birds). The most potentially replicated species was the East African mole rat *Tachyoryctes splendens*, for which 12 synonyms were identified.

S2.2 Additional information for trait compilation

Correlations among closely related traits




Figure S2.2: (a) Body mass versus body length and (b) longevity versus age at sexual maturity in amphibians. Pearson's correlation coefficient was 0.71 in (a) and 0.55 in (b) (taxonomic order was considered in these coefficients).




Figure S2.3: Generation length versus longevity data in (a) mammals and (b) birds. Pearson's correlation coefficient was 0.74 in (a) and 0.70 in (b).

S2.3 Cutting distribution maps by altitudinal limits




Figure S2.4: Availability of altitudinal limits across species. Upper and lower altitudinal limits were extracted from the IUCN Red List (IUCN, 2020).




Figure S2.5: Range sizes before versus after cutting the range maps by altitudinal limits.

S2.4 Impact of taxonomic corrections on trait coverage





Figure S2.6: Trait coverage when no taxonomic correction is applied before matching sources by binomial names, versus when I applied the described procedure. The identification of synonymous names resulted in an increase in trait coverage in most cases.

S2.5 Assemblage-level median, mean and standard deviation of trait completeness (maps)


(a) Mammals




Median completeness
(birds and mammals):

- NA
- 80 – 90%
- 90 – 91%
- 91 – 92%
- 92 – 93%
- 93 – 94%
- 94 – 95%
- 95 – 96%
- 96 – 97%
- 97 – 98%
- 98 – 99%
- 99 – 100%

(b) Birds



(c) Amphibians



Median completeness
(herptiles):

- NA
- 0 – 10%
- 10 – 20%
- 20 – 30%
- 30 – 40%
- 40 – 50%
- 50 – 60%
- 60 – 70%
- 70 – 80%
- 80 – 90%
- 90 – 100%

(d) Reptiles





Figure S2.7: Spatial distribution of assemblage-level median trait completeness in each class. Note that the color breaks differ for mammals and birds and for herptiles.


(a) Mammals




Mean completeness
(birds and mammals):

- NA
- 80 – 90%
- 90 – 91%
- 91 – 92%
- 92 – 93%
- 93 – 94%
- 94 – 95%
- 95 – 96%
- 96 – 97%
- 97 – 98%
- 98 – 99%
- 99 – 100%

(b) Birds



(c) Amphibians



Mean completeness
(herptiles):

- NA
- 0 – 10%
- 10 – 20%
- 20 – 30%
- 30 – 40%
- 40 – 50%
- 50 – 60%
- 60 – 70%
- 70 – 80%
- 80 – 90%
- 90 – 100%

(d) Reptiles





Figure S2.8: Spatial distribution of assemblage-level mean trait completeness in each class. Note that the color breaks differ for mammals and birds and for herptiles.


(a) Mammals




Standard deviation
of completeness
(birds and mammals):

- NA
- 0 – 5%
- 5 – 10%
- 10 – 15%
- 15 – 20%
- 20 – 25%
- 25 – 30%

(b) Birds



(c) Amphibians



Standard deviation
of completeness
(herptiles):

- NA
- 0 – 10%
- 10 – 15%
- 15 – 20%
- 20 – 25%
- 25 – 30%
- 30 – 35%
- 35 – 40%
- 40 – 50%
- 50 – 60%
- 60 – 75%

(d) Reptiles




Figure S2.9: Spatial distribution of assemblage-level standard deviation of trait completeness in each class. Note that the color breaks differ for mammals and birds and for herptiles.




Figure S2.10: Assemblage-level species richness against standard deviation in trait completeness in each class.

S2.6 Phylogenetic patterns in trait completeness




Figure S2.11: Within-family median trait completeness in mammals.





Figure S2.12: Within-family median trait completeness in birds.






Figure S2.14: Within-family species richness against the within-family standard deviation of trait completeness.

S2.7 Poisson model summaries

Table S2.1: Coefficients of the model investigating whether species range size explained the number of sampled traits. Class was added as an interacting predictor. The reference level for class is mammals. The model was fitted using a Poisson error distribution.

	Estimate	Std. Error	z value	Pr(> z)
Intercept	1.678	0.022	76.809	< 2e - 16
log Range Size	0.015	0.002	8.086	6.16e - 16
Class Birds	-0.092	0.028	-3.350	0.000809
Class Amphibians	-0.689	0.029	-24.099	< 2e - 16
Class Reptiles	-0.872	0.027	-31.856	< 2e - 16
log Range Size:Class Birds	0.003	0.002	1.415	0.157
log Range Size:Class Amphibians	0.017	0.003	6.427	1.30e - 10
log Range Size:Class Reptiles	0.026	0.002	11.159	< 2e - 16

Table S2.2: Coefficients of the model investigating whether species range size explained the number of sampled traits, using range maps not cut by altitudinal limits. Class was added as an interacting predictor. The reference level for class is mammals. The model was fitted using a Poisson error distribution.

	Estimate	Std. Error	z value	Pr(> z)
Intercept	1.665	0.023	72.070	< 2e - 16
log Range Size	0.015	0.002	8.167	3.16e - 16
Class Birds	-0.110	0.029	-3.763	0.0002
Class Amphibians	-0.700	0.030	-23.721	< 2e - 16
Class Reptiles	-0.928	0.029	-32.403	< 2e - 16
log Range Size:Class Birds	0.004	0.002	1.840	0.066
log Range Size:Class Amphibians	0.018	0.003	6.564	5.24e - 11
log Range Size:Class Reptiles	0.031	0.002	12.630	< 2e - 16




Figure S2.15: Relationship between number of sampled traits and geographical range size using distribution maps not cut by altitudinal limits. The model was fitted using a Poisson error distribution. Class was added as a predictor interacting with range size. Rates of increase were not significantly different for mammals and birds, but differed for reptiles and amphibians, with steeper rates of increase for reptiles overall.

S2.8 Spatial model summaries

Table S2.3: Spatial model summary for amphibians. The spatial model was fitted to explain assemblage-level median completeness with species richness. Biogeographic realm was added as an interacting explanatory variable.

	Estimate	Std. Error	z value	Pr(> z)
Intercept - Realm: Afrotropic	0.0738	0.0064	11.4908	0
log(Species richness)	-0.0025	0.0017	-1.4261	0.1538
Realm: Australasia	-0.0109	0.0095	-1.1453	0.2521
Realm: Indo-Malay	0.0455	0.0119	3.8294	0.0001
Realm: Nearctic	0.0441	0.0082	5.3905	0.000000
Realm: Neotropic	-0.0377	0.0083	-4.5538	0.00001
Realm: Palearctic	0.0047	0.0067	0.6992	0.4844
log(Species richness):Australasia	0.0018	0.0038	0.4789	0.6320
log(Species richness):Indo-Malay	-0.0147	0.0039	-3.7294	0.0002
log(Species richness):Nearctic	-0.0097	0.0030	-3.2003	0.0014
log(Species richness):Neotropic	0.0144	0.0026	5.6454	0.000000
log(Species richness):Palearctic	0.0109	0.0029	3.7358	0.0002

Table S2.4: Spatial model summary for reptiles. The spatial model was fitted to explain assemblage-level median completeness with species richness. Biogeographic realm was added as an interacting explanatory variable.

	Estimate	Std. Error	z value	Pr(> z)
Intercept - Realm: Afrotropic	0.2001	0.0144	13.9349	0
log(Species richness)	-0.0316	0.0031	-10.0547	0
Realm: Australasia	-0.1284	0.0189	-6.7851	0
Realm: Indo-Malay	-0.0453	0.0263	-1.7215	0.0852
Realm: Nearctic	-0.0788	0.0140	-5.6366	0.000000
Realm: Neotropic	-0.0932	0.0145	-6.4425	0
Realm: Palearctic	-0.1030	0.0131	-7.8787	0
log(Species richness):Australasia	0.0386	0.0046	8.4019	0
log(Species richness):Indo-Malay	0.0124	0.0061	2.0397	0.0414
log(Species richness):Nearctic	0.0346	0.0038	9.1601	0
log(Species richness):Neotropic	0.0220	0.0034	6.4231	0
log(Species richness):Palearctic	0.0286	0.0033	8.6153	0

S2.9 Trait coverage and taxonomic matching

Here, I briefly explore the robustness of my work to taxonomic uncertainty by comparing trait coverage obtained with the procedure I developed for taxonomic matching against trait coverage obtained when extracting synonyms from class-specific sources, which could potentially contain more taxonomic information than the ITIS or the IUCN Red List, notably for herptiles. I corrected the datasets for taxonomy again using the ‘rangeBuilder’ R package (Rabosky et al., 2016), which allows to extract accepted names from class-specific sources. Overall, my results are robust to the use of a different taxonomic backbone (Figure S2.16); the main conclusions of this work are likely unaffected by taxonomic uncertainty.




Figure S2.16: Comparison of trait coverage among datasets corrected for taxonomy in different ways: using the described procedure (purple bars); using datasets corrected for taxonomy with the ‘rangeBuilder’ package (yellow bars; Rabosky et al. (2016)); and using datasets uncorrected for taxonomy (green bars).

Appendix 2: Supporting information for Chapter 3

S3.1 Land-use categories in PREDICTS and sample sizes

Table S3.1: Land-use categories in the PREDICTS database. See Hudson et al. (2014, 2017) for more details.

Land-use category	Definition
Primary vegetation	Native vegetation, undisturbed since its establishment under current climatic conditions. No known alterations due to human activities or to extreme natural events.
Mature secondary vegetation	Vegetation recovering after complete destruction of primary vegetation & where succession is near complete – the structure approaches that of primary vegetation.
Intermediate secondary vegetation	Vegetation recovering after complete destruction of primary vegetation at a mid-successional stage.
Young secondary vegetation	Vegetation recovering after complete destruction of primary vegetation at an early successional stage.
Plantation forest	Previously cleared areas planted with crop trees or shrubs grown and harvested for human consumption or for commercial purposes (includes wood, fruit, oil, biofuel, rubber, etc).
Pasture	Areas grazed by livestock, permanently or regularly. Can be improved through cultivation techniques.
Cropland	Previously cleared areas planted with herbaceous crops and harvested for human or animal consumption (including animal feed and crops used in the food industry), or for commercial purposes (e.g., crops grown for the textile industry).
Urban	Previously cleared areas built up by humans. Vegetation is managed for civic or personal purposes.




Figure S3.1: Number of sites in each land use and land-use intensity for which FRic and FDis were calculated, across all species. The number of sites for FRic can be smaller than the number of sites for FDis because FRic could not necessarily be computed for all assemblages in which FDis was estimated (in assemblages where species richness was 2 or 3, FRic could not be computed).




Figure S3.2: Number of sites in each land use, land-use intensity and for which FRic and FDis were calculated, within each class. The number of sites for FRic can be smaller than the number of sites for FDis because FRic could not necessarily be computed for all assemblages in which FDis was estimated (in assemblages where species richness was 2 or 3, FRic could not be computed).

S3.2 Trait data & imputation of missing trait values

I used the compilation of trait data across terrestrial vertebrates from Chapter 2. I selected seven traits that were available at least for a subset of the species in each class: body mass, trophic level, lifespan, litter/clutch size, diel activity, habitat breadth and use of artificial habitats. These traits related to life-history, habitat specialisation and use of geographical space (for instance, habitat breadth is a significant predictor of geographical range size in all classes, Figure S3.3).




Figure S3.3: Relationship between habitat breadth and geographical range size across species in each class. The derivation of geographical range sizes is described in Chapter 2.

Trait-data coverage was highly variable among classes and traits, with important geographical and phylogenetic biases in trait data for reptiles and amphibians (Chapter 2; Figure S3.4, Figure S3.5). To obtain complete species-trait datasets, I imputed missing trait values. Further, in order to assess the sensitivity of the models to variation in imputed values, I imputed the missing trait values eight independent times. This allowed me to assess the congruence of the model predictions when using the different imputed trait datasets in the analyses.




Figure S3.4: Trait coverage for the vertebrate species sampled in the PREDICTS database. For a given trait, coverage is calculated as the percentage of species for which an estimate was available.




Figure S3.5: Distribution of trait completeness across the vertebrate species sampled in the PREDICTS database. For a given species, trait completeness is calculated as the proportion of traits for which an estimate was available. Dashed lines represent the median trait completeness.

S3.2.1 Choice of imputation technique

There exist several imputation techniques (Debastiani et al., 2021; Johnson et al., 2021; Penone et al., 2014), such as K-nearest neighbour (Troyanskaya et al., 2001), multivariate imputation by chained equations (Buuren and Oudshoorn, 2007), random forest algorithms (implementable in R with the ‘missforest’ function from the ‘missForest’ package; Stekhoven, 2016; Stekhoven and Bühlmann, 2012), and phylogenetic imputations (implementable in R with PhyloPars, Bruggeman et al., 2009). Penone et al. (2014) assessed the performance of these four techniques and showed that ‘missforest’ and ‘PhyloPars’ performed better when traits were phylogenetically conserved, and when the species phylogenetic position was included as a predictor of missing trait values. ‘PhyloPars’ can only handle continuous data, while ‘missforest’ is compatible with mixed-type (including categorical) data. When no phylogenetic information was included, ‘mice’ was found to be the best method, with fast imputations of mixed-type data (Penone et al., 2014). Therefore, to assess whether ‘missforest’ or ‘mice’ was more appropriate here, I measured the phylogenetic signal in the traits. For continuous traits, I used Pagel’s λ (Pagel, 1999), and for categorical traits I used Borges’ δ (Borges et al., 2018). Strong phylogenetic signals would indicate that traits are phylogenetically conserved, and hence ‘missforest’ would be the most suited approach for imputing missing trait values, with the inclusion of species’ phylogenetic positions as a predictor.

S3.2.2 Phylogenetic signal in traits

Across all classes, similar traits were used for calculating functional diversity metrics: body mass, litter/clutch size, lifespan (using different proxies in different vertebrate classes: generation length for birds and mammals, longevity for reptiles, and age at sexual maturity for amphibians), trophic level, diel activity, habitat breadth and use of artificial habitats. In addition, I included some class-specific traits for the imputations, as certain class-specific traits could be useful predictors of other traits (such as body length for instance in amphibians (Santini et al., 2018)). Table S3.2 details the traits that were included for the imputations in each class and the phylogenetic signal for each of these traits. Continuous traits were log-10 transformed before assessing Pagel’s λ to improve normality. Pagel’s λ was estimated using the ‘phylosig’ function of the ‘phytools’ package (Revell, 2012), and Borges’ δ was assessed using code provided by Borges et al. (2018), available at : https://github.com/mrborges23/delta_statistic. To test for the significance of δ , I generated null distributions of δ for each categorical trait by randomising trait vectors 50 times, and calculating δ for each randomised vector – following the guidelines proposed by Borges et al. (2018). I then tested whether the observed medians were greater than the null distributions using one-sided Wilcoxon rank sum tests.

I used class-specific phylogenies to estimate phylogenetic signal, all downloaded on 13th April 2020. Trees from Faurby et al. (2018, 2020) were used for mammals (downloaded from <https://zenodo.org/record/3690867#.Xyc5wyhKhPZ>). For amphibians, birds and reptiles (squamates only), I downloaded trees from <https://data.vertlife.org/>. Trees were from Jetz et al. (2012) for birds, from Jetz and Pyron (2018) for amphibians and from Tonini et al. (2016) for squamates. For each class, I downloaded a distribution of 1,000 trees, from which I obtained consensus trees to estimate phylogenetic signal (to that end, I used the TreeAnnotator programme of the BEAST software (Bouckaert et al., 2014)).

Table S3.2: Phylogenetic signal in continuous and categorical traits. BM: body mass; BL: body length; GL: generation length; MA: age at sexual maturity; ML: maximum longevity; L: longevity; LCS: litter/clutch size; HB: habitat breadth; TL: trophic level; DA: diel activity; UA: use of artificial habitats. Continuous traits were log-10 transformed to improve normality before estimating Pagel's λ – except for habitat breadth which was square-rooted. A star indicates a significant signal (p-value<0.05 for the log-likelihood ratio test in the case of λ ; and a significant difference from the simulated null distribution of δ for categorical traits). 'NA' indicates traits that were not considered for a given class. All traits showed significant phylogenetic signal, with signals for BM, BL, L, GL, MA and LCS being particularly strong (above 0.8) across the four classes.

Class	Pagel's λ								Borges' δ		
	BM	BL	GL	MA	ML	L	LCS	HB	TL	DA	UA
Amphibians	0.98*	0.94*	NA	0.85*	0.82*	NA	0.93*	0.99*	18*	3.4*	4.5*
Birds	0.99*	NA	0.97*	NA	NA	NA	0.95*	0.60*	13*	32e3*	1.8*
Mammals	0.99*	NA	0.97	NA	NA	NA	0.99	0.71	26*	17*	1.3*
Reptiles	1.0*	NA	NA	NA	0.94*	0.98*	1.0*	0.52*	6.3*	6.4*	1.4*

S3.2.3 Implementation of ‘missforest’ imputations

As phylogenetic signals were strong in many categorical and continuous traits (Table S3.2), I imputed missing trait values using random forest algorithms, as implemented in R with ‘missforest’ from the ‘missForest’ package (Stekhoven, 2016; Stekhoven and Bühlmann, 2012). Another advantage of ‘missforest’ was that, being a non-parametric approach, no prior assumption about data distribution was required. The data were therefore not transformed prior to imputations. In addition, Penone et al. (2014) showed that including phylogenetic information did not decrease the accuracy of imputations for traits that were less phylogenetically conserved, such as habitat breadth in this work.

Phylogenetic relationships were included as additional predictors in the form of phylogenetic eigenvectors (Diniz-Filho et al., 2012), extracted from the phylogenies using the ‘PVR’ package (Santos, 2018). Following Penone et al. (2014), I included the first ten phylogenetic eigenvectors as additional predictors of missing trait values in each class, enough to minimise imputation error. As not all species were represented in the phylogenies, I also added taxonomic order as a predictor for all species. All traits in Table S3.2 were included in the imputations. Tuning parameters of ‘missforest’ were set to ten maximum iterations and to

one hundred trees grown in each forest.

S3.3 Degree of multicollinearity among traits

Multicollinearity among traits can be problematic when calculating functional diversity indices (Cadotte et al., 2011). After imputing missing trait values and before estimating functional metrics, I assessed whether the degree of multicollinearity among categorical and continuous traits was problematically high. To this end, I used generalised variance inflation factors (Fox and Monette, 1992). Given a regression model, variance inflation factors quantify the overestimation in the variance of estimated regression coefficients due to multicollinearity among the predictors. A GVIF value of 5 or 10 is commonly used as a threshold to select out collinear predictors (Dormann et al., 2013). I used the ‘stepwise.vif’ function of the ‘Rnalytica’ package (<https://github.com/awsm-research/Rnalytica>), with a threshold of 5, to determine the GVIF of each trait. I used the imputed traits from the 8th imputation iteration to assess whether multicollinearity was problematically high. Continuous traits were log-10 transformed (except for habitat breadth which was square-rooted). The degree of multicollinearity among traits was not detected to be problematically high, as all traits had a GVIF value below 2 (Table S3.3). As such, all seven traits were included in the calculation of the functional diversity indices.

Table S3.3: Variance inflation factors across considered (imputed) traits.

Trait	GVIF
Diel activity	1.1
Trophic level	1.3
Use of artificial habitats	1.4
Body mass (log10)	1.5
Habitat breadth (square-root)	1.5
Litter/clutch size (log10)	1.6
Lifespan proxy (log10)	1.7

S3.4 Imputation performance




Figure S3.6: Distribution of continuous traits considered in the calculation of the functional diversity metrics (shown as density plots), before and after missing value imputations, in each class and for the species occurring in the PREDICTS database. All traits were log10-transformed, except habitat breadth, which was square-rooted.




Figure S3.7: Frequency distribution for categorical traits considered in the calculation of the functional diversity metrics (shown as % of total species in each category) before and after missing value imputations, in each class, for the species occurring in the PREDICTS database.




Figure S3.8: Out-of-bag imputation errors for the continuous traits (a) and categorical traits (b). For continuous traits, the Normalised Root Mean Squared Error (NRMSE) was obtained by dividing the Mean Squared Error (MSE) by the variance of the known trait distribution, then square-rooting the result. The MSE was returned for each trait by the ‘missForest’ function (‘missForest’ package, Stekhoven, 2016; Stekhoven and Bühlmann, 2012) and corresponds to an out-of-bag error. For categorical traits, the error was estimated as the out-of-bag proportion of falsely classified traits.

S3.5 Functional loss and functional gain

Across all vertebrates, I estimated functional loss and functional gain using 84 studies for the tropical subset and 39 studies for the temperate subset (51,514 and 30,470 pairwise comparisons between sites respectively, Table S3.4). Because of this large number of pairwise comparisons, I did not develop a null modelling approach (if I used 100 randomisations per pair of sites, I would need to compute functional loss and functional gain for more than 8 million pairs, which would be very computationally demanding). I grouped mature, intermediate and young secondary vegetation together in this analysis. I could not estimate the effects in all land uses (for instance, sample sizes for tropical urban sites were too small; Figure S3.1).

Within classes, I used 18 tropical studies and 1 temperate study for amphibians; 38 tropical studies and 21 temperate studies for birds; 28 tropical studies and 9 temperate studies for mammals; and 11 tropical studies and 7 temperate studies for reptiles. As sample sizes differed among pairs of land uses and land-use intensities (Table S3.5), I was not able to estimate all effects, notably for the intensely-used land uses.

To calculate functional loss and functional gain, the Gower distance matrix (obtained from the global species by trait dataset for the species occurring in PREDICTS) was first subsetted to the species occurring in a given pair of sites. Cailliez corrections were applied when the distance matrix was not Euclidian (Cailliez corrections consist of applying the smallest positive constant to the distances so as to make them Euclidian (Cailliez, 1983); ‘ade4’ R package (Dray et al., 2007)). I then performed a principal coordinates analysis on the (corrected) Gower distance matrix, retaining the first two axes to reduce the computational load in the calculation of convex hulls. Sites that contained fewer than three functionally different species were excluded (the computation of a convex hull requiring more species in the assemblage than PCoA axes). Then I estimated the hypervolume of trait space occupied by each assemblage of a given pair, as well as the volume of the shared trait space (intersection), from which I derived functional loss and functional gain.

Table S3.4: Sample sizes (number of pairs of sites) for the calculation of functional loss and functional gain across all species.

Region	Pairs	Minimal use	Light use	Intense use
Temperate	PV-PV	7626	22546	492
	PV-SV	511	72	–
	PV-PF	9	166	–
	PV-PA	8	40	–
	PV-CR	150	–	–
	PV-UR	6306	1197	7
Tropical	PV-PV	8547	4016	16722
	PV-SV	6584	1124	9713
	PV-PF	580	1378	–
	PV-PA	36	20	22
	PV-CR	1700	1088	–
	PV-UR	–	–	–

Table S3.5: Sample sizes (number of pairs of sites) for the calculation of functional loss and functional gain within each class.

Class	Region	Pair of land uses	Minimal use	Light use	Intense use
Amphibians	Temperate	PV/PV	–	45	–
Amphibians	Temperate	PV/SV	8	70	–
Amphibians	Temperate	PV/AGR	3	–	–
Amphibians	Temperate	PV/UR	6	100	–
Amphibians	Tropical	PV/PV	501	241	307
Amphibians	Tropical	PV/SV	838	–	90
Amphibians	Tropical	PV/PF	422	91	–
Amphibians	Tropical	PV/AGR	1	3	1
Birds	Temperate	PV/PV	7,382	19,300	491
Birds	Temperate	PV/SV	150	1	–
Birds	Temperate	PV/PF	9	166	–
Birds	Temperate	PV/AGR	145	40	–
Birds	Temperate	PV/UR	6,300	992	–
Birds	Tropical	PV/PV	5,059	3,117	9,014
Birds	Tropical	PV/SV	3,491	1,058	5,225
Birds	Tropical	PV/PF	156	994	–
Birds	Tropical	PV/AGR	1,626	1,085	–
Mammals	Temperate	PV/PV	110	3,030	–
Mammals	Temperate	PV/SV	25	–	–
Mammals	Temperate	PV/AGR	5	–	–
Mammals	Temperate	PV/UR	–	105	7
Mammals	Tropical	PV/PV	1,989	637	64
Mammals	Tropical	PV/SV	230	65	8
Mammals	Tropical	PV/PF	2	–	–
Mammals	Tropical	PV/AGR	109	20	21
Reptiles	Temperate	PV/PV	132	2	1
Reptiles	Temperate	PV/SV	250	1	–
Reptiles	Temperate	PV/AGR	5	–	–
Reptiles	Tropical	PV/PV	989	137	5,140
Reptiles	Tropical	PV/SV	1,760	1	3,456
Reptiles	Tropical	PV/PF	–	190	–

S3.6 Diagnostic plots




Figure S3.9: Diagnostic plots for Model 1a, obtained using the 'performance' R package (Lüdecke et al., 2021).




Figure S3.10: Diagnostic plots for **Model 1b**, obtained using the ‘performance’ R package (Lüdecke et al., 2021).




Figure S3.11: Diagnostic plots for Model 2a, obtained using the ‘performance’ R package (Lüdecke et al., 2021).




Figure S3.12: Diagnostic plots for Model 2b, obtained using the ‘performance’ R package (Lüdecke et al., 2021).




Figure S3.13: Diagnostic plots for Model 3, obtained using the ‘DHARMA’ R package (Hartig, 2021).




Figure S3.14: Diagnostic plots for Model 4a, obtained using the ‘performance’ R package (Lüdecke et al., 2021).




Figure S3.15: Diagnostic plots for Model 4b, obtained using the ‘performance’ R package (Lüdecke et al., 2021).




Figure S3.16: Diagnostic plots for Model 5a, obtained using the ‘performance’ R package (Lüdecke et al., 2021).




Figure S3.17: Diagnostic plots for **Model 5b**, obtained using the ‘performance’ R package (Lüdecke et al., 2021).

S3.7 Model robustness




Figure S3.18: Effects of land use, land-use intensity and region on FRic (a) and FDis (b) across vertebrates, for the subset of species with complete trait data (i.e., excluding species with less than 100% trait completeness). Effects are plotted as a % difference relative to the reference land use (primary vegetation, PV). For FRic, I fitted **Model 1a**, which included the effects of land use, land-use intensity and region, as well as interactions between land use and land-use intensity and between land use and region. For FDis, I fitted **Model 1b**, which included effects of land use, land-use intensity and region, and interactions between land use and land-use intensity. Error bars represent 95% confidence intervals. MSV: mature secondary vegetation; ISV: intermediate secondary vegetation; YSV: young secondary vegetation; PF: plantation forest; PA: pasture; CR: cropland; UR: urban. Effects for Intense use in MSV could not be estimated as there weren't enough sampled sites.




Figure S3.19: Effects of land use, land-use intensity and region on FRic (a) and FDis (b), for the subset of species with complete trait data (i.e., excluding species with less than 100% trait completeness), with geographical range size as an additional trait considered in the calculation of functional diversity metrics. Effects are plotted as a % difference relative to the reference land use (primary vegetation, PV). For FRic, I fitted **Model 1a**, which included the effects of land use, land-use intensity and region, as well as interactions between land use and land-use intensity and between land use and region. For FDis, I fitted **Model 1b**, which included effects of land use, land-use intensity and region, and interactions between land use and land-use intensity. Error bars represent 95% confidence intervals. MSV: mature secondary vegetation; ISV: intermediate secondary vegetation; YSV: young secondary vegetation; PF: plantation forest; PA: pasture; CR: cropland; UR: urban. Effects for Intense use in MSV could not be estimated as there weren't enough sampled sites.




Figure S3.20: Effects of land use and land-use intensity on FRic (a) and FDis (b), obtained when calculating FRic and FDis with each set of imputed traits (eight in total). Effects are plotted as a % difference relative to the reference land use (primary vegetation, PV). For FRic, I fitted **Model 1a, which included the effects of land use, land-use intensity and region, as well as interactions between land use and land-use intensity and between land use and region. For FDis, I fitted **Model 1b**, which included effects of land use, land-use intensity and region, and interactions between land use and land-use intensity. Error bars represent 95% confidence intervals. MSV: mature secondary vegetation; ISV: intermediate secondary vegetation; YSV: young secondary vegetation; PF: plantation forest; PA: pasture; CR: cropland; UR: urban. Effects for Intense use in MSV could not be estimated as there weren't enough sampled sites.**




Figure S3.21: Effects of land use and land-use intensity on FRic (a) and FDis (b), obtained when re-sampling primary vegetation sites twenty independent times. I fixed the sample size for primary vegetation sites at 50. For FRic, we fitted **Model 1a**, which included the effects of land use, land-use intensity and region, as well as interactions between land use and land-use intensity and between land use and region. For FDis, I fitted **Model 1b**, which included effects of land use, land-use intensity and region, and interactions between land use and land-use intensity. Error bars represent 95% confidence intervals. MSV: mature secondary vegetation; ISV: intermediate secondary vegetation; YSV: young secondary vegetation; PF: plantation forest; PA: pasture; CR: cropland; UR: urban. Effects for Intense use in MSV could not be estimated as there weren't enough sampled sites.





Figure S3.22: Effects of land use, land-use intensity and taxonomic class on FRic (a) and FDis (c), for the subset of species with complete trait data (i.e., excluding species with less than 100% trait completeness). Effects are plotted as a % difference relative to the reference land use (primary vegetation, PV). I did not include the effects of region here as sample sizes were not large enough for some classes. For FRic, the model included the effects of land use, land-use intensity and class, and interactions between land use and land-use intensity as well as land use and class. For FDis, the model included an additional interaction between land-use intensity and class. Error bars represent 95% confidence intervals. SV: secondary vegetation; PF: plantation forest; PA: pasture; CR: cropland; UR: urban. Effects for reptiles in urban land uses could not be estimated as there weren't enough sampled sites.

— complete and imputed trait data — complete trait data subset Use intensity • Minimal use ▲ Light use * Intense use

(a) FRic



(b) FDis




Figure S3.23: Effects of land use, region, land-use intensity and taxonomic class on FRic and FDis obtained with the imputed trait data (black points) or with the complete data subsets (red points). Effects are plotted as a % difference relative to the reference land use (primary vegetation, PV). For FRic, I fitted Model 2a, and I fitted Model 2b for FDis. Error bars represent 95% confidence intervals. SV: secondary vegetation; PF: plantation forest; AGR: agricultural (cropland and pasture); UR: urban. Effects for reptiles in urban land uses could not be estimated as there weren't enough sampled sites.




Figure S3.24: Effects of land use, region, land-use intensity and taxonomic class on FRic and FDis, obtained when calculating FRic and FDis with each set of imputed traits (eight in total). Effects are plotted as a % difference relative to the reference land use (primary vegetation, PV). For FRic, I fitted Model 2a, and Model 2b for FDis. Error bars represent 95% confidence intervals. SV: secondary vegetation; PF: plantation forest; AGR: agricultural (cropland and pasture); UR: urban. Effects for reptiles in urban land uses could not be estimated as there weren't enough sampled sites.

S3.8 Model robustness – time since land-use conversion

Time since land-use conversion could have important impacts on assemblage composition and thus, on local functional diversity. I did not investigate these effects because PREDICTS contained data on time since land-use conversion only for about 22% of the sites, considerably reducing samples sizes. Here, I investigated whether the results are likely robust to the inclusion of time since land-use conversion using the subset of sites for which time since land-use conversion was provided. To this end, I found the best-fitting models explaining FRic and FDis, using backwards stepwise selection, starting with complete models that included the effects of land use, time since land-use conversion, region, land-use intensity (for FRic only) and all two-way interactions among these predictors.

- For FRic, the best-fitting model included the main effects of land use and time since land-use conversion, but no interaction between these predictors. The model's summary (Table S3.6) showed that time since conversion had a significant negative effect on FRic, but the relationship between FRic and time since land-use conversion was similar in different land uses (as there were no interactions between land use and time since conversion retained in the best-fitting model, such that the slopes were similar in different land uses, and so the rate at which FRic decreased with time was similar in different land uses). The intercept was only different for urban land uses (significantly lower). Thus, based on this data subset, I expect time since land-use conversion to have a similar effect in different land-use types.

Table S3.6: Summary of the model explaining FRic by land use and time since land-use conversion, fitted on the subset of data for which there are information on time since land-use conversion.

	Estimate	Std. Error	t value
Intercept: Primary vegetation	1.156	0.073	15.921
Mature secondary vegetation	0.178	0.093	1.907
Intermediate secondary vegetation	0.018	0.072	0.249
Young secondary vegetation	-0.078	0.051	-1.532
Plantation forest	-0.018	0.082	-0.224
Pasture	-0.005	0.093	-0.054
Cropland	0.133	0.152	0.875
Urban	-0.316	0.133	-2.368
Time since conversion (log years)	-0.094	0.021	-4.566

I then compared this model's predictions with a simpler model that did not account for time since land-use conversion ($\text{FRic} \sim \text{Land use}$). The predictions (Figure S3.25) showed that including time since land-use

conversion did not bias the results, as I found a similar significant effect with both models in urban land uses, and elsewhere the effects were congruent. Thus, given this data subset, I argue that the results are robust to the inclusion of time since land-use conversion.




Figure S3.25: Effects of land use on FRic for the model that includes time since land-use conversion (blue points) versus the model that does not take time since land-use conversion into account (red points).

- For FDis, the best-fitting model included the main effects of land use, time since land-use conversion as well as interactions between land use and time since land-use conversion (I did not consider land-use intensity in the starting model because of sample size issues). Nevertheless, the main effect of time since land-use conversion was not significant (Table S3.7), and the relationship between time since land-use conversion and FDis was also not significant in most disturbed land uses (except for plantation forest). Thus, I argue the available data do not allow us to properly investigate the relationship between time since land-use conversion and FDis.

Table S3.7: Summary of the model explaining FDis by land use and time since land-use conversion, fitted on the subset of data for which I have information on time since land-use conversion.

	Estimate	Std. Error	t value
Intercept: Primary vegetation	0.366	0.011	32.219
Mature secondary vegetation	0.032	0.055	0.577
Intermediate secondary vegetation	-0.015	0.050	-0.298
Young secondary vegetation	0.020	0.015	1.386
Plantation forest	0.074	0.023	3.213
Pasture	-0.017	0.048	-0.346
Cropland	-0.013	0.042	-0.317
Urban	0.031	0.054	0.573
Time since conversion (years, log)	-0.004	0.004	-1.186
Mature secondary vegetation:Time since conversion (log years)	-0.005	0.015	-0.335
Intermediate secondary vegetation:Time since conversion (log years)	0.011	0.016	0.650
Young secondary vegetation:Time since conversion (log years)	-0.008	0.007	-1.170
Plantation forest:Time since conversion (log years)	-0.023	0.007	-3.077
Pasture:Time since conversion (log years)	0.010	0.015	0.688
Cropland:Time since conversion (log years)	0.007	0.012	0.620
Urban:Time since conversion (log years)	-0.016	0.022	-0.714

Appendix 3: Supporting information for Chapter 4

S4.1 Compiling diet information

For mammals and birds, diet information was obtained from the EltonTraits database (Wilman et al., 2014). Before processing the data, the taxonomy was aligned to that of the trait datasets complied in Chapter 2. Primary diet – that is, the diet inferred from the combination of food items that represent more than 50% of species consumption – was directly available for birds, but not for mammals. For both classes, diet was described as the percent use of different food items (namely: invertebrates, vertebrates – either ectotherms, endotherms, fish or unknown –, carrion, fruit, nectar, seed or other plant material). In order to have a consistent classification scheme across mammals and birds, I chose not to use the provided primary diet for birds, and instead I applied my own procedure to infer primary diet from recorded food items across birds and mammals. I first grouped the different vertebrate food items together with carrion to create a single ‘vertebrate’ food item category. I then used the percent uses of the food items to infer primary diet, classifying species’ primary diet into the following categories: vertebrate eaters; invertebrate eaters; plant/seed eaters; fruit/nectar eaters; and omnivores [these categories are similar to those employed for birds’ primary diet in EltonTraits]. When all food items had a percent use below (or equal to) 50% percent, species were classified as omnivores. For each species, I calculated diet breadth as the number of consumed food items (regardless of the percent use of those items; I kept vertebrate food items grouped together in the calculation of diet breadth, such that I did not count carrion as a separate food item).

For amphibians, diet information was partly extracted from the AmphiBIO database (Oliveira et al., 2017), and partly compiled from the literature (see [next section](#)). In AmphiBIO, diet information was recorded as the consumption of six food items (leaves, flowers, seeds, fruit, arthropods and vertebrates), but the percent use of these items was not recorded (only whether they were consumed or not). From AmphiBIO, I classified amphibians into the different diet categories, [depending on the combinations of consumed food](#)

items.

To increase diet data coverage for amphibians, I compiled data from published papers and from the grey literature, targeting species occurring in the PREDICTS database (which I used for inferring land-use responses). I was able to collect diet information for an additional 108 amphibians from 26 published sources (all found to be invertebrate eaters; see below for the list of sources). For reptiles, there was no readily available diet information (except for trophic level information, see Etard et al. (2020)). Thus, I collected diet data from the literature, again specifically targeting reptiles occurring in the PREDICTS database. From the literature, I added diet information for 239 reptiles. Finally, diet breadth was calculated across amphibians and reptiles from the recorded food items. The compiled diet data are available at: https://figshare.com/articles/Reptile_Diet_csv/12024309 (DOI: 10.6084/m9.figshare.12024309.v1) and https://figshare.com/articles/Untitled_Item/12024312 (DOI: 10.6084/m9.figshare.12024312.v4).

Complementary data sources for amphibians

- Aguilar-López & Pineda (2013). An exotic species of earthworm preyed by *Craugastor rhodopis*. Herpetology Notes, 6: 335-336.
- Arroyo, S. B., Serrano-Cardozo, V. H., & Ramírez-Pinilla, M. P. (2008). Diet, microhabitat and time of activity in a Pristimantis (Anura, Strabomantidae) assemblage. Phylomedusa: Journal of Herpetology, 7(2), 109-119.
- Loc Barragán, J. A. & Woolrich, G. (2016). *Smilisca baudinii*. Diet. Mesoamerican Herpetology, 3(3).
- Berry, P. Y., & Bullock, J. A. (1962). The Food of the Common Malayan Toad, *Bufo melanostictus* Schneider. Copeia, 1962(4), 736–741.
- Blommers-Schlösser, R. M. A. (1975). A unique case of mating behaviour in a Malagasy tree frog, *Gephyromantis liber* (Peracca, 1893), with observations on the larval development (Amphibia, Ranidae). Beaufortia, 23(296), 15–25.
- Cappo, M. (1986). Frogs as predators of organisms of aquatic origin in the Magela Creek System, Northern Territory. MSc thesis. Dept of Zoology, University of Adelaide. <https://hdl.handle.net/2440/110846>.
- Costello, J.M. (2013). Differences in Morphology and Behavior in Green Frogs (*Lithobates clamitans*) from Urban and Rural Sites in New York and New Jersey. PhD thesis. City University of New York.

- Brito de Carvalho, C. & Freitas, E. & Faria, R. & Batista, R. & Batista, C. & Coelho, W. & Bocchiglieri, A. (2008). Natural history of *Leptodactylus mystacinus* and *Leptodactylus fuscus* (Anura: Leptodactylidae) in the Cerrado of Central Brazil. *Biota Neotropica*, 8.
- Batista, R. & Brito de Carvalho, C. & Freitas, E.B. & Franco, S.C. & Batista, C.C. & Coelho, Wellington & Faria, Renato. (2011). Diet of *Rhinella schneideri* (Werner, 1894) (Anura: Bufonidae) in the Cerrado, Central Brazil. *Herpetology Notes*, 4:17-21.
- Fonseca-Pérez, K. A., Molina, C., & Tárano, Z. (2017). Diet of *Dendropsophus microcephalus* and *Scarthyla vigilans* (Anura: Hylidae) at a locality in north-western Venezuela with notes on microhabitat occupation. *Papéis Avulsos De Zoologia*, 57(7), 93-104.
- García-R, J.C. & Posso-Gómez, C. E. & Cárdenas-Henao, H. (2015). Diet of direct-developing frogs (Anura: Craugastoridae: Pristimantis) from the Andes of western Colombia. *Acta Biológica Colombiana*, 20:79-87.
- Zumbado-Ulato, H. & Acosta-Chaves, V. (2012). *Cochranella granulosa* (Granular glass frog) feeding behaviour. *Herpetological review*.
- Yap, C.H. & Jafaar, I. (2012). Feeding analysis of *Hylarana cf. labialis*, *Leptobrchiun hendricksoni*, and *Occidozyga laevis* (Amphibia: Anura) from a lowland dipterocarp forest in Kedah, Malaysia. *Herpetological Review*, 43(1).
- Schwenk, K. (2000). Feeding: Form, function and evolution in tetrapod vertebrates. Academic Press.
- Sabagh, L.T., Ferreira V.L., & Rocha C.F. (2010). Living together, sometimes feeding in a similar way: the case of the syntopic hylid frogs *Hypsiboas raniceps* and *Scinax acuminatus* (Anura: Hylidae) in the Pantanal of Miranda, Mato Grosso do Sul State, Brazil. *Brazilian Journal of Biology*, 70(4):955-9.
- Luría-Manzano R. & Ramírez-Bautista A. (2017). Diet comparison between rainforest and cave populations of *Craugastor alfredi* (Anura: Craugastoridae): does diet vary in contrasting habitats? *Journal of Natural History*, 51:39-40, 2345-2354.
- Martínez-Coronel, M. & Pérez-Gutiérrez, M. (2011). Diet composition of *Craugastor lineatus* (Anura: Craugastoridae) of Chiapas, Mexico. *Acta Zoológica Mexicana* 27(2):215-230.
- McAlister, W. H. (1963). Evidence of mild toxicity in the saliva of the hognose snake (Heterodon). *Herpetologica*, 19:132-137.

- Moreno-Barbosa, S. & Hoyos, J. (2014). Ontogeny of the diet in Anurans (Amphibia) collected at La Vieja river basin in the departamento of Quindio (Colombia). *Caldasia*, 36 (365-372).
- Piatti, L. & Souza, F. (2011). Diet and resource partitioning among anurans in irrigated rice fields in Pantanal, Brazil. *Brazilian journal of biology*, 71 (653-61).
- Savini, C. O. K., Chuang, M. F., & Ishida, C. (2004). Diet Selection in the Green Paddy Frog (*Rana erythraea*). International field biology course 2004.
- Simon, M. P., & Toft, C. A. (1991). Diet Specialization in Small Vertebrates: Mite-Eating in Frogs. *Oikos*, 61(2):263–278.
- Sugai, J.L.M.M., Terra, J.S., & Ferreira, V.L. (2012): Diet of *Leptodactylus fuscus* (Amphibia: Anura: Leptodactylidae) in the Pantanal of Miranda river, Brazil. *Biota Neotropica*, 12: 99-104.
- Toft, C. A. (1981). Feeding Ecology of Panamanian Litter Anurans: Patterns in Diet and Foraging Mode. *Journal of Herpetology*, 15(2):139–144.
- Vences, M., Glaw, F. & Zapp, C.(1999). Stomach content analysis in Malagasy frogs of the genera *Tomopterna*, *Aglyptodactylus*, *Boophis* and *Mantidactylus*. *Herpetozoa*, 11(3/4):109-116.
- Yap, C.H. & Jaafar, I. (2011). Stomach Content Analysis of Tropical Forest Toads *Ingerophrynus parvus* and *Phrynooides aspera* (ANURA : BUFONIDAE) from Kedah, Malaysia. Taxonomist and Ecologist Conference 2011.

Complementary data sources for reptiles

The 148 sources are listed in the dataset available from https://figshare.com/articles/Reptile_Diet_csv/12024309 (DOI: 10.6084/m9.figshare.12024309.v1).

S4.2 Imputing missing trait values

S4.2.1 Trait data coverage




Figure S4.1: Trait coverage, including coverage for diet information, calculated as the proportion of species for which trait values are not missing. The dashed line represents 50% coverage.

S4.2.2 Phylogenetic signal in traits

I measured the phylogenetic signal in traits using Pagel's λ (for continuous traits) and Borges' δ (for categorical traits). I found evidence of phylogenetic conservatism in all traits.

Table S4.1: Phylogenetic signal in continuous and categorical traits. BM: body mass; BL: body length; GL: generation length; MA: age at sexual maturity; ML: maximum longevity; L: longevity; LCS: litter/clutch size; HB: habitat breadth; DA: diel activity; UA: use of artificial habitats. Continuous traits were log-10 transformed to improve normality before estimating Pagel's λ – except for habitat breadth which was square-rooted. A star indicates a significant signal ($P<0.05$ for the log-likelihood ratio test in the case of λ ; and a significant difference from the simulated null distribution of δ for categorical traits). ‘NA’ indicates traits that were not considered for a given class.

Class	Pagel's λ										Borges' δ		
	BM	BL	GL	MA	ML	L	LCS	HB	Diet breadth	Diet	DA	UA	
Amphibians	0.98*	0.94*	NA	0.85*	0.82*	NA	0.93*	0.99*	0.61*	3.4*	3.4*	4.5*	
Birds	0.99*	NA	0.97*	NA	NA	NA	0.95*	0.60*	0.72*	6.4*	32e3*	1.8*	
Mammals	0.99*	NA	0.97	NA	NA	NA	0.99	0.71		26*	52*	1.3*	
Reptiles	1.0*	NA	NA	NA	0.94*	0.98*	1.0*	0.52*	0.84*	2.2*	6.4*	1.4*	

S4.2.3 Implementation of missing value imputations

I imputed missing trait values using random forest algorithms, as implemented in R with missForest (Stekhoven, 2016; Stekhoven and Bühlmann, 2012). Phylogenetic relationships were included as additional predictors in the form of phylogenetic eigenvectors (Diniz-Filho et al., 2012), extracted from the phylogenies using the PVR package (Santos, 2018). I used class-specific phylogenetic trees from <https://zenodo.org/record/3690867#.Xyc5wyhKhPZ> for mammals (Phylacine 1.2; Faurby et al. (2018, 2020)); and from <https://data.vertlife.org/> for amphibians (Jetz & Pyron 2018), birds (Jetz et al. 2012) and squamates (Tonini et al. 2016). For each class, I used a consensus tree obtained with the TreeAnnotator programme of the BEAST software (Bouckaert et al. 2014), from an available distribution of 1000 trees.

Following Penone et al. (2014), I included the first ten phylogenetic eigenvectors as additional predictors of missing trait values in each class. As not all species were represented in the phylogenies, I also added taxonomic order as a predictor for all species. All traits in Table S4.1 were included in the imputations. Habitat & diet breadth were considered as categorical variables for the imputations (and so, discretised), in order to ensure that only integer estimates were obtained from the imputations for these traits. Tuning parameters of ‘missforest’ were set to ten maximum iterations and to one hundred trees grown in each forest.

S4.2.4 Imputation error




Figure S4.2: Out-of-bag estimation of imputation errors for the traits included in the analyses. (a) For the categorical traits, I show the proportion of falsely classified traits ('PFC', out-of-bag estimates); (b) For the continuous traits, I calculate the normalised root-mean-squared error (NRMSE), from the out-of-bag mean square error that I divide by the standard deviation of the known trait distribution. The lower the NRMSE, the lower the imputation error, with values close to 0 indicating low imputation error and values close to 1 tending to indicate high imputation error.

S4.3 Land-use types in PREDICTS and sample sizes (number of sampled sites across classes)

Table S4.2: Land-use categories in the PREDICTS database. See Hudson et al. (2014, 2017) for more details.

Land-use category	Definition
Primary vegetation	Native vegetation, undisturbed since its establishment under current climatic conditions. No known alterations due to human activities or to extreme natural events.
Mature secondary vegetation	Vegetation recovering after complete destruction of primary vegetation & where succession is near complete – the structure approaches that of primary vegetation.
Intermediate secondary vegetation	Vegetation recovering after complete destruction of primary vegetation at a mid-successional stage.
Young secondary vegetation	Vegetation recovering after complete destruction of primary vegetation at an early successional stage.
Plantation forest	Previously cleared areas planted with crop trees or shrubs grown and harvested for human consumption or for commercial purposes (includes wood, fruit, oil, biofuel, rubber, etc).
Pasture	Areas grazed by livestock, permanently or regularly. Can be improved through cultivation techniques.
Cropland	Previously cleared areas planted with herbaceous crops and harvested for human or animal consumption (including animal feed and crops used in the food industry), or for commercial purposes (e.g., crops grown for the textile industry).
Urban	Previously cleared areas built up by humans. Vegetation is managed for civic or personal purposes.




Figure S4.3: Sample sizes (number of PREDICTS sites) for the different land-use types, in each class.

S4.4 Land-use responses: multicollinearity checks among the models' explanatory variables

Table S4.3: Generalised Variance Inflation Factors among the candidate explanatory variables for the mixed-effects model fitted across amphibians, prior to the exclusion of diet. The model aimed at investigating the effects of land use, land-use intensity and the species-level ecological characteristics on species occurrence probability.

Predictor	GVIF
Diel activity	1.7
Lifespan proxy (\log_{10})	1.8
Specialisation	1.8
Range area (\log_{10})	2.0
Body mass (\log_{10})	2.0
Land use	2.0
Litter/clutch size (\log_{10})	2.5
Land-use intensity	2.6
Habitat breadth (square-root)	3.2
Diet breadth (square-root)	22.8
Primary diet	23.6

Table S4.4: Generalised Variance Inflation Factors among the explanatory variables for the mixed-effects model fitted across amphibians (after the exclusion of diet), looking at the effects of land use, land-use intensity and the species-level ecological characteristics on species occurrence probability.

Predictor	GVIF
Diel activity	1.6
Diet breadth (sqrt)	1.7
Lifespan proxy (\log_{10})	1.8
Specialisation	1.8
Range area (\log_{10})	1.9
Land use	2.0
Body mass (\log_{10})	2.0
Litter/clutch size (\log_{10})	2.4
Land-use intensity	2.6
Habitat breadth (square-root)	3.1

Table S4.5: Generalised Variance Inflation Factors among the explanatory variables for the mixed-effects model fitted across birds, looking at the effects of land use, land-use intensity and the species-level ecological characteristics on species occurrence probability.

Predictor	GVIF
Diel activity	1.1
Land use	1.2
Land-use intensity	1.2
Litter/clutch size (\log_{10})	1.3
Range area (\log_{10})	1.4
Diet breadth (square-root)	1.5
Specialisation	1.6
Lifespan proxy (\log_{10})	1.7
Habitat breadth (square-root)	1.8
Body mass (\log_{10})	1.9
Primary diet	2.3

Table S4.6: Generalised Variance Inflation Factors among the explanatory variables for the mixed-effects model fitted across mammals, looking at the effects of land use, land-use intensity and the species-level ecological characteristics on species occurrence probability.

Predictor	GVIF
Diel activity	1.2
Range area (\log_{10})	1.2
Specialisation	1.4
Land-use intensity	1.4
Diet breadth (square-root)	1.7
Land use	1.8
Habitat breadth (square-root)	1.8
Litter/clutch size (\log_{10})	2.7
Body mass (\log_{10})	3.0
Lifespan proxy (\log_{10})	3.4
Primary diet	4.4

Table S4.7: Generalised Variance Inflation Factors among the candidate explanatory variables for the mixed-effects model fitted across reptiles, prior to the exclusion of diet. The model aimed at investigating the effects of land use, land-use intensity and the species-level ecological characteristics on species occurrence probability.

Predictor	GVIF
Diel activity	1.1
Specialisation	1.3
Range area (\log_{10})	1.3
Habitat breadth (square-root)	1.6
Lifespan proxy (\log_{10})	1.9
Litter/clutch size (\log_{10})	2.8
Land use	3.2
Land-use intensity	3.5
Body mass (\log_{10})	3.9
Diet breadth (square-root)	5.8
Primary diet	9.9

Table S4.8: Generalised Variance Inflation Factors among the explanatory variables for the mixed-effects model fitted across reptiles (*after the exclusion of diet*), looking at the effects of land use, land-use intensity and the species-level ecological characteristics on species occurrence probability.

Predictor	GVIF
Diel activity	1.1
Diet breadth (square-root)	1.2
Specialisation	1.2
Range area (\log_{10})	1.3
Habitat breadth (square-root)	1.6
Lifespan proxy (\log_{10})	1.9
Litter/clutch size (\log_{10})	2.7
Land use	3.2
Body mass (\log_{10})	3.2
Land-use intensity	3.4

S4.5 Implementing Climate-niche Factor Analysis across terrestrial vertebrates

S4.5.1 Historical climate data: groups of intercorrelated variables




Figure S4.4: Groups of intercorrelated climatic variables **using a cutoff of 0.65 for Pearson's correlation coefficient**, obtained using the ‘removeCollinearity’ R function (‘virtualspecies’ package, Leroy et al. (2015)).

S4.5.2 CENFA estimation and resolution

I estimated climate-change sensitivity across terrestrial vertebrates with the CENFA framework using three different resolutions for the species distribution files and the climatic variables: 50 km², 10 km² and 5 km². Indeed, the finer the resolution, the better **species distribution** is likely to be captured, particularly for narrow-ranging species (Figure S4.5). When working with coarser resolutions, the actual geographical distribution of a narrow-ranging species might be overestimated (Figure S4.5), such that the climatic niche breadth of the species might also be overestimated, and consequently its climate-change sensitivity **might be** underestimated. However, finer resolutions are more computationally demanding, which can be limiting when working across several thousand species.

Thus, I looked for a resolution that provided the best estimations of climate-change sensitivity for the narrow-ranging species while requiring acceptable computational load. With a resolution of 50 km^2 , climate-change sensitivity tended to be overestimated for a larger number of narrow-ranging species than at 10 km^2 (Figure S4.6); and at 10 km^2 , climate-change sensitivity tended to be overestimated for a larger number of narrow-ranging species than at 5 km^2 (Figure S4.6). Below 5 km^2 , I deemed the computational load not acceptable for a global estimation of climate-change sensitivity across terrestrial vertebrates.

Hence, I chose to work with a resolution of 5 km^2 . At this resolution, there were still some narrow-ranging species for which sensitivity was likely overestimated (Figure S4.6). To prevent any impact of these species on the analyses, I removed species with the smallest geographical range areas, using a conservative threshold of 100 km^2 for geographical range area (Figure S4.6).




Figure S4.5: Possible impact of resolution on estimated geographical range area. I represent a virtual distribution for a species (orange shape). The species distribution is more accurately captured at finer resolutions (e.g., 5 km^2) than at coarser resolutions (e.g., 10 km^2 or 50 km^2). A possible consequence is that coarser resolutions can tend to disproportionately overestimate the geographical range area of narrow-ranging species, because the aggregation of grid cells where the species is found to be present can artificially augment the amount of occupied area at coarser resolutions, and relatively more so if the species is narrow-ranging.




Figure S4.6: Estimated climate-change sensitivity estimations at three different resolutions (50 km^2 , 10 km^2 and 5 km^2) against geographical range area (estimated at 1 km^2). With the red dashed lines, I highlight the range areas that correspond to the surface area of one grid cell (i.e., $2,500 \text{ km}^2$, 100 km^2 and 25 km^2 respectively). Climate-change sensitivity was estimated using the CENFA framework (Rinnan and Lawler, 2019). I chose to work at a resolution of 5 km^2 and I excluded species whose range area was $\leq 100 \text{ km}^2$ (blue dashed line), that is, species whose distribution could intersect up to four grid cells at a resolution of 5 km^2 .

S4.6 Climate-change sensitivity models: multicollinearity checks among models' explanatory variables

Table S4.9: Generalised Variance Inflation Factors among the candidate explanatory variables for the phylogenetic least-square regression fitted across amphibians, prior to the exclusion of diet breadth. The model aimed at investigating the association between the species-level ecological characteristics and species climate-change sensitivity.

Predictor	GVIF
Diel activity	1.1
Lifespan proxy (\log_{10})	1.2
Range area (\log_{10})	1.3
Body mass (\log_{10})	1.4
Litter/clutch size (\log_{10})	1.5
Specialisation	1.6
Habitat breadth (square-root)	1.9
Primary diet	17.1
Diet breadth (square-root)	17.1

Table S4.10: Generalised Variance Inflation Factors among the explanatory variables for the phylogenetic least-square regression fitted across amphibians, after excluding diet breadth. The model aimed at investigating the association between the species-level ecological characteristics and species climate-change sensitivity.

Predictor	GVIF
Diel activity	1.1
Primary diet	1.1
Lifespan proxy (\log_{10})	1.2
Range area (\log_{10})	1.3
Body mass (\log_{10})	1.4
Litter/clutch size (\log_{10})	1.5
Specialisation	1.6
Habitat breadth (square-root)	1.9

Table S4.11: Generalised Variance Inflation Factors among the explanatory variables for the phylogenetic least-square regression fitted across birds. The model aimed at investigating the association between the species-level ecological characteristics and species climate-change sensitivity.

Predictor	GVIF
Diel activity	1.1
Range area (\log_{10})	1.2
Litter/clutch size (\log_{10})	1.3
Diet breadth (square-root)	1.5
Specialisation	1.6
Habitat breadth (square-root)	1.8
Lifespan proxy (\log_{10})	1.9
Body mass (\log_{10})	2.0
Primary diet	2.1

Table S4.12: Generalised Variance Inflation Factors among the explanatory variables for the phylogenetic least-square regression fitted across mammals. The model aimed at investigating the association between the species-level ecological characteristics and species climate-change sensitivity.

Predictor	GVIF
Range area (\log_{10})	1.2
Diel activity	1.3
Specialisation	1.3
Habitat breadth (square-root)	1.5
Diet breadth (square-root)	1.6
Body mass(\log_{10})	2.3
Litter/clutch size (\log_{10})	2.4
Primary diet	2.7
Lifespan proxy (\log_{10})	3.0

Table S4.13: Generalised Variance Inflation Factors among the explanatory variables for the phylogenetic least-square regression fitted across reptiles. The model aimed at investigating the association between the species-level ecological characteristics and species climate-change sensitivity.

Predictor	GVIF
Diel activity	1.1
Range area(\log_{10})	1.2
Specialisation	1.4
Habitat breadth (square-root)	1.5
Lifespan proxy (\log_{10})	1.6
Litter/clutch size (\log_{10})	2.0
Body mass (\log_{10})	2.9
Diet breadth (square-root)	2.9
Primary diet	3.6

S4.7 Land-use responses: occurrence probability predictions from the partial models for artificial habitat use and diel activity




Figure S4.7: Predicted occurrence probability as a function of land use, land-use intensity, artificial habitat use and their interactions, for each class of terrestrial vertebrates (median $\pm 95\%$ confidence interval; the predictions are rescaled with reference to minimally-used primary vegetation). The predictions were obtained from the partial models fitted in each class for artificial habitat use. Effects could not be estimated for urban reptiles, as there weren't any sampled sites. Primary: primary vegetation; Secondary: secondary vegetation; plantation: plantation forest; agricultural: cropland and pasture.




Figure S4.8: Predicted occurrence probability as a function of land use, land-use intensity, diel activity and their interactions, for each class of terrestrial vertebrates (median $\pm 95\%$ confidence interval; the predictions are rescaled with reference to minimally-used primary vegetation). The predictions were obtained from the partial models fitted in each class for diel activity. Effects could not be estimated for urban reptiles, as there weren't any sampled sites. Error bars are large for non-nocturnal urban mammals because there were very few sampled species (only five). Primary: primary vegetation; Secondary: secondary vegetation; plantation: plantation forest; agricultural: cropland and pasture.

S4.8 Land-use responses: diagnostic plots for the full models




Figure S4.9: Diagnostic plots for the mixed-effects model fitted on amphibians, looking at the effects of land use, land-use intensity and the species-level ecological characteristics on species occurrence probability (full model). The diagnostic plots were obtained from the ‘DHARMA’ R package (Hartig, 2021).




Figure S4.10: Diagnostic plots for the mixed-effects model fitted on birds, looking at the effects of land use, land-use intensity and the species-level ecological characteristics on species occurrence probability (full model). The diagnostic plots were obtained from the ‘DHARMA’ R package (Hartig, 2021).




Figure S4.11: Diagnostic plots for the mixed-effects model fitted on mammals, looking at the effects of land use, land-use intensity and the species-level ecological characteristics on species occurrence probability (full model). The diagnostic plots were obtained from the ‘DHARMA’ R package (Hartig, 2021).




Figure S4.12: Diagnostic plots for the mixed-effects model fitted on reptiles, looking at the effects of land use, land-use intensity and the species-level ecological characteristics on species occurrence probability (full model). The diagnostic plots were obtained from the 'DHARMA' R package (Hartig, 2021).

S4.9 Climate-change sensitivity: model summaries and diagnostic plots

S4.9.1 Summaries & diagnostic plots for models fitted on species with range area $>100 \text{ km}^2$

Table S4.14: Summary for the PGLS model fitted on amphibians, investigating associations between species-level ecological characteristics and climate-change sensitivity, excluding species whose range area was $\leq 100 \text{ km}^2$ (sample size: n=4,537).

	Estimate	Std. Error	t value	Pr(> t)
Intercept	1.15	0.21	5.49	< 0.001
$\log_{10}(\text{Body mass})$	-1.97	0.61	-3.22	0.001
$\log_{10}(\text{Body mass})^2$	-0.26	0.42	-0.60	0.55
$\log_{10}(\text{Body mass})^3$	0.46	0.37	1.24	0.22
$\log_{10}(\text{Lifespan proxy})$	-0.21	0.59	-0.36	0.72
$\log_{10}(\text{Lifespan proxy})^2$	-0.14	0.43	-0.32	0.75
$\log_{10}(\text{Lifespan proxy})^3$	0.58	0.35	1.66	0.10
$\log_{10}(\text{Litter/clutch size})$	1.59	0.54	2.96	0.003
$\log_{10}(\text{Litter/clutch size})^2$	-0.06	0.38	-0.16	0.87
$\log_{10}(\text{Litter/clutch size})^3$	-0.74	0.31	-2.37	0.02
$\log_{10}(\text{Range area})$	-26.60	0.34	-77.15	< 0.001
$\log_{10}(\text{Range area})^2$	4.27	0.29	14.57	< 0.001
$\log_{10}(\text{Range area})^3$	-1.65	0.28	-5.96	< 0.001
square-root(Habitat breadth)	-2.26	0.43	-5.32	< 0.001
square-root(Habitat breadth) ²	0.81	0.30	2.67	0.01
square-root(Habitat breadth) ³	-0.59	0.28	-2.10	0.04
Specialisation: Natural habitat specialist	0.02	0.01	1.85	0.06
Diel activity: Non-nocturnal	0.04	0.01	3.28	0.001
Primary diet: Omnivore	0.01	0.03	0.29	0.77
Primary diet: Plants/seeds	0.04	0.13	0.31	0.76
Primary diet: Vertebrates	0.13	0.15	0.87	0.39




Figure S4.13: Diagnostic plots for the PGLS model fitted on amphibians, investigating associations between species-level ecological characteristics and climate-change sensitivity, excluding species whose range area was $\leq 100 \text{ km}^2$ (sample size: n=4,537).

Table S4.15: Summary for the PGLS model fitted on birds, investigating associations between species-level ecological characteristics and climate-change sensitivity, excluding species whose range area was $\leq 100 \text{ km}^2$ (sample size: n=10,198).

	Estimate	Std. Error	t value	Pr(> t)
Intercept	0.64	0.08	7.91	< 0.001
$\log_{10}(\text{Body mass})$	2.24	0.67	3.36	0.001
$\log_{10}(\text{Body mass})^2$	0.16	0.42	0.37	0.71
$\log_{10}(\text{Body mass})^3$	-0.22	0.37	-0.60	0.55
$\log_{10}(\text{Lifespan proxy})$	-0.23	0.59	-0.38	0.70
$\log_{10}(\text{Lifespan proxy})^2$	-0.84	0.39	-2.16	0.03
$\log_{10}(\text{Lifespan proxy})^3$	-0.10	0.28	-0.36	0.72
$\log_{10}(\text{Litter/clutch size})$	3.72	0.39	9.46	< 0.001
$\log_{10}(\text{Litter/clutch size})^2$	-0.42	0.33	-1.26	0.21
$\log_{10}(\text{Litter/clutch size})^3$	-0.40	0.27	-1.47	0.14
$\log_{10}(\text{Range area})$	-30.69	0.27	-113.09	< 0.001
$\log_{10}(\text{Range area})^2$	7.22	0.24	29.92	< 0.001
$\log_{10}(\text{Range area})^3$	-2.73	0.23	-11.74	< 0.001
square-root(Habitat breadth)	0.86	0.33	2.59	0.01
square-root(Habitat breadth) ²	-0.89	0.24	-3.63	< 0.001
square-root(Habitat breadth) ³	-0.22	0.23	-0.95	0.34
square-root(Diet breadth)	-0.50	0.31	-1.64	0.10
square-root(Diet breadth) ²	-0.11	0.25	-0.44	0.66
square-root(Diet breadth) ³	0.32	0.24	1.36	0.18
Specialisation: Natural habitat specialist	0.06	0.01	10.13	< 0.001
Diel activity: Non-nocturnal	-0.02	0.04	-0.66	0.51
Primary diet: Invertebrates	0.06	0.01	5.53	< 0.001
Primary diet: Omnivores	0.02	0.01	2.09	0.04
Primary diet: Plants/seeds	0.06	0.01	4.69	< 0.001
Primary diet: Vertebrates	0.01	0.02	0.83	0.41




Figure S4.14: Diagnostic plots for the PGLS model fitted on birds, investigating associations between species-level ecological characteristics and climate-change sensitivity, excluding species whose range area was $\leq 100 \text{ km}^2$ (sample size: n=10,198).

Table S4.16: Summary for the PGLS model fitted on mammals, investigating associations between species-level ecological characteristics and climate-change sensitivity, excluding species whose range area was $\leq 100 \text{ km}^2$ (sample size: n=4,712).

	Estimate	Std. Error	t value	Pr(> t)
Intercept	0.84	0.16	5.37	< 0.001
$\log_{10}(\text{Body mass})$	-4.62	0.94	-4.93	< 0.001
$\log_{10}(\text{Body mass})^2$	0.40	0.56	0.72	0.47
$\log_{10}(\text{Body mass})^3$	0.59	0.44	1.33	0.18
$\log_{10}(\text{Lifespan proxy})$	1.60	1.03	1.55	0.12
$\log_{10}(\text{Lifespan proxy})^2$	-0.79	0.49	-1.60	0.11
$\log_{10}(\text{Lifespan proxy})^3$	-0.15	0.43	-0.35	0.73
$\log_{10}(\text{Litter/clutch size})$	3.29	0.71	4.63	< 0.001
$\log_{10}(\text{Litter/clutch size})^2$	0.06	0.42	0.14	0.89
$\log_{10}(\text{Litter/clutch size})^3$	-0.16	0.33	-0.47	0.64
$\log_{10}(\text{Range area})$	-24.17	0.31	-78.21	< 0.001
$\log_{10}(\text{Range area})^2$	4.15	0.28	15.09	< 0.001
$\log_{10}(\text{Range area})^3$	-0.90	0.26	-3.45	0.001
square-root(Habitat breadth)	-1.24	0.34	-3.60	< 0.001
square-root(Habitat breadth) ²	0.22	0.27	0.82	0.41
square-root(Habitat breadth) ³	-0.03	0.26	-0.10	0.92
square-root(Diet breadth)	-1.21	0.47	-2.55	0.01
square-root(Diet breadth) ²	0.33	0.36	0.91	0.36
square-root(Diet breadth) ³	0.11	0.34	0.33	0.74
Specialisation: Natural habitat specialist	0.04	0.01	3.22	0.001
Diel activity: Non-nocturnal	0.003	0.01	0.23	0.82
Primary diet: Invertebrates	-0.02	0.03	-0.62	0.54
Primary diet: Omnivores	-0.02	0.03	-0.77	0.44
Primary diet: Plants/seeds	0.03	0.02	1.32	0.19
Primary diet: Vertebrates	-0.04	0.04	-0.98	0.33




Figure S4.15: Diagnostic plots for the PGLS model fitted on mammals, investigating associations between species-level ecological characteristics and climate-change sensitivity, excluding species whose range area was $\leq 100 \text{ km}^2$ (sample size: n=4,712).

Table S4.17: Summary for the PGLS model fitted on reptiles, looking at the effects of species-level ecological characteristics on climate-change sensitivity, excluding species whose range area was $\leq 100 \text{ km}^2$ (sample size: n=7,330).

	Estimate	Std. Error	t value	Pr(> t)
Intercept	1.02	0.13	7.70	< 0.001
$\log_{10}(\text{Body mass})$	-4.62	0.62	-7.47	< 0.001
$\log_{10}(\text{Body mass})^2$	0.97	0.41	2.35	0.02
$\log_{10}(\text{Body mass})^3$	-0.96	0.33	-2.90	0.004
$\log_{10}(\text{Lifespan proxy})$	0.89	0.51	1.74	0.08
$\log_{10}(\text{Lifespan proxy})^2$	-1.27	0.39	-3.27	0.001
$\log_{10}(\text{Lifespan proxy})^3$	-0.29	0.33	-0.88	0.38
$\log_{10}(\text{Litter/clutch size})$	0.84	0.68	1.24	0.22
$\log_{10}(\text{Litter/clutch size})^2$	-0.34	0.45	-0.76	0.45
$\log_{10}(\text{Litter/clutch size})^3$	0.26	0.35	0.74	0.46
$\log_{10}(\text{Range area})$	-31.88	0.31	-101.48	< 0.001
$\log_{10}(\text{Range area})^2$	5.15	0.27	19.03	< 0.001
$\log_{10}(\text{Range area})^3$	-1.97	0.26	-7.53	< 0.001
square-root(Habitat breadth)	0.56	0.35	1.59	0.11
square-root(Habitat breadth) ²	-0.67	0.27	-2.49	0.01
square-root(Habitat breadth) ³	0.05	0.26	0.21	0.83
square-root(Diet breadth)	-0.33	0.37	-0.90	0.37
square-root(Diet breadth) ²	-0.02	0.27	-0.08	0.94
square-root(Diet breadth) ³	0.31	0.26	1.22	0.22
Specialisation: Natural habitat specialist	0.10	0.01	8.50	< 0.001
Diel activity: Non-nocturnal	-0.02	0.01	-1.88	0.06
Primary diet: Omnivore	-0.03	0.03	-0.92	0.36
Primary diet: Vertebrates	-0.02	0.02	-1.05	0.30




Figure S4.16: Diagnostic plots for the PGLS model fitted on reptiles, investigating associations between species-level ecological characteristics and climate-change sensitivity, excluding species whose range area was $\leq 100 \text{ km}^2$ (sample size: n=7,330).

S4.9.2 Summaries for the PGLS models fitted on all species (including those with range area $\leq 100 \text{ km}^2$)

Table S4.18: Summary for the PGLS model fitted on amphibians, investigating associations between species-level ecological characteristics and climate-change sensitivity, including species whose range area was $\leq 100 \text{ km}^2$ (sample size: n=5,197).

	Estimate	Std. Error	t value	Pr(> t)
Intercept	1.23	0.20	6.01	< 0.001
$\log_{10}(\text{Body mass})$	-1.70	0.74	-2.29	0.02
$\log_{10}(\text{Body mass})^2$	-0.56	0.52	-1.09	0.28
$\log_{10}(\text{Body mass})^3$	0.12	0.44	0.27	0.78
$\log_{10}(\text{Lifespan proxy})$	0.46	0.72	0.64	0.52
$\log_{10}(\text{Lifespan proxy})^2$	0.21	0.53	0.40	0.69
$\log_{10}(\text{Lifespan proxy})^3$	0.50	0.43	1.17	0.24
$\log_{10}(\text{Litter/clutch size})$	0.95	0.64	1.48	0.14
$\log_{10}(\text{Litter/clutch size})^2$	-0.34	0.45	-0.74	0.46
$\log_{10}(\text{Litter/clutch size})^3$	-1.07	0.38	-2.79	0.01
$\log_{10}(\text{Range area})$	-28.60	0.42	-67.90	< 0.001
$\log_{10}(\text{Range area})^2$	-3.57	0.37	-9.76	< 0.001
$\log_{10}(\text{Range area})^3$	8.78	0.34	25.56	< 0.001
square-root(Habitat breadth)	-2.55	0.52	-4.89	< 0.001
square-root(Habitat breadth) ²	0.46	0.38	1.21	0.23
square-root(Habitat breadth) ³	-1.26	0.35	-3.62	< 0.001
Specialisation: Natural habitat specialist	0.02	0.01	1.55	0.12
Diel activity: Non-nocturnal	0.03	0.01	2.46	0.01
Primary diet: Omnivore	0.001	0.04	0.02	0.99
Primary diet: Plants/seeds	-0.12	0.15	-0.80	0.42
Primary diet: Vertebrates	0.20	0.18	1.14	0.26

Table S4.19: Summary for the PGLS model fitted on birds, investigating associations between species-level ecological characteristics and climate-change sensitivity, including species whose range area was $\leq 100 \text{ km}^2$ (sample size: n=10,340).

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	0.64	0.08	7.62	< 0.001
$\log_{10}(\text{Body mass})$	2.07	0.70	2.94	0.003
$\log_{10}(\text{Body mass})^2$	0.26	0.45	0.57	0.57
$\log_{10}(\text{Body mass})^3$	-0.15	0.39	-0.39	0.70
$\log_{10}(\text{Lifespan proxy})$	-0.47	0.62	-0.76	0.45
$\log_{10}(\text{Lifespan proxy})^2$	-0.67	0.41	-1.64	0.10
$\log_{10}(\text{Lifespan proxy})^3$	-0.07	0.29	-0.24	0.81
$\log_{10}(\text{Litter/clutch size})$	3.25	0.41	7.84	< 0.001
$\log_{10}(\text{Litter/clutch size})^2$	-0.64	0.35	-1.82	0.07
$\log_{10}(\text{Litter/clutch size})^3$	-0.30	0.29	-1.04	0.30
$\log_{10}(\text{Range area})$	-33.48	0.29	-116.96	< 0.001
$\log_{10}(\text{Range area})^2$	6.56	0.25	25.72	< 0.001
$\log_{10}(\text{Range area})^3$	1.56	0.24	6.39	< 0.001
square-root(Habitat breadth)	1.01	0.35	2.88	0.004
square-root(Habitat breadth) ²	-1.22	0.26	-4.75	< 0.001
square-root(Habitat breadth) ³	-0.38	0.24	-1.59	0.11
square-root(Diet breadth)	-0.44	0.32	-1.38	0.17
square-root(Diet breadth) ²	-0.27	0.26	-1.04	0.30
square-root(Diet breadth) ³	0.43	0.25	1.73	0.08
Specialisation: Natural habitat specialist	0.06	0.01	9.75	< 0.001
Diel activity: Non-nocturnal	-0.01	0.04	-0.35	0.73
Primary diet: Invertebrates	0.06	0.01	6.03	< 0.001
Primary diet: Omnivore	0.03	0.01	2.61	0.01
Primary diet: Plants/seeds	0.07	0.01	5.13	< 0.001
Primary diet: Vertebrates	0.02	0.02	0.85	0.39

Table S4.20: Summary for the PGLS model fitted on mammals, investigating associations between species-level ecological characteristics and climate-change sensitivity, including species whose range area was $\leq 100 \text{ km}^2$ (sample size: n=4,841).

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	0.86	0.15	5.68	< 0.001
$\log_{10}(\text{Body mass})$	-4.57	0.97	-4.69	< 0.001
$\log_{10}(\text{Body mass})^2$	0.37	0.59	0.62	0.53
$\log_{10}(\text{Body mass})^3$	0.67	0.47	1.44	0.15
$\log_{10}(\text{Lifespan proxy})$	1.50	1.09	1.38	0.17
$\log_{10}(\text{Lifespan proxy})^2$	-0.93	0.52	-1.79	0.07
$\log_{10}(\text{Lifespan proxy})^3$	-0.05	0.46	-0.11	0.91
$\log_{10}(\text{Litter/clutch size})$	3.28	0.75	4.35	< 0.001
$\log_{10}(\text{Litter/clutch size})^2$	0.19	0.44	0.43	0.67
$\log_{10}(\text{Litter/clutch size})^3$	-0.35	0.35	-0.99	0.32
$\log_{10}(\text{Range area})$	-26.32	0.34	-78.41	< 0.001
$\log_{10}(\text{Range area})^2$	2.16	0.30	7.26	< 0.001
$\log_{10}(\text{Range area})^3$	3.80	0.29	13.28	< 0.001
square-root(Habitat breadth)	-1.24	0.37	-3.31	0.001
square-root(Habitat breadth) ²	0.10	0.29	0.34	0.73
square-root(Habitat breadth) ³	-0.15	0.28	-0.55	0.58
square-root(Diet breadth)	-1.57	0.50	-3.12	0.002
square-root(Diet breadth) ²	0.34	0.38	0.88	0.38
square-root(Diet breadth) ³	-0.03	0.36	-0.09	0.93
Specialisation: Natural habitat specialist	0.04	0.01	3.55	< 0.001
Diel activity: Non-nocturnal	-0.001	0.02	-0.07	0.95
Primary diet: Invertebrates	-0.03	0.03	-0.97	0.33
Primary diet: Omnivore	-0.03	0.03	-0.96	0.34
Primary diet: Plants/seeds	0.02	0.03	0.83	0.41
Primary diet: Vertebrates	-0.06	0.04	-1.59	0.11

Table S4.21: Summary for the PGLS model fitted on reptiles, investigating associations between species-level ecological characteristics and climate-change sensitivity, including species whose range area was $\leq 100 \text{ km}^2$ (sample size: n=7,945).

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	1.08	0.14	7.79	< 0.001
$\log_{10}(\text{Body mass})$	-4.04	0.72	-5.64	< 0.001
$\log_{10}(\text{Body mass})^2$	0.22	0.48	0.46	0.64
$\log_{10}(\text{Body mass})^3$	-0.84	0.39	-2.19	0.03
$\log_{10}(\text{Lifespan proxy})$	0.72	0.61	1.18	0.24
$\log_{10}(\text{Lifespan proxy})^2$	-1.42	0.46	-3.07	0.002
$\log_{10}(\text{Lifespan proxy})^3$	-0.30	0.39	-0.77	0.44
$\log_{10}(\text{Litter/clutch size})$	1.23	0.80	1.54	0.12
$\log_{10}(\text{Litter/clutch size})^2$	-0.21	0.53	-0.40	0.69
$\log_{10}(\text{Litter/clutch size})^3$	-0.23	0.41	-0.56	0.58
$\log_{10}(\text{Range area})$	-32.18	0.37	-87.40	< 0.001
$\log_{10}(\text{Range area})^2$	-6.51	0.32	-20.05	< 0.001
$\log_{10}(\text{Range area})^3$	9.84	0.31	31.28	< 0.001
square-root(Habitat breadth)	1.05	0.42	2.53	0.01
square-root(Habitat breadth) ²	-0.91	0.32	-2.83	0.005
square-root(Habitat breadth) ³	-0.06	0.30	-0.20	0.84
square-root(Diet breadth)	-0.52	0.43	-1.20	0.23
square-root(Diet breadth) ²	0.02	0.32	0.07	0.95
square-root(Diet breadth) ³	0.24	0.31	0.77	0.44
Specialisation: Natural habitat specialist	0.10	0.01	7.07	< 0.001
Diel activity: Non-nocturnal	-0.02	0.01	-1.76	0.08
Primary diet: Omnivore	-0.01	0.04	-0.18	0.85
Primary diet: Vertebrates	-0.01	0.02	-0.64	0.52

S4.9.3 Effects for the PGLS models fitted on all species, against effects for the PGLS models fitted on species whose range area was $>100 \text{ km}^2$




Figure S4.17: Effects for the PGLS models investigating associations between species-level ecological characteristics and climate-change sensitivity, either fitted on all species (y-axis), or fitted on the species whose range area was $>100 \text{ km}^2$ (x-axis). Overall, the estimates from both sets of models were congruent, except for those estimated for geographical range area (I show the effects for range area separately from the effects of other characteristics). Across all classes, the relationship between sensitivity and geographical range area was reversed between the two sets of models. I found that sensitivity was positively associated with geographical range area when including all species, likely because of the underestimation of climate-change sensitivity for the most narrow-ranging species when working with a resolution of 5 km^2 (see Figure S4.6). The dashed line is the identity line ($y=x$).

S4.10 Validations on complete trait data subsets

Table S4.22: Summary of the effects of the ecological characteristics (except for diet) on (a) species' responses to disturbed land uses ('within land-use type' effects) and on (b) species climate-change sensitivity, for each class of terrestrial vertebrates, from the models fitted on empirical trait values (excluding all imputed values). The symbol  indicates where a characteristic has a significant negative effect on occurrence probability within a disturbed land-use type (within any of the land-use intensities), or where the characteristic renders species significantly more sensitive to climate change. A  indicates a significantly positive effect of a characteristic on occurrence probability in a land-use type (within any of the land-use intensities), or significantly lower sensitivity to climate change. For the land-use effects, I report 'within land-use type effects' here, that is, within a disturbed land use whether there were significant differences in occurrence probability among species with different trait values. These effects were derived from the interactive terms of the full models.

(a) "Within land-use type" effects of the species-level characteristics on occurrence probability - <=> more sensitive; + <=> less sensitive															Predictors					
Secondary vegetation				Plantation forest				Agricultural				Urban				Habitat affinity	Specialisation	Resource use	Life-history	
																				
0	-	0	0	0	-	0	0	0	-	-	+	+	-	0	NA	More narrowly-distributed	Habitat affinity	Specialisation	Resource use	Life-history
0	-	0	0	-	-	0	0	-	-	0	-	-	-	0	NA	Smaller habitat breadth				
-	-	0	0	0	-	0	0	-	-	-	+	0	-	0	NA	Natural habitat specialist	Specialisation	Resource use	Life-history	
0	+	+	0	+	+	(+)	+	0	+	+	0	0	0	0	NA	Non-nocturnal				
-	0	(+)	0	(-)	0	+	0	-	0	(+)	+	-	-	0	NA	Narrower diet breadth	Resource use			
NA	(+)	+	NA	NA	+	+	NA	NA	-	0	NA	NA	+	0	NA	Smaller body mass				
NA	0	0	-	NA	-	(-)	0	NA	-	(-)	0	NA	0	-	NA	Smaller litter/clutch size	Life-history			
NA	0	0	-	NA	(-)	0	0	NA	+	0	+	NA	-	0	NA	Shorter-lived				




Figure S4.18: Predicted occurrence probability as a function of land use, land-use intensity, diet and their interactions in each class. The predictions were obtained from the partial models fitted in each class for diet, estimated using empirical trait data subsets (i.e., excluding imputed trait values). Empty plots are drawn where there were no data for a diet category in given class (e.g., amphibian fruit/nectar eaters). Effects could not be estimated for urban reptiles, as well as for urban vertebrate, fruit/nectar and plant/seed eaters for mammals because there were no sampled species. The predictions are rescaled with reference to minimally-used primary vegetation. Primary: primary vegetation; Secondary: secondary vegetation; plantation: plantation forest; agricultural: cropland and pasture.

Table S4.23: Summary for the PGLS model fitted on amphibians, using the empirical trait data subset (i.e., excluding imputed trait values), looking at the association between the species-level ecological characteristics and climate-change sensitivity.

	Estimate	Std. Error	t value	Pr(> t)
Intercept	1.825	0.342	5.344	< 0.001
$\log_{10}(\text{Body mass})$	-0.030	0.035	-0.875	0.385
$\log_{10}(\text{Lifespan proxy})$	0.180	0.103	1.742	0.086
$\log_{10}(\text{Litter/clutch size})$	0.052	0.032	1.639	0.106
$\log_{10}(\text{Range area})$	-0.261	0.024	-10.888	< 0.001
square-root(Habitat breadth)	-0.032	0.030	-1.078	0.285
square-root(Diet breadth)	-0.068	0.107	-0.635	0.527
Specialisation: Natural habitat specialist	-0.006	0.059	-0.095	0.925
Diel activity: Non-nocturnal	-0.009	0.044	-0.205	0.839

Table S4.24: Summary for the PGLS model fitted on birds, using the empirical trait data subset (i.e., excluding imputed traits values), looking at the association between the species-level ecological characteristics and climate-change sensitivity.

	Estimate	Std. Error	t value	Pr(> t)
Intercept	1.551	0.143	10.875	< 0.001
$\log_{10}(\text{Body mass})$	0.020	0.011	1.759	0.079
$\log_{10}(\text{Lifespan proxy})$	0.047	0.036	1.308	0.191
$\log_{10}(\text{Litter/clutch size})$	0.212	0.022	9.487	< 0.001
$\log_{10}(\text{Range area})$	-0.230	0.003	-73.130	< 0.001
square-root(Habitat breadth)	-0.011	0.005	-2.319	0.020
square-root(Diet breadth)	0.015	0.011	1.319	0.187
Specialisation: Natural habitat specialist	0.044	0.007	6.586	< 0.001
Diel activity: Non-nocturnal	0.001	0.058	0.019	0.984
Primary diet: Invertebrates	0.047	0.014	3.330	0.001
Primary diet: Omnivore	0.016	0.014	1.144	0.253
Primary diet: Plants/seeds	0.048	0.016	2.993	0.003
Primary diet: Vertebrates	-0.005	0.022	-0.247	0.805

Table S4.25: Summary for the PGLS model fitted on mammals, using the empirical trait data subset (i.e., excluding imputed trait values), looking at the association between the species-level ecological characteristics and climate-change sensitivity.

	Estimate	Std. Error	t value	Pr(> t)
Intercept	2.194	0.187	11.705	< 0.001
$\log_{10}(\text{Body mass})$	-0.047	0.012	-3.865	< 0.001
$\log_{10}(\text{Lifespan proxy})$	0.065	0.046	1.416	0.157
$\log_{10}(\text{Litter/clutch size})$	0.147	0.034	4.293	< 0.001
$\log_{10}(\text{Range area})$	-0.269	0.005	-54.621	< 0.001
square-root(Habitat breadth)	-0.042	0.010	-4.320	< 0.001
square-root(Diet breadth)	-0.030	0.020	-1.487	0.137
Specialisation: Natural habitat specialist	0.016	0.013	1.235	0.217
Diel activity: Non-nocturnal	0.012	0.015	0.792	0.428
Primary diet: Invertebrates	-0.038	0.031	-1.220	0.223
Primary diet: Omnivores	-0.031	0.027	-1.148	0.251
Primary diet: Plants/seeds	0.034	0.027	1.298	0.195
Primary diet: Vertebrates	-0.039	0.040	-0.995	0.320

Table S4.26: Summary for the PGLS model fitted on reptiles, using the empirical trait data subset (i.e., excluding imputed trait values), looking at the association between the species-level ecological characteristics and climate-change sensitivity.

	Estimate	Std. Error	t value	Pr(> t)
Intercept	1.269	0.522	2.433	0.018
$\log_{10}(\text{Body mass})$	-0.033	0.058	-0.570	0.571
$\log_{10}(\text{Lifespan proxy})$	0.015	0.130	0.119	0.905
$\log_{10}(\text{Litter/clutch size})$	0.095	0.128	0.739	0.463
$\log_{10}(\text{Range area})$	-0.093	0.049	-1.878	0.066
square-root(Habitat breadth)	-0.094	0.045	-2.097	0.041
square-root(Diet breadth)	0.067	0.112	0.595	0.554
Specialisation: Natural habitat specialist	0.221	0.075	2.936	0.005
Diel activity: Non-nocturnal	-0.118	0.080	-1.486	0.143

Appendix 4: Supporting information for Chapter 5

Table S5.1: Number of species for which RMR data was collected in each class & data sources, initial RMR coverage for PREDICTS species and phylogenetic signal in RMR.

Class	RMR data	Coverage for PREDICTS species	Phylogenetic signal (Pagel's λ , $\pm 95\%$ CI)
Amphibians	126 species from Stark et al. (2020)	16/379 species (4%)	0.89 (0.86-0.91)
Birds	719 species from McNab (2009) Fristoe et al. (2015) Londoño et al. (2015) Stark et al. (2020)	317/3129 species (10%)	0.97 (0.95-0.98)
Mammals	685 species from PanTHERIA (Jones et al., 2009) Fristoe et al. (2015) Stark et al. (2020)	148/556 species (27%)	0.99 (0.98-0.99)
Reptiles	173 species from Stark et al. (2020)	24/329 species (7.3%)	0.90 (0.86-0.92)

$$\begin{aligned} \log(\text{Abundance}) = & \text{LU} + \text{LUI} + \text{TG} + \log(\text{BM}) + \\ & \text{LU:LUI} + \text{LU:TG} + \text{LU:log(BM)} + \text{LUI:TG} + \text{LUI:log(BM)} + \text{TG:log(BM)} + \\ & \text{LU:TG:log(BM)} + \text{LUI:TG:log(BM)} \end{aligned}$$

Figure S5.1: Formula for the abundance model, used to understand the role of shifts in the body mass of species on observed changes in tRMR (see main text, ‘Disentangling the effects of body mass and abundance on tRMR’). I fitted a model to explain changes in species abundance (given presence) by land use, land-use intensity, trophic group, body mass and their interactions. The model included all two-way interactions among these predictors. To account for potential differences in the slope of the relationship between abundance and body mass among the different trophic groups, I also included two three-way interactions in the model (among land use, trophic group and body mass; and among land-use intensity, trophic group and body mass). Random effects included study, site and species identity. LU: land use; LUI: land-use intensity; TG: trophic group; BM: body mass.




Figure S5.2: Effects of land use, land-use intensity, trophic group, body mass and their interactions on assemblage-level total abundance, estimated from the model specified in Figure S5.1. The predictions are rescaled with reference to minimally used primary vegetation, considered to be the undisturbed baseline. Primary: primary vegetation; secondary: secondary vegetation; plantation: plantation forest. For visualisation purposes, I plotted the predictions for three body mass levels (but body mass was considered as a continuous variable in the model).





Figure S5.3: Diagnostic plots (qq-plot and residual distribution) for the linear mixed-effects model looking at the effects of land use, land-use intensity, trophic group and their interactions on assemblage-level total RMR. The diagnostic plots were obtained with the ‘performance’ R package (Lüdecke et al., 2021).

(a) Coefficient estimates



(b) Standard error estimates




Figure S5.4: Estimates for the models looking at the effects of land use, land-use intensity, trophic group and their interactions on assemblage-level total RMR. I plotted the estimates from the model fitted on the empirical and imputed RMR values (presented in the main text) on the y-axis, and the estimates from the model fitted on the empirical RMR values only on the x-axis.




Figure S5.5: Effects of land use, land-use intensity, trophic level and their interactions on assemblage-level total RMR, estimated from the model fitted on the empirical and imputed RMR values (presented in the main text) and from the model fitted on the empirical values only. The predictions are rescaled with reference to minimally used primary vegetation, considered to be the undisturbed baseline. Primary: primary vegetation; secondary: secondary vegetation; plantation: plantation forest.




Figure S5.6: Diagnostic plots for the generalised mixed-effects model looking at the effects of land use, land-use intensity, trophic group, residual RMR and their interactions on species' probability of occurrence. The diagnostic plots were obtained with the 'DHARMA' R package (Hartig, 2021).





Figure S5.7: Model's coefficients from the occurrence model fitted using the 'lme4' package (Bates et al., 2015) against coefficients from the model fitted using a Bayesian framework with the 'MCMCglmm' package (Hadfield, 2010). The models were fitted to investigate the effects of land use, land-use intensity, trophic group and residual RMR on species occurrence probability.

(a) Slope of the relationships between occurrence probability (log-odds) and residual RMR



(b) Predicted effects of residual RMR on occurrence probability (minimal use intensity)




Figure S5.8: (a) Slope estimates for the relationship between residual RMR and occurrence probability in each land-use type and for the three levels of land-use intensity, from the model fitted using the empirical RMR values (i.e., excluding imputed RMR values). The black horizontal line indicates the slope for the reference level (primary vegetation) for minimal land-use intensity. The grey dashed line marks 0. Error bars are 95% confidence intervals. (b) Effect of residual RMR on species probability of occurrence within each trophic level and for each land use-type. I plotted the predictions for minimal land-use intensity only. Solid lines represent significant relationships. Primary: primary vegetation; secondary: secondary vegetation; plantation: plantation forest.

Bibliography for the Appendices

- Bates, D., Mächler, M., Bolker, B. M., and Walker, S. C. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*. DOI: 10.18637/jss.v067.i01.
- Borges, R., Machado, J. P., Gomes, C., Rocha, A. P., and Antunes, A. (2018). Measuring phylogenetic signal between categorical traits and phylogenies. *Bioinformatics*. DOI: 10.1093/bioinformatics/bty800.
- Bouckaert, R., Heled, J., Kühnert, D., Vaughan, T., Wu, C. H., Xie, D., Suchard, M. A., Rambaut, A., and Drummond, A. J. (2014). BEAST 2: A Software Platform for Bayesian Evolutionary Analysis. *PLoS Computational Biology*. DOI: 10.1371/journal.pcbi.1003537.
- Bruggeman, J., Heringa, J., and Brandt, B. W. (2009). PhyloPars: Estimation of missing parameter values using phylogeny. *Nucleic Acids Research*. DOI: 10.1093/nar/gkp370.
- Buuren, S van and Oudshoorn, C. G. M. (2007). MICE: multivariate imputation by chained equations. *R package version*.
- Cadotte, M. W., Carscadden, K., and Mirochnick, N. (2011). Beyond species: Functional diversity and the maintenance of ecological processes and services. *Journal of Applied Ecology*. DOI: 10.1111/j.1365-2664.2011.02048.x.
- Cailliez, F. (1983). The analytical solution of the additive constant problem. *Psychometrika*. DOI: 10.1007/BF02294026.
- Chamberlain, S. (2018). *rredlist: 'IUCN' Red List Client*. R package version 0.5.0.
- Chamberlain, S. A. and Szöcs, E. (2013). taxize : taxonomic search and retrieval in R [version 2; referees: 3 approved]. *F1000Research*. DOI: 10.12688/f1000research.2-191.v2.
- Debastiani, V. J., Bastazini, V. A. G., and Pillar, D (2021). Ecological Informatics Using phylogenetic information to impute missing functional trait values in ecological databases. 63.April. DOI: 10.1016/j.ecoinf.2021.101315.
- Diniz-Filho, J. A. F., Bini, L. M., Rangel, T. F., Morales-Castilla, I., Olalla-Tárraga, M. Á., Rodríguez, M. Á., and Hawkins, B. A. (2012). On the selection of phylogenetic eigenvectors for ecological analyses. *Ecography*. DOI: 10.1111/j.1600-0587.2011.06949.x.
- Dormann, C. F., Elith, J., Bacher, S., Buchmann, C., Carl, G., Carré, G., Marquéz, J. R., Gruber, B., Lafourcade, B., Leitão, P. J., Münkemüller, T., McClean, C., Osborne, P. E., Reineking, B., Schröder, B., Skidmore, A. K., Zurell, D., and Lautenbach, S. (2013). Collinearity: A review of methods to deal with it and a simulation study evaluating their performance. *Ecography*. DOI: 10.1111/j.1600-0587.2012.07348.x.

- Dray, S., Dufour, A.-B., and Chessel, D. (2007). The ade4 Package – II: Two-Table and K-Table Methods. *R News*, 7.2, 47–52.
- Etard, A., Morrill, S., and Newbold, T. (2020). Global gaps in trait data for terrestrial vertebrates. *Global Ecology and Biogeography*, November 2019, 1–16. DOI: 10.1111/geb.13184.
- Faurby, S., Davis, M., Pedersen, R., Schowanek, S. D., Antonelli1, A., and Svenning, J.-C. (2018). PHYLACINE 1.2: The Phylogenetic Atlas of Mammal Macroecology. *Ecology*, 99.11, 2626–2626. DOI: 10.1002/ecy.2443.
- Faurby, S., Pedersen, R., Davis, M., Schowanek, S. D., Jarvie, S., Antonelli, A., and Svenning, J.-C. (2020). PHYLACINE 1.2.1: An update to the Phylogenetic Atlas of Mammal Macroecology. DOI: 10.5281/zenodo.3690867.
- Fox, J. and Monette, G. (1992). Generalized collinearity diagnostics. *Journal of the American Statistical Association*. DOI: 10.1080/01621459.1992.10475190.
- Fristoe, T. S., Burger, J. R., Balk, M. A., Khaliq, I., Hof, C., and Brown, J. H. (2015). Metabolic heat production and thermal conductance are mass-independent adaptations to thermal environment in birds and mammals. *Proceedings of the National Academy of Sciences of the United States of America*, 112.52, 15934–15939. DOI: 10.1073/pnas.1521662112.
- Hadfield, J. D. (2010). MCMC Methods for Multi-Response Generalized Linear Mixed Models: The MCMCglmm R Package. *Journal of Statistical Software*, 33.2, 1–22.
- Hartig, F. (2021). *DHARMA: Residual Diagnostics for Hierarchical (Multi-Level / Mixed) Regression Models*. R package version 0.4.1.
- Hudson, L. N. et al. (2014). The PREDICTS database: A global database of how local terrestrial biodiversity responds to human impacts. *Ecology and Evolution*. DOI: 10.1002/ece3.1303.
- (2017). The database of the PREDICTS (Projecting Responses of Ecological Diversity In Changing Terrestrial Systems) project. *Ecology and Evolution*. DOI: 10.1002/ece3.2579.
- IUCN (2020). The IUCN Red List of Threatened Species. <http://www.iucnredlist.org>. Version 2020.1.
- Jetz, W., Thomas, G. H., Joy, J. B., Hartmann, K., and Mooers, A. O. (2012). The global diversity of birds in space and time. *Nature*. DOI: 10.1038/nature11631.
- Jetz, W. and Pyron, R. A. (2018). The interplay of past diversification and evolutionary isolation with present imperilment across the amphibian tree of life. *Nature Ecology and Evolution*. DOI: 10.1038/s41559-018-0515-5.
- Johnson, T. F., Isaac, N. J., Paviolo, A., and González-Suárez, M. (2021). Handling missing values in trait data. *Global Ecology and Biogeography*. DOI: 10.1111/geb.13185.
- Jones, K. E. et al. (2009). PanTHERIA: a species-level database of life history, ecology, and geography of extant and recently extinct mammals. *Ecology*. DOI: 10.1890/08-1494.1.
- Lüdecke, D., Ben-Shachar, M. S., Patil, I., Waggoner, P., and Makowski, D. (2021). performance: An R Package for Assessment, Comparison and Testing of Statistical Models. *Journal of Open Source Software*, 6.60, 3139. DOI: 10.21105/joss.03139.
- Leroy, B., Meynard, C. N., Bellard, C., and Courchamp, F. (2015). virtualspecies, an R package to generate virtual species distributions. *Ecography*. DOI: 10.1111/ecog.01388.

- Londoño, G. A., Chappell, M. A., Castañeda, M. d. R., Jankowski, J. E., and Robinson, S. K. (2015). Basal metabolism in tropical birds: Latitude, altitude, and the 'pace of life'. *Functional Ecology*, 29.3, 338–346. DOI: 10.1111/1365-2435.12348.
- McNab, B. K. (2009). Ecological factors affect the level and scaling of avian BMR. *Comparative Biochemistry and Physiology - A Molecular and Integrative Physiology*, 152.1, 22–45. DOI: 10.1016/j.cbpa.2008.08.021.
- Oliveira, B. F., São-Pedro, V. A., Santos-Barrera, G., Penone, C., and Costa, G. C. (2017). AmphiBIO, a global database for amphibian ecological traits. *Scientific Data*. DOI: 10.1038/sdata.2017.123.
- Pagel, M. (1999). Inferring the historical patterns of biological evolution. *Nature*. DOI: 10.1038/44766.
- Penone, C., Davidson, A. D., Shoemaker, K. T., Di Marco, M., Rondinini, C., Brooks, T. M., Young, B. E., Graham, C. H., and Costa, G. C. (2014). Imputation of missing data in life-history trait datasets: Which approach performs the best? *Methods in Ecology and Evolution*. DOI: 10.1111/2041-210X.12232.
- Rabosky, A. D., Cox, C., Rabosky, D., Title, P., Holmes, I., Feldman, A., and McGuire, J. (2016). Coral snakes predict the evolution of mimicry across New World snakes. *Nature Communications*, 7, 11484.
- Revell, L. J. (2012). phytools: An R package for phylogenetic comparative biology (and other things). *Methods in Ecology and Evolution*, 3, 217–223.
- Rinnan, D. S. and Lawler, J. (2019). Climate-niche factor analysis: a spatial approach to quantifying species vulnerability to climate change. *Ecography*, 42.9, 1494–1503. DOI: <https://doi.org/10.1111/ecog.03937>.
- Santini, L., Benítez-López, A., Ficetola, G. F., and Huijbregts, M. A. J. (2018). Length–mass allometries in amphibians. *Integrative Zoology*, 13.1, 36–45. DOI: 10.1111/1749-4877.12268.
- Santos, T. (2018). Package ‘PVR’. Phylogenetic Eigenvectors Regression and Phylogenetic Signal-Representation Curve.
- Stark, G., Pincheira-Donoso, D., and Meiri, S. (2020). No evidence for the ‘rate-of-living’ theory across the tetrapod tree of life. *Global Ecology and Biogeography*, 29.5, 857–884. DOI: 10.1111/geb.13069.
- Stekhoven, D. J. (2016). Nonparametric Missing Value Imputation using Random Forest. *R Package version 1.4*. DOI: 10.1093/bioinformatics/btr597.
- Stekhoven, D. J. and Bühlmann, P. (2012). Missforest-Non-parametric missing value imputation for mixed-type data. *Bioinformatics*. DOI: 10.1093/bioinformatics/btr597.
- Tonini, J. F. R., Beard, K. H., Ferreira, R. B., Jetz, W., and Pyron, R. A. (2016). Fully-sampled phylogenies of squamates reveal evolutionary patterns in threat status. *Biological Conservation*. DOI: 10.1016/j.biocon.2016.03.039.
- Troyanskaya, O., Cantor, M., Sherlock, G., Brown, P., Hastie, T., Tibshirani, R., Botstein, D., and Altman, R. B. (2001). Missing value estimation methods for DNA microarrays. *Bioinformatics*. DOI: 10.1093/bioinformatics/17.6.520.
- Wilman, H., Belmaker, J., Simpson, J., Rosa, C. de la, Rivadeneira, M. M., and Jetz, W. (2014). EltonTraits 1.0: Species-level foraging attributes of the world’s birds and mammals. *Ecology*. DOI: 10.1890/13-1917.1.