

MARANDA 2025 KCSE PREDICTION



EXPECTED EXAM 1



CHEMISTRY

233/3

PAPER 3 (PRACTICAL)

TIME: 2¼ HOURS

NAME.....

SCHOOL..... SIGN.....

INDEX NO..... ADM NO.....

Kenya Certificate of Secondary Education.

INSTRUCTIONS TO CANDIDATES

- Write your name in the spaces provided above.
- Answer all the questions in the spaces provided
- All workings must be clearly shown where necessary

FOR EXAMINERS USE ONLY

QUESTION	MAXIMUM SCORE	CANDIDATE'S SCORES
1	20	
2	11	
3	09	
TOTAL SCORE	40	

QUESTIONS

1. You are provided with:

- 0.30g solid A, magnesium metal
- Hydrochloric acid, solution B
- 0.15M sodium carbonate, solution C
- Methyl orange indicator

You are required to determine:

- The enthalpy change, ΔH per mole for the reaction between magnesium and excess hydrochloric acid.
- The concentration in moles per litre of hydrochloric acid, solution B.

PROCEDURE I

- i. Using a burette, place 50.0cm³ of Solution B in a 100 ml plastic beaker.
- ii. Measure the temperature of solution B after every 30 seconds. Record your readings in Table 1.
- iii. At the 90th second, add **ALL** of solid A provided to the beaker, stir gently with the thermometer and continue measuring and recording the temperature after every 30 seconds. Complete Table


1. **Retain the content in the beaker for use in procedure II.**

Table 1

Time (S)	0	30	60	90	120	150	180	210	240	270
temperature (°C)				X						

(3 marks)

- a) On the grid provided, plot a graph of temperature (vertical axis) against time. **(3 marks)**



b) Determine the change in temperature, ΔT , for the complete reaction. Show the working on the graph.

$\Delta T = \dots\dots\dots$

(1mark)

- c) Calculate the heat change for the reaction. (Assume density of solution = 1.0g/cm^3 , specific capacity of the solution = $4.2\text{ kJ g}^{-1}\text{ K}^{-1}$). **(1mark)**

.....

- d) The relative atomic mass of magnesium is 24.0. Calculate the enthalpy change, ΔH , for the reaction per mole of magnesium. **(1mark)**

.....

PROCEDURE II

- i. Rinse the burette and fill it with 0.15M sodium carbonate, solution C.
- ii. Transfer **ALL** the solutions in the beaker from procedure I into a 250ml volumetric flask. Rinse both the beaker and thermometer with distilled water and add the rinsing to the volumetric flask. Add more distilled water to the volumetric flask up to the mark. Label this as solution T.
- iii. Pipette 25cm^3 of solution T into a 250ml conical flask and add 2 drops of methyl orange indicator.
- iv. Titrate solution T in the conical flask with the sodium carbonate, solution C and record the results in Table 2.
- v. Repeat steps (iii) and (iv) and complete Table 2.

Table 2

Titre	1	2	3
Final burette reading (cm^3)			
Initial Burette reading (cm^3)			
Volume of solution C (cm^3)			

(4 marks)

- a) Calculate the average volume of the 0.15M sodium carbonate, solution C used. **(1 mark)**

.....
.....
.....
b) Calculate the number of moles of :

i. Sodium carbonate used. (1 mark)

.....
.....
.....
ii. Hydrochloric acid in the 25.0 cm³ of solution T. (1 mark)

.....
.....
.....
iii. Hydrochloric acid in the 25.0 cm³ of solution T. (1 mark)

.....
.....
.....
iv. Hydrochloric acid that reacted with magnesium metal. (1 mark)

.....
.....
.....
v. Total number of moles of hydrochloric acid in the 50.0 cm³, solution B. (1 mark)

.....
.....
.....
c) Determine the concentration of hydrochloric acid in moles per litre, in solution B. (1 mark)

c)

Observation	Inferences
(1mark)	(1mark)

To the third portion, add 2M ammonia solution dropwise until in excess.

d)

Observation	Inferences
(1mark)	(½ mark)

To the fourth portion, add 2 - 3 drops of aqueous potassium iodide solution.

Observation	Inferences
(½ mark)	(½ mark)

e) To the fifth portion, add 2 - 3 drops of barium nitrate solution.

Observation	Inferences
(1mark)	(1mark)

THIS IS THE LAST PRINTED PAGE