

INTRO TO DATA SCIENCE LECTURE 6: LOGISTIC REGRESSION

Jason Dolatshahi Data Scientist, EveryScreen Media RECAP 2

LAST TIME:

- LINEAR REGRESSION
- POLYNOMIAL REGRESSION
- REGULARIZATION

QUESTIONS?

AGENDA 3

- I. LOGISTIC REGRESSION
- **II. OUTCOME VARIABLES**
- III. ERROR TERMS
- IV. INTERPRETING RESULTS

EXERCISES:

IMPLEMENTING A LOGISTIC FIT IN R

INTRO TO DATA SCIENCE

I. LOGISTIC REGRESSION

LOGISTIC REGRESSION

	continuous	categorical
supervised	???	???
unsupervised	???	???

LOGISTIC REGRESSION

supervised
unsupervisedregression
dimension reductionclassification
clustering

Q: What is **logistic regression**?

Q: What is **logistic regression**?

A: A generalization of the linear regression model to *classification* problems.

In linear regression, we used a set of covariates to predict the value of a (continuous) outcome variable.

In linear regression, we used a set of covariates to predict the value of a (continuous) outcome variable.


In logistic regression, we use a set of covariates to predict *probabilities* of (binary) class membership.


In linear regression, we used a set of covariates to predict the value of a (continuous) outcome variable.

In logistic regression, we use a set of covariates to predict *probabilities* of (binary) class membership.

These probabilities are then mapped to *class labels*, thus solving the classification problem.

PROBABILITIES 12






NOTE

Probability predictions look like this.

value of independent variable



value of independent variable

The logistic regression model is an *extension* of the linear regression model, with a couple of important differences.

LOGISTIC REGRESSION

The logistic regression model is an *extension* of the linear regression model, with a couple of important differences.

The first difference is in the outcome variable.

LOGISTIC REGRESSION

The logistic regression model is an *extension* of the linear regression model, with a couple of important differences.

The first difference is in the outcome variable.

The second difference is in the error term.

INTRO TO DATA SCIENCE

II. OUTCOME VARIABLES

OUTCOME VARIABLES 18

The key variable in any regression problem is the **conditional mean** of the outcome variable y given the value of the covariate x:

The key variable in any regression problem is the **conditional mean** of the outcome variable y given the value of the covariate x:

In linear regression, we assume that this conditional mean is a linear function taking values in $(-\infty, +\infty)$:

$$E(y|x) = \alpha + \beta x$$

OUTCOME VARIABLES

In logistic regression, we've seen that the conditional mean of the outcome variable takes values only in the unit interval [0, 1].

OUTCOME VARIABLES

In logistic regression, we've seen that the conditional mean of the outcome variable takes values only in the unit interval [0, 1].

The first step in extending the linear regression model to logistic regression is to map the outcome variable $E(y \mid x)$ into the unit interval.

In logistic regression, we've seen that the conditional mean of the outcome variable takes values only in the unit interval [0, 1].

The first step in extending the linear regression model to logistic regression is to map the outcome variable $E(y \mid x)$ into the unit interval.

Q: How do we do this?

THE LOGISTIC FUNCTION


A: By using a transformation called the **logistic function**:

$$E(y|x) = \pi(x) = \frac{e^{\alpha + \beta x}}{1 + e^{\alpha + \beta x}}$$

A: By using a transformation called the **logistic function**:

$$E(y|x) = \pi(x) = \frac{e^{\alpha + \beta x}}{1 + e^{\alpha + \beta x}}$$


We've already seen what this looks like:



A: By using a transformation called the **logistic function**:

$$E(y|x) = \pi(x) = \frac{e^{\alpha + \beta x}}{1 + e^{\alpha + \beta x}}$$

We've already seen what this looks like:



NOTE

For any value of x, y is in the interval [0, 1]

This is a nonlinear transformation!

The **logit function** is an important transformation of the logistic function. Notice that it returns the linear model!

$$g(x) = ln(\frac{\pi(x)}{1-\pi(x)}) = \alpha + \beta x$$

The **logit function** is an important transformation of the logistic function. Notice that it returns the linear model!

$$g(x) = \ln(\frac{\pi(x)}{1 - \pi(x)}) = \alpha + \beta x$$

The logit function is also called the log-odds function.

The **logit function** is an important transformation of the logistic function. Notice that it returns the linear model!

$$g(x) = ln(\frac{\pi(x)}{1-\pi(x)}) = \alpha + \beta x$$

NOTE

This name hints at its usefulness in interpreting our results.

We will see why shortly.

The logit function is also called the log-odds function.

INTRO TO DATA SCIENCE

III. ERROR TERMS

The second difference between linear regression and the logistic regression model is in the error term.

The second difference between linear regression and the logistic regression model is in the error term.

One of the key assumptions of linear regression is that the error terms follow independent Gaussian distributions with zero mean and constant variance:

$$\epsilon \sim N(0, \sigma^2)$$

In logistic regression, the outcome variable can take only two values: 0 or 1.

In logistic regression, the outcome variable can take only two values: 0 or 1.

It's easy to show from this that instead of following a Gaussian distribution, the error term in logistic regression follows a Bernoulli distribution:

$$\epsilon \sim B(0, \pi(1-\pi))$$

In logistic regression, the outcome variable can take only two values: 0 or 1.

It's easy to show from this that instead of following a Gaussian distribution, the error term in logistic regression follows a Bernoulli distribution:

$$\epsilon \sim B(0, \pi(1-\pi))$$

This is the same

This is the same distribution followed by a coin toss.

Think about why this makes sense!

AN ASIDE: GLM

These two key differences define the logistic regression model, and they also lead us to a kind of unification of regression techniques called **generalized linear models**.

These two key differences define the logistic regression model, and they also lead us to a kind of unification of regression techniques called **generalized linear models**.

Briefly, GLMs generalize the distribution of the error term, and allow the conditional mean of the response variable to be related to the linear model by a **link function**.

AN ASIDE: GLM

In the present case, the error term follows a Bernoulli distribution, and the logit is the link function that connects us to the linear predictor.

In the present case, the error term follows a Bernoulli distribution, and the logit is the link function that connects us to the linear predictor.

$$g(x) = ln(\frac{\pi(x)}{1-\pi(x)}) = \alpha + \beta x$$

In the present case, the error term follows a Bernoulli distribution, and the logit is the link function that connects us to the linear predictor.

$$g(x) = \ln(\frac{\pi(x)}{1 - \pi(x)}) = \alpha + \beta x$$

Since the Bernoulli distribution and the logit function share a common parameter π , we say that the logit is the **canonical link function** for the Bernoulli distribution.

In the present case, the error term follows a Bernoulli distribute the logit is the link function that connects us to the linear

$$g(x) = \ln(\frac{\pi(x)}{1 - \pi(x)}) = \alpha + \beta x$$

Since the Bernoulli distribution and the logit function share a common parameter π , we say that the logit is the **canonical link function** for the Bernoulli distribution.

INTRO TO DATA SCIENCE

IV. INTERPRETING RESULTS

In linear regression, the parameter β represents the change in the response variable for a unit change in the covariate.

In linear regression, the parameter β represents the change in the response variable for a unit change in the covariate.

In logistic regression, β represents the change in the logit function for a unit change in the covariate.

In linear regression, the parameter β represents the change in the response variable for a unit change in the covariate.

In logistic regression, β represents the change in the logit function for a unit change in the covariate.

Interpreting this change in the logit function requires another definition first.

The **odds** of an event are given by the ratio of the probability of the event by its complement:

$$O(x=1) = \frac{\pi(1)}{(1-\pi(1))}$$

The **odds** of an event are given by the ratio of the probability of the event by its complement:

$$O(x=1) = \frac{\pi(1)}{(1-\pi(1))}$$

The **odds ratio** of a binary event is given by the odds of the event divided by the odds of its complement:

$$OR = \frac{O(x=1)}{O(x=0)} = \frac{\pi(1)/[1-\pi(1)]}{\pi(0)/[1-\pi(0)]}$$

Substituting the definition of $\pi(x)$ into this equation yields (after some algebra),

$$OR = e^{\beta}$$

Substituting the definition of $\pi(x)$ into this equation yields (after some algebra),

$$OR = e^{\beta}$$

This simple relationship between the odds ratio and the parameter β is what makes logistic regression such a powerful tool.

Q: So how do we interpret this?

Q: So how do we interpret this?

A: The odds ratio of a binary event gives the increase in likelihood of an outcome if the event occurs.

Suppose we are interested in mobile purchase behavior. Let y be a class label denoting purchase/no purchase, and let x denote a mobile OS (for example, iOS).

Suppose we are interested in mobile purchase behavior. Let y be a class label denoting purchase/no purchase, and let x denote a mobile OS (for example, iOS).

In this case, an odds ratio of 2 (eg, $\beta = \log(2)$) indicates that a purchase is twice as likely for an iOS user as for a non-iOS user.

INTRO TO DATA SCIENCE

EX: LOGISTIC REGRESSION