

QFT

Adwait

Quantum Field Theory

Canonical Quantisation.

For Quantising fields using second quantisation, we follow the algorithm.

step 1:

Get the Lagrangian Density \mathcal{L}

step 2:

Get the momentum and Hamiltonian density

$$\Pi^\nu = \frac{\partial \mathcal{L}}{\partial(\partial_\nu \phi)} \mathcal{H} = \Pi^0 \partial_0 \phi - \mathcal{L}$$

step 3:

$$[\hat{\phi}(x^\nu), \hat{\Pi}^0(y^\nu)] = i\delta^3(x^\nu - y^\nu)$$

Step 4:

Expand the field in plane waves and get other operators. Use Wick ordering to prevent infinities.

Free Scalar field:

step I:

$$\mathcal{L} = \frac{1}{2}((\partial_\nu \phi)^2 - m^2 \phi^2)$$

Step II:

$$\Pi^\nu = \partial^\nu \phi$$

$$\mathcal{H} = \frac{1}{2}(\dot{\phi}^2 + (\nabla\phi)^2 + m^2\phi^2)$$

Step III:

Do the above

Step IV:

$$\hat{\phi}(x^\nu) = \int \frac{d^4p}{(2\pi)^4} (\hat{a}(k_\nu) e^{ik_\nu x^\nu} + \hat{a}^\dagger(k_\nu) e^{-ik_\nu x^\nu}) = \int \frac{d^3p}{(2\pi)^3 \sqrt{2E_p}} (\hat{a}_p e^{ip \cdot x} + \hat{a}_p^\dagger e^{-ip \cdot x})$$

Where $E_p = \sqrt{p^2 + m^2}$

$$\mathcal{H} = \hat{T}\mathcal{H} = \int d^3p E_p \hat{a}_p^\dagger \hat{a}_p$$