

# MA2102: LINEAR ALGEBRA

## Lecture 27: Diagonalization

3rd November 2020

Indian Institute of Science Education & Research Kolkata



Suppose  $T : V \rightarrow V$  is a linear map. In order to compute trace and determinants of  $T^k$ , it definitely helps if there exist a basis  $\beta$  such that  $[T]_\beta$  is a diagonal matrix. In that case,

$$\text{trace}(T^k) = \text{trace}([T^k]_\beta) = \text{trace}([T]_\beta)^k,$$

which is easy to compute. Similarly, determinant of  $T$  becomes the product of the diagonal entries of  $[T]_\beta$ . As determinant is multiplicative,

$$\det(T^k) = \det([T^k]_\beta) = \det([T]_\beta)^k.$$

**Definition [Diagonalizable]** A matrix  $A \in M_n(F)$  is called **diagonalizable** if  $A$  is similar to a diagonal matrix, i.e., there exists an invertible matrix  $C \in M_n(F)$  such that  $CAC^{-1}$  is diagonal.

A linear map  $T : V \rightarrow V$  is called **diagonalizable** if  $[T]_\beta$  is diagonal for some basis  $\beta$ .

**Remark** If  $\gamma$  is another basis of  $V$ , then  $[T]_\gamma$ , being similar to  $[T]_\beta$ , is diagonalizable as a matrix. Conversely, suppose  $[T]_\gamma$  is diagonalizable for some basis  $\gamma$ , then  $Q[T]_\gamma Q^{-1}$  is diagonal for some matrix  $Q$ . Use  $Q$  to define a basis  $\beta$  such that  $[T]_\beta$  is diagonal. Thus,  $T : V \rightarrow V$  is diagonalizable if  $[T]_\gamma$  is diagonalizable for any basis  $\gamma$ .

If such a basis  $\beta$  exists, then write  $\beta = \{v_1, \dots, v_n\}$  and

$$[T]_\beta = D(\lambda_1, \dots, \lambda_n).$$

In other words,  $v_j \neq 0$  and  $T(v_j) = \lambda_j v_j$ .

**Definition [Eigenvector]** A non-zero vector  $v \in V$  is called an **eigenvector** of  $T : V \rightarrow V$  if  $T(v) = \lambda v$  for some scalar  $\lambda \in F$ . The scalar  $\lambda$  is called the **eigenvalue** of the eigenvector  $v$ .

**Definition** [Eigenvector] A non-zero vector  $v \in F^n$  is called an **eigenvector** of  $A \in M_n(F)$  if  $Av = \lambda v$  for some  $\lambda \in F$ . The scalar  $\lambda$  is called the **eigenvalue** of  $A$  corresponding to the eigenvector  $v$ .

**Remark** Eigenvalues are also called proper values, latent values, spectral values or characteristic values. 'Eigen' is a German word meaning *own*.

Geometrically, if  $v$  is an eigenvector of  $T$ , then consider the 1-dimensional subspace  $L$  spanned by  $v$ . It follows that  $T(L) \subseteq L$ , where equality is achieved if and only if the eigenvalue is non-zero.

**Example** (1) Any non-zero vector  $v$  in the null space of  $T : V \rightarrow V$  is an eigenvector with eigenvalue 0.

(2) Any non-zero vector  $v$  is an eigenvector of  $I_V : V \rightarrow V$  with eigenvalue 1.

**Theorem** [Characterization of eigenvalues]

*The following conditions are equivalent:*

- (a)  $\lambda$  is an eigenvalue for  $T$ ;
- (b)  $T - \lambda I_V$  is not invertible;
- (c)  $\det(T - \lambda I_V) = 0$ .

**Proof.**

(a)  $\Rightarrow$  (b): There exists  $v \neq 0$  such that  $T(v) = \lambda v$ , whence  $v \in N(T - \lambda I_V)$ . Thus,  $T - \lambda I_V$  is not invertible.

(b)  $\Rightarrow$  (c): If  $T - \lambda I_V$  is not invertible as a linear map, then  $[T - \lambda I_V]_\beta$  is not invertible as a matrix, whence  $\det([T]_\beta - \lambda I_n) = 0$ .

(c)  $\Rightarrow$  (a): The assumption implies that  $T - \lambda I_V$  is not invertible, i.e., it is not of full rank. Thus, there exists a non-zero  $v$  such that  $(T - \lambda I_V)(v) = 0$ . □

Observe that  $\lambda$  is an eigenvalue of  $T$  if and only if  $\lambda$  is an eigenvalue of  $[T]_\beta$  for any ordered basis  $\beta$ .

**Example** (1) If  $A \in M_n(F)$  is upper triangular, then  $\lambda$  is an eigenvalue if and only if  $\lambda = a_{jj}$  for some  $j \in \{1, \dots, n\}$ . Note that  $\det(A)$  is the product of the eigenvalues.

(2) If  $v \in V$  satisfies  $T(v) = \lambda v$ , then  $T^k(v) = \lambda^k v$ . Thus, eigenvectors of  $T$  are always eigenvectors of  $T^k$  for  $k \in \mathbb{N}$ . More generally, let  $p(x) = a_0 + a_1x + \dots + a_kx^k$  be a polynomial of degree  $k$ . It follows that  $v$  is an eigenvector of  $p(T)$  since

$$p(T)(v) = (a_0I_V + a_1T + \dots + a_kT^k)(v) = (a_0 + a_1\lambda + \dots + a_k\lambda^k)v.$$

(3) Let  $T : V \rightarrow V$  be linear and  $V$  be finite dimensional. Then  $T$  is invertible if and only if  $0$  is not an eigenvalue of  $T$ .

(4) Recall that  $[T^*]_{\beta^*} = [T]_{\beta}^t$  for  $T : V \rightarrow V$  and any basis  $\beta$  of  $V$ . Thus,

$$\det([T^*]_{\beta^*} - \lambda I_n) = \det([T]_{\beta}^t - \lambda I_n) = \det([T]_{\beta} - \lambda I_n).$$

Therefore, eigenvalues of  $T$  and  $T^*$  coincide.

Note that not all linear maps admit eigenvalues. For instance, let  $R_{\theta}$  denotes the counter-clockwise rotation on the plane by an angle  $\theta$ , i.e.,

$$R_{\theta} : \mathbb{R}^2 \rightarrow \mathbb{R}^2, \quad \begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}.$$

Let  $\theta \neq 0, \pi$ , i.e.,  $R_{\theta}$  is not  $\pm I_2$ . If  $(a, b)$  is an eigenvector with *real* eigenvalue  $\lambda$ , then

$$a \cos \theta - b \sin \theta = \lambda a, \quad a \sin \theta + b \cos \theta = \lambda b.$$

Squaring both equations and adding, we obtain  $a^2 + b^2 = \lambda^2(a^2 + b^2)$ , whence  $\lambda = \pm 1$  as  $a^2 + b^2 \neq 0$ . **Show that  $\lambda = \pm 1$  is not possible.**

Consider the complexified map

$$R_\theta : \mathbb{C}^2 \rightarrow \mathbb{C}^2, \begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}.$$

By condition (c) of the Theorem, we solve for

$$0 = (\cos \theta - \lambda)(\cos \theta - \lambda) + \sin^2 \theta = \lambda^2 - 2\lambda \cos \theta + 1.$$

This indeed has two complex roots although it has no real roots.

**Remark** The map  $R_\theta$  is an isometry and the eigenvalues (real or complex) of any isometry must have absolute value 1. From geometry, we see that if  $v$  is an eigenvector, then only rotation by 0 or  $\pi$  maps  $v$  to  $v$  or  $-v$  respectively.