



Introduction to Scientific Computation

Lecture 2

Fall 2026

Solving Linear Systems


How to solve this ?

$$Ax = b$$

How to solve this ?

$$Ax = b$$


Direct methods



$$x = A^{-1}b$$

Exact solution,
except for rounding errors,
fixed number of computational steps

Iterative methods



$$x_t = f(x_{t-1}, A, b)$$

Often work better in practice,
due to care about numerical stability,
iteratively improve solution estimate

Direct methods

If all linear equations are uncoupled:

If all linear equations are uncoupled:

$$A = \begin{bmatrix} a_{11} & & & \\ & a_{22} & & \\ & & \ddots & \\ & & & a_{nn} \end{bmatrix}$$

If all linear equations are uncoupled:

$$A = \begin{bmatrix} a_{11} & & & \\ & a_{22} & & \\ & & \ddots & \\ & & & a_{nn} \end{bmatrix}$$

Solution is obvious

If all linear equations are uncoupled:

$$A = \begin{bmatrix} a_{11} & & & \\ & a_{22} & & \\ & & \ddots & \\ & & & a_{nn} \end{bmatrix}$$

Solution is obvious

$$x_i = \frac{b_i}{a_{ii}}$$

If all linear equations are uncoupled:

$$A = \begin{bmatrix} a_{11} & & & \\ & a_{22} & & \\ & & \ddots & \\ & & & a_{nn} \end{bmatrix}$$

Solution is obvious

$$x_i = \frac{b_i}{a_{ii}}$$

Can we diagonalize every matrix?

If all linear equations are uncoupled:

$$A = \begin{bmatrix} a_{11} & & & \\ & a_{22} & & \\ & & \ddots & \\ & & & a_{nn} \end{bmatrix}$$

Solution is obvious

$$x_i = \frac{b_i}{a_{ii}}$$

Can we diagonalize every matrix?

$$P^{-1}AP$$

If all linear equations are uncoupled:

$$A = \begin{bmatrix} a_{11} & & & \\ & a_{22} & & \\ & & \ddots & \\ & & & a_{nn} \end{bmatrix}$$

Solution is obvious

$$x_i = \frac{b_i}{a_{ii}}$$

Can we diagonalize every matrix?

Keyword: Jordan Normal Form

$$P^{-1}AP$$

$$A = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$

Upper triangular matrix:

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ & a_{22} & \ddots & \vdots \\ & & \ddots & \vdots \\ & & & a_{nn} \end{bmatrix}$$

$$a_{ij} = 0, \forall i > j$$

$$a_{nn}x_n = b_n, \implies x_n = \frac{b_n}{a_{nn}}$$

$$a_{n-1,n-1}x_{n-1,n-1} = b_{n-1} - a_{n-1,n}x_n, \implies x_{n-1} = \frac{b_{n-1} - a_{n-1,n}x_n}{a_{n-1,n-1}}$$

Upper triangular matrix:

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ & a_{22} & \ddots & \vdots \\ & & \ddots & \vdots \\ & & & a_{nn} \end{bmatrix}$$

$$a_{ij} = 0, \forall i > j$$

$$a_{nn}x_n = b_n, \implies x_n = \frac{b_n}{a_{nn}}$$

$$a_{n-1,n-1}x_{n-1,n-1} = b_{n-1} - a_{n-1,n}x_n, \implies x_{n-1} = \frac{b_{n-1} - a_{n-1,n}x_n}{a_{n-1,n-1}}$$

Quiz: What is the final complexity?

Upper triangular matrix:

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ & a_{22} & \ddots & \vdots \\ & & \ddots & \vdots \\ & & & a_{nn} \end{bmatrix}$$

$$a_{ij} = 0, \forall i > j$$

$$a_{nn}x_n = b_n, \implies x_n = \frac{b_n}{a_{nn}}$$

$$a_{n-1,n-1}x_{n-1,n-1} = b_{n-1} - a_{n-1,n}x_n, \implies x_{n-1} = \frac{b_{n-1} - a_{n-1,n}x_n}{a_{n-1,n-1}}$$

Quiz: What is the final complexity?

Answer: $O(N^2)$

Gaussian Elimination

Exact solution,
except for rounding errors

$$A = LU$$

$$Ly = b$$

$$Ux = y$$

$$PA = LU$$

Not unique

Problems when A is singular

Solution is LUP - additional permutation matrix which aims to maximize absolute values of diagonal elements

Finding LU (LUP) has complexity $O(N^3)$

Special case: SPD matrices

Symmetric positive definite: if $A = A^T$ and $x^T A x > 0, \forall x \neq 0$

$$A = LL^*$$

Two times less memory than LU

QR decomposition

$$A = QR$$

$$Q^T Q = I$$

R is upper (right) triangular

Roughly twice more arithmetic operations than LU $\left(\frac{4n^3}{3} \text{ vs. } \frac{2n^3}{3} \right)$ but unconditional stability

Methods: modified Gram-Schmidt, Householder reflections, or Givens rotations

Conditioning

Relative error: $\frac{||x - \hat{x}||}{||x||}$

$$r = b - A\hat{x} = Ax - A\hat{x} = A(x - \hat{x})$$

$$x - \hat{x} = A^{-1}r \implies ||x - \hat{x}|| = ||A^{-1}r|| \leq ||A^{-1}|| ||r||$$

$$\frac{||x - \hat{x}||}{||x||} \leq ||A^{-1}|| ||r|| \frac{||A||}{||b||}$$

Note:


$$A\hat{x} = b + \Delta b \implies ||\Delta b|| = ||r||$$

Condition number: $k(A) = ||A|| ||A^{-1}||$

$$\frac{||x - \hat{x}||}{||x||} \leq k(A) \frac{||\Delta b||}{||b||}$$

the relative error in the solution is bounded by the condition number of the matrix A times the relative error in the residual.

Iterative methods




Stationary methods

$$Mx_t = Nx_{t-1} + b, A = M - N$$

We need initial guess for the solution: x_0

Error is: $e_0 = x - x_0$ 

We do not know the exact solution
(if we know, why do we care?) and
cannot compute

$r_0 = Ae_0 = b - Ax_0$  We can compute

$$x = x_0 + e_0 = x_0 + A^{-1}r_0$$

Set $M\tilde{e} = r_k$

Preconditioner: M

Then $x_t = x_{t-1} + \tilde{e}$

$$A = D + E + F$$

Jacobi: $M = D$

Gauss-Seidel: $M = D + E$

SOR: $M = D + \frac{1}{\omega}E$

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 & 0 \\ a_{21} & 0 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \dots & 0 \end{bmatrix} E$$

$$+ \begin{bmatrix} 0 & a_{12} & \dots & a_{1n} \\ 0 & 0 & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 0 \end{bmatrix} F$$

$$+ \begin{bmatrix} a_{11} & 0 & \dots & 0 \\ 0 & a_{22} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & a_{nn} \end{bmatrix} D$$

Nonstationary methods

$$x_t = x_{t-1} + \alpha_t p_t$$

Gradient descent (SPD only)

$$x_t = x_{t-1} + \alpha_t p_t$$

or

$$M \tilde{e} = r_t$$

$$p_k = r_k$$

$$M = \alpha_t I$$

Cast as an optimization problem via the least squares formulation, where the residual becomes the gradient (zero residual = minimum).

Gradient descent (SPD only)

$$x_t = x_{t-1} + \alpha_t p_t$$

$$p_k = r_k$$

or

$$M \tilde{e} = r_t$$

$$M = \alpha_t I$$

Conjugate gradient (SPD only)

$$x_t = x_{t-1} + \alpha_t p_t$$

$$\alpha_t = \frac{p_t^T r_t}{p_t^T A p_t}$$

$$p_t = r_t - \sum_{i < t} \frac{p_i^T A r_t}{p_i^T A p_i} p_i$$

Avoids oscillations by making steps in the direction orthogonal to all the previous ones.