## Advanced Robotics

Melnikov E. R., Markeeva L. B., Usvyatsov M. R.  ${\it April~13,~2015}$ 

## 1 Part 1

## 1.1 Problem 1




Figure 1: Problem 1 description

1. 
$${}_{1}^{0}R = \begin{pmatrix} \cos(\alpha) & \sin(\alpha) & 0 \\ -\sin(\alpha) & \cos(\alpha) & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$${}_{0}P = \begin{pmatrix} 0 \\ 0 \\ 8 \end{pmatrix}$$

$$\text{Thus} \ _{1}^{0}T = \begin{pmatrix} \cos(\alpha) & \sin(\alpha) & 0 & 0 \\ -\sin(\alpha) & \cos(\alpha) & 0 & 0 \\ 0 & 0 & 1 & 8 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

$$2. \ _{1}^{2}R = \begin{pmatrix} \cos(\alpha) & \sin(\alpha) & 0 \\ -\sin(\alpha) & \cos(\alpha) & 0 \\ 0 & 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ -1 & 0 & 0 \end{pmatrix} = \begin{pmatrix} 0 & \sin(\alpha) & \cos(\alpha) \\ 0 & \cos(\alpha & -\sin(\alpha) \\ -1 & 0 & 0 \end{pmatrix}$$

$$\text{We know, that} \ \ _{B}^{A}R = \begin{pmatrix} (^{B}_{A}R)^{T}, & \text{thus} \ \ _{2}^{1}R = \begin{pmatrix} 0 & \sin(\alpha) & \cos(\alpha) \\ 0 & \cos(\alpha & -\sin(\alpha) \\ -1 & 0 & 0 \end{pmatrix}^{T} = \begin{pmatrix} 0 & 0 & -1 \\ \sin(\alpha) & \cos(\alpha) & 0 \\ \cos(\alpha) & -\sin(\alpha) & 0 \end{pmatrix}$$

$$\text{$^{1}P = \begin{pmatrix} 5 \cdot \cos(\alpha) \\ -5 \cdot \sin(\alpha) \\ 0 \end{pmatrix}$$

$$\text{Thus} \ _{2}^{1}T = \begin{pmatrix} 0 & 0 & -1 & 5 \cdot \cos(\alpha) \\ \sin(\alpha) & \cos(\alpha) & 0 & -5 \cdot \sin(\alpha) \\ \cos(\alpha) & -\sin(\alpha) & 0 & 0 \end{pmatrix}$$

$$\text{Thus} \ _{2}^{1}T = \begin{pmatrix} 0 & 0 & -1 & 5 \cdot \cos(\alpha) \\ \sin(\alpha) & \cos(\alpha) & 0 & -5 \cdot \sin(\alpha) \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

## 1.2 Problem 2




Figure 2: Problem 2 description

1. Let us define 
$$\alpha = \arctan(0.5)$$

$$/ \cos(\alpha) - \sin(\alpha) = 0$$

$${}^{2}_{1}R = \left( \begin{array}{ccc} \cos(\alpha) & -\sin(\alpha) & 0 \\ \sin(\alpha) & \cos(\alpha) & 0 \\ 0 & 0 & 1 \end{array} \right) \cdot \left( \begin{array}{ccc} 0 & 0 & -1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{array} \right) = \left( \begin{array}{ccc} 0 & -\sin(\alpha) & -\cos(\alpha) \\ 0 & \cos(\alpha) & -\sin(\alpha) \\ 1 & 0 & 0 \end{array} \right)$$

$${}^{2}P = \left( \begin{array}{c} 2 \\ 8 \cdot \cos(\alpha) \\ 8 \cdot \sin(\alpha) \end{array} \right)$$

Thus 
$${}_{2}^{1}T = \begin{pmatrix} 0 & -\sin(\alpha) & -\cos(\alpha) & 2\\ 0 & \cos(\alpha) & -\sin(\alpha) & 8 \cdot \cos(\alpha)\\ 1 & 0 & 0 & 8 \cdot \sin(\alpha)\\ 0 & 0 & 0 & 1 \end{pmatrix}$$

$$2. \ _{2}^{3}R = \left(\begin{array}{ccc} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 1 \end{array}\right) \cdot \left(\begin{array}{ccc} 1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{array}\right) = \left(\begin{array}{ccc} 0 & 0 & -1 \\ -1 & 0 & 0 \\ 0 & 1 & 0 \end{array}\right)$$

$${}^{3}P = \left(\begin{array}{c} 2\\0\\4\sqrt{5} \end{array}\right)$$

Thus 
$${}_{2}^{3}T = \begin{pmatrix} 0 & 0 & -1 & 2 \\ -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 4\sqrt{5} \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

$$3. \ _{1}^{3}R = \left(\begin{array}{ccc} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{array}\right) \cdot \left(\begin{array}{ccc} \frac{\sqrt{2}}{2} & -\frac{\sqrt{2}}{2} & 0 \\ \frac{\sqrt{2}}{2} & \frac{\sqrt{2}}{2} & 0 \\ 0 & 0 & 1 \end{array}\right) = \left(\begin{array}{ccc} \frac{\sqrt{2}}{2} & -\frac{\sqrt{2}}{2} & 0 \\ -\frac{\sqrt{2}}{2} & -\frac{\sqrt{2}}{2} & 0 \\ 0 & 0 & -1 \end{array}\right)$$

$$^{3}P = \left( \begin{array}{c} 4 \cdot \sin(\alpha) \\ 8 \cdot \sin(\alpha) \\ 0 \end{array} \right)$$

Thus 
$${}_{1}^{3}T = \begin{pmatrix} \frac{\sqrt{2}}{2} & -\frac{\sqrt{2}}{2} & 0 & 4 \cdot \sin(\alpha) \\ -\frac{\sqrt{2}}{2} & -\frac{\sqrt{2}}{2} & 0 & 8 \cdot \sin(\alpha) \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$