





Exploring the World of Science

University of Michigan Science Olympiad 2021 Invitational Tournament

Detector Building C

Test length: 50 Minutes

Team name: KEY

Student names: KEY

Detector Building C - Detector Building C - University of Michigan Div C - 02-20-2021

This test consists of 40 questions and you will have 50 minutes to complete it.

The Tiebreakers for this test will be: Question 19, Question 25, and Question 7 in that order.

If you experience technical difficulties during the test:

Immediately contact the event supervisor through the classroom feature on Scilympiad, stating clearly what issue you are having.


If your work is not saving/submitting, take screenshots of your answers on Scilympiad and submit them to this google form (https://docs.google.com/forms/d/19cRQLafN7EARRS7tZ18HOCt4B1F-4fYOSliON1kro/viewform?edit_requested=true). Try to stay within your allotted 50 minutes.

This test contains various types of questions, make sure you read the entire question!

1. (1.00 pts) Which law is Gay Lussac known for?
A) If a gas's pressure decreases, then so does its temperature if the mass and volume of the gas are held constant.
O B) If a gas's temperature decreases, then its pressure increases if the mass and volume of the gas are held constant.
O C) If a gas's temperature increases, then so does its pressure if the mass of the gas is held constant.
O D) If a gas's temperature increases, then so does its pressure if the size of the gas is held constant.
2. (1.00 pts) A famous physicist with the first name Daniel is known for his work developing a temperature scale. Which one is it?
A) The Fahrenheit Scale
O B) The Kelvin Scale
C) The Celsius Scale
Op) The Rankine Scale
3. (2.00 pts) What is 0 degrees Fahrenheit in Kelvin? (Nearest whole number, do not include units)
255
4. (1.00 pts) What is the coldest temperature possible?
O A) Total zero
B) Absolute zero
O C) Absolute freezing
O D) Complete zero
O E) The critical point
5. (1.00 pts) What is an NTC thermistor?
A thermally sensitive resistor whose resistance distinctively decreases (a precise and predictable change) as the temperature at the core increases.

C) A resistor that changes colors based on the temperature at its core.
O D) A thermistor that changes size significantly based on the temperature at its core.
O E) A thermally sensitive resistor whose resistance decreases (by subtle margins that require estimations) as the temperature at the core decreases.
6. (1.00 pts) What is not true about the Light Emitting Diode?
A) When buying an LED bulb, you will notice they cost less than incandescent bulbs.
O B) They hold the title for producing the longest-lasting and most efficient lighting available today.
O C) They work great with solar powered systems since they consume low amounts of energy.
O D) They can be made very compact so they can fit anywhere.
O E) Display screens can be made out of them.
7. (1.00 pts) Tie Breaker 3 How much more energy is wasted in an incandescent than an LED?
O B) 40% of power goes to heat in an incandescent and 5% goes to heat in an LED
C) 80% of power goes to heat in an incandescent and 0% goes to heat in an LED
O D) 30% of power goes to heat in an incandescent and 80% goes to heat in an LED
8. (1.00 pts) What is the measurement of light based off of (where a gram – the measurement of mass – is "based off of" 1 cubic cm of water at 4 degrees Celcius)? This is the SI unit of luminous intensity.
O A) 1 Small LED
O B) 1 Florescent Lightbulb
C) 1 Candle
O D) 1 Spark
O E) 1 Firework
9. (1.00 pts) In a substance, temperature is the average energy of the moving particles.
O A) Potential
O B) Frictional
O C) Thermal
C) Thermal D) Nuclear
O D) Nuclear
O D) Nuclear
○ D) Nuclear ■ E) Kinetic
 ○ D) Nuclear ○ E) Kinetic 10. (1.00 pts) Which of the following are ways to increase water's boiling point?
 ○ D) Nuclear ● E) Kinetic 10. (1.00 pts) Which of the following are ways to increase water's boiling point? ○ A) Adding more water
 ○ D) Nuclear ○ E) Kinetic 10. (1.00 pts) Which of the following are ways to increase water's boiling point? ○ A) Adding more water ○ B) Removing some water

11. (2.00 pts) Calculate the Total Resistance of this circuit where r is 2 Ohms. Round to the nearest whole number and do not include units. (Free Response)




12. (1.00 pts)

Two cylindrical resistors are made from the same material and have equal diameters but the first resistor (resistor A) has length L and the second resistor (resistor B) has length 2L. I the current is the same through both, compare the voltages across them. (VA is voltage across resistor A and VB is voltage across resistor B)

- O A) VA > VB
- B) _{VA = VB}
- © C) VA < VB</p>


13. (3.00 pts)

In a circuit there are two identical batteries with voltage V = 9 Volts and each battery is in series with a resistor Rb = 7.87 Ohms. Another resistor, Re is rated at 9.06 Ohms. What is the magnitude of current through Re? Round to the nearest hundredth place and do not include units. (Free Response)




0.69

14. (2.00 pts) Four identical bulbs are placed in a circuit. If D burns out (no current flows through D), what will happen to the brightness of bulb C?



- A) C gets brighter
- O B) C gets dimmer
- O C) C stays the same

15. (2.00 pts) Four identical bulbs are placed in a circuit. If D burns out (no current flows through D), what will happen to the Voltage across C?



- A) V stays the same
- O B) V decreases
- O C) V increases

16. (1.00 pts) What is NOT a practical use for an LED?

- O A) Can be used to measure speed
- O B) Can be used in remotes to transmit information
- O C) Can be used to check if code is working
- O) Can be used to store energy

17. (1.00 pts) What can an LED do? Select all that apply.

(Mark ALL correct answers)

- $\ \square$ A) Cause a drop in current
- B) Cause a drop in voltage
- C) Cause current to only flow in one direction


□ D) Store energy that can be used later by the circuit
Store energy as light that can be used later by the circuit
18. (1.00 pts) In a DC circuit, what does adding resistors before an LED do?
A) Makes the LED dimmer
O B) Makes the LED brighter
C) LED brightness will remain the same
19. (2.00 pts) Tie Breaker 1 What is the Steinhart-Hart equation?
Converts a reading of resistance to a value measured in Kelvin
O B) Converts a reading of resistance to a value measured in Celsius
O C) Converts a reading of current to a value measured in Kelvin
O D) Converts a reading of voltage to a temperature value
20. (2.00 pts) What happens at 0 Kelvin?
O A) Water freezes
O B) Things start to melt
O C) Water boils
All molecular motion stops
21. (1.00 pts) There is a thermometer that consists of a bulb containing a liquid that expands into a capillary tube. The current liquid is replaced with another (same volume) that expands more fo the same rise in temperature. The new thermometer will have
21. (1.00 pts) There is a thermometer that consists of a bulb containing a liquid that expands into a capillary tube. The current liquid is replaced with another (same volume) that expands more fo the same rise in temperature. The new thermometer will have
21. (1.00 pts) There is a thermometer that consists of a bulb containing a liquid that expands into a capillary tube. The current liquid is replaced with another (same volume) that expands more for the same rise in temperature. The new thermometer will have O A) More sensitivity and a larger range
21. (1.00 pts) There is a thermometer that consists of a bulb containing a liquid that expands into a capillary tube. The current liquid is replaced with another (same volume) that expands more fo the same rise in temperature. The new thermometer will have
21. (1.00 pts) There is a thermometer that consists of a bulb containing a liquid that expands into a capillary tube. The current liquid is replaced with another (same volume) that expands more for the same rise in temperature. The new thermometer will have A) More sensitivity and a larger range B) Less sensitivity and a larger range
21. (1.00 pts) There is a thermometer that consists of a bulb containing a liquid that expands into a capillary tube. The current liquid is replaced with another (same volume) that expands more for the same rise in temperature. The new thermometer will have One is a thermometer that consists of a bulb containing a liquid that expands into a capillary tube. The current liquid is replaced with another (same volume) that expands more for the same rise in temperature. The new thermometer will have One is a sensitivity and a larger range One is a sensitivity and a larger range One is a sensitivity and less range
21. (1.00 pts) There is a thermometer that consists of a bulb containing a liquid that expands into a capillary tube. The current liquid is replaced with another (same volume) that expands more for the same rise in temperature. The new thermometer will have One is a thermometer that consists of a bulb containing a liquid that expands into a capillary tube. The current liquid is replaced with another (same volume) that expands more for the same rise in temperature. The new thermometer will have One is a sensitivity and a larger range One is a sensitivity and a larger range One is a sensitivity and less range
21. (1.00 pts) There is a thermometer that consists of a bulb containing a liquid that expands into a capillary tube. The current liquid is replaced with another (same volume) that expands more for the same rise in temperature. The new thermometer will have A) More sensitivity and a larger range B) Less sensitivity and a larger range C) More sensitivity and less range D) Less sensitivity and less range 10 Less sensitivity and less range
21. (1.00 pts) There is a thermometer that consists of a bulb containing a liquid that expands into a capillary tube. The current liquid is replaced with another (same volume) that expands more for the same rise in temperature. The new thermometer will have A) More sensitivity and a larger range B) Less sensitivity and a larger range C) More sensitivity and less range D) Less sensitivity and less range 10 D) Less sensitivity and less range
21. (1.00 pts) There is a thermometer that consists of a bulb containing a liquid that expands into a capillary tube. The current liquid is replaced with another (same volume) that expands more for the same rise in temperature. The new thermometer will have O A) More sensitivity and a larger range O B) Less sensitivity and a larger range O C) More sensitivity and less range O D) Less sensitivity and less range 22. (1.00 pts) In a liquid-in-glass thermometer, temperature shown increases. What is constant?
21. (1.00 pts) There is a thermometer that consists of a bulb containing a liquid that expands into a capillary tube. The current liquid is replaced with another (same volume) that expands more for the same rise in temperature. The new thermometer will have O A) More sensitivity and a larger range O B) Less sensitivity and less range O D) Less sensitivity and less range 22. (1.00 pts) In a liquid-in-glass thermometer, temperature shown increases. What is constant? A) Mass of liquid O B) Density of liquid O C) Volume of liquid
21. (1.00 pts) There is a thermometer that consists of a bulb containing a liquid that expands into a capillary tube. The current liquid is replaced with another (same volume) that expands more for the same rise in temperature. The new thermometer will have O A) More sensitivity and a larger range O B) Less sensitivity and a larger range O C) More sensitivity and less range O D) Less sensitivity and less range 22. (1.00 pts) In a liquid-in-glass thermometer, temperature shown increases. What is constant?
21. (1.00 pts) There is a thermometer that consists of a bulb containing a liquid that expands into a capillary tube. The current liquid is replaced with another (same volume) that expands more for the same rise in temperature. The new thermometer will have O A) More sensitivity and a larger range O B) Less sensitivity and less range O D) Less sensitivity and less range 22. (1.00 pts) In a liquid-in-glass thermometer, temperature shown increases. What is constant? A) Mass of liquid O B) Density of liquid O C) Volume of liquid
21. (1.00 pts) There is a thermometer that consists of a bulb containing a liquid that expands into a capillary tube. The current liquid is replaced with another (same volume) that expands more for the same rise in temperature. The new thermometer will have O A) More sensitivity and a larger range O B) Less sensitivity and less range O D) Less sensitivity and less range 22. (1.00 pts) In a liquid-in-glass thermometer, temperature shown increases. What is constant? A) Mass of liquid O B) Density of liquid O C) Volume of liquid
21. (1.00 pts) There is a thermometer that consists of a bulb containing a liquid that expands into a capillary tube. The current liquid is replaced with another (same volume) that expands more for the same rise in temperature. The new thermometer will have A) More sensitivity and a larger range B) Less sensitivity and less range C) More sensitivity and less range D) Less sensitivity and less range 12. (1.00 pts) In a liquid-in-glass thermometer, temperature shown increases. What is constant? A) Mass of liquid B) Density of liquid C) Volume of liquid D) The kinetic energy of the molecules
21. (1.00 pts) There is a thermometer that consists of a bulb containing a liquid that expands into a capillary tube. The current liquid is replaced with another (same volume) that expands more for the same rise in temperature. The new thermometer will have O A) More sensitivity and a larger range O B) Less sensitivity and less range O D) Less sensitivity and less range 22. (1.00 pts) In a liquid-in-glass thermometer, temperature shown increases. What is constant? A) Mass of liquid O D) The kinetic energy of the molecules

C) Thermocouple Thermometer D) Solid Core Thermometer
24. (1.00 pts)
All thermometers must be calibrated to ensure they function properly. If you must do this without another thermometer, you may need a substance of a temperature known to you so you can compare it with the reading of the thermometer. What is needed:
O A) A high fixed-point temperature
O B) A low fixed-point temperature
C) A low and high fixed-point temperature
O D) No fixed-point temperatures are required for calibration
25. (1.00 pts) Tie Breaker 2 How does an Infrared Thermometer work?
Collects infrared light from molecules and converts it to heat.
O B) The beam allows the sensor to feel the surface, kind of like it was an extension.
C) It measures the heat of the infrared beam shot from the thermometer.
O D) It measures the heat in between the detector and object and makes a calculation based on distance.
26. (3.00 pts) Cold objects (above 0 K) will always
A) Emit light
O B) Cool other objects
O C) Contract
O D) Expand
27. (1.00 pts) What state of matter do modern thermometers rely on the expansion and contraction of to calculate the temperature?
O A) Gas
O B) Solid
C) Fluid
O D) Some combination of them
28. (1.00 pts) What is not a type of resistance varying resistor (variable resistor)?
O A) Humistor
O B) Photoresistor
C) Potentiometer
Grid resistor
20 (4 00 de) What adaptively and a self-bate a selection of the self-bate and self-bat
29. (1.00 pts) What color would you use to calibrate a color sensor? (Use all lower case letters)
white

30. (1.00 pts) What does LED stand for? (Use all lower case letters)

light emitting diode

31. (2.00 pts) Four identical bulbs are placed in a circuit. If A burns out (no current flows through A), what will happen to the Voltage across C?



- O A) V stays the same
- B) V decreases
- O C) V increases

32. (3.00 pts)

There is a battery, LED, and a resistor wired in series in a circuit. What is the needed resistance of the resistor to prevent the LED from burning out? The ideal current for the LED is 28mA, VLED is 2.2V, and VBattery is 9V. Round to the nearest whole number if needed and do not include units.


243

33. (1.00 pts)

You have many LEDs, all rated for 1.5V and 2.0mA, but you do not have any resistors. With a 9V battery, what is the minimum number of LEDs you would have to use to ensure you do not burn out your LEDs?

6

34. (2.00 pts) Calculate the voltage of the battery if the current through the battery is 3 Amps. r = 2 Ohms. Do not include units or decimals. (Free Response)



25 /4 00 mtm) Enter the three generally assented body temperatures considered healthy. No desimals
35. (1.00 pts) Enter the three generally accepted body temperatures considered healthy. No decimals.
97 98 99
36. (1.00 pts) Daniel Fahrenheit was responsible for inventing which thermometer?
O A) Alcohol-in-glass thermometer
B) Mercury-in-glass thermometer
O C) Solid Core thermometer
O D) Digital thermometer
37. (2.00 pts) What was the earliest known thermometer called?
O A) Thermometer
B) Thermoscope
O C) Temperature Detector
O D) Temperature Finder
O E) Detector
38. (1.00 pts) What was the first standard temperature scale?
O A) Celsius
O B) Kelvin
O C) Henry
D) Fahrenheit
39. (10.00 pts) In the next few questions, you will answer the following: If you were competing in person, how would you go about making the detector?
Part 1:
Describe the physical components of how you would build your detector. Include describing electrical components and how you would make your thermometer probe.
Expected Answer: +10 (Based on specificity) for describing how to construct the thermometer and protect the thermistor from water. +2 for mentioning a circuit board, +2 for connecting it to a thermistor, +2 for including LEDs, and up to +4 if water proofing of thermometer tip is adequate (scaling could be: epoxy/resins that harden to become a plastic sea would be considered very reliable (+4), ceran wrap has resulted in some leaks so somewhat reliable(+2.5), and tape doesn't really work so not very reliable(+1)). Example of one wit full points: To make the detector we would start off with an Arduino circuit board. Connected to the board is an NTC thermistor, located at the tip of a 3D printed probe. To ensure that water does not enter the tip of the probe we will coat the tip and the gap between it and the body of the probe with epoxy so that water does not enter and interfere with the electronics. Also, we will use LEDs with our Arduino to signal the temperature range.

40. (8.00 pts) Part 2:

Describe the process you would use to convert the information from the sensor to temperature (without using the Steinhart-Hart equation or program libraries). Be specific!	
Expected Answer: +8 total (Based on specificity) for describing the process of creating a model to convert resistance from thermistor to readable temperature. +2 for measuring resistance output when putting probe into different temperatures of liquid, +2 for creating graph and making a line of best fit, +4 for using the line of best fit formula to convert resistance at any value to a temperature. Example: We would use our probe in water of various temperatures (about 10 different ones) including 0 Celsius and 100 Celsius and down the resistance value at each temperature. Then, we would graph our resistance values and the temperature at which they were recorded and create a line of best fit for tidata points. Then, we would code our microcontroller to use the formula for the line of best fit to convert resistance readings to temperature in Celsius.	d write
Congratulations on completing the University of Michigan 2021 Invitational Detector Building test!	
If you have any questions or concerns pertaining to this event, please email tec.umichscioly@umich.edu, and we will try to get back to you as soon as we can.	

© 2021 - powered by Scilympiad (https://scilympiad.com) | Tournaments (/umso/Home/Subscribers)

Terms of Use (/ToS.html) | **Privacy (/Privacy.html)** | Cookies Policy (/CookiePolicy.html) | Support (/umso/Support) | Contact (/umso/Home/Contact)