

Age/Gender :- 35 Year (M)
Referred By :- Dr. GAURAV ANAND "M.B.B.S" "D.T.C.D"

SEROLOGY EXAMINATION

Investigations	Observed Value	Unit	Reference - Range
Malaria Pf/Pv Rapid Card	Negative		Negative
Plasmodium Vivax (PV)	Negative		Negative
Plasmodium Falciparum (PF)			

Method/Technology:- Card/Strip:- J. Mitra & Co. Pvt. Ltd. Batch:- 042228 Expiry:- August 2025

Interpretation:-

Malaria is caused by the malarial parasite belonging to genus plasmodium. Malaria or Plasmodium has following types : Plasmodium vivax (Tertian Malaria) is the most common in all present in the tropics, subtropics, and the temperate regions, Plasmodium malariae (Quartan Malaria) found in asia and south America. Most commonly seen in tropical Africa, Plasmodium falciparum (malignant tertian malaria) is found in tropical and subtropical area, Plasmodium ovale (ovale tertian malaria) found in subtropical and temperate area.

Once the malarial parasite enters the body through blood, then they enter the liver where they mature. In the liver after several days mature parasite enters the bloodstream and then in RBCs. In the RBCs, the parasite multiply in 48 to 72 hours and then RBCs rupture and then continue to infect other RBCs. During this period of RBCs infection, patient develops symptoms and last for 2 to 3 days.

Continue To Next Page