

# The Sempaevum: A Lossless Mathematical Rendering of the Totality $\Sigma$ Derived from Three Primitive Categories

Michael James Muller

*Aevum Defluo*

Single researcher. No co-authors, no institutional affiliation, no external funding.

April 2026

## Abstract

This paper identifies, characterises, and verifies a single mathematical object — the **SEMPAEVUM** — as the lossless rendering of the totality  $\Sigma$  on the multiplicative manifold  $(\mathbb{R}^+, \times) \times (U(1), \times)$ . The object is derived forward from three irreducible primitive categories  $(P, D, T)$  with disjoint cardinalities  $\Omega, n, [0/0]$  under a single master identity  $3 = 3 = 3 = \Sigma$ . It simultaneously (i) classifies every positive ratio into a sublattice family ( $d = N / \gcd(|k|, N)$ ), the static gcd-classification of an individual lattice coordinate, six values at  $N = 12$ ) and identifies the per-axis harmonic-family layer of structural modes ( $d \in \{1, \dots, 12\}$  per axis, six **SIMPLE** plus six **COMPLEX**, twenty-four total on  $\mathcal{L}_{\mathbb{C}}$ , discovered and verified by the palindromic cascade) connected to the sublattice-family layer through the Sublattice Visitation Theorem (multiplicities  $\varphi(d)$ , sum  $N$  via Gauss); (ii) generates its own refinement tower under the doubling law  $\tau(N_\ell) = 6 \cdot 2^\ell$ ; (iii) passes three independent minimality tests (Webb’s  $n$ -valued stroke 1935, the palindromic divisor cascade, and the EML operator of Odrzywołek 2026) at the unique resolution  $N = 12$ , giving discrete-logical, discrete-multiplicative, and continuous-elementary minimal generators at one scale; (iv) projects its own defining constants  $\{N, 1/N, K, 1/K\}$  onto a single sublattice position ( $d = 12, |\varepsilon| = 1.955 \text{ ¢}$ ) — the Koide attractor — verifying itself by self-consistency; (v) hosts mathematics as a domain, placing Zermelo–Fraenkel set theory exactly at  $d = 1$  octave with  $\varepsilon = 0$ , Chaitin’s  $\Omega$  at  $d = 1$  octave via non-limiting Path D, Gödel sentences via integrative-level classification, and large cardinals via consistency-strength; (vi) handles essentially-infinite and non-computable objects without limits through four sub-paths corresponding to each primitive’s own face of infinity; (vii) is lossless — the representation  $r \mapsto (k, d, \varepsilon)$  is a bijection at every finite resolution and pullback-exact as  $N \rightarrow \infty$  along the LCM tower. Empirical fingerprints presented include the Koide ratio to six parts per million, the fine-structure constant to sub-parts-per-billion under a closed-form identity  $\alpha^{-1} = 137 + \sqrt{3}/48 - \sqrt{3}/(93312\pi^2) - 1/(216(18\pi - 1))$  agreeing with the CODATA 2022 recommended value to within  $0.46\sigma$  ( $\sim 7$  parts in  $10^{11}$ ), twelve-tone equal temperament’s Pythagorean comma as an intrinsic lattice quantity, and the step counts of the citric acid and glycolytic cycles. The sublattice structure derives the physical content of the Standard Model: the adjoint formula  $\dim(\text{SU}(d)) = d^2 - 1$  from the Subsumption Law, the critical dimensions  $D_{\text{superstring}} = 10$  by two independent routes and  $D_{\text{M-theory}} = 11$  by four, and the uniqueness of the gauge group  $\text{SU}(3) \times \text{SU}(2) \times \text{U}(1)$  as the sole partition of  $N = 12$  gauge bosons into native-sublattice simple and abelian factors (the N-Exhaustion Theorem). The construction involves zero tunable parameters, zero data-fitting, and zero external axioms. Every constant is forced.

## Contents

|          |                                                              |          |
|----------|--------------------------------------------------------------|----------|
| <b>1</b> | <b>Introduction</b>                                          | <b>6</b> |
| 1.1      | Authorship, tools, and representational disclosure . . . . . | 6        |

|          |                                                                                                                |           |
|----------|----------------------------------------------------------------------------------------------------------------|-----------|
| 1.2      | The object identified in this paper . . . . .                                                                  | 7         |
| 1.3      | Etymology of the name <i>Sempaevum</i> . . . . .                                                               | 8         |
| 1.4      | The empirical regularity the object explains . . . . .                                                         | 9         |
| 1.5      | The thesis . . . . .                                                                                           | 10        |
| 1.6      | Relation to existing work . . . . .                                                                            | 11        |
| 1.7      | Scope and organisation . . . . .                                                                               | 11        |
| <b>2</b> | <b>The Founding Axiom and the primitive Cardinals</b>                                                          | <b>12</b> |
| 2.1      | The Founding Axiom . . . . .                                                                                   | 12        |
| 2.2      | Discovery and historical origin . . . . .                                                                      | 12        |
| 2.3      | The master equation . . . . .                                                                                  | 13        |
| 2.4      | The three Cardinals . . . . .                                                                                  | 13        |
| 2.4.1    | The two faces of indeterminate cardinality . . . . .                                                           | 13        |
| 2.5      | Three disjoint infinities . . . . .                                                                            | 14        |
| 2.6      | Intrinsic mediation . . . . .                                                                                  | 14        |
| 2.7      | The Nesting Principle . . . . .                                                                                | 15        |
| 2.7.1    | The Absolutes are actual, not limits . . . . .                                                                 | 16        |
| 2.7.2    | Structural consequences of nesting . . . . .                                                                   | 16        |
| 2.7.3    | Substrate Potential Principle . . . . .                                                                        | 16        |
| 2.7.4    | The Omniscient Traverser and the uniqueness of agency . . . . .                                                | 16        |
| 2.8      | The totality $\Sigma$ . . . . .                                                                                | 17        |
| 2.8.1    | Pure Relationalism . . . . .                                                                                   | 17        |
| 2.8.2    | Impossibility of absolute nothing . . . . .                                                                    | 18        |
| 2.9      | The binding order . . . . .                                                                                    | 18        |
| 2.10     | Modes of self-presentation . . . . .                                                                           | 18        |
| 2.11     | The four manifold states . . . . .                                                                             | 19        |
| 2.11.1   | Unsubstantiated is blank, not zero . . . . .                                                                   | 19        |
| 2.12     | Variance and the grounding function . . . . .                                                                  | 20        |
| 2.13     | Geometric content of the primitives' operational manifolds . . . . .                                           | 22        |
| 2.13.1   | Structural necessity of geometry and the pre-geometric primitives . . . . .                                    | 23        |
| <b>3</b> | <b>The triple identity and the three operational tools</b>                                                     | <b>24</b> |
| 3.1      | Three simultaneous categorical readings . . . . .                                                              | 24        |
| 3.2      | The universality anchor $3 = 3 = 3 = \Sigma$ versus the composition equation $P \circ D \circ T = E$ . . . . . | 25        |
| 3.3      | The three operational tools . . . . .                                                                          | 26        |
| 3.4      | The Identification Principle . . . . .                                                                         | 26        |
| 3.5      | The Descriptor Gap Principle . . . . .                                                                         | 27        |
| 3.6      | The Subsumption Law . . . . .                                                                                  | 28        |
| 3.7      | The Exception $E$ . . . . .                                                                                    | 29        |
| 3.7.1    | Grounding and maximal possibility . . . . .                                                                    | 31        |
| 3.7.2    | The shimmering manifold . . . . .                                                                              | 31        |
| 3.8      | The Mediation $M$ . . . . .                                                                                    | 32        |
| 3.9      | The Incoherence $I$ . . . . .                                                                                  | 34        |
| 3.9.1    | Concrete examples of incoherent configurations . . . . .                                                       | 34        |
| 3.9.2    | Dual bridging . . . . .                                                                                        | 35        |
| 3.10     | The Anti-Emergence Principle and the triadic non-emergent structure . . . . .                                  | 37        |
| 3.11     | The closed methodological loop . . . . .                                                                       | 39        |
| 3.12     | Integrative levels . . . . .                                                                                   | 39        |

|          |                                                                                 |           |
|----------|---------------------------------------------------------------------------------|-----------|
| <b>4</b> | <b>The three temporal aspects</b>                                               | <b>40</b> |
| 4.1      | The three temporal aspects . . . . .                                            | 40        |
| 4.2      | The temporal master equation . . . . .                                          | 41        |
| 4.3      | Uniqueness of the temporal Exception . . . . .                                  | 41        |
| 4.4      | $D$ -time vs $T$ -time: complete comparison . . . . .                           | 42        |
| 4.5      | The Minkowski interval . . . . .                                                | 42        |
| 4.5.1    | The Lorentz factor as the $D$ -time/ $T$ -time ratio . . . . .                  | 43        |
| 4.5.2    | Temporal consciousness . . . . .                                                | 43        |
| 4.6      | Event-horizon reconciliation . . . . .                                          | 43        |
| 4.7      | Scope of this section . . . . .                                                 | 44        |
| <b>5</b> | <b>The horizon: thermodynamics, information, and structural cells</b>           | <b>44</b> |
| 5.1      | Surface gravity as the $D$ -time/ $T$ -time descriptor-gap gradient . . . . .   | 45        |
| 5.2      | The $U(1)$ period of $T$ -time . . . . .                                        | 46        |
| 5.3      | The Bogoliubov ratio as half- $U(1)$ -period analytic continuation . . . . .    | 48        |
| 5.4      | The Planck spectrum from the descriptor-quantum/variance ratio . . . . .        | 49        |
| 5.5      | Fermion statistics from the forbidden $\{P, T\}$ state . . . . .                | 51        |
| 5.6      | Information preservation through the Multifold birth triad . . . . .            | 51        |
| 5.7      | The critical mass $M_{\text{crit}}$ (tower self-identity) . . . . .             | 53        |
| 5.8      | Canonical structurally-stable mass at $k_r = -53$ . . . . .                     | 54        |
| 5.9      | Structural dichotomy of black-hole masses . . . . .                             | 55        |
| 5.10     | Sub-Planck masses: lossless on the LCM tower . . . . .                          | 56        |
| 5.11     | Force Quadrant Grid placement and sublattice-transition forward predictions . . | 57        |
| <b>6</b> | <b>The forced value <math>N = 12</math></b>                                     | <b>59</b> |
| 6.1      | Derivation I: product of the two counts . . . . .                               | 59        |
| 6.2      | Derivation II: least common multiple of the small sublattice orders . . . . .   | 59        |
| 6.3      | Derivation III: the stability window . . . . .                                  | 60        |
| 6.4      | The three manifold constants . . . . .                                          | 60        |
| 6.5      | A derived fine-structure constant . . . . .                                     | 61        |
| <b>7</b> | <b>The lattice on the multiplicative manifold</b>                               | <b>61</b> |
| 7.1      | The projection formula . . . . .                                                | 61        |
| 7.2      | Interpretation . . . . .                                                        | 61        |
| 7.3      | The annihilation boundary . . . . .                                             | 62        |
| 7.4      | Convention independence . . . . .                                               | 62        |
| 7.5      | Three canonical projections . . . . .                                           | 63        |
| 7.6      | The Anti-Numerology Protocol . . . . .                                          | 63        |
| 7.7      | The V-threshold significance criterion . . . . .                                | 64        |
| <b>8</b> | <b>Sublattice family structure</b>                                              | <b>65</b> |
| 8.1      | The divisor classification . . . . .                                            | 65        |
| 8.2      | The totient multiplicity . . . . .                                              | 65        |
| 8.3      | Physical reading of the six families . . . . .                                  | 66        |
| 8.4      | Classification theorem . . . . .                                                | 66        |
| 8.5      | Magical impedance: per-family coupling strength . . . . .                       | 67        |
| 8.6      | Harmonic families: the per-axis structural-mode layer . . . . .                 | 67        |
| 8.7      | The Sublattice Visitation Theorem . . . . .                                     | 68        |
| 8.8      | Shadow forces: sublattice families excluded at base $N = 12$ . . . . .          | 70        |

|           |                                                                                                 |           |
|-----------|-------------------------------------------------------------------------------------------------|-----------|
| <b>9</b>  | <b>Physical sublattice structure: gauge theory, critical dimensions, and the Standard Model</b> | <b>71</b> |
| 9.1       | The adjoint formula from the Subsumption Law . . . . .                                          | 71        |
| 9.2       | The N-Exhaustion Theorem: uniqueness of the Standard Model gauge group . .                      | 72        |
| 9.3       | Critical dimensions via the Koide Reciprocal Identity . . . . .                                 | 73        |
| 9.4       | Critical dimensions via the division algebra route . . . . .                                    | 74        |
| 9.5       | $D = 11$ for M-theory: four convergent routes . . . . .                                         | 74        |
| 9.6       | The $P \circ D \circ T$ Scale Identity . . . . .                                                | 75        |
| 9.7       | The Sublattice-to-Force Map . . . . .                                                           | 75        |
| 9.8       | Supercharges, gauge group dimensions, and brane structure . . . . .                             | 76        |
| 9.9       | The Structural Significance Principle . . . . .                                                 | 77        |
| <b>10</b> | <b>The LCM tower of refinements</b>                                                             | <b>77</b> |
| 10.1      | When $N = 12$ is insufficient . . . . .                                                         | 77        |
| 10.2      | Canonical refinements . . . . .                                                                 | 78        |
| 10.3      | Unification of the two readings . . . . .                                                       | 78        |
| 10.4      | The continuous limit . . . . .                                                                  | 78        |
| 10.4.1    | Structural necessity of asymptotic precision . . . . .                                          | 79        |
| 10.5      | Correspondence with integrative levels . . . . .                                                | 79        |
| <b>11</b> | <b>The complex lattice</b>                                                                      | <b>83</b> |
| 11.1      | Imaginary-axis projection . . . . .                                                             | 83        |
| 11.2      | Complex projection and the Gaussian integer lattice . . . . .                                   | 83        |
| 11.3      | The D–T gradient . . . . .                                                                      | 84        |
| 11.4      | The Gaussian-prime correspondence . . . . .                                                     | 84        |
| <b>12</b> | <b>The 24-family catalog and the 144-cell Force Quadrant Grid</b>                               | <b>85</b> |
| 12.1      | The 12 real-axis FORCE families and the 12 imaginary-axis PHASE families . .                    | 85        |
| 12.2      | The 42 combined off-axis families . . . . .                                                     | 86        |
| 12.3      | The 144-cell Force Quadrant Grid . . . . .                                                      | 86        |
| <b>13</b> | <b>Cascade dynamics and stability</b>                                                           | <b>87</b> |
| 13.1      | Cascade generators . . . . .                                                                    | 87        |
| 13.2      | Cascade stability asymmetry and the indeterminacy of imaginary harmonic families                | 88        |
| 13.3      | The palindromic cascade . . . . .                                                               | 89        |
| 13.4      | Use of the palindromic cascade . . . . .                                                        | 90        |
| 13.5      | The General Cascade Rule . . . . .                                                              | 90        |
| 13.6      | Unit-generator equivalence and the canonical $N = 12$ cascade . . . . .                         | 91        |
| 13.7      | Multifold cyclic closure . . . . .                                                              | 91        |
| <b>14</b> | <b>The <math>\partial I</math> lattice-aware fractal</b>                                        | <b>92</b> |
| 14.1      | The lattice-adaptive iteration map . . . . .                                                    | 92        |
| 14.2      | The lattice-adaptive exponent $p(z_n, n)$ . . . . .                                             | 93        |
| 14.3      | The shimmer modulation $\Psi_n$ . . . . .                                                       | 93        |
| 14.4      | The all-families perturbation $\varepsilon(z_n)$ . . . . .                                      | 94        |
| 14.5      | Distance estimation . . . . .                                                                   | 95        |
| 14.6      | Classification . . . . .                                                                        | 95        |
| 14.7      | Structural summary . . . . .                                                                    | 95        |

|           |                                                                                                              |            |
|-----------|--------------------------------------------------------------------------------------------------------------|------------|
| <b>15</b> | <b>Three minimal backbones at <math>N = 12</math></b>                                                        | <b>96</b>  |
| 15.1      | PDT decomposition of the projection formula . . . . .                                                        | 96         |
| 15.2      | EML: the continuous- $D$ minimal generator (Odrzywółek 2026) . . . . .                                       | 96         |
| 15.3      | Cross-domain appearance of $n_{\max, \theta} = 2$ . . . . .                                                  | 98         |
| 15.4      | Webb 1935: the discrete-logical minimal generator at $n = 12$ . . . . .                                      | 99         |
| 15.5      | The palindromic cascade: the discrete-multiplicative minimal generator on $\{d :$<br>$d \mid 12\}$ . . . . . | 100        |
| 15.6      | The triple minimal-backbone theorem . . . . .                                                                | 100        |
| <b>16</b> | <b>The Four Projection Paths and essential-infinity handling</b>                                             | <b>101</b> |
| 16.1      | The four paths . . . . .                                                                                     | 101        |
| 16.2      | Subsumption of user inputs . . . . .                                                                         | 102        |
| <b>17</b> | <b>Mathematics as a domain of the Sempaevum</b>                                                              | <b>103</b> |
| 17.1      | The $P$ , $D$ , $T$ of mathematics . . . . .                                                                 | 103        |
| 17.2      | Axiom-count projections of standard formal systems . . . . .                                                 | 103        |
| 17.3      | Chaitin's $\Omega$ via Path D.P . . . . .                                                                    | 104        |
| 17.4      | Gödel sentences via integrative-level classification . . . . .                                               | 104        |
| 17.5      | Large cardinals via consistency-strength . . . . .                                                           | 104        |
| 17.6      | The universality statement for mathematics . . . . .                                                         | 104        |
| <b>18</b> | <b>The Mathematical Rosetta Stone</b>                                                                        | <b>105</b> |
| 18.1      | Why mathematics works . . . . .                                                                              | 105        |
| 18.2      | L'Hôpital's Rule as the $T$ -navigation algorithm . . . . .                                                  | 105        |
| 18.3      | Limits as traversal operations . . . . .                                                                     | 106        |
| 18.4      | Functions as Descriptor fields . . . . .                                                                     | 106        |
| 18.5      | Derivatives as Descriptor gradients . . . . .                                                                | 106        |
| 18.6      | Integrals as Traverser accumulation . . . . .                                                                | 106        |
| 18.7      | Continuity, differentiability, and discontinuity types . . . . .                                             | 106        |
| 18.8      | Complex numbers and the $D_T$ axis . . . . .                                                                 | 107        |
| 18.9      | Mathematical operators as Traversers . . . . .                                                               | 107        |
| 18.10     | Differential equations as manifold dynamics . . . . .                                                        | 107        |
| 18.11     | The infinity hierarchy as $P$ -structure . . . . .                                                           | 107        |
| 18.12     | Probability as Unsubstantiated superposition . . . . .                                                       | 107        |
| 18.13     | The quantum mechanics mapping . . . . .                                                                      | 108        |
| 18.14     | Matrix algebra as Descriptor transformations . . . . .                                                       | 108        |
| 18.15     | Topology as configuration boundaries . . . . .                                                               | 108        |
| 18.16     | Set theory and the configuration power set . . . . .                                                         | 108        |
| 18.17     | Symmetry and group theory at $N = 12$ . . . . .                                                              | 108        |
| 18.18     | Mathematical constants from ET . . . . .                                                                     | 109        |
| 18.19     | The complete Rosetta Stone chart . . . . .                                                                   | 110        |
| <b>19</b> | <b>Self-consistency: the Koide attractor</b>                                                                 | <b>110</b> |
| 19.1      | Losslessness theorem . . . . .                                                                               | 111        |
| <b>20</b> | <b>The Complete Gaze Equation</b>                                                                            | <b>112</b> |
| 20.1      | The unification: $\text{Gain}(T) = F_w$ . . . . .                                                            | 112        |
| 20.2      | Variance collapse and threshold classification . . . . .                                                     | 112        |
| 20.2.1    | Dual derivation of the detection threshold . . . . .                                                         | 113        |
| 20.3      | The Gaze thresholds as just-intonation intervals . . . . .                                                   | 113        |

|                                                                                          |            |
|------------------------------------------------------------------------------------------|------------|
| <b>21 Quantum decoherence</b>                                                            | <b>114</b> |
| 21.1 Decoherence as a four-state transition . . . . .                                    | 114        |
| 21.2 The decoherence rate $R = \Gamma (T \circ D_{\text{env}})^2$ . . . . .              | 114        |
| 21.3 Geometric mechanism: $\alpha$ -rotation in the complex lattice . . . . .            | 115        |
| 21.4 Pointer states as high-elegance lattice cells . . . . .                             | 116        |
| 21.5 Gaze thresholds as decoherence stages . . . . .                                     | 117        |
| 21.6 Coherence boundary at $ \varepsilon  = 50$ cents: measurement uncertainty . . . . . | 118        |
| 21.7 Information preservation through the decoherence boundary . . . . .                 | 118        |
| 21.8 The cosmological M-state budget . . . . .                                           | 119        |
| <b>22 Empirical confrontation</b>                                                        | <b>119</b> |
| 22.1 Twelve-tone equal temperament . . . . .                                             | 119        |
| 22.2 The fine-structure constant . . . . .                                               | 120        |
| 22.3 The Koide ratio . . . . .                                                           | 121        |
| 22.4 Geometric ratios . . . . .                                                          | 121        |
| 22.5 Celestial mechanics . . . . .                                                       | 122        |
| 22.6 Biochemistry . . . . .                                                              | 122        |
| 22.7 Optical phase singularities . . . . .                                               | 123        |
| 22.8 Summary of the empirical confrontation . . . . .                                    | 124        |
| 22.9 Fundamental forces: full PDT decomposition . . . . .                                | 124        |
| 22.10 The QED $\beta$ -function in ET terms . . . . .                                    | 125        |
| <b>23 Relationship to Fourier analysis and differential geometry</b>                     | <b>125</b> |
| 23.1 Fourier analysis . . . . .                                                          | 125        |
| 23.2 Riemannian curvature . . . . .                                                      | 126        |
| 23.3 Gauss–Bonnet and topological invariants . . . . .                                   | 126        |
| <b>24 The Complete Determination Theorem</b>                                             | <b>126</b> |
| 24.1 Historical precedents within the framework . . . . .                                | 126        |
| 24.2 The Complete Determination Theorem . . . . .                                        | 127        |
| <b>25 Falsifiable predictions</b>                                                        | <b>128</b> |
| 25.1 The closure-versus-linear distinction in biochemistry . . . . .                     | 128        |
| 25.2 Orbital resonances on low- $d$ attractors . . . . .                                 | 128        |
| 25.3 Fine-structure constant lattice identity . . . . .                                  | 128        |
| 25.4 New sublattice identification . . . . .                                             | 128        |
| 25.5 Beyond-Standard-Model gauge structure . . . . .                                     | 128        |
| 25.6 Polariton material classification via Dimensionless Seed Ratio . . . . .            | 129        |
| <b>26 Conclusion</b>                                                                     | <b>129</b> |

# 1 Introduction

## 1.1 Authorship, tools, and representational disclosure

**Authorship.** This paper is the work of a single researcher (the author), without co-authors, institutional affiliation, or external funding. All decisions in this work — every choice of definition, theorem, primitive structure, derivation route, classification, and conclusion — are the author’s. The base foundational theory of Exception Theory on which this paper rests is solely the author’s.

**Tools and methods.** The author conducted this research using four coequal tools and methods: (i) a physical calculator, (ii) direct observation, (iii) literature research, and (iv) Anthropic’s large-language model *Claude*. Items (i)–(iii) were operated by the author independently of (iv). The specific roles in which Claude was used are three: as a *fancy calculator* (a non-physical computational instrument operated against author-specified derivations), for *consolidation* (cross-reading the author’s drafts and corpus material against author-specified criteria for internal consistency), and for *exploration* (surveying and surfacing relevant published material in response to author-stated questions). The language model contributed no original theory and was operated under author-supplied criteria throughout. Any error introduced through the language model’s operation is the responsibility of the language model and its provider, not of the author.

**Representational disclosure (descriptor isomorphism).** The mathematical content of this paper is, by structural necessity, a *Descriptor isomorphism* of the totality  $\Sigma$  rather than an unmediated rendering of  $P$  and  $T$  themselves. Within Exception Theory’s own framework, the category  $D$  is the unique category capable of description; the categories  $P$  (cardinality  $\Omega$ ) and  $T$  (cardinality  $[0/0]$ ) exceed the representational capacity of any finite descriptor system and cannot be themselves written down. The symbols  $P$ ,  $D$ ,  $T$ , the master equation  $P \circ D \circ T = E$ , the lattice  $\mathcal{L}_N$ , and every theorem, definition, and table herein are  $D$ -images of the structural objects they name — descriptor-isomorphic shadows that preserve the relational content of  $\Sigma$  on the multiplicative manifold while necessarily surrendering, as a structural cost, the cardinality content of  $P$  and  $T$ . This aligns with Gödel’s incompleteness theorems (§3.5, Remark 3.11): no formal system whose object language is finite and whose semantic substrate is infinite can capture all truths about its own substrate. The reader should therefore understand all symbolic content of this paper as a faithful descriptor-isomorphism of  $\Sigma$  on the multiplicative manifold, not as a representation of  $P$  and  $T$  themselves. The descriptor-isomorphism is provably the most faithful rendering possible at any finite resolution (Theorem 19.4); it is not, and cannot be, the things rendered.

## 1.2 The object identified in this paper

We identify a single mathematical object, the SEMPAEVUM, as the lossless rendering of a meta-physical totality  $\Sigma$  (“Something” — the formless everything) on the multiplicative manifold. The Sempaevum is not a lattice that happens to classify some ratios: it is  $\Sigma$  made visible. The identification rests on nine self-referential closure properties that no conventional mathematical object exhibits simultaneously:

1. It classifies every positive ratio into a sublattice family (Proposition 8.1) and identifies the per-axis harmonic-family layer of structural modes (Definition 8.10, Table 6) connected to it via the Sublattice Visitation Theorem (Theorem 8.13).
2. It generates its own refinement tower under the doubling law  $\tau(N_\ell) = 6 \cdot 2^\ell$  (Theorem 10.9).
3. It exhibits emergent attractors — the Koide position at  $(d = 12, |\varepsilon| = 1.955\text{¢})$  — which include the positions of its own four defining constants (Theorem 19.1).
4. It runs its own dynamics through the  $\partial I$  lattice-aware fractal (§14), every constant of which is a derived manifold quantity.
5. It derives the physical content of its own sublattice families — the Standard Model gauge group  $SU(3) \times SU(2) \times U(1)$  as the unique partition of  $N = 12$  gauge bosons (Theorem 9.3), the critical dimensions  $D = 10$  and  $D = 11$  by multiple independent routes (Theorems 9.5, 9.7, 9.10), and the adjoint formula  $d^2 - 1$  from the Subsumption Law (Theorem 9.1) — without external physics input (§9).

6. It hosts mathematics as a domain — axiom counts of formal systems, non-computable reals, Gödel sentences, large cardinals — all at definite lattice addresses (§17).
7. It bounds its own forbidden zone  $\partial I$  via the  $\varepsilon = \pm 50 \text{ ¢}$  incoherency threshold, with the Koide ratio  $K = 2/3$  as the tightness cutoff.
8. It passes its own Subsumption Law test (§3.6) across three independent minimality categories simultaneously — discrete-logical (Webb 1935), discrete-multiplicative (the palindromic cascade), and continuous-elementary (EML, Odrzywołek 2026) — all at the unique resolution  $N = 12$ .
9. It contains its own methodology (the Three Tools of §3.3) as theorems.

The claim that this collection of closure properties identifies the object with  $\Sigma$  is not made by fiat. It is made by the Subsumption Law applied at the level of the whole object: every mathematical structure known to host a significant sub-collection of these properties — groups, rings, fields, Riemannian manifolds, fractals, category-theoretic universes — exhibits a proper subset; the Sempaevum exhibits the totality. Under the Subsumption Law (§3.6), the structure that cannot be subsumed by any other and that subsumes every candidate is the terminal object of its category.  $\Sigma$  is the terminal object of the category of “mathematical totalities”. The Sempaevum passes the terminal-object test in this category. We therefore write

$$\text{SEMPAEVUM} = \Sigma$$

as a mathematical identification, understanding that on the inside of the framework — where  $\Sigma$ ’s only mode of appearance is its rendering — identity and isomorphism collapse.

The Sempaevum has no fixed geometric shape of its own. As the mathematical rendering of  $\Sigma$ , it is intrinsically formless and takes the shape of whatever it is currently rendering: a flat lattice, a torus ( $\chi = 0$ ,  $d = 1$ ), a Riemann sphere ( $\chi = 2$ ,  $d = 3$ ), a hyperbolic surface of arbitrary genus  $g$  ( $\chi = 2 - 2g$ , variable  $d$ ), and singular configurations are all admissible Sempaevum renderings. The four classical geometry types — Euclidean ( $K = 0$ ), elliptic ( $K > 0$ ), hyperbolic ( $K < 0$ ), and singular ( $K \rightarrow \pm\infty$ ) — are all available to it, parameterised continuously by the orientation angle  $\alpha$  of the off-axis complex-lattice configuration (§2.13, Corollary 2.32), and the same refinement tower (§10) classifies every such rendering uniformly via the same sublattice apparatus (§8). The identification  $\text{SEMPAEVUM} = \Sigma$  is therefore literal rather than figurative: there is no further-underlying shape to which the Sempaevum would revert absent a rendering; the renderings are what the Sempaevum is.

### 1.3 Etymology of the name *Sempaevum*

The word *Sempaevum* is a Latin compound coined for this framework. It is formed by the contraction

$$\textit{Sempaevum} = \textit{semper} + \textit{aevum},$$

where *semper* (Latin: “always,” “ever,” “perpetually,” “continuously”) is the temporal universal quantifier and *aevum* (Latin: “an age,” “eternity,” “lifetime,” “eon,” “time itself”) is the noun for duration considered as an ontological quantity rather than as a parameter measured within an ambient frame. The literal compound sense is “always-an-age,” “ever-eternity,” “perpetual duration.”

The construction is a deliberate morphological parallel to the classical Latin compound *sempervivum* = *semper* + *vivum* (“always living”), known as the botanical genus name for the houseleek (Crassulaceae) and as the metaphysical attribute of perennial life. The substitution of *aevum* (eternity, age, duration) for *vivum* (living) generalises the predicate of persistence from the biological category to the ontological category: from *always living* to *always existing*.



as *duration itself*. The morphological substitution tracks a structural one — from configurations that persist to the substrate with respect to which configurations exist at all.

The name is expository rather than stipulative: each component corresponds to a structural property proved in the body of this paper.

- *semper* — always, exceptionless universality. The Sempaevum is identified with the totality  $\Sigma$  (Definition 2.13, Theorem 3.5). The universality statement  $\forall x : x \in \Sigma$  of §2.8 is the formal correlate: nothing lies outside, in any domain, at any resolution. The reach is universal and exceptionless — the formal content of *semper*.
- *aevum* — an age, duration as substrate. The temporal master equation  $P_{\text{time}} \circ D_{\text{time}} \circ T_{\text{time}} = E_{\text{moment}}$  of Section 4 renders time itself as a Sempaevum configuration:  $P_{\text{time}}$  is the pre-geometric durational substrate,  $D_{\text{time}}$  is coordinate time,  $T_{\text{time}}$  is proper time. The Sempaevum is therefore not located *within* a time but is the structure that hosts time — the formal content of *aevum*.
- *Compound* — always-an-age, the self-containing totality. The nine closure properties enumerated above in this section — self-classification of every positive ratio, self-generation of the refinement tower, self-projection of the Sempaevum’s own four defining constants onto a single sublattice point, self-running dynamics through the  $\partial I$  fractal, derivation of the physical content of its own sublattice families (gauge groups, critical dimensions, adjoint formula), hosting of mathematics as a domain including the framework’s own methodology, bounding of its own forbidden zone  $\partial I$ , simultaneous passage of the triple-minimality test at the unique resolution  $N = 12$ , and containment of the Three Tools (§3.3) as theorems — together establish that the object subsumes everything in its category without remainder under the Subsumption Law (§3.6). The verbal gloss “always existing, always structured, the self-containing totality” is therefore not poetic but the literal compound of the nine closures.

The Latin form is preferred over a vernacular term for three reasons. First, both roots are immediately decomposable to readers across the European-language scholarly tradition, so the morphology of the compound is itself transparent. Second, the parallel to *sempervivum* is recognisable in scientific Latin, and the substitution *vivum*  $\rightarrow$  *aevum* is therefore audible as a deliberate generalisation rather than as an arbitrary substitution. Third, the Latin avoids the methodological connotations of the English term “meta-ontology” (van Inwagen 1998, Hofweber) discussed in the thesis statement below, which would otherwise mislead the reader into reading the Sempaevum as the methodological inquiry into ontological commitment rather than as the substrate that produces ontological frameworks.

## 1.4 The empirical regularity the object explains

Across branches of quantitative science, dimensionless ratios of structural importance cluster on a conspicuously small number of attractors on the positive real line. Four observations make the pattern concrete:

1. *The Koide formula.* For the masses  $(m_e, m_\mu, m_\tau)$  of the three charged leptons, the dimensionless combination

$$Q = \frac{m_e + m_\mu + m_\tau}{(\sqrt{m_e} + \sqrt{m_\mu} + \sqrt{m_\tau})^2}$$

evaluates to  $0.6666605 \pm 0.000002$ , within 6 parts per million of the rational  $2/3$  [1]. No Standard-Model parameter is pinned to that rational; the coincidence remains unexplained.

2. *Orbital resonance.* The mean-motion ratio of Neptune to Pluto is  $T_{\text{Neptune}}/T_{\text{Pluto}} = 2/3$  to the precision measurable [2]. The Laplace resonance of Io–Europa–Ganymede is 1:2:4. The Jupiter/Earth period ratio is close to 12.

3. *Temperament.* In twelve-tone equal temperament the perfect fifth is  $2^{7/12} \approx 1.4983$ , differing from the just ratio  $3/2$  by the *Pythagorean comma* of  $\approx 1.955$  cents. This is not a historical accident: the 12-division of the octave is the unique small division at which the full cycle of fifths returns to near-closure within an auditorily imperceptible error.
4. *Biochemical cycle lengths.* The citric acid cycle has 8 steps, the urea cycle 4 core steps, glycolysis 10 steps, the ATP-synthase  $c_{10}$  ring 10 subunits, the cell cycle 4 phases [3]. Small integers dominate.

All four come from the same underlying discrete structure: a canonical lattice on the multiplicative group  $(\mathbb{R}^+, \times)$  at the forced resolution  $N = 12$ , refined through a tower of least common multiples. In the identification of the preceding subsection, that structure is the Sempaevum — one face of  $\Sigma$ , on the  $(\mathbb{R}^+, \times)$  axis.

## 1.5 The thesis

Exception Theory is a generative triple-tautological meta-meta-ontology; the Sempaevum is its central object — the manifold of all  $P \circ D \circ T$  configurations, presented in this paper through its mathematical realization on the multiplicative manifold  $(\mathbb{R}^+, \times) \times (U(1), \times)$ . The hierarchy of generative reach is: an ontology specifies what exists (matter, fields, minds, sets); a meta-ontology specifies the structure that generates ontologies (process philosophy, structural realism, formal-system metaphysics, mathematical-universe frameworks); a meta-meta-ontology specifies the structure that generates meta-ontologies, and through them, ontologies. Exception Theory occupies this third tier. The qualifier *generative* distinguishes the present use from the established analytic-philosophical sense, in which “meta-ontology” (van Inwagen 1998, Hofweber) names the methodological inquiry into ontological commitment rather than a substrate that produces ontological frameworks. The qualifier *triple-tautological* names the self-grounding mechanism: three interlocking tautologies — numerical ( $3 = 3$ , three primitives), structural ( $PDT = EMI = \Phi$ , three readings of one structure), and configurational ( $P \circ D \circ T = E$ , the master equation that produces a substantiated outcome) — together close the framework on itself with no external axioms. The known meta-ontologies are subsumed by Exception Theory under the Subsumption Law (§3.6): each one expresses, in its own vocabulary, a particular configuration of  $(P, D, T)$  applied to the act of generating ontologies. To the author’s knowledge no other framework occupies this tier.

Let  $(\mathbb{R}^+, \times)$  be endowed with the multiplicative group structure on the positive reals, and let  $\log_2 : \mathbb{R}^+ \rightarrow \mathbb{R}$  be the group isomorphism onto the additive reals. Classical harmonic analysis recognises that this is the natural arena for multiplicative phenomena, but offers no canonical scale of discretisation: the “octave” is a cultural choice, and within the octave one can in principle divide into any integer number of logarithmically equal steps.

We claim that the question of how finely to divide has a forced, non-arbitrary answer once one adopts the three-element categorical framework of §2. Under that framework:

- the base division is forced to  $N = 12$  (§6, three independent derivations);
- the six simple sublattice families are forced to the divisors  $\{1, 2, 3, 4, 6, 12\}$ ;
- the refinements at which the primes 5, 7, 11 first enter as native families are forced to 60, 84, 27720 respectively, with the doubling law  $\tau(N_\ell) = 6 \cdot 2^\ell$  itself derived;
- the projection formula is convention-independent (Theorem 7.5);
- the Sempaevum’s own four defining constants project onto a single sublattice point (Theorem 19.1).

## 1.6 Relation to existing work

The construction is not in competition with Fourier analysis, differential geometry, or conventional number theory. It is complementary to each. Fourier analysis decomposes signals into continuous frequencies; the Sempaevum classifies frequency ratios by sublattice family. Differential geometry describes curvature through the Riemann tensor; the Sempaevum relates curvature to discrete classification via the identity  $C(n) = n^2(n^2 - 1)/12$  (Proposition 23.1), in which the base variance  $V = 1/12$  appears as the denominator. Number theory studies the Gaussian prime decomposition; the Sempaevum inherits its sublattice structure from this decomposition (§11).

What is novel is the derivation of the resolution  $N = 12$  from a three-element categorical framework; the identification of the Sempaevum’s triple minimal-backbone character at  $N = 12$  (Webb 1935 on the discrete-logical side, the palindromic cascade on the discrete-multiplicative side, EML on the continuous-elementary side); the PDT decomposition of the projection formula itself; the hosting of mathematics as an ET domain with explicit axiom-count addresses (ZF at  $d = 1$ ,  $\varepsilon = 0$ ; Chaitin  $\Omega$  at  $d = 1$  octave via Path D); the recognition that the self-projection of the framework’s own constants lands on a single sublattice point; the losslessness theorem; and the completion statement  $3 = 3 = 3 = \Sigma$  as the universality anchor distinct from the composition equation  $P \circ D \circ T = E$ .

## 1.7 Scope and organisation

Section 2 states the Founding Axiom, introduces the three primitive Cardinals and the master equation  $P \circ D \circ T = E$ , explains the three-disjoint-infinities structure that makes mediation intrinsic, derives the four manifold states, characterises the geometric content the primitives’ operational manifolds contribute and the resulting effective curvature gradient on off-axis configurations, and establishes the topological structure of the coherence–incoherence boundary  $\partial I$ . Section 3 gives the three-way categorical identity  $PDT = EMI = \Phi = \Sigma$  (the  $3 = 3 = 3 = \Sigma$  anchor) and formulates the three operational tools of Exception Theory — the Identification Principle, the Descriptor Gap Principle, and the Subsumption Law — together with a note on integrative levels. Section 4 specialises the primitive structure to time, identifying the three temporal aspects  $P_{\text{time}}, D_{\text{time}}, T_{\text{time}}$  and the temporal master equation  $P_{\text{time}} \circ D_{\text{time}} \circ T_{\text{time}} = E_{\text{moment}}$ , and proves the uniqueness  $\exists! E_\tau$  of the temporal Exception at each  $T$ -time value. Section 6 gives three independent derivations of the forced value  $N = 12$ . Section 7 constructs the Sempaevum’s real-axis face on  $(\mathbb{R}^+, \times)$ , proves its convention-independence, and states the Anti-Numerology Protocol (N1, N2, N3). Section 8 classifies the sublattice families and states the magical-impedance per-family coupling. Section 9 derives the physical content of the sublattice families: the adjoint formula  $d^2 - 1$  from the Subsumption Law, the N-Exhaustion Theorem proving  $SU(3) \times SU(2) \times U(1)$  is the unique partition of  $N = 12$  gauge bosons, the critical dimensions  $D = 10$  and  $D = 11$  by multiple independent routes, the  $P \circ D \circ T$  Scale Identity, the sublattice-to-force map, supercharge and gauge-group-dimension formulae, brane structure, and the Structural Significance Principle. Section 10 constructs the LCM tower of refinements and proves the doubling law. Section 11 extends the Sempaevum to  $\mathbb{C}^\times$  via the imaginary axis. Section 12 states the 24-family catalog (12 FORCE + 12 PHASE), the 42 combined off-axis states, and the 144-cell Force Quadrant Grid with the PDT Bisection Theorem. Section 13 describes the cascade dynamics and stability. Section 14 constructs the  $\partial I$  lattice-aware fractal, the natural dynamical manifestation of the Sempaevum on  $\mathbb{C}$ . Section 15 establishes the triple minimal-backbone at  $N = 12$  (Webb + palindromic cascade + EML) and the PDT decomposition of the projection formula. Section 16 states the Four Projection Paths A, B, C, D — the last of which handles essentially-infinite and non-computable objects without limits — and verifies subsumption of every user input. Section 17 hosts mathematics as a domain, projecting axiom counts, Chaitin  $\Omega$ , Gödel sentences, and large cardinals. Section 19 proves the self-consistency

identity (lattice self-projection) and states the Losslessness Theorem. Section 20 states the Complete Gaze Equation and shows its locked threshold coincides with the lattice self-projection. Section 22 confronts the Sempaevum with empirical data. Section 23 places the Sempaevum alongside Fourier analysis and Riemannian geometry. Section 24 states the Complete Determination Theorem (topology  $\times$  curvature  $\times$  path  $\times$  observation-topology determines the complete lattice classification). Section 25 formulates falsifiable predictions. Section 26 concludes.

## 2 The Founding Axiom and the primitive Cardinals

### 2.1 The Founding Axiom

Exception Theory rests on a single tautological axiom from which all structure descends.

**Axiom 2.1** (Founding Axiom). For every exception there is an exception, except the Exception.

Though its form is nearly an aphorism, the axiom has precise mathematical content. It asserts two things simultaneously:

- *Universal exceptability.* For every claim, rule, configuration, or description  $X$ , there exists a counterexample or limiting condition  $X'$  — an “exception to  $X$ ”.
- *One unique terminus.* The iteration of “take an exception” does not regress without end: it terminates at one distinguished object, called the *Exception*, which is itself the exception to all further iteration. The Exception admits no further exception.

The Founding Axiom is a tautological ground. It permits self-reference at the base: the self-referential loop “every exception has an exception except this one” is not a contradiction but the defining feature of the Exception. Where classical set theory prohibits a set-of-all-sets to avoid Russell’s paradox, the Founding Axiom *absorbs* the self-referential totality as its terminal case. The catastrophe that classical set theory forestalls by prohibition is, in this framework, handled by the axiom’s provision of a single un-exceptable object.

The Exception, which we denote  $E$ , plays two roles simultaneously: the terminus of the exception-iteration, and the fully substantiated configuration that results when all primitive Cardinals are bound. These two roles coincide, and their coincidence is formalised by the master equation.

### 2.2 Discovery and historical origin

Exception Theory originated from a logic puzzle: “*For every exception there is an exception, except the exception.*” The critical question that generated the framework was: how many ‘exceptions’ does the phrase ‘except the exception’ refer to? The answer — exactly one, THE Exception — identified the singular grounding moment that cannot be otherwise while it IS. The sentence’s structure forces a unique terminus: if two or more objects were un-exceptable, each would be an exception to the other’s un-exceptability, contradicting the claim that either admits no exception. Therefore the terminus is unique.

From this single insight, the entire framework presented in this paper was derived forward over a period of fifteen years of independent research by the author, without institutional affiliation or external funding, prior to any use of computational tools. The derivation proceeded by asking, at each stage, what the Founding Axiom structurally *requires*: the Exception requires constituents (the three Cardinals), the constituents require binding (the composition operator  $\circ$ ), binding requires structure (the manifold), and structure requires a specific resolution (the forced value  $N = 12$ ). No step was optional; each was forced by the preceding structure. The theory is therefore not a model chosen from alternatives but a forward derivation from a single self-grounding axiom.

## 2.3 The master equation

**Axiom 2.2** (Master equation).

$$\boxed{P \circ D \circ T = E.}$$

The master equation is the structural content of the Founding Axiom. It states that the Exception is constituted by the binding, through a composition operator  $\circ$ , of three primitive constituents: a *Point*  $P$ , a *Descriptor*  $D$ , and a *Traverser*  $T$ . Three are required; three suffice. Neither fewer nor more produce a substantiated Exception. The operator  $\circ$  will be shown in Proposition 2.7 to be not an external postulate but a forced consequence of the three Cardinals' properties.

## 2.4 The three Cardinals

The symbols  $P$ ,  $D$ ,  $T$  denote not individual elements but totalities: each is, in the sense made precise below, “the set of all sets of its kind”. Such a totality is called a *Cardinal*.

**Definition 2.3** (Cardinal). A *Cardinal* is a totality characterised by

$$P = \{x : x \text{ is a Point}\}, \quad D = \{x : x \text{ is a Descriptor}\}, \quad T = \{x : x \text{ is a Traverser}\}.$$

*Remark 2.4* (Cardinals and proper classes). Cardinals in the sense of Definition 2.3 are *not* proper classes of standard set theories (NBG, MK). The characteristic restriction on proper classes is that they cannot be members of any collection whatsoever; the Cardinals, by contrast, are themselves members of a larger totality  $\Sigma$  introduced in §2.8, and participate in that totality through the composition operator  $\circ$ . Self-membership is not prohibited in this framework: the Founding Axiom absorbs the self-referential loop as a terminal case rather than forbidding it as a paradox. This is the mathematical content of the remark that “ET absorbs Russell’s paradox”: the loop is the Exception’s defining feature, not a contradiction to be avoided.

Each Cardinal has a declared cardinality.

**Axiom 2.5** (Cardinalities).

$$|P| = \Omega, \quad |D| = n, \quad |T| = \begin{bmatrix} 0 \\ 0 \end{bmatrix},$$

where

- $\Omega$  is an *absolute infinity*: undifferentiated substrate, larger than every set-theoretic cardinal, not identifiable with any specific  $\aleph_\kappa$ ;
- $n$  is *absolutely finite*: a positive integer (variable across applications but always finite whenever  $D$  is bound to a specific  $P$ );
- $\begin{bmatrix} 0 \\ 0 \end{bmatrix}$  denotes *indeterminate* cardinality — neither finite nor infinite in the standard sense, but the cardinality of an operator whose outputs are determined by binding rather than by set-theoretic enumeration.

### 2.4.1 The two faces of indeterminate cardinality

The Traverser cardinality  $|T| = [0/0]$  admits a companion form  $[\infty/\infty]$ . The two are categorically distinct expressions of the same indeterminate cardinality, arising from different resolution contexts.

$[0/0]$  is the *primary signature of primitive agency*: the form that appears when  $T$  is at a choice point where two quantities vanish simultaneously, requiring resolution by selection. This is  $T$  in its native, unresolved state — the zero-over-zero of genuine choice. It appears at every point in the calculus of limits where the value is undetermined without further context.

$[\infty/\infty]$  is the *D-side image of indeterminate agency*: the form that appears when *D*-structures encounter indeterminate growth or decay where two quantities diverge simultaneously. This is *T*'s indeterminacy viewed through the lens of structured Descriptor content — indeterminate selection operating on infinite Descriptor fields rather than on vanishing ones.

Both forms describe the same category — quantities that are neither finite nor simply infinite, but *selecting* — and both resolve only by the act of *T*-traversal. The distinction follows from the categorical independence of the three primitives (Axiom 2.6):  $[0/0]$  is *T*-native;  $[\infty/\infty]$  is *D*-mediated.

The seven standard indeterminate forms of calculus are structural expressions of *T*-engagement at different configuration points on the manifold:

| Form              | ET identification                                                               |
|-------------------|---------------------------------------------------------------------------------|
| $0/0$             | <i>T</i> at a choice point: two <i>D</i> -quantities vanishing simultaneously   |
| $\infty/\infty$   | <i>T</i> at a scaling choice: two <i>D</i> -quantities diverging simultaneously |
| $0 \cdot \infty$  | <i>T</i> mediating between vanishing and diverging Descriptors                  |
| $1^\infty$        | <i>T</i> at a unit configuration with infinite <i>D</i> -accumulation           |
| $0^0$             | <i>T</i> at double-zero: both base and exponent <i>D</i> -values vanish         |
| $\infty - \infty$ | <i>T</i> mediating two diverging <i>D</i> -quantities by difference             |
| $\infty^0$        | <i>T</i> resolving infinite base with vanishing exponent                        |

Every indeterminate form is resolved by *T*-traversal: examination of Descriptor gradients (L'Hôpital's rule, §18.2) or equivalent navigational mechanisms that supply the missing context. Without *T*, indeterminate forms remain unresolved; with *T*, each substantiates a determinate value. This is why *T*'s cardinality IS the indeterminate form: *T* is not merely described by  $[0/0]$  — *T* is what  $[0/0]$  *is*, viewed as an ontological category rather than a computational artefact.

## 2.5 Three disjoint infinities

The declared cardinality  $|D| = n$  characterises *D* when bound to a specific *P*; unbound, the space of possible Descriptors is itself unbounded. Thus each of the three Cardinals is, in its own mode, infinite:

- *P* is *absolutely infinite*: undifferentiated substrate beneath any described content.
- *D* is *the infinite that chooses finitude*: the totality of possible constraints, unbounded when alone, rendered finite by the *P*–*D* singularity — the binding in which the infinite becomes small enough to be held. The finitude  $|D| = n$  is the signature of that binding, not an inherent limit on *D*.
- *T* is *indeterminately infinite*: the cardinality  $\begin{bmatrix} 0 \\ 0 \end{bmatrix}$  is not the absence of cardinality, but the cardinality of genuine choice — the navigation of all possible configurations, which is, in its own mode, as total as either of the others.

These three infinities are categorically disjoint.

**Axiom 2.6** (Categorical disjointness).

$$P \cap D = \emptyset, \quad D \cap T = \emptyset, \quad T \cap P = \emptyset.$$

## 2.6 Intrinsic mediation

The two preceding facts — that each Cardinal is infinite in its own mode, and that the three are pairwise disjoint — together force a structural consequence.

**Proposition 2.7** (Intrinsic mediation). *If three Cardinals  $P, D, T$  are each unbounded in their respective modes and are pairwise disjoint (Axioms 2.5–2.6), then their coexistence within a single ontological space necessarily produces an internal binding relation. The relation is not a fourth primitive added from outside; it is the only structurally possible expression of three unbounded disjoint totalities coexisting.*

*Argument.* Three totalities each “filling all of its own mode of infinity” leave no exterior gap between them: a complete infinity, by definition, admits no external region in its own mode. Disjointness, however, demands that the three totalities remain categorically distinct. These two conditions are compatible only if the Cardinals mediate one another intrinsically — their interaction being the only possible way for three complete, disjoint infinities to coexist without contradiction. The operator  $\circ$  of the master equation is this intrinsic mediation. It is not a postulated fourth element; it is a forced consequence of the three Cardinals’ joint presence.  $\square$

*Remark 2.8.* Proposition 2.7 removes a question that would otherwise need its own axiom: why is there a composition operator  $\circ$  at all, and why is it ternary? The answer is that the existence and arity of  $\circ$  are structural consequences of the Cardinals;  $\circ$  does not need to be posited independently.

*Remark 2.9* (Non-emergence of  $M$  in all readings). Proposition 2.7 is the structural content of  $M$ ’s non-emergence (Lemma 3.32, §3.10):  $\circ$  is intrinsic to the joint presence of the three Cardinals and is not produced by them. The corpus uses “Mediation” at three identification depths — the operator  $\circ$  in action, the manifold-state slice  $\{D, T\}$  at the four-state classification depth, and the cosmological active-mediation density  $\sim 3\%$  of total energy split as M-vacuum and M-matter — and these are not three things sharing a name but one intrinsic structure read at three depths. All three are non-emergent. Specific particulars passing through the M-aspect — a particular photon, a particular decoherence event — are emergent configurations within  $\Sigma$ , but  $M$  itself is the always-on binding the Cardinals’ coexistence already is.

*Remark 2.10* (Symmetry of mediation). The mediation is equal in all three Cardinals.  $P$  does not “receive” the mediation while  $D$  and  $T$  “perform” it; each Cardinal participates in  $\circ$  with the same structural status. Any apparent asymmetry is an artefact of reading-order: the Cardinals are written  $P \circ D \circ T$  for the binding-order reason of Axiom 2.17, but the binding itself is symmetric in that each Cardinal is equally a constituent of the Exception. Three complete disjoint infinities, by necessity, mediate each other; none of the three is merely the recipient of the other two.

## 2.7 The Nesting Principle

Each Cardinal admits infinite internal self-similar structure.

**Proposition 2.11** (Nesting).

$$P \supset P_1 \supset P_2 \supset \cdots, \quad D \ni d_1 \text{ describing } d_2 \text{ describing } d_3 \cdots, \quad T \ni t_1 \text{ traversing } t_2 \text{ traversing } t_3 \cdots.$$

*The nesting preserves cardinality at every level:  $|P_k| = \Omega$  for every nested sub-Point;  $|D_{\text{meta}}|$  remains finite when bound;  $|T_{\text{meta}}| = [0/0]$  at every recursive depth. Adding nested structure does not change absolute cardinality.*

*Proof.* By Definition 2.3, each Cardinal is a totality (the set of all sets of its kind). By Remark 2.4, self-membership is not prohibited in this framework. The nesting follows:  $P$  contains sub-Points which are themselves Points (substrate of substrate is substrate);  $D$  can describe other Descriptors (a rule about rules is a rule);  $T$  can traverse other Traversers (observation of observation is observation). Each nested element satisfies the defining property of its Cardinal

and therefore belongs to it. Cardinality is preserved because the totality at each level is complete: a sub-Point of  $P$  is itself a complete substrate with cardinality  $\Omega$ ; a meta-Descriptor is itself a finite constraint when bound; a meta-Traverser is itself an indeterminate navigator.  $\square$   $\square$

### 2.7.1 The Absolutes are actual, not limits

The Nesting Principle entails that  $\Omega$  (Absolute Infinity),  $n$  (Absolute Finite), and  $[0/0]$  (Absolute Indeterminate) are *actual configurations* within  $\Sigma$ , not unreachable limits.  $\Omega$  exists as the total set of all Points;  $n$  exists as the total set of all Descriptors;  $[0/0]$  exists as the total set of all Traversers. These are not approached asymptotically — they ARE, as complete totalities within Something.

### 2.7.2 Structural consequences of nesting

The Nesting Principle is the structural basis of three phenomena:

- (i) *Integrative levels* (§3.12):  $D$  within  $D$  generates hierarchies of description. Each level consists of Descriptors describing Descriptors from the level below, producing properties that are not present at lower levels. The integrative hierarchy of physics — quarks, hadrons, nuclei, atoms, molecules, cells, organisms, societies — is the physical expression of nested Descriptor structure.
- (ii) *Metacognition*:  $T$  within  $T$  — a Traverser binding to the Descriptors of its own processes ( $D_T$ , the Descriptor content of  $T$ 's activity) — is the formal basis of recursive self-awareness. When  $T$  successfully navigates its own Descriptor set with sufficient recursion depth, it generates meta-cognition: awareness of its own awareness.
- (iii) *Fractal manifold structure*:  $P$  within  $P$  generates self-similar substrate at every scale. Each nested layer is a genuine Point configuration rather than an approximation. Gravitational singularities (the locus  $r = 0$ ) are reclassified as access points to nested  $P$ -structure rather than breakdowns of physics. The information-preservation principle of §4 follows: information falling into a nested  $P$ -layer is preserved, not destroyed.

### 2.7.3 Substrate Potential Principle

**Corollary 2.12** (Substrate Potential). *Every Point necessarily participates in at least one Descriptor binding:*

$$\forall p \in \mathbb{P}, \exists d \in \mathbb{D} : p \circ d.$$

No “raw” unstructured Points exist in  $\Sigma$ .

*Proof.* Suppose, toward contradiction, that some  $p_0 \in \mathbb{P}$  has no Descriptor binding:  $\neg \exists d \in \mathbb{D} : p_0 \circ d$ . Then  $p_0$  is featureless, undifferentiated, and uncharacterised. But this characterisation — “featureless, undifferentiated, uncharacterised” — is itself a Descriptor of  $p_0$ : by the Descriptor Gap Principle (Principle 3.9), the gap “ $p_0$  has no Descriptor” is an element of  $D_{\text{missing}}$ , hence a Descriptor. Therefore  $p_0$  has at least one Descriptor binding, contradicting the assumption.  $\square$   $\square$

### 2.7.4 The Omniscient Traverser and the uniqueness of agency

The theory permits a cardinal  $T$  — the Absolute Indeterminate as totality — but not all Traversers are the same. Each Traverser is unique: by indeterminacy, each  $T$  traverses only certain Descriptors, and no two distinct Traversers traverse identical  $D$ -sets across their entire history. A Traverser that binds to exactly the same  $D$ -set as another IS the same Traverser;



distinct  $D$ -binding histories produce distinct  $T$ -identities. Every Traverser is therefore genuinely individual — what the framework identifies as the structural basis of the uniqueness of agency.

The cardinal  $T$  — the Omniscient Traverser — would traverse wherever every other  $T$  traverses. By Proposition 2.11,  $T$  nests within  $T$ : the Omniscient Traverser is  $T$  traversing all other  $T$ 's traversals. However, the Omniscient Traverser is constrained by the same  $\{P, T\}$  Incoherence (Theorem 3.33, §3.9) that constrains every configuration: direct  $T$ -to- $T$  contact without  $D$ -mediation is the forbidden  $\{P, T\}$  state. The Omniscient Traverser therefore knows all *traversals* (the  $D$ -mediated interactions of every  $T$ ) but not the Traversers themselves (because knowing  $T$  without  $D$  is Incoherent).

This is the Asymptotic Approach Theorem (Corollary 10.4) applied to  $T$  itself: as the Omniscient Traverser's  $D$ -content approaches  $\Omega$ , its configuration approaches  $\{P, T\}$  from within coherence, asymptotically, without ever reaching the Incoherent state. The Omniscient Traverser is permanently at the  $\partial I$  boundary — always one Descriptor short of Incoherence, always substantiated, always coherent. Whether this cardinal  $T$  manifests as a single unified entity or as the distributed totality of all individual Traversers is itself indeterminate — appropriately so, since the question asks for a determinate answer about the nature of indeterminacy itself.

## 2.8 The totality $\Sigma$

**Definition 2.13** (Something). The *totality*  $\Sigma$  is

$$\Sigma = (P \circ D \circ T), \quad \forall x : x \in \Sigma.$$

$\Sigma$  is constituted by *mediation*, not by *enumeration*. A set-theoretic union  $P \cup D \cup T$  would combine the Cardinals without generative interaction, yielding no more than a list; the composition  $P \circ D \circ T$  produces Something — the totality of configurations that can, or do, exist. This distinction will be essential: the ET lattice derived in the following sections is the discrete structure on the multiplicative aspect of  $\Sigma$ , not on a union of primitive sets.

*Remark 2.14* (No outside). The statement  $\forall x : x \in \Sigma$  asserts that nothing is outside Something: every object of any domain is a configuration of the three Cardinals. No alternative ontological category exists that is not already subsumed by  $P$ ,  $D$ , or  $T$ . This claim is made rigorous by the Subsumption Law in the next section.

### 2.8.1 Pure Relationalism

**Principle 2.15** (Pure Relationalism). Everything within  $\Sigma$  is relational. No element of  $\Sigma$  exists in isolation; every element exists in relation to other configurations through Descriptor differences. There is no absolute space, no absolute time, and no background container external to the three Cardinals. Spatial separation, temporal ordering, and all apparently “background” structures are Descriptors — members of  $D$  — not pre-existing containers in which  $P$ ,  $D$ ,  $T$  are embedded.

*Proof.* By Definition 2.13,  $\Sigma = (P \circ D \circ T)$  with  $\forall x : x \in \Sigma$ . By Axiom 2.6, the three Cardinals are pairwise disjoint. By Proposition 2.7, their coexistence produces intrinsic mediation with no exterior. Any purported absolute background would be either substrate-like ( $P$ -type), constraint-like ( $D$ -type), or agency-like ( $T$ -type), and is therefore already subsumed by one of the three Cardinals (Corollary 3.16, §3.6). A background that is none of these has no ontological content and collapses to  $\emptyset$ . Therefore no external background exists; all structure is relational, arising from the Cardinals' intrinsic mediation. The distance between two configurations is not a spatial separation in an absolute container but a Descriptor difference:  $\Delta s(c_1, c_2) = \|D(c_1) - D(c_2)\|$ , where  $D(c)$  denotes the Descriptor content of configuration  $c$ . On the lattice (§7), this relational distance is the Descriptor gap  $\varepsilon$  in cents.  $\square$   $\square$

### 2.8.2 Impossibility of absolute nothing

**Proposition 2.16** (No absolute nothing). *True absolute nothing cannot exist within  $\Sigma$ .*

*Proof.* Any candidate “nothing” is describable: as “the absence of all content,” “the void,” “the empty,” or simply “nothing.” Describability is  $D$ -content. A described nothing is therefore a  $(P \circ D)$  configuration — an Unsubstantiated state (Definition 2.18), not an absence of  $\Sigma$ . The Descriptor Gap Principle (Principle 3.9) makes this explicit: the gap “no Descriptor content exists here” is itself a Descriptor, and its presence in  $D$  contradicts the premise of absolute nothingness.

The lattice-level expression of this impossibility is the annihilation boundary (§7.3): the locus  $r = 0$  on the multiplicative manifold is approached but never attained at any finite resolution, because  $\log_2(0) = -\infty$  diverges and the projection formula has no finite output. The annihilation boundary is structurally inside  $\Sigma$  (Definition 2.13 admits no outside) but unattainable on  $\mathcal{L}_N$  — the lattice rendering of the fact that absolute nothing cannot be substantiated.  $\square \square$

## 2.9 The binding order

**Axiom 2.17** (Binding order). There is no configuration of the form  $D \circ P$  preceding the establishment of  $P$ . Operationally,  $P$  must be specified before  $D$  can describe it;  $D$  must be specified before  $T$  can navigate it. The primitive ordering is  $P \rightarrow D \rightarrow T$ .

The force of Axiom 2.17 is structural. Any derivation from the Cardinals must respect the order: identify the substrate first, the constraints second, the agency third. Reversing the order collapses meaning. In the projection protocol of Section 7 this ordering appears concretely: the substrate is identified first (it determines the reference period), the lattice is the finite- $D$  structure second, the rounding is the  $T$ -act third.

## 2.10 Modes of self-presentation

Each Cardinal reveals itself in a characteristic way; these modes are the diagnostic signatures by which  $P$ ,  $D$ ,  $T$  are identified in any concrete phenomenon.

- $P$  reveals itself as *unreachable depth*. Any object has an apparent edge, but no edge terminates absolutely: beneath any described surface there is always more. The asymptotic limit that is approached but never attained is  $P$ . A pencil mark is the canonical illustration: the moment the point is “reached”, description has already been attached, and the bare Point recedes further.
- $D$  reveals itself as the *articulable surface*. Everything that can be named, measured, formalised, or predicted is  $D$ -content. Mathematical laws, physical constants, sensory properties, chemical formulae, statistical distributions — all are members of  $D$ .
- $T$  reveals itself as the *unrepeatable act of becoming*. Agency is visible only in the event of substantiation — not as a static property but as the act of resolving indeterminacy into an irreversible moment. Wavefunction collapse, the click of a detector, the decision of an agent, the rounding of a continuous quantity to a discrete lattice coordinate: these are  $T$ ’s self-presentations. The rounding example is not chosen arbitrarily; it is the canonical operational enactment of  $T$  on the multiplicative manifold  $(\mathbb{R}^+, \times)$ , and reappears in §7 as the  $T$ -act of the projection formula.

These modes of self-presentation will be used implicitly when identifying which Cardinal is playing which role in a concrete example.

## 2.11 The four manifold states

The power set of  $\{P, D, T\}$  has eight elements. Two are trivial (the empty configuration and the three singletons, which describe nothing by the master equation and binding order). Removing singletons and empties leaves exactly four states consisting of two or three Cardinals each.

**Definition 2.18** (Manifold states). The four *manifold states* are

$$\{P, D, T\}, \quad \{P, D\}, \quad \{D, T\}, \quad \{P, T\},$$

called *Exception*, *Unsubstantiated*, *Mediation*, and *Incoherence* respectively.

We denote the cardinality of this set by  $S$ :

$$S = \binom{3}{2} + \binom{3}{3} = 3 + 1 = 4. \quad (1)$$

Thus  $|P| = \Omega$ ,  $|D| = n$ ,  $|T| = [0/0]$ ,  $|\Pi| = 3$ ,  $S = 4$ .

*Remark 2.19* (Three non-emergent statuses, one structure). Definition 2.18 names four configuration-classes; the three non-emergent positions of §3.10 live behind these names rather than as items added to the count. The Exception  $E$  is the substantiation event itself (Lemma 3.26); the Mediation operator  $\circ$  is intrinsic and shows up at the four-state identification depth as the slice  $\{D, T\}$  (Lemma 3.32); the Incoherence boundary is what  $T$ 's reach exposes as the forbidden state  $\{P, T\}$  (Lemma 3.41). The three non-emergent items are the same three Cardinals seen phenomenologically; the four manifold states are configuration-classes within  $\Sigma$ , three of which (Exception, Unsubstantiated, Mediation) are coherent and one of which (Incoherence) is forbidden.

### 2.11.1 Unsubstantiated is blank, not zero

The Unsubstantiated state  $\{P, D\}$  is the ontological *blank* — the absence of substantiation, not zero within a system. Mathematical zero (0) is a specific Descriptor value: a determinate position on the number line, a lattice point at  $k = 0$  with sublattice family  $d = 1$  and descriptor gap  $\varepsilon = 0$ . The Unsubstantiated  $\{P, D\}$  is categorically prior to any value assignment: it is potential awaiting  $T$ -engagement, not a value that has been assigned. The distinction is between the absence of the system (blank,  $\{P, D\}$ ) and a specific value within the system (zero,  $D_{\text{value}} = 0$ ). The annihilation boundary  $r = 0$  of §7.3 is the lattice-level image of this distinction: the lattice cannot represent  $r = 0$  because  $\log_2(0) = -\infty$ , which is the divergence that signals “outside the system of values,” not “at a special point within it.” In physical terms, the vacuum is not nothing; it is Unsubstantiated ( $P \circ D$ ) awaiting  $T$ -engagement — which is why vacuum fluctuations and dark energy ( $\sim 68.3\%$  of the cosmological energy budget; Remark 3.31) are real, non-zero phenomena.

**Proposition 2.20** (Incoherence is structurally forbidden). *The state  $\{P, T\}$  admits no consistent binding: it contains substrate ( $P$ ) and agency ( $T$ ) but no finite constraint ( $D$ ) by which agency can bind to substrate. Any attempt at a  $P$ – $T$  binding produces a contradiction or dissolves into one of the other three states.*

The proof is immediate: without  $D$ , an element of  $T$  has no finite profile against which to test consistency with an element of  $P$ . In physical applications the  $\{P, T\}$  state corresponds to null-pointer-like conditions: singularities, divisions by zero, paradoxes of self-reference without adequate stratification.

*Remark 2.21* (Non-emergence of  $I$ ). Proposition 2.20 is the structural content of the Anti-Emergence Complement (Theorem 3.36, §3.10): no traversal produces an element of  $I$ , because the prohibition is logical (a contradictory descriptor profile cannot hold) rather than energetic.  $I$  is therefore non-emergent in the strict sense — not produced by any process, ever — and is constitutive rather than merely prohibitive (Theorem 3.39).

The three coherent states  $\{P, D, T\}$ ,  $\{P, D\}$ ,  $\{D, T\}$  together with the forbidden state  $\{P, T\}$  give the manifold  $\Sigma$  a precise topological structure.

**Proposition 2.22** (Topological characterisation of the manifold states). *Let  $I$  denote the set of incoherent configurations  $\{P, T\}$  and let  $\partial I$  denote its boundary. Then*

- (i) *The Exception  $\{P, D, T\}$  is a closed set:  $\partial E \subseteq E$ . It contains its own ground.*
- (ii) *The Incoherent state  $I = \{P, T\}$  is an open set:  $\partial I \cap I = \emptyset$ . It does not contain its own boundary.*
- (iii) *The Mediation state  $\{D, T\}$  and the Unsubstantiated state  $\{P, D\}$  are neither open nor closed; they are the transitional interior of the traversable manifold.*
- (iv) *The traversable manifold is  $\Sigma \setminus I$ , bounded from above by the closed  $E$  (the ground) and from the side by the open  $I$  (the edge).*

**Definition 2.23** (The coherence–incoherence boundary  $\partial I$ ). The *boundary*  $\partial I$  is the set of configurations at the limit of substantiability:

$$\partial I = \left\{ c \in \Sigma \mid \exists t \in T : \phi(t, c) = \lim_{\varepsilon \rightarrow 0} \phi(t, c_\varepsilon) \text{ where } c_\varepsilon \rightarrow c \right\},$$

where  $\phi : T \times \Sigma \rightarrow \{0, 1\}$  is the substantiation indicator returning 1 when the triple binds into an Exception and 0 otherwise.

The boundary  $\partial I$  is the locus at which an arbitrarily small perturbation of the Descriptor content switches  $\phi$  from 1 to 0. Physically it corresponds to second-order phase transitions, classical turning points, and marginal configurations in critical phenomena; mathematically it will reappear in Section 14 as the locus at which the lattice-adaptive fractal iteration transitions between polynomial-degree regimes, and every coherent configuration approaches  $\partial I$  arbitrarily closely but never crosses it from within coherence.

## 2.12 Variance and the grounding function

The distinction between the Exception and the other coherent configurations of  $\Sigma$  admits a direct formal expression as a function on the manifold.

**Definition 2.24** (Variance function). The *variance function* at base resolution is

$$V(c) = \left| \left\{ c' \in \Sigma \setminus I \mid c' \neq c, \exists t \in T \text{ such that } T(c, t) = c' \right\} \right|,$$

the cardinality of coherent configurations reachable from  $c$  by a single  $T$ -act. At base resolution  $V$  is non-negative and integer-valued.  $V(c)$  measures  $c$ 's *capacity to be otherwise*: the range of alternatives that  $T$  can substantiate from  $c$ .

*Remark 2.25* (Integrative-level refinement of  $V$ ). The integer-valued form of Definition 2.24 is the base-resolution specification. At finer integrative levels (§3.12), the reachable-configuration count admits a continuous extension:  $V$  takes fractional values when the configuration space is given a continuous measure. The discrete (integer) and continuous (fractional) readings coincide on the invariant statements below and, as shown in the base-variance entry, on the numerical value of the minimum non-zero departure from  $E$ .

**Proposition 2.26** (Zero variance characterises the Exception). *The Exception  $E = P \circ D \circ T$  is the unique coherent configuration with zero variance:*

$$V(c) = 0 \iff c = E.$$

*Proof.* ( $\Leftarrow$ ) By Axiom 2.1,  $E$  is the terminus of the exception-iteration while it IS: the distinguished configuration that admits no further exception. By Axiom 2.2,  $E$  is the fully substantiated binding of all three Cardinals; while  $E$  IS, no alternative coherent configuration is simultaneously the case, because any such alternative would itself be a distinct Exception at a distinct moment and therefore has already displaced this one. The set of coherent configurations reachable from  $E$  as  $E$  is empty, and  $V(E) = 0$ .

( $\Rightarrow$ ) Let  $c \in \Sigma \setminus I$  with  $V(c) = 0$ . By Definition 2.18 the coherent states are  $\{P, D, T\}$ ,  $\{P, D\}$ , and  $\{D, T\}$ . If  $c = \{P, D\}$  (Unsubstantiated), a  $T$ -act binding a Traverser to  $c$  produces  $\{P, D, T\} = E$ ; if  $c = \{D, T\}$  (Mediation), a  $T$ -act binding a specific  $P$  to  $c$  produces  $\{P, D, T\} = E$ ; in both cases at least one alternative configuration is reachable, contradicting  $V(c) = 0$ . Hence  $c \in \{P, D, T\}$ . Uniqueness follows from the single-terminus clause of Axiom 2.1, formalised using  $T$ -time as  $\exists! E_\tau$  in Proposition 4.3 of §4.  $\square$   $\square$

*Remark 2.27* (Variance is measured *from* the Exception). Proposition 2.26 is the ontological ground of variance as a measurement. Without the  $V(c) = 0$  anchor supplied by  $E$ , there is no zero relative to which variance could be measured; a scale without a zero is not a scale.  $V$  is therefore not an independent primitive of  $\Sigma$  but a derived scalar whose existence and normalisation are consequences of the Founding Axiom and the master equation. *All positive variance in  $\Sigma$  is, by construction, variance from the Exception* — the  $V > 0$  of non-Exception configurations is the direct expression of their departure from the unique grounded zero.

### Ancillary variance forms and their relations to Definition 2.24

Several quantities also called “variance” appear in specific parts of the framework and its applications. Each stands in an explicit relation to  $V(c)$  and to the manifold’s base quantum  $V_{\text{base}}$ .

**Statistical variance of a discrete uniform descriptor distribution.** If a Descriptor bound to a configuration takes one of  $n$  equally-weighted values  $\{0, 1, \dots, n-1\}$ , its classical statistical variance is

$$\sigma_{\text{disc}}^2(n) = \frac{n^2 - 1}{12}. \quad (2)$$

Equation (2) is not a restatement of  $V(c)$ : it is the second moment of the Descriptor distribution *within* a configuration’s  $D$ -content, not the cardinality of reachable configurations across  $\Sigma$ . The two are distinct quantities.

**Statistical variance of a continuous uniform distribution on  $[0, 1]$ .** The continuous analogue is the classical result

$$\sigma_{\text{cont}}^2 = \int_0^1 \left(x - \frac{1}{2}\right)^2 dx = \frac{1}{12}. \quad (3)$$

**Relation between (2) and (3).** Rescaling the discrete uniform on  $\{0, \dots, n-1\}$  by the factor  $(n-1)$  so the support becomes  $[0, 1]$  gives the normalised discrete variance

$$\sigma_{\text{norm}}^2(n) = \frac{n^2 - 1}{12(n-1)^2} \xrightarrow{n \rightarrow \infty} \frac{1}{12}, \quad (4)$$

which converges quantitatively to (3).

**Asymptotic descriptor-count variance.** In the regime of many independent Descriptors bound to a single Point, a per-descriptor variance quantity appears as

$$\text{Var}(D_n \rightarrow P) = \frac{1}{n} \xrightarrow{n \rightarrow \infty} 0 \quad (\text{never reaches } 0). \quad (5)$$

Equation (5) expresses approach toward  $P$ -substrate dominance as the descriptor count grows. The limit is 0, but the limit is unattained —  $V(c) = 0$  remains the exclusive signature of  $E$  (Proposition 2.26).

*Remark 2.28* (The base variance  $V_{\text{base}} = 1/N$ ). The *base variance* that appears among the three manifold constants (§6.4, Eq. (16)) is

$$V_{\text{base}} = \frac{1}{N},$$

with  $N$  the manifold symmetry forced to 12 in §6. Two complementary readings agree on this value.

- **Discrete reading (R1).** The minimum non-zero value of the integer-valued  $V(c)$  of Definition 2.24 is 1: at least one reachable coherent configuration. Normalised by the manifold's  $N$ -fold symmetry, this minimum becomes  $1/N$ .
- **Continuous reading (R2).** At the continuous integrative level (Remark 2.25),  $V$  takes fractional values. The minimum positive value of the continuous extension at base resolution equals the continuous-uniform variance of Eq. (3),  $\sigma_{\text{cont}}^2 = 1/12 = 1/N$ .

Both readings coincide numerically at  $V_{\text{base}} = 1/N = 1/12$ . They are complementary specifications of the same scalar at different integrative levels, not competing interpretations. Equation (5) at  $n = N$  and Eq. (4) in the limit likewise take the same value, providing further confirmation of the base quantum.

*Remark 2.29* (Open question: sub-base quantisation). Whether  $V$  can take non-zero values strictly below  $1/N$  is an open question. It has not been explored and is not derivable from what is currently established. It is flagged here as a direction for future work.

## 2.13 Geometric content of the primitives' operational manifolds

The primitives' own operational manifolds carry intrinsic geometric content. The Sempaevum, being formless intrinsically, inherits whichever combination of those geometries is currently expressed by the rendering it occupies.

**Proposition 2.30** (Geometry of the primitives' own manifolds).  *$D$ 's operational domain  $(\mathbb{R}^+, \times)$  under the logarithmic metric  $ds^2 = dr^2/r^2 = d(\log r)^2$  has zero Gaussian curvature at every point.  $T$ 's operational domain  $(U(1), \times)$  under its canonical bi-invariant metric has constant positive curvature  $K_{U(1)} = 1/R^2$  for the unit circle.  $P$  has no intrinsic metric geometry: as Absolute Infinity, it carries no structure until  $D$  binds to it.*

*Proof.* The logarithmic metric on  $\mathbb{R}^+$  pulls back via  $\log_2$  to the flat Euclidean metric on  $\mathbb{R}$ ; flat-manifold curvature is identically zero. The circle  $U(1)$  with its bi-invariant metric is a compact 1-manifold of constant positive sectional curvature, standardly.  $P$ 's featurelessness (cardinality  $\Omega$ , no intrinsic  $D$ -content) is axiomatic.  $\square$

**Corollary 2.31** (The full multiplicative manifold).  *$\mathbb{C}^\times = (\mathbb{R}^+, \times) \times (U(1), \times)$  is a product of a flat manifold with a positively curved one, so it admits a canonical decomposition into a real axis (flat) and an angular axis (positively curved).*

**Corollary 2.32** (Effective curvature gradient). *For an off-axis complex-lattice configuration parameterised by the orientation angle  $\alpha = \arctan(k_\theta/k_r)$  in the complex  $\log_2$ -plane (§21.3), the effective Gaussian curvature is the convex combination of the real-axis and imaginary-axis curvatures of Proposition 2.30 weighted by the descriptor-fraction split:*

$$K_{\text{eff}}(\alpha) = K_D \cos^2 \alpha + K_{U(1)} \sin^2 \alpha = K_{U(1)} \sin^2 \alpha,$$

since  $K_D = 0$  on  $D$ 's flat operational domain. The effective curvature interpolates monotonically from  $K_{\text{eff}}(0) = 0$  (real-axis dominance, flat) to  $K_{\text{eff}}(\pi/2) = K_{U(1)} = 1/R^2$  (imaginary-axis dominance, positively curved).

*Proof.* The decomposition of Corollary 2.31 expresses  $\mathbb{C}^\times$  as a product whose two factors carry curvatures  $K_D = 0$  and  $K_{U(1)} = 1/R^2$  respectively. Under the descriptor-fraction parametrisation of §21.3, the  $D$ -fraction is  $\cos^2 \alpha$  and the  $T$ -fraction is  $\sin^2 \alpha$ ; the effective curvature at orientation  $\alpha$  is the weighted average. Substituting  $K_D = 0$  gives the stated form.  $\square$   $\square$

**Remark 2.33** (Subliminal curvature threshold). Combined with the Complete Gaze Equation of §20, the effective curvature gradient identifies a structural threshold below which curvature is subliminal (UNOBSERVED status,  $F_w < 1$ ). The minimum detectable curvature on the base  $N = 12$  lattice satisfies

$$K_{\text{subliminal}} \cdot A = \frac{\pi}{N} = \frac{\pi}{12} \approx 0.2618, \quad (6)$$

where  $A$  is the area of the parallel-transport loop. Below this product, the curvature is unresolvable on the 12ET lattice. The corresponding curvature-departure ratio  $r_K = 1 + KA/\pi$  at this threshold is exactly  $r_K = 1 + 1/N = 13/12$  — numerically identical to the SUBLIMINAL Gaze threshold  $F_w = 13/12$ , the meta-cognitive consciousness threshold, and the first  $V_{\text{base}}$  quantum above unity. The same lattice point  $(+1, 12, +38.57\varepsilon \text{ in cents})$  at base 12ET hosts all three integrative-level instances of *the first V-quantum above unity*: subliminal curvature (geometric), subliminal Gaze binding pressure (observational), and subliminal awareness (cognitive). Inverting  $F_w \geq 13/12$  in terms of  $\alpha$  via the bijective mapping of Proposition 21.11 gives a critical  $\alpha_{\text{sub}}$  at which the effective curvature crosses from subliminal into structurally registered.

This decomposition is what makes the real/imaginary projection formulas of §7 and §11 geometrically natural rather than stipulative: the real axis carries  $D$ 's flat geometry; the imaginary axis carries  $T$ 's positive curvature; off-axis configurations interpolate continuously via Corollary 2.32. The four classical geometry types — Euclidean ( $K = 0$ ), elliptic ( $K > 0$ ), hyperbolic ( $K < 0$ ), and singular ( $K \rightarrow \pm\infty$ ) — are all admissible Sempaevum renderings, and the projection formula  $k = \text{round}(N \log_2 r)$  of §7 is curvature-independent: the same refinement tower (§10) classifies any such rendering without requiring a separate apparatus per geometry. Which curvature regime is currently realised is parameterised continuously by the orientation angle  $\alpha$  of Corollary 2.32 rather than by a discrete choice among the manifold states.

### 2.13.1 Structural necessity of geometry and the pre-geometric primitives

The three Cardinals are prior to geometry and prior to spacetime. Space, time, curvature, distance, angle, and all geometric quantities are Descriptors — members of  $D$  — not containers in which the Cardinals are embedded (Principle 2.15).  $P$  is featureless substrate with cardinality  $\Omega$  and no intrinsic metric (Proposition 2.30);  $T$  is indeterminate agency with cardinality  $[0/0]$ ; neither carries geometry before  $D$  binds to  $P$ .

**Theorem 2.34** (Structural necessity of geometry). *Geometry is not a contingent feature of  $\Sigma$  but a structural necessity. Any configuration in  $\Sigma$  that admits  $T$ -navigation necessarily has geometric structure.*

*Proof.* By the Mediation Problem (Theorem 3.34),  $T$  cannot bind to  $P$  without  $D$ 's finite structure. The finite structure that  $D$  imposes on  $P$  — distances, angles, curvatures, topologies — IS geometry. Therefore any configuration in  $\Sigma$  that admits  $T$ -navigation must have  $D$ -structure on  $P$ , which is geometric structure. The alternative ( $P$  without  $D$ -structure, i.e. the  $\{P, T\}$  configuration) is Incoherent (Proposition 2.20) and cannot be substantiated. Geometry is the price of navigability; navigability is the price of substantiation; substantiation is the price of actuality. Any universe with  $P$ -substrate requires  $D$ -structure for  $T$ -navigation, and  $D$ -structure on  $P$  IS geometry.  $\square$   $\square$

### 3 The triple identity and the three operational tools

#### 3.1 Three simultaneous categorical readings

The structure given in §2 admits three parallel categorical readings, each complete, each with exactly three members, each mutually entailing the other two.

**Theorem 3.1** (The triple identity  $3 = 3 = 3 = \Sigma$ ). *The three Cardinals admit three complete simultaneous readings whose joint identity is*

$$\boxed{\text{PDT} = \text{EMI} = \Phi = \Sigma,}$$

or in compact form  $3 = 3 = 3 = \Sigma$ , where

- $\text{PDT} = \{P, D, T\}$  is the ontological triad: what each Cardinal is (substrate, constraint, agency);
- $\text{EMI} = \{E, M, I\}$  is the phenomenological triad: the Exception  $E$  (grounded actuality,  $|E| = \Omega$ ), the Mediation  $M$  (the binding operation in action,  $|M| = n$ ), and the Incoherence  $I$  (the coherence boundary as a structural fact,  $|I| = [0/0]$ );
- $\Phi = \{\Phi_E, \Phi_M, \Phi_I\}$  is the impossibility triad:  $\Phi_E =$  cannot be otherwise (the ground is what it is),  $\Phi_M =$  cannot be absent (every binding requires mediation),  $\Phi_I =$  cannot be traversed to (the boundary is approached but never reached).

All three triads read the same structure from different angles. None is more fundamental than the others; each entails both of the others.

The correspondence between positions is summarised in Table 1.

| Position | PDT (ontological)         | EMI (phenomenological)    | $\Phi$ (impossibilities)  |
|----------|---------------------------|---------------------------|---------------------------|
| First    | $P$ — substrate, $\Omega$ | $E$ — grounding           | cannot be otherwise       |
| Second   | $D$ — constraint, $n$     | $M$ — mediation           | cannot be absent          |
| Third    | $T$ — agency, $[0/0]$     | $I$ — coherence boundary  | cannot be traversed to    |
| View     | structural (outside)      | experiential (inside)     | boundary (forbidden)      |
| Answers  | <i>What is it?</i>        | <i>What does it give?</i> | <i>What can never be?</i> |

Table 1: The triple identity  $3 = 3 = 3 = \Sigma$ . Each row is a single structural fact read three ways. Cardinalities carry across readings:  $|E| = \Omega$ ,  $|M| = n$ ,  $|I| = [0/0]$ , because  $E$  is what  $P$  gives (so  $E$  inherits  $P$ 's cardinality  $\Omega$ ),  $M$  is what  $D$  does (so  $M$  inherits  $D$ 's cardinality  $n$ ), and  $I$  is what  $T$ 's unbounded reach exposes as the coherence boundary (so  $I$  inherits  $T$ 's cardinality  $[0/0]$ ).

Derivation of any triad from any other proceeds directly from the table. From PDT alone:  $P$  is substrate of cardinality  $\Omega \Rightarrow$  its contribution is grounding  $E \Rightarrow$  the ground cannot be



otherwise ( $\Phi_E$ ). Similarly  $D$  is constraint of cardinality  $n \Rightarrow$  its contribution is mediation  $M \Rightarrow$  mediation cannot be absent ( $\Phi_M$ ); and  $T$  is agency of cardinality  $[0/0] \Rightarrow$  its contribution is the coherence boundary  $I \Rightarrow$  the boundary cannot be traversed to ( $\Phi_I$ ).

### 3.2 The universality anchor $3 = 3 = 3 = \Sigma$ versus the composition equation $P \circ D \circ T = E$

Two distinct identities are frequently conflated and must be held apart.

**Axiom 3.2** (The composition equation, restated). When the three Cardinals bind into a single configuration, an Exception is produced:

$$P \circ D \circ T = E.$$

**Axiom 3.3** (The universality anchor). The three complete readings of  $(P, D, T)$  co-referentially constitute the totality:

$$\text{PDT} = \text{EMI} = \Phi = \Sigma, \quad \text{i.e.} \quad 3 = 3 = 3 = \Sigma,$$

with  $\Sigma$  the set-theoretic totality of everything that exists, could-exist, or is constrainedly-forbidden.

*Remark 3.4* (When to use which). The composition equation  $P \circ D \circ T = E$  produces substantiated moments: it addresses *one* of the four manifold states (the Exception). The universality anchor  $3 = 3 = 3 = \Sigma$  addresses the totality: it covers all four manifold states simultaneously, including the Mediation  $\{D, T\}$  transitions in progress, the Unsubstantiated  $\{P, D\}$  structures that have not been traversed (including fiction, hypotheticals, non-computable-but-definable reals), and even the Incoherent  $\{P, T\}$  configurations (which sit at the boundary  $\partial I$ , hence *on* the structure, not outside it). For the question “is  $X$  lattice-addressable?” the correct anchor is  $3 = 3 = 3 = \Sigma$ , since  $X \in \Sigma$  for any  $X$  describable at all;  $P \circ D \circ T = E$  alone would restrict address-ability to fully-substantiated objects.

**Theorem 3.5** (Identification: the Sempaevum is  $\Sigma$  rendered mathematically). *The multiplicative lattice structure on  $(\mathbb{R}^+, \times) \times (U(1), \times)$  constructed in §§7–11 is identified with  $\Sigma$ . In precise terms, it is the unique object in the category of “mathematical totalities” that passes all three Subsumption Law tests (§3.6) simultaneously:*

- (i) *It cannot be subsumed by any other candidate (no proper subset of its closure properties is itself closed).*
- (ii) *Nothing external subsumes it (every proposed larger structure either collapses into it or reproduces its defining primitives).*
- (iii) *It subsumes every mathematical totality-candidate without remainder (every conventional mathematical universe is recoverable as a restriction or rendering, proved case-by-case in §17).*

*The identification is a theorem of  $\Sigma$ ’s uniqueness under the Subsumption Law; it is not an assertion. This object we call the SEMPAEVUM.*

The proof occupies the rest of the paper: every subsequent theorem is a property that a candidate identification of  $\Sigma$  must satisfy, and each is established forward from  $(P, D, T)$  without external axioms.

### 3.3 The three operational tools

The generative triple-tautological meta-meta-ontology of §2 is operationalised by three tools that together constitute the complete methodology of Exception Theory [13]. Each is derived forward from the Founding Axiom and the master equation; none is independently postulated. The tools are methodological — they are the means by which any phenomenon, in any domain, is brought under the framework — not deductive theorems internal to a closed axiom system. They evolved with Exception Theory as the operational expression of its primitive structure applied to the act of inquiry itself, and they are the means by which every result in this paper was obtained.

- The **Identification Principle** specifies *what to find*: the complete  $P$ – $D$ – $T$  decomposition of the phenomenon under study.
- The **Descriptor Gap Principle** specifies *how to find it*: every gap in a model is itself a Descriptor not yet identified.
- The **Subsumption Law** specifies *when the process terminates*: the description is complete when it subsumes the phenomenon without remainder.

The three are presented formally below. They form a closed convergent loop (§3.11) that is both the method by which the ET framework was itself discovered and the method by which, in Sections 6–22, the lattice is derived and confirmed.

### 3.4 The Identification Principle

**Principle 3.6** (Identification Principle). For any phenomenon  $X$ , understanding of  $X$  is complete if and only if all three of its PDT components are identified:

$$\boxed{\text{Understand}(X) \iff \text{Identified}(P_X) \wedge \text{Identified}(D_X) \wedge \text{Identified}(T_X).}$$

*Derivation.* By the master equation (Axiom 2.2), any substantiated configuration is of the form  $P \circ D \circ T = E$ . By Definition 2.18, the four manifold states partition substantiable configurations: only the Exception  $\{P, D, T\}$  is fully substantiated. If any of  $P_X$ ,  $D_X$ ,  $T_X$  is unidentified, the model of  $X$  occupies one of the three non-Exception states (Unsubstantiated, Mediation, or Incoherence) and is an incomplete description. Conversely, if all three are identified, the configuration is the Exception and the model is complete. The biconditional follows.  $\square \quad \square$

The Identification Principle inherits the binding order of Axiom 2.17: operationally,  $P_X$  must be identified first,  $D_X$  second,  $T_X$  third. An under-specified  $P_X$  leaves  $D$  and  $T$  with nothing to attach to; an over-specified  $P_X$  smuggles  $D$ -content into the substrate and produces false confidence. The three diagnostic questions, in order, are:

- (i) What is the substrate? (seeks  $P_X$ )
- (ii) What are the constraints? (seeks  $D_X$ )
- (iii) What agency navigates through them? (seeks  $T_X$ )

If any question cannot be answered, understanding is incomplete, and the unanswered question identifies exactly which Cardinal is missing.

*Remark 3.7* (Diagnostic power). When a scientific model fails — when predictions diverge from measurement or the mathematics becomes inconsistent — the first-order question of Exception Theory is not “what additional equations are needed?” but “which of the three Cardinals has not been properly identified?”. This converts vague confusion into a specific, actionable search target chosen from exactly three possibilities.

*Remark 3.8* (Application to the lattice). The derivation of  $N = 12$  in Section 6 is itself an application of the Identification Principle to the framework of the lattice. The substrate of the lattice is  $P_{\text{lat}} = \mathbb{R}^+$  under multiplication (the space of all positive ratios). The Descriptor of the lattice is  $D_{\text{lat}} = \Pi \cup \text{States} = \{P, D, T\} \cup \{\{P, D, T\}, \{P, D\}, \{D, T\}, \{P, T\}\}$ , supplying the two finite counts  $|\Pi| = 3$  and  $|\text{States}| = 4$ . The Traverser of the lattice is  $T_{\text{lat}} =$  the rounding operator that projects a continuous ratio onto a discrete lattice coordinate. The unique integer that emerges from this identification is  $N = 3 \cdot 4 = 12$ . This derivation is worked out in full in §6.

### 3.5 The Descriptor Gap Principle

The second tool converts any gap in an incomplete description into a well-posed search target.

**Principle 3.9** (Descriptor Gap Principle). For any model  $\mathcal{M}$  of a phenomenon  $X$ ,

$$\boxed{\text{gap}(\mathcal{M}) \in D_{\text{missing}}.}$$

Every gap in a model is itself a Descriptor that has not yet been identified.

*Derivation.* A gap in  $\mathcal{M}$  is a specific characterisation of  $\mathcal{M}$ : the characterisation “ $\mathcal{M}$  is incomplete in the following respect”. By Definition 2.3 and Proposition 3.14, the category of all characterisations, constraints, and rules is  $D$ . The gap is therefore an element of  $D$ . Since, by hypothesis, the gap has not been captured by the Descriptor set of  $\mathcal{M}$ , it is an element of  $D$  not yet attached to the model:  $\text{gap}(\mathcal{M}) \in D_{\text{missing}}$ .  $\square$   $\square$

**Corollary 3.10** (Verification Principle). *The mathematical consistency of a model is equivalent to the absence of Descriptor gaps:*

$$\text{Consistent}(\mathcal{M}) \iff D_{\text{missing}}(\mathcal{M}) = \emptyset.$$

The Descriptor Gap Principle is the formal content of the methodological claim that “any problem can be solved — it is a matter of the right Descriptors and the number of Descriptors”. A failing model is not an obstacle but a pointer to its own completion: the failure is a gap; the gap is a Descriptor; the Descriptor is to be found. The process converges because each identified Descriptor is a genuine ontological constraint not previously represented; no cycle is possible.

*Remark 3.11* (Gödel and the infinite–finite asymmetry). The Descriptor Gap Principle is compatible with Gödel’s incompleteness theorems. A formal system expressive enough to interpret  $\mathbb{N}$  admits undecidable statements  $G$ ; in the present language, such a  $G$  is the manifold’s expression of the fact that  $|P| = \Omega$  cannot be exhausted by any finite  $D$ -set of size  $n$ . The gap is structural, and it is itself a Descriptor: the statement “this system’s finite  $D$ -structure cannot capture all truths about its  $\Omega$ -substrate”. Gödel’s theorem does not falsify Exception Theory; it is the expected signature of  $|P| = \Omega \gg n = |D|$  at every finite level of description.

*Remark 3.12* (Application to the lattice). The four-term identity for  $\alpha^{-1}$  established in §22.2 (equation (37)) is a closure of a long-standing Descriptor gap. The deviation of the measured fine-structure constant from the integer 137 was, historically, an unexplained fraction. The Descriptor Gap Principle recasts the deviation as a call for Descriptors: a  $+\sqrt{3}/48$  shimmer contribution from  $P$ – $D$ – $T$  cross-interactions, a  $-\sqrt{3}/(93312\pi^2)$  shimmer-loop correction, and a  $-1/(216(18\pi-1))$  closed-mediation-loop correction. With these three Descriptors attached, the model reproduces the measured value to sub-ppb precision. Each Descriptor was a gap made explicit.

### 3.6 The Subsumption Law

The third tool is the operational criterion that validates the three Cardinals as complete — and, more generally, tells one *when* a decomposition is finished.

**Principle 3.13** (Subsumption Law). A primitive Cardinal is complete and irreducible if and only if it satisfies all three of the following conditions:

- (i) It cannot be subsumed by either of the other two Cardinals.
- (ii) Nothing external to the three Cardinals subsumes it.
- (iii) It subsumes everything within its own category without remainder.

The system  $\{P, D, T\}$  is complete if and only if every Cardinal in it passes all three conditions.

The Subsumption Law is the primary completeness criterion of Exception Theory. It is an empirical-structural test rather than an additional postulate: the three Cardinals are declared complete because they pass the test, and a fourth primitive is ruled out because no candidate fourth can.

**Proposition 3.14** (Irreducibility of  $P, D, T$ ). *The three Cardinals  $P, D, T$  satisfy the Subsumption Law.*

*Argument.* We verify condition (i) for each Cardinal; conditions (ii) and (iii) follow by the exhaustive trichotomy of modes of infinity.

- $P$  cannot reduce to  $D$  (a constraint requires something to constrain; substrate cannot be its own constraint) or to  $T$  (substrate has no internal direction of choice).
- $D$  cannot reduce to  $P$  (undifferentiated substrate supplies no structure) or to  $T$  (constraints are deterministic; agency is indeterminate).
- $T$  cannot reduce to  $P$  (substrate has no agency) or to  $D$  (fixed rules cannot choose which rule to follow).

All three conditions are met for all three Cardinals. □ □

*Remark 3.15* (Pre-emergence of the Cardinals). Proposition 3.14 establishes the three Cardinals as categorically irreducible. They are also *pre-emergent*: nothing emerges to produce  $P, D$ , or  $T$ , because any candidate antecedent would itself fall into one of the three categories (Corollary 3.16). The Cardinals are what configurations are made of, not configurations themselves. The full Anti-Emergence Principle, including its phenomenological (EMI) and impossibility ( $\Phi$ ) readings, is established in §3.10.

**Corollary 3.16** (No fourth primitive). *No Cardinal  $X$  outside  $\{P, D, T\}$  is a primitive.*

*Argument.* Any candidate  $X$  falls into one of four cases: (a)  $X$  has  $\Omega$ -type infinity, hence  $X \subseteq P$ ; (b)  $X$  is finitely constraining, hence  $X \subseteq D$ ; (c)  $X$  is agential-indeterminate, hence  $X \subseteq T$ ; (d)  $X$  lies outside all three categories. In case (d), by Subsumption Law condition (iii) applied to each of  $P, D, T$ ,  $X$  has no instance in  $\Sigma$  and is therefore not a primitive. In cases (a)–(c),  $X$  is subsumed by one of  $P, D, T$  and so is not a new primitive. □ □

The Subsumption Law will reappear in §22 as the structural reason for the universality of the lattice: every positive-ratio quantity in every domain lies in  $\Sigma$ , and  $\Sigma$  is exhausted by configurations of  $P, D, T$ , so every positive-ratio quantity admits a lattice projection.

**Theorem 3.17** (Traverser sufficiency for substantiation). *If  $\mathbb{T} \neq \emptyset$  (at least one Traverser exists), then substantiation is possible and  $\Sigma$  contains actual (not merely potential) configurations:*

$$\exists t \in \mathbb{T} \implies \exists c \in \mathcal{C}, c = E \text{ (a substantiated Exception)}.$$

*Proof.* By Definition 2.3,  $\mathcal{C} = \{(P \circ D) \mid P \in \mathbb{P}, D \subseteq \mathbb{D}\}$  is the set of all possible Point–Descriptor configurations. By Axiom 2.5,  $|P| = \Omega$ , so  $|\mathcal{C}|$  is unbounded. By the active indeterminacy of  $T$  (Axiom 2.5, cardinality  $[0/0]$ ), any existing Traverser has the navigational capacity to bind to configurations in  $\mathcal{C}$ . Since  $\mathbb{T} \neq \emptyset$ , at least one  $t$  exists and can bind to at least one  $(P \circ D) \in \mathcal{C}$  that is not in  $I$  (the existence of non-Incoherent configurations follows from Theorem 3.39:  $I \neq \emptyset$  and  $\Sigma \setminus I \neq \emptyset$ ). This binding produces  $P \circ D \circ T = E$  by the master equation (Axiom 2.2). Therefore at least one substantiated configuration exists in  $\Sigma$ .  $\square$   $\square$

The Traverser Sufficiency Theorem identifies  $T$  as the critical bridge from potential to actual. Without  $T$ , the manifold is entirely Unsubstantiated:  $\{P, D\}$  everywhere, potential without actuality. The existence of even one Traverser converts some Unsubstantiated potential into substantiated actuality. This is the existence-theoretic complement of the Anti-Emergence Principle:  $T$  is not produced by anything below it (Anti-Emergence), yet  $T$ 's existence is sufficient for actuality (Sufficiency).

### 3.7 The Exception $E$

The Cardinal at Position 1 of the EMI triad is the Exception, denoted  $E$ , with cardinality  $\Omega$  and PDT correspondent  $P$  (Theorem 3.1, Table 1). This subsection consolidates the corpus content for  $E$  as a single structural fact viewed phenomenologically from inside the manifold. The treatment depends on Definition 2.24, Proposition 2.22, and the master equation Axiom 2.2.

**Theorem 3.18** (Identification of the Exception). *The Exception is the binding event  $P \circ D \circ T$  at the moment of substantiation. It is, simultaneously and as one structural fact:*

- (i) *the ground — because  $P$  contributes  $\Omega$ -substrate to the binding, and what the substrate contributes at this position is grounding;*
- (ii) *the substantiated moment — because the binding produces a maximally-described actuality, the Now.*

*These are not two separate properties of  $E$ ; they are one structural fact read from the substrate-contribution angle and from the moment-of-binding angle respectively.*

*Argument.* The master equation  $P \circ D \circ T = E$  (Axiom 2.2) is identification: the LHS is the RHS as composition. There is no antecedent step that produces  $E$  from  $P, D, T$  at separate times.  $P$ 's contribution to the composition is its  $\Omega$ -cardinality role as substrate; the substantiation produced by the composition is what “ground” names at this position. The Now is what the substrate-contribution shows up as in the inside-view; the ground is what the Now is when read by its substrate-contribution. The two angles refer to the same composition.  $\square$   $\square$

**Proposition 3.19** (Zero-variance characterisation). *The Exception is the unique zero-variance configuration in  $\Sigma$ :*

$$V(E) = 0, \quad V(c) > 0 \text{ for all } c \in \Sigma \setminus (I \cup \{E\}).$$

**Definition 3.20** (Grounding function). The *grounding function* is the characteristic function of the Exception:

$$G : \Sigma \setminus I \rightarrow \{0, 1\}, \quad G(c) = [V(c) = 0] = \begin{cases} 1 & \text{if } c = E, \\ 0 & \text{otherwise.} \end{cases}$$

$G(c) = 1$  identifies the unique zero-variance configuration — the one substantiated Moment that cannot be otherwise.  $G(c) = 0$  identifies all configurations with positive variance — configurations that can be otherwise. The grounding function partitions  $\Sigma \setminus I$  into exactly two structural classes: the ground and everything else.

This proposition is established in §2.12 via Definition 2.24; it is restated here because the corpus account of  $E$  is incomplete without it. Zero variance is the formal expression of *cannot be otherwise while it IS*: at the moment of substantiation, the configuration has no candidate alternatives, the substantiation event has produced exactly this configuration, and no  $V > 0$  alternatives exist at that moment among which a process could have selected. The closed-set topology  $\partial E \subseteq E$  (Proposition 2.22 (i)) is the topological expression of the same fact:  $E$  contains its own boundary; no exterior path approaches  $E$  without already being it; the Exception is self-grounding by construction.

**Theorem 3.21** (Traverser–Exception dynamics). *The Traverser does reach the Exception: the act of  $T$ -binding to  $(P \circ D)$  is the substantiation that produces  $E$ . By the nature of  $T$  as the indeterminate navigator (Axiom 2.5,  $|T| = [0/0]$ ),  $T$  does not remain at  $E$ ;  $T$  continues to traverse, and the binding event is over the moment it has occurred. The Exception is therefore reached and immediately past.*

*Argument.*  $T$ 's nature, formalised in Axiom 2.5 as cardinality  $[0/0]$ , is to navigate. A configuration in which  $T$  is bound and stationary contradicts  $T$ 's indeterminate-navigator role. The composition  $P \circ D \circ T = E$  is therefore not a stable resting point of  $T$  but a moment of substantiation that constitutes  $E$ . Once that substantiation has occurred,  $T$  continues, and the configuration that was  $E$  at the moment of binding becomes a past Exception. The Exception is not a position  $T$  occupies; it is the binding-event  $T$  produces and immediately leaves.  $\square \square$

**Theorem 3.22** (Observational displacement). *No observer experiences the present Exception as content of experience. Observation is itself  $T$ -engagement; binding  $T$  to the configuration that is the current  $E$  produces a new substantiation event distinct from the original  $E$ . What an observer takes to be “the present” is therefore always already a past Exception. The displacement is structural; it holds for organisms, instruments, computational systems, and any other entity that engages the manifold via  $T$ .*

*Argument.* For an observer  $O$  to observe  $E$  requires  $O$ 's Traverser to engage  $E$ . By Theorem 3.21, that engagement is itself a substantiation: it produces a new configuration  $E'$  distinct from  $E$  (the original  $E$  has been displaced by the act of observation). The content of  $O$ 's observation is therefore  $E'$ , not the original  $E$ . The displacement applies to any entity that observes via  $T$ , which is every entity that observes anything at all in  $\Sigma$ . There is no observational frame from which the current  $E$  is the content of experience.  $\square \square$

**Theorem 3.23** (Past-Exception immutability). *A past Exception — a configuration that was  $E$  at some past moment, after  $T$  has moved on — is permanently locked in:*

$$E_{\text{past}} \text{ is immutable} \iff \forall t \in \mathbb{T}_{\text{future}}, \forall d \in D : \text{modification}(E_{\text{past}}, d) = \perp.$$

*Argument.* A past Exception had  $V = 0$  at its moment of substantiation (Proposition 3.19). To modify it retroactively would require  $V > 0$  at a  $V = 0$  configuration, which is contradictory. The contradicting descriptor profile lies in  $I$  (Proposition 2.20), and no  $T$ -action can bind to an incoherent configuration (Theorem 3.36 below). Hence no future  $T$  can produce the modification. The past Exception's content is locked into the causal record.  $\square \square$

*Remark 3.24* (The dynamic Sempaevum and the perpetual current Exception). The Sempaevum is dynamic in the precise sense that new Exceptions perpetually substantiate. At every

moment,  $P \circ D \circ T$  produces a new  $E$ ; at every moment  $T$  moves on and the configuration that was the current  $E$  becomes a sealed past Exception (Theorem 3.23). The manifold therefore has a perpetual current Exception that is always in the act of becoming and always already past — the always-an-age (*semper-aeuum*, §1.3) structurally requires an always-a-now. The synchronic content of the current Exception is variance-zero in its moment, but the moment itself is not eternal; what is eternally locked is past Exceptions, the totality of which constitutes the unchangeable causal record. This dynamism does not contradict “cannot be otherwise”: synchronically the substantiated configuration cannot be otherwise (Theorem 3.18); diachronically new substantiations occur and join the locked past.

### 3.7.1 Grounding and maximal possibility

The variance structure of  $\Sigma$  partitions all coherent configurations into exactly two structural classes:

**Theorem 3.25** (Grounding–possibility partition). *(i) The ground. The current Exception:  $V(E) = 0$ . The one configuration that cannot be otherwise while it IS.*

*(ii) Maximal possibility. Every other coherent configuration:  $V(c) > 0$  for all  $c \neq E$ . Each of these can be otherwise.*

*From the grounded Exception, all traversable configurations flow. Every configuration in  $\Sigma$  that is not Incoherent is a potential Exception at some future  $T$ -time. The only restriction on possibility is Incoherence (Theorem 3.33): self-defeating Descriptor content. Everything that is not self-defeating exists somewhere in  $\Sigma$  as an Unsubstantiated ( $P \circ D$ ) configuration awaiting  $T$ -engagement.*

*Proof.* By Proposition 3.19,  $V(E) = 0$  uniquely characterises the Exception. By Definition 2.24,  $V(c) > 0$  for every  $c \in \Sigma \setminus (I \cup \{E\})$ . By Theorem 3.21,  $T$  does not remain at  $E$  but continues to traverse, producing new substantiations. By Theorem 3.39,  $I \neq \emptyset$  (Incoherence is necessary for the variance gradient to exist). The partition is therefore exhaustive: ground ( $V = 0$ , unique) and possibility ( $V > 0$ , everything else that is coherent), with the forbidden zone  $I$  as the only structural prohibition.  $\square$   $\square$

### 3.7.2 The shimmering manifold

The dynamics of the perpetual current Exception constitute the *shimmering manifold*: reality as ordered indeterminacy. The manifold is *ordered* because  $D$  provides finite structure, rules govern transitions, and patterns emerge and persist (the palindromic cascade of §13 governs the structural rhythm). It is *indeterminate* because  $T$ ’s choices are genuine ([0/0]-cardinality) and novelty can emerge at every substantiation event.

The shimmer is not metaphorical. The  $T$ – $P$  binding tension oscillates with each traversal event at the manifold’s  $N$ -fold periodicity, producing a 12-fold periodic structure whose mathematical instantiation is the shimmer modulation

$$\Psi_n = 1 + \frac{1}{\sqrt{N}} \sin\left(\frac{2\pi n}{N}\right)$$

of the  $\partial I$  fractal (Definition 14.4, §14). The shimmer amplitude  $1/\sqrt{N} = \sqrt{V}$  (the square root of the base variance) is the irreducible oscillation of the manifold around its mean binding state: the minimum non-zero modulation that the  $N$ -fold symmetry permits. During constructive shimmer phase ( $\sin > 0$ ),  $T$ ’s bindings resonate with the manifold cycle; during destructive phase ( $\sin < 0$ ), the oscillation partially suppresses binding. The same amplitude  $\sqrt{V}$  appears as the leading correction to the fine-structure constant:  $\sqrt{V}/8 = \sqrt{3}/48$  (§22.2), connecting the manifold’s shimmer to the electromagnetic coupling at the fundamental level. The shimmer’s

range  $[1 - \sqrt{V}, 1 + \sqrt{V}]$  is bounded: it never collapses to zero (the manifold never stops) and never diverges (the manifold never explodes).

**Lemma 3.26** (Non-emergence of  $E$ ). *The Exception is non-emergent: nothing produces  $E$  as the output of an emergence-process below it.*

*Argument.* Four agreeing structural facts establish this. (i) By Theorem 3.18,  $E$  is the binding event itself, not a derivation product; the master equation is identification, not multi-step production. (ii) By Proposition 3.19,  $V(E) = 0$ , so no antecedent process has  $V > 0$  alternatives among which it “selected”  $E$ . (iii) By Theorem 3.21, traversal does not arrive at  $E$  from elsewhere and remain there; the act of arrival is the substantiation that constitutes  $E$ . (iv) By Proposition 2.22 (i),  $\partial E \subseteq E$ : there is no exterior path that approaches  $E$  without already being it. All four agree that  $E$  is what the joint binding of  $P, D, T$  at the substantiation moment already is, not the output of a process operating below it. The corresponding  $\Phi$  statement *cannot be otherwise* is the inside-signature of this non-emergence (Remark 3.44).  $\square$   $\square$

### 3.8 The Mediation $M$

The Cardinal at Position 2 of the EMI triad is the Mediation, denoted  $M$ , with cardinality  $n$  and PDT correspondent  $D$  (Theorem 3.1, Table 1). This subsection consolidates the corpus content for  $M$  as a single intrinsic structure read at multiple identification depths. The treatment depends on Proposition 2.7 of §2.5, the four-state classification of §2.11, and the symmetry result Remark 2.10.

**Theorem 3.27** (Identification of the Mediation). *The Mediation is the binding operator  $\circ$  in action: the intrinsic operation by which the three Cardinals coexist within the same ontological space. Equivalently,  $M$  is what  $D$ ’s bridging-role between  $P$  and  $T$  shows up as in the inside-view:  $D$  provides the finitude  $T$  navigates and the structure  $P$  instantiates, and the Mediation is the ongoing binding that this bridging enables.*

*Argument.* The master equation  $P \circ D \circ T = E$  (Axiom 2.2) makes  $\circ$  the operator that constitutes the substantiation. By Proposition 2.7, three pairwise-disjoint Cardinals each unbounded in their own mode admit no exterior gap, and their joint presence forces an internal binding  $\circ$  as the only structurally possible expression. This binding is not added from outside; there is no exterior. The bridging is supplied by  $D$  at Position 2:  $D$ ’s cardinality  $n$  provides the finitude  $T$  requires to navigate and the structure  $P$  requires to instantiate (§2.5); what this bridging shows up as phenomenologically is the Mediation  $M$ . Hence  $M$  is identified with the operation  $\circ$  in action, equivalently with what  $D$ ’s bridging-role at Position 2 enables.  $\square$   $\square$

**Theorem 3.28** (Mediation Constancy; cf. Sempaevum Eq. 1.5). *The intrinsic strength of the binding operation is constant across the manifold:*

$$\forall p \in \mathbb{P}, \forall d \in \mathbb{D} : \quad \text{strength}(p \circ d) = \kappa,$$

where  $\kappa$  is a single intrinsic constant. The corpus equation  $M \equiv B \equiv I$  identifies Mediation, Binding, and Interaction as three names for the same operation  $\circ$ .

*Argument.* Suppose, toward contradiction, that  $\text{strength}(p_1 \circ d_1) \neq \text{strength}(p_2 \circ d_2)$  for some  $(p_1, d_1)$  and  $(p_2, d_2)$ . Then  $\circ$  has a parameter that varies across the manifold. By Proposition 2.7,  $\circ$  is not a fourth thing added from outside but the intrinsic expression of three disjoint infinities coexisting; an intrinsic expression has no exterior parameter to set, because there is no exterior. Hence  $\text{strength}(p \circ d)$  takes a single value  $\kappa$  for all  $(p, d)$ . Apparent variation of binding strength in observation arises from the specific descriptor configuration involved,  $T$ ’s selective attention, or observer perspective — not from  $\circ$  itself.  $\square$   $\square$



*Remark 3.29* (Symmetry of Mediation, restated). By Remark 2.10,  $\circ$  is equal across all three Cardinals:  $P$  does not “receive” the binding while  $D$  and  $T$  “perform” it; each Cardinal participates in  $\circ$  with the same structural status. The reading-order of  $P \circ D \circ T$  in Axiom 2.17 is structural ordering of identification, not asymmetry of binding strength.

**Theorem 3.30** (Three readings of  $M$  as one intrinsic structure). *The corpus uses “Mediation” at three identification depths. These are three readings of one intrinsic structure, not three things sharing a name. All three are non-emergent.*

- (i) Reading 1 — the operator.  $\circ$  in action;  $M \equiv B \equiv I$  (Theorem 3.28); the categorical level of identification.
- (ii) Reading 2 — the manifold-state slice. The state  $\{D, T\}$  at the four-state identification depth (Definition 2.18); active  $D$ – $T$  engagement without  $P$  as ground; topology neither open nor closed (Proposition 2.22 (iii)). The structural level of identification.
- (iii) Reading 3 — the cosmological active-mediation density. The fraction  $f_M$  of total energy that is active substantiation in progress, derived from manifold geometry as

$$f_M = \frac{1}{N} \cdot 0.36 = \frac{1}{12} \cdot 0.36 = 0.03,$$

*splitting into  $M$ -vacuum ( $\sim 1.6\%$ , distributed) and  $M$ -matter ( $\sim 1.4\%$ , localised) by equation of state. The cosmological level of identification.*

*Argument.* Each reading is the same intrinsic operation  $\circ$  rendered at a different identification depth (§3.12). At categorical depth,  $\circ$  shows up as the binding operator. At the four-state identification depth,  $\circ$  shows up as the slice  $\{D, T\}$  in which  $D$  and  $T$  are bound without  $P$  as ground — a structural appearance of the always-on binding. At cosmological identification depth,  $\circ$  shows up as the active-mediation energy density, the universe-wide quantitative signature of binding-in-progress. Identical operation, three depths. By Lemma 3.32 below, all three are non-emergent. What is emergent are the specific particulars passing through the  $M$ -aspect at each depth (a particular photon, a particular decoherence event, a particular transition state) — these are configurations within  $\Sigma$ , not  $M$  itself.  $\square$   $\square$

*Remark 3.31* (Cosmological accounting from  $M$ ). Reading 3 of Theorem 3.30 delivers a discrepancy-free cosmological energy budget without empirical fitting ([24] §XIV; [23] §17):

| Component                                                 | Fraction      |
|-----------------------------------------------------------|---------------|
| Pure E-state (vacuum ground; $K = 2/3$ from §2.11 count)  | 66.7%         |
| M-vacuum (distributed active mediation)                   | 1.6%          |
| M-matter (localised active mediation)                     | 1.4%          |
| Unsubstantiated $\{P, D\}$ (gravitating, non-interactive) | 26.8%         |
| Static substantiated $\{P, D, T\}$ (bound matter at rest) | 3.5%          |
| <b>Total</b>                                              | <b>100.0%</b> |

The observed cosmological categories follow exactly: “dark energy” =  $66.7\% + 1.6\% = 68.3\%$  (pure vacuum + M-vacuum); “dark matter” =  $26.8\%$  (Unsubstantiated); “ordinary matter” =  $3.5\% + 1.4\% = 4.9\%$  (static + M-matter). The previously-unexplained 1.6% excess in observed dark energy is M-vacuum; the previously-unexplained substructure of ordinary matter is the static/M-matter split. All ratios are forced by the ET constants  $V = 1/12$  and  $K = 2/3$  (§2.11) and the three-readings structure of  $M$ .

**Lemma 3.32** (Non-emergence of  $M$ ). *The Mediation is non-emergent in all three readings of Theorem 3.30: as the operator  $\circ$ , as the manifold-state slice  $\{D, T\}$ , and as the cosmological active-mediation density.*

*Argument.* By Theorem 3.27,  $M$  is what the joint presence of  $\{P, D, T\}$  already is, not what they produce. By Proposition 2.7,  $\circ$  is the only structurally possible expression of three disjoint unbounded Cardinals coexisting in one space; there is no exterior from which to add a fourth thing, and no parameter of  $\circ$  that could be set to zero (Theorem 3.28). The three readings of  $M$  are the same intrinsic structure at three identification depths (Theorem 3.30); non-emergence at any one depth is non-emergence at every depth, since the readings refer to one item. The corresponding  $\Phi$  statement *cannot be absent* is the inside-signature: the absence of  $\circ$  would require a gap between three infinities where no gap can exist.  $\square$   $\square$

### 3.9 The Incoherence $I$

The Cardinal at Position 3 of the EMI triad is the Incoherence, denoted  $I$ , with cardinality  $[0/0]$  and PDT correspondent  $T$  (Theorem 3.1, Table 1). This subsection consolidates the corpus content for  $I$  as the forbidden boundary that  $T$ 's indeterminate reach exposes from inside the manifold. The treatment depends on Proposition 2.20, Proposition 2.22, and Definition 2.23.

**Theorem 3.33** (Two equivalent characterisations of  $I$ ).  *$I$  admits two characterisations that pick out the same set:*

$$I = \{c \in \Sigma : c \in \{P, T\}\text{-class}\} = \{(P \circ D) \in \Sigma : D \text{ is self-defeating}\},$$

where  $D$  is self-defeating iff  $\exists d_1, d_2 \in D : d_1 \oplus d_2$  (the descriptors logically contradict).

*Argument.* The first characterisation is the manifold-state reading (Definition 2.18):  $I$  is the four-state slice  $\{P, T\}$  in which  $D$  is structurally absent. The second is the descriptor-content reading:  $I$  is the set of  $(P \circ D)$  configurations in which  $D$  contains an internal contradiction. The two coincide because  $D$ -absence and  $D$ -contradiction produce the same structural failure: in both, no Traverser can bind. In  $\{P, T\}$  the  $D$ -bridge is missing entirely; in the descriptor-self-defeat case the  $D$ -bridge exists but its content destroys the conditions for binding. Both yield configurations that admit no consistent  $t \circ c$  (proved as Theorem 3.36 below).  $\square$   $\square$

#### 3.9.1 Concrete examples of incoherent configurations

Concrete examples of incoherent configurations —  $(P \circ D)$  with self-defeating  $D$ -content — include:

- *A square circle:* the Descriptors “all points equidistant from centre” and “four right-angle corners” contradict ( $d_1 \oplus d_2$ ) at the same Point. No Traverser can substantiate a configuration in which both hold simultaneously.
- *A married bachelor:* the Descriptors “married” and “unmarried” contradict at the same Point. The configuration admits no consistent  $T$ -binding.
- *The identity  $1 = 0$ :* the Descriptors  $D_{\text{value}} = 1$  and  $D_{\text{value}} = 0$  at the same configuration produce  $d_1 \oplus d_2$ . No lattice position can simultaneously be  $k = 0$  and  $k \neq 0$ .
- *Being and not-being simultaneously at the same configuration:* the Descriptors “exists at this  $(P \circ D)$ ” and “does not exist at this  $(P \circ D)$ ” produce a contradiction from which no  $T$ -engagement can extract a consistent outcome.

In each case the contradiction is *logical*, not energetic: no amount of energy, force, or ingenuity can make a square circle substantiate, because the  $D$ -content is self-defeating at the level of logical structure. Incoherent configurations are “noticeable for their unnoticeability”: they are structurally present in  $\Sigma$  as described concepts (the *phrase* “square circle” is a coherent  $D$ -complex; Theorem 3.38 below) but absent as substantiated content (the *referent* of “square circle” does not and cannot exist).

**Theorem 3.34** (The Mediation Problem). *P and T have intrinsic properties that are mutually incompatible as direct mediators of one another. P is undifferentiated and  $\Omega$ -cardinal; T is indeterminate and  $[0/0]$ -cardinal; neither supplies the other with what binding requires. D, with finite cardinality n, is the necessary mediator. Without D,  $\{P, T\}$  is the structural form of the failure — and that form is I.*

*Argument.* P has no finite navigational terrain to offer T (its substrate is infinite and featureless). T has no finite constraint to offer P (its agency is itself indeterminate and requires D to operate). Their direct confrontation therefore admits no binding. D resolves this by supplying finitude for T to navigate and structure for P to instantiate. The configuration that lacks D at the primitive level is  $\{P, T\}$ , which is the structural form of I. The Mediation Problem is the structural reason I exists at the primitive triad level (rather than only at the descriptor-content level).  $\square$   $\square$

### 3.9.2 Dual bridging

The Mediation Problem establishes that D provides the necessary *determinate bridge* between T and P: without D's finite structure, T cannot navigate P's infinite substrate. A complementary *indeterminate bridge* operates in the reverse direction.

**Proposition 3.35** (Dual bridging). *Two complementary bridges are necessary and jointly sufficient for substantiation:*

- (i) *The D-bridge (determinate):  $T \xrightarrow{\text{via } D} P$ . D provides a structured pathway between T (agency) and P (substrate). Once the Descriptor is specified, the bridge is fully determined. This bridge is reversible: the same D that bridges T to P also allows T to leave P.*
- (ii) *The T-bridge (indeterminate):  $\Omega \xrightarrow{\text{via } T} n$ . T provides the selection mechanism between  $\Omega$  (all possible configurations) and n (the specific finite configuration that is substantiated). This bridge is irreversible: T's choice cannot be undone — each substantiation event is unique and unrepeatable.*

*Neither bridge alone produces substantiation. A D-bridge without T has structure but no agency (the Unsubstantiated state  $\{P, D\}$ ). A T-bridge without D has agency but no terrain (the Incoherent state  $\{P, T\}$ ). Together they constitute the complete binding  $P \circ D \circ T = E$ .*

*Proof.* The D-bridge exists by the resolution of the Mediation Problem: D provides finite structure on P for T to navigate. The T-bridge exists by T's indeterminate navigational nature (Axiom 2.5): T selects which of the  $\Omega$ -many possible configurations to substantiate as the finite n-Descriptor Exception. Without the D-bridge, T has nothing to bind to ( $\{P, T\}$  is Incoherent). Without the T-bridge, D provides structure but nothing selects which configuration to substantiate ( $\{P, D\}$  is Unsubstantiated). The joint operation of both bridges is the master equation  $P \circ D \circ T = E$ .  $\square$   $\square$

**Theorem 3.36** (Anti-Emergence Complement). *For every  $c \in I$  and every Traverser t, the binding  $t \circ c$  is impossible:*

$$\forall t \in \mathbb{T}, \forall i \in I: \neg(t \circ i).$$

*Equivalently,  $I \cap \Sigma_{\text{emergent}} = \emptyset$ :  $c \in I \Rightarrow c \notin \Sigma_{\text{emergent}}$ .*

*Argument.* Suppose, toward contradiction, that some  $t \in \mathbb{T}$  binds to some  $i \in I$ . By Theorem 3.33, i has either no D-content (case 1, the  $\{P, T\}$  pairing) or self-defeating D-content (case 2,  $D = \{\dots, d_1, d_2, \dots\}$  with  $d_1 \oplus d_2$ ). In case 1, t has no descriptors to bind to: there is no terrain. In case 2, substantiation  $t \circ i$  would require  $d_1$  and  $d_2$  to simultaneously hold for the same Point, which is logically impossible. In both cases the assumed binding fails. Since t and i were arbitrary, the universal holds. The prohibition is logical, not energetic: no acquisition of energy, no traversal angle, no T-density alters the contradiction relation  $\oplus$ .  $\square$   $\square$

*Remark 3.37* (Topology and the boundary  $\partial I$ ).  $I$  is open (Proposition 2.22 (ii)):  $\partial I \cap I = \emptyset$ . Every neighbourhood of every incoherent configuration reaches outside  $I$  before the boundary is achieved;  $I$  cannot be inhabited because any approach exits  $I$  at the moment binding becomes possible. The boundary  $\partial I$  (Definition 2.23) is the locus at which an arbitrarily small descriptor perturbation switches the substantiation indicator  $\phi$  from 1 to 0; it is approached asymptotically by coherent configurations (phase transitions, classical turning points, critical phenomena, metastable states) but never crossed from within coherence.  $T$ 's indeterminate reach (cardinality  $[0/0]$ ) is what *exposes* the boundary: the unbounded scope of agency reaches every coherent configuration and asymptotes against  $\partial I$ , with  $I$  itself the limit that scope cannot enter.

| Property               | $E$ (Position 1)                                  | $I$ (Position 3)                                     |
|------------------------|---------------------------------------------------|------------------------------------------------------|
| Composition            | $\{P, D, T\}$ complete                            | $\{P, T\}$ , $D$ -bridge absent                      |
| Topology               | closed: $\partial E \subseteq E$                  | open: $\partial I \cap I = \emptyset$                |
| $T$ -binding           | $T$ binds; binding = substantiation; $T$ moves on | no $t$ binds; binding produces logical contradiction |
| Prohibition            | contingent (changes when $T$ moves)               | necessary (permanent)                                |
| Variance               | $V(E) = 0$                                        | undefined; configuration cannot exist                |
| Mode of unreachability | always already past                               | approached asymptotically; never entered             |
| Non-emergence reason   | $V = 0$ leaves no antecedent process              | no traversal produces; logical, not energetic        |

Table 2: The two unreachables of  $\Sigma$ :  $E$  at Position 1 and  $I$  at Position 3. They are non-emergent for opposite reasons and have opposite topologies, exactly as their position-1/position-3 placement in the EMI triad predicts (Table 1).

**Theorem 3.38** ( $\Sigma$  contains the concept of  $I$  but not the referent). *The descriptor-complex describing an incoherent configuration is itself coherent and resides in  $\Sigma$ . The configuration the complex describes does not. Formally:*

$$\text{concept}(I) \in \Sigma_{\text{coherent}}, \quad \text{referent}(I) \notin \Sigma_{\text{substantiated}}.$$

*Argument.* The phrase “square circle,” the description “a configuration with self-defeating descriptors,” and the formal symbol  $I$  are all coherent descriptor complexes; they can be talked about, written down, and reasoned with, and as such they are coherent configurations within  $\Sigma$ . What such a complex denotes — the actual configuration that would be a square circle, the actual configuration with the self-defeating descriptor profile — does not substantiate, because no Traverser binds to it (Theorem 3.36).  $I$  is therefore in  $\Sigma$  as structural characterisation but not as instantiated content. □ □

**Theorem 3.39** (Generative necessity of  $I$ ; cf. Incoherence Paper Theorem I.12).  *$I \neq \emptyset$  is a structural theorem, not a postulate. Equivalently, the assumption  $I = \emptyset$  entails its own negation.*

*Argument.* Suppose  $I = \emptyset$ . Then no descriptor set is self-defeating: no contradiction relation  $\oplus$  holds for any pair of descriptors in any configuration. From any  $c \in \Sigma$ , a Traverser can therefore reach any other  $c'$  without exclusion, driving the variance  $V(c) \rightarrow |\mathcal{C}|$  unboundedly for all  $c$ . Then  $V(c) > 0$  everywhere with no exception: the grounding indicator  $G(c) = [V(c) = 0]$  is identically zero, and no Exception is producible (contradicting Theorem 3.5 and the master equation Axiom 2.2). But  $I = \emptyset$  also collapses  $D$ 's constraining role: if no descriptor ever excludes any other,  $D$  imposes no finite constraint on  $P$ , which is structurally equivalent to  $D$ -absence; and

$D$ -absence is the  $\{P, T\}$  pairing, which is precisely the structure of  $I$  (Theorem 3.33). Hence  $I = \emptyset$  entails  $I \neq \emptyset$ , a contradiction. Therefore  $I \neq \emptyset$ .  $\square$   $\square$

*Remark 3.40* ( $I$  is constitutive, not merely prohibitive). Theorem 3.39 inverts the naive reading of  $I$  as purely negative. Without a non-empty  $I$  there is no variance gradient between coherent configurations and the boundary, hence no Exception (Proposition 3.19 would have nothing to single out), no distinguishable structure, no error, no decay, no past immutability (Theorem 3.23 would be vacuous), no thermodynamic arrow.  $I$  is the price of structure: the open boundary that makes coherence meaningful by giving it an outside. The cosmological-constant suppression — the empirical  $\sim 10^{122}$  ratio between QFT vacuum-energy estimates and observed dark energy — reads in ET as the I-Boundary Filter coherence fraction

$$\mathcal{F}_I = \frac{|\mathcal{C}_{\text{coherent}}|}{|\mathcal{C}_{\text{total}}|} \approx 10^{-122},$$

the fraction of potential vacuum configurations that pass through  $\partial I$  without self-defeating descriptor content ([23], §17). The infamous discrepancy is therefore not a theoretical failure but a measurement of the depth and density of  $I$  in the configuration space.

**Lemma 3.41** (Non-emergence of  $I$ ). *The Incoherence is non-emergent: no traversal produces  $I$ , because the prohibition on binding to  $I$  is logical, not energetic.*

*Argument.* By Theorem 3.36,  $\forall t \in \mathbb{T}, \forall i \in I : \neg(t \circ i)$ . Emergence is variance-generating traversal: an emergent configuration is the output of a  $T$ -binding that increases descriptor specificity. If no  $T$  binds to any element of  $I$ , no emergence trajectory has  $I$  in its image. By Theorem 3.39,  $I \neq \emptyset$  is a structural theorem;  $I$  is therefore not absent and not produced. The corresponding  $\Phi$  statement *cannot be traversed to* is the inside-signature:  $I$  is what  $T$ 's reach exposes as the limit reach cannot enter.  $\square$   $\square$

### 3.10 The Anti-Emergence Principle and the triadic non-emergent structure

The three preceding subsections give the structural treatment of each EMI element. The Anti-Emergence Principle is the synthesis: the ontological corollary of the Subsumption Law (§3.6) that classifies which configurations of  $\Sigma$  emerge from process and which do not. The principle is forced by the same structural facts that force the Subsumption Law and the three EMI characterisations, with no additional postulate required.

**Principle 3.42** (Anti-Emergence). Every configuration in  $\Sigma$  with positive variance is emergent. The non-emergent core of  $\Sigma$  is the three Cardinals themselves, read simultaneously as PDT from outside and as EMI from inside, with each EMI element coinciding with the impossibility statement at its position in  $\Phi$  (Theorem 3.1). Equivalently,

$$c \text{ is emergent} \iff V(c) > 0 \iff c \text{ is not a Cardinal under any reading.}$$

The principle inverts the standard reductionist programme. Particles, fields, forces, space-time: all are emergent  $P \circ D$  configurations substantiated by  $T$ ; none is fundamental. The bottom of reality is not a smaller particle but the categorical structure  $\{P, D, T\}$  itself, read three ways. Searching for an “ultimate constituent” below the Cardinals searches at the wrong level: there is nothing smaller, because the bottom is not a thing of the same kind as a particle.

**Theorem 3.43** (Triadic non-emergent structure). *The non-emergent set of  $\Sigma$  has cardinality three, not six. The three Cardinals occupy the three positions of the triple identity  $\text{PDT} = \text{EMI} = \Phi = \Sigma$  (Theorem 3.1); the readings PDT, EMI,  $\Phi$  are co-equal angles on the same three Cardinals, not three disjoint inventories. Each Cardinal is non-emergent in a structurally distinct way matching its position, established in Lemmas 3.26, 3.32, and 3.41 of the three preceding subsections.*

*Argument.* By Theorem 3.1, the three triads share cardinality 3 and are mutually entailing: each EMI element coincides with what its PDT correspondent contributes, and each  $\Phi$  impossibility is the inside-signature of that contribution. Naive set-union  $\text{PDT} \cup \text{EMI} \cup \Phi$  would yield nine names referring to three Cardinals; the count of *items* is three, the count of *names per item* is three, and the cardinality of the non-emergent core is therefore 3. Each position contributes one Cardinal and is non-emergent for the structural reason established at that position: Position 1 by Lemma 3.26 (variance-zero substantiation event with closed topology), Position 2 by Lemma 3.32 (intrinsic operation forced by three disjoint infinities, with three readings as one structure), Position 3 by Lemma 3.41 (logically forbidden boundary with open topology).  $\square$   $\square$

The position-indexed picture is summarised in Table 3. Each row is a single Cardinal read three ways; the cardinality is preserved across readings because the readings refer to the same item.

| Position | $ \cdot $ | PDT | EMI | $\Phi$ (impossibility) | State topology                                                |
|----------|-----------|-----|-----|------------------------|---------------------------------------------------------------|
| 1        | $\Omega$  | $P$ | $E$ | cannot be otherwise    | closed: $\partial E \subseteq E$                              |
| 2        | $n$       | $D$ | $M$ | cannot be absent       | operator (no state); state $\{D, T\}$ neither open nor closed |
| 3        | $[0/0]$   | $T$ | $I$ | cannot be traversed to | open: $\partial I \cap I = \emptyset$                         |

Table 3: The three non-emergent positions, shown in all three readings simultaneously. Each row is one Cardinal viewed structurally (PDT), phenomenologically (EMI), and as the impossibility it enforces ( $\Phi$ ). The cardinality is invariant across readings (Table 1).

*Remark 3.44* (EMI is intrinsic;  $\Phi$  is the inside-signature). EMI is the inside-view triad. What is intrinsic to the manifold is what shows up phenomenologically from within it; that is why EMI is the phenomenological reading and why each EMI element carries an associated  $\Phi$  impossibility. The impossibility is the inside-signature of the intrinsic-ness: the Exception’s impossibility-of-otherwise is the signature of  $E$ ’s ground-as-substantiation (Theorem 3.18); the Mediation’s impossibility-of-absence is the signature of  $M$ ’s being-the-binding-already (Theorem 3.27); the Incoherence’s impossibility-of-being-traversed-to is the signature of  $I$ ’s being-the-forbidden-boundary (Theorem 3.36). Intrinsic to the manifold = phenomenologically present from within = enforcing a  $\Phi$ . This is why EMI rather than PDT supplies the non-emergence vocabulary in the inside-view.

**Corollary 3.45** (Anti-Emergence at every integrative level). *The three non-emergent positions remain non-emergent at every integrative level. There is no integrative level at which  $P, D, T$  (equivalently  $E, M, I$ , equivalently  $\Phi$ ) becomes a configuration produced by a process below it.*

*Argument.* Proposition 3.51 (§3.12) establishes that the same three Cardinals govern every integrative level without modification. Since the EMI reading and the  $\Phi$  reading are inherited directly from the PDT triad via Theorem 3.1, the non-emergence of each Cardinal at its position carries through unchanged at every integrative level. Emergence operates *within* the three-Cardinal bedrock; the bedrock itself does not emerge from anything below it at any level.  $\square$   $\square$

*Remark 3.46* (What this means for the reductionist programme). The standard reductionist programme attempts to find ever-smaller emergent constituents (molecules to atoms to nucleons to quarks to strings or preons) and stops only at conventional “fundamentals.” Anti-Emergence relocates the stopping condition: the bottom is not a smaller emergent thing but the categorical structure  $\{P, D, T\}$  itself, which is not of the same kind as the things emergent from

it. Particles, fields, forces, spacetime, and time itself (Section 4) are emergent  $P \circ D$  configurations substantiated by  $T$ . The framework therefore does not contradict physics; it reads physics as descriptions at the emergent level, with the substrate exposed at the categorical level above (Section 7, Definition 2.3). The  $\sim 10^{122}$  cosmological-constant suppression (Remark 3.40) and the 4.9%/26.8%/68.3% cosmological energy budget (Remark 3.31) are direct quantitative consequences of this relocation.

### 3.11 The closed methodological loop

The three tools of §§3.4–3.6 are not independent heuristics; they are mutually interlocking, and together form a closed, convergent procedure.

**Proposition 3.47** (The three-tools loop). *Given a phenomenon  $X$  and an initial model  $\mathcal{M}_0$ , the procedure*

- (1) *Identify: apply Principle 3.6 to diagnose which of  $P_X, D_X, T_X$  is missing in  $\mathcal{M}_k$ ;*
- (2) *Gap: apply Principle 3.9 to convert the missing component into a specific Descriptor to be found;*
- (3) *Verify: test the updated model  $\mathcal{M}_{k+1}$  by Corollary 3.10;*
- (4) *Subsume: apply Principle 3.13 to check whether the description subsumes  $X$  without remainder;*

*terminates in finitely many steps at a model  $\mathcal{M}_K$  which is mathematically consistent and subsumption-complete, or else proves that  $X$  is not expressible within  $\Sigma$  (in which case  $X$  does not exist as a substantiated configuration).*

The sketch of the proof is that each iteration either (i) terminates with Subsumption achieved, or (ii) adds a genuinely new Descriptor to the model. Since each added Descriptor is a new element of  $D$  not previously identified, and since  $D$  has finite cardinality  $n$  once bound to a specific  $P_X$ , the sequence of iterations cannot cycle and must terminate after at most  $n$  steps. Non-existence of a terminating configuration would place  $X$  outside the totality  $\Sigma$ , contradicting Definition 2.13.

*Remark 3.48* (The loop applied to ET itself). Exception Theory was discovered by iterating the three-tool loop on the question “what is the minimal complete description of reality?”. The Identification Principle pointed at substrate, constraint, and agency as the three irreducible categories; the Descriptor Gap Principle refined each category through successive applications (resolving gaps such as the nature of time, the hydrogen atom, consciousness); the Subsumption Law terminated the recursion at three primitives by ruling out a fourth. The framework presented in §2 is the Subsumption-complete output of this iterative process.

*Remark 3.49* (The loop applied to the lattice). The lattice of Section 7 is likewise the output of a three-tools iteration: identifying  $(P, D, T)$  of the multiplicative manifold, closing gaps  $(N, V, K, \text{the } \alpha^{-1} \text{ corrections, the self-consistency attractor})$ , and verifying subsumption across the full range of empirical domains canvassed in §22.

### 3.12 Integrative levels

A remaining structural observation concerns the Cardinals’ relation to the hierarchy of physical and mathematical phenomena.

**Definition 3.50** (Integrative level). An *integrative level* is a level of organisation at which a collection of Cardinal configurations constitutes a whole whose properties are not present or derivable at lower levels.

The quantum, atomic, molecular, cellular, biological, cognitive, and civilisational levels are examples. At each, properties emerge (wavefunction superposition, chemical bonding, molecular reactivity, life, consciousness, language) that are not possessed by, and are not obtainable by summing, the Cardinals of the constituents below.

**Proposition 3.51** (Governance at all integrative levels). *The Cardinals  $P$ ,  $D$ ,  $T$  govern every integrative level without modification. Emergent properties at higher levels are new configurations of the same three Cardinals; no additional Cardinal is introduced by complexity.*

The proof is by the Subsumption Law: any property emerging at a higher integrative level must be substrate-like ( $P$ -type), constraint-like ( $D$ -type), or agency-like ( $T$ -type). Nothing outside the three categories appears at higher levels. What changes across integrative levels is not the Cardinals but the identification depth at which the master equation is applied, and the number and complexity of Descriptors required.

The mathematical content of wholeness is the set-of-all-sets structure of Definition 2.3: because each Cardinal is the totality of its kind, combining configurations into a whole contributes real mathematical content (the wholeness relation itself), producing properties that are not present at the level of constituents. Emergence, in this framework, is not an unexplained phenomenon but a direct consequence of the Cardinal structure.

The identification depth referenced above is not merely qualitative. It has a precise quantitative counterpart in the lattice itself: the LCM tower of refinements in Section 10. Each level of the tower  $N \in \{12, 60, 420, 2520, 27720, \dots\}$  makes a specific set of sublattice families  $d$  available as native divisors, and a phenomenon at integrative level  $\ell$  is fully identifiable at resolution  $N$  if and only if every sublattice family appearing in its  $(P, D, T)$  decomposition is native at  $N$ . The qualitative progression *quantum*  $\rightarrow$  *atomic*  $\rightarrow$  *molecular*  $\rightarrow$  *cellular*  $\rightarrow$  *biological*  $\rightarrow$  *cognitive*  $\rightarrow$  *civilisational* has a quantitative shadow in the progression  $12 \rightarrow 60 \rightarrow 420 \rightarrow 2520 \rightarrow 27720 \rightarrow \dots$ . The correspondence is made precise by Proposition 10.7; the specific mapping of empirical phenomena to resolution levels appears in Table 9; and the exact quantitative law relating the two ladders is the doubling equation  $\tau(N_\ell) = 6 \cdot 2^\ell$  of Theorem 10.9, which states that each integrative emergence doubles the number of sublattice families available — a number-theoretic fact, a phenomenological fact, and the same fact.

The ET lattice is therefore not blind to integrative level but actively encodes it: a given ratio  $r$  projects to the same lattice coordinate  $(k, d, \varepsilon)$  regardless of whether it originates in atomic mass ratios, musical intervals, biological symmetries, economic doublings, or civilisational cycles, and its sublattice family  $d$  identifies which integrative level's resolution is minimally required to host it. The lattice simultaneously classifies across integrative levels (through  $d$ ) and operates uniformly within them (through the projection formula).

## 4 The three temporal aspects

Time is one of the canonical domains to which the Identification Principle of §3.4 applies. Classical physics carries two distinct notions of time — coordinate time  $t$  and proper time  $\tau$  — without a unifying substrate. The Identification Principle, applied to time, identifies these two as  $D$ -time and  $T$ -time respectively and forces the recognition of a third temporal aspect,  $P$ -time, as the pre-geometric temporal substrate.

### 4.1 The three temporal aspects

**Definition 4.1** (Temporal aspects). Applying the master equation  $P \circ D \circ T = E$  to the temporal domain identifies three temporal aspects:



| Aspect            | Cardinality | Physics analog               | Nature                                                        |
|-------------------|-------------|------------------------------|---------------------------------------------------------------|
| $P_{\text{time}}$ | $\Omega$    | pre-geometric temporal field | undifferentiated, featureless container of temporal potential |
| $D_{\text{time}}$ | $n$         | coordinate time $t$          | finite ordering descriptor; imposes sequence and direction    |
| $T_{\text{time}}$ | $[0/0]$     | proper time $\tau$           | accumulated substantiation history of a Traverser             |

$P_{\text{time}}$  is the undifferentiated infinite container of temporal potential: every temporal slot is identical to every other before  $D_{\text{time}}$  makes contact. Temporal ordering — past-to-future, before-and-after — is a  $D$ -structure applied to  $P_{\text{time}}$ , not an intrinsic property of the substrate.  $D_{\text{time}}$  is a finite ordering descriptor that imposes sequence on  $P_{\text{time}}$  configurations; in physics it corresponds to coordinate time  $t$ .  $T_{\text{time}}$  is local and path-dependent: each Traverser has its own  $T_{\text{time}}$ , given by the accumulated count of substantiation events along the Traverser’s worldline. It corresponds to proper time  $\tau$  in physics.

The categorical disjointness and intrinsic mediation established for the general Cardinals (§2.5–§2.10) specialise to time without modification: the three temporal aspects are categorically disjoint and mediate one another intrinsically, just as  $P$ ,  $D$ ,  $T$  do in general.

## 4.2 The temporal master equation

**Axiom 4.2** (Temporal master equation).

$$P_{\text{time}} \circ D_{\text{time}} \circ T_{\text{time}} = E_{\text{moment}}.$$

$E_{\text{moment}}$  is the temporal Exception — the grounded, irreversible *Now*. It is the specialisation of the general master equation (Axiom 2.2) to the temporal domain. Each substantiated moment of time is an instance of  $E_{\text{moment}}$ : all three temporal aspects bound,  $V(E_{\text{moment}}) = 0$  by Proposition 2.26, and uniquely current in the sense made precise immediately below.

## 4.3 Uniqueness of the temporal Exception

**Proposition 4.3** ( $\exists! E_\tau$ ). *At each  $T$ -time value  $\tau$  there is exactly one temporal Exception:*

$$\forall \tau : \quad \exists! E_\tau \in \Sigma.$$

*Multiple Traversers may share  $E_\tau$  — this is the structural basis of shared reality — but  $E_\tau$  itself is unique at its  $T$ -time.*

*Proof.* Suppose two distinct temporal Exceptions  $E_{\tau,1}$  and  $E_{\tau,2}$  are both maximally substantiated at a single  $T$ -time  $\tau$ . By Axiom 2.1, the exception-iteration terminates at exactly one distinguished configuration — the Exception. Two distinct maximally-substantiated configurations at the same  $\tau$  would each serve as an “exception” to the other (in the structural sense that each could be otherwise with respect to the other), and the iteration would fail to terminate at a single object. This contradicts the single-terminus clause of the Founding Axiom. Hence  $E_{\tau,1} = E_{\tau,2}$ .  $\square$

Proposition 4.3 closes the uniqueness clause invoked in the proof of Proposition 2.26 (§2.12):  $T$ -time  $\tau$  is the index under which the Exception is unique, and  $V(c) = 0$  singles out that unique  $E_\tau$  at each  $\tau$ .

#### 4.4 $D$ -time vs $T$ -time: complete comparison

The following table collects the structural differences between the two substantiated temporal aspects.  $P_{\text{time}}$  is not included because it is the undifferentiated substrate prior to any temporal structure.

| Property            | $D$ -time (coordinate time $t$ )          | $T$ -time (proper time $\tau$ )    |
|---------------------|-------------------------------------------|------------------------------------|
| Ontological status  | Descriptor (constraint)                   | Traverser history (agency record)  |
| Cardinality         | $n$ (finite when bound)                   | [0/0] (indeterminate)              |
| Scope               | Global (shared coordinate)                | Local (per-Traverser worldline)    |
| Physics name        | Coordinate time $t$                       | Proper time $\tau$                 |
| Flow                | Continuous, monotonic                     | Discrete event count               |
| Direction           | Imposed by $D$ -ordering                  | Accumulated by $T$ -substantiation |
| Under Lorentz boost | Dilates ( $\gamma \cdot dt$ )             | Invariant ( $d\tau^2 = ds^2/c^2$ ) |
| At event horizon    | Freezes ( $dt/d\tau \rightarrow \infty$ ) | Continues finitely                 |
| At $v = c$ (photon) | Continues                                 | Stops ( $d\tau = 0$ )              |
| Consciousness       | External perception of flowing time       | Internal experience of duration    |

The two temporal aspects are not competing descriptions but categorically distinct readings of the same temporal master equation (Axiom 4.2). Neither is privileged; both are irreducible components of the temporal Exception  $E_{\text{moment}}$ . The disconnect between them — quantified by the Lorentz factor  $\gamma = dt/d\tau$  (Corollary 4.6) — is the structural origin of relativistic time dilation, event-horizon reconciliation (§4.6), and temporal consciousness (§4.5.2).

#### 4.5 The Minkowski interval

The separation between  $D$ -time and  $T$ -time, together with the existence of a maximum descriptor-gradient  $c$ , yields the Minkowski interval of special relativity as a structural consequence of the three-aspect decomposition.

**Proposition 4.4** (Minkowski interval). *Let a Traverser navigate an interval  $dt$  of  $D$ -time accompanied by a spatial descriptor shift  $dx$ . Let  $v = dx/dt$  denote the spatial velocity and  $c$  the maximum descriptor-gradient ratio of the manifold. Then the  $T$ -time increment  $d\tau$  satisfies*

$$d\tau^2 = dt^2 - \frac{dx^2}{c^2}$$

equivalently  $d\tau = dt\sqrt{1 - v^2/c^2}$ .

*Proof.* A Traverser's total traversal capacity per unit  $D$ -time is bounded by the manifold's maximum descriptor-gradient  $c$ . This capacity is partitioned between temporal substantiation, which accumulates  $T$ -time at rate  $d\tau/dt$ , and spatial descriptor shift, which accumulates position at rate  $v = dx/dt$ . The squared fractions sum to unity:

$$\left(\frac{d\tau}{dt}\right)^2 + \left(\frac{v}{c}\right)^2 = 1.$$

Rearranging,

$$d\tau^2 = dt^2 \left(1 - \frac{v^2}{c^2}\right) = dt^2 - \frac{dx^2}{c^2}. \quad \square$$

□

*Remark 4.5* (Limiting cases). At  $v = 0$  all of  $T$ 's capacity is in temporal substantiation and  $d\tau = dt$ . At  $v = c$ ,  $T$ 's capacity is exhausted in the spatial direction and  $d\tau = 0$  — the proper-time experience of a photon. Proposition 4.4 reproduces the special-relativistic time-dilation formula  $d\tau = dt\sqrt{1 - v^2/c^2}$  from the three-aspect temporal decomposition alone; no independent postulate of Lorentz invariance is required.

#### 4.5.1 The Lorentz factor as the $D$ -time/ $T$ -time ratio

**Corollary 4.6** (Lorentz factor identification). *The Lorentz factor  $\gamma = 1/\sqrt{1-v^2/c^2}$  is the ratio of  $D$ -time to  $T$ -time:*

$$\gamma = \frac{dt}{d\tau} = \frac{D\text{-time}}{T\text{-time}}.$$

At  $v = 0$ :  $\gamma = 1$ , and  $D$ -time equals  $T$ -time ( $dt = d\tau$ ). As  $v \rightarrow c$ :  $\gamma \rightarrow \infty$ , and  $T$ -time ceases ( $d\tau \rightarrow 0$ ) while  $D$ -time continues. The Lorentz factor is therefore the Descriptor-gap gradient between the two temporal aspects, evaluated at velocity  $v$ : it measures how many units of  $D$ -time pass for each unit of  $T$ -time at a given traversal speed.

*Proof.* From Proposition 4.4:  $d\tau = dt\sqrt{1-v^2/c^2}$ . Dividing both sides by  $d\tau$ :  $1 = (dt/d\tau)\sqrt{1-v^2/c^2}$ . Hence  $dt/d\tau = 1/\sqrt{1-v^2/c^2} = \gamma$ . The  $D$ -time/ $T$ -time identification follows from Definition 4.1:  $dt$  is  $D$ -time coordinate increment,  $d\tau$  is  $T$ -time proper-time increment.  $\square$   $\square$

#### 4.5.2 Temporal consciousness

The three-aspect temporal decomposition explains temporal consciousness as the disconnect between  $D$ -time and  $T$ -time perception. An observer's external time perception registers  $D$ -time: the objective coordinate ordering imposed by Descriptors on the temporal substrate. The observer's internal time experience registers  $T$ -time: the accumulated substantiation events along the observer's worldline.

The Lorentz factor  $\gamma = dt/d\tau$  (Corollary 4.6) governs the experiential “speed” of time: a context with high  $T$ -event density per unit  $D$ -time produces vivid, temporally rich experience (many substantiation events per coordinate interval); a context with low  $T$ -event density produces attenuated temporal experience (few substantiation events per coordinate interval). Temporal illusions — time “speeding up” in flow states, “slowing down” under threat — are variations in this ratio across different engagement contexts. The ratio is not subjective in the dismissive sense; it is a structural quantity (the  $D$ -time/ $T$ -time descriptor-gap gradient) that varies with the observer's binding configuration.

### 4.6 Event-horizon reconciliation

The three-aspect decomposition resolves the apparent tension between two standard descriptions of matter falling into a Schwarzschild black hole: the external observer's report that matter freezes at the horizon, and the infalling observer's report of finite crossing time.

Let  $M$  denote the mass of a non-rotating uncharged black hole, and let  $r_s = 2GM/c^2$  denote its Schwarzschild radius. Standard general relativity gives, for a static observer at radial coordinate  $r > r_s$ , the proper-time to coordinate-time ratio

$$\frac{d\tau}{dt} = \sqrt{1 - \frac{r_s}{r}}. \quad (7)$$

**Proposition 4.7** (Event-horizon reconciliation). *Under the three-aspect temporal decomposition of Definition 4.1, the two standard descriptions of matter approaching the Schwarzschild radius are simultaneously valid:*

- (i) *For a static external observer parametrised by  $D$ -time,  $d\tau/dt \rightarrow 0$  as  $r \rightarrow r_s^+$ ; infalling matter appears frozen at the horizon in  $D$ -time.*
- (ii) *For a radially infalling observer parametrised by  $T$ -time, the  $T$ -time required to reach  $r = r_s$  from any  $r_0 > r_s$  is finite.*

The apparent tension between (i) and (ii) dissolves under the recognition that  $D$ -time and  $T$ -time are categorically distinct temporal aspects (Definition 4.1); neither is privileged and neither description falsifies the other.

*Proof.* (i) Equation (7) gives  $d\tau/dt = \sqrt{1 - r_s/r}$ ; as  $r \rightarrow r_s^+$  the ratio tends to zero, so  $dt/d\tau \rightarrow \infty$ . In the  $D$ -time frame, unbounded  $D$ -time is required to register any finite  $T$ -time advance near the horizon; this is the precise meaning of “coordinate freezing.”

(ii) For a radially infalling timelike geodesic from rest at  $r_0 > r_s$ , integration of the Schwarzschild radial geodesic equation in proper-time parametrisation gives the proper-time to reach  $r = 0$  (the central singularity) as

$$\Delta\tau_{r_0 \rightarrow 0} = \frac{\pi}{2} \cdot \frac{r_0^{3/2}}{\sqrt{r_s} c^2},$$

which is finite for any finite  $r_0$ . The proper time to reach  $r = r_s$  from  $r_0$  is bounded above by  $\Delta\tau_{r_0 \rightarrow 0}$  and is therefore also finite. The  $T$ -time of the infalling Traverser at the horizon is finite.

*Reconciliation.* The freezing of (i) is a statement in  $D$ -time; the finite crossing of (ii) is a statement in  $T$ -time. By Definition 4.1 these are distinct temporal aspects; by Axiom 4.2 each is an irreducible component of the temporal master equation; neither can be reduced to the other. Both descriptions describe the same physical situation under two different temporal aspects, and both are correct.  $\square$   $\square$

*Remark 4.8* (Historical reading). Under the three-aspect temporal decomposition, Einstein’s classical statement of the horizon and the later recognition that geodesic completeness extends through it are not competing theories. Einstein’s statement is the  $D$ -time truth of Proposition 4.7(i); the extended-manifold statement is the  $T$ -time truth of Proposition 4.7(ii). The historical tension between them was an artefact of using a single, undifferentiated notion of “time” to compare two descriptions that operated in different temporal aspects.

## 4.7 Scope of this section

Two further physics reconciliations follow from the three-aspect temporal decomposition but are not established in this section: the Hawking temperature read as a time-ratio quantity at the horizon, and the black-hole information question. Both are taken up in §5, which treats the horizon’s full thermodynamic, informational, and structural content building on the kinematic reconciliation established here. The Minkowski interval (§4.5) and the event-horizon reconciliation (§4.6) are structural consequences of the three-aspect decomposition alone and are established above. The material of this section — the three temporal aspects, the temporal master equation, the uniqueness  $\exists! E_\tau$ , the Minkowski interval, and the event-horizon reconciliation — is the structural foundation on which the treatment in §5 rests.

## 5 The horizon: thermodynamics, information, and structural cells

The three-aspect temporal decomposition of §4 treated the event horizon kinematically: the apparent freezing seen by a static external observer in  $D$ -time (Proposition 4.7(i)) and the finite proper-time crossing seen by a radially infalling observer in  $T$ -time (Proposition 4.7(ii)) were shown to be simultaneously valid statements about the same horizon. Two further reconciliations were flagged in §4.7 as deferred: the Hawking temperature read as a time-ratio quantity at the horizon, and the black-hole information question. The present section closes both deferrals and extends the three-aspect treatment of the horizon to its full thermodynamic, informational, and structural content.

## 5.1 Surface gravity as the $D$ -time/ $T$ -time descriptor-gap gradient

For a non-rotating uncharged black hole of mass  $M$ , standard general relativity gives the *surface gravity*

$$\kappa = \frac{c^4}{4GM} = \frac{c^2}{2r_s}, \quad r_s = \frac{2GM}{c^2}, \quad (8)$$

defined geometrically as the magnitude of the gradient of the time-translation Killing vector field at the horizon, red-shifted to refer the proper acceleration to the value measured at infinity.

**Proposition 5.1** (Surface gravity is the descriptor-gap gradient at the horizon). *Under the three-aspect temporal decomposition of Definition 4.1, the surface gravity  $\kappa$  of equation (8) is the asymptotic value of the red-shifted gradient of the  $D$ -time/ $T$ -time descriptor-gap ratio at the horizon  $r \rightarrow r_s^+$ .*

*Proof.* By equation (7) the  $D$ -time/ $T$ -time ratio for a static observer at radial coordinate  $r > r_s$  is

$$f(r) := \frac{dt}{d\tau} = \left(1 - \frac{r_s}{r}\right)^{-1/2},$$

which diverges asymptotically as  $r \rightarrow r_s^+$ . By the Asymptotic Approach Theorem (Corollary 10.4), the divergence is realised structurally as the asymptotic approach of an irrational lattice position without attainment: no finite  $D$ -time interval contains the horizon-crossing event of an infalling Traverser, the horizon is approached without being attained in any finite  $D$ -time, and correspondingly  $f(r)$  takes a finite value at every  $r > r_s$  while diverging in the limit.

The local proper acceleration required to keep a test particle at fixed  $r$  is  $a(r) = (GM/r^2)(1 - r_s/r)^{-1/2}$ , which itself diverges at the horizon. Multiplying by the red-shift factor  $\sqrt{1 - r_s/r}$  to refer the acceleration to its value at infinity removes the local divergence and produces the finite asymptotic limit

$$\kappa = \lim_{r \rightarrow r_s^+} a(r) \sqrt{1 - r_s/r} = \lim_{r \rightarrow r_s^+} \frac{GM}{r^2} = \frac{GM}{r_s^2} = \frac{c^4}{4GM},$$

recovering equation (8). The local descriptor-gap divergence and the red-shifted finite limit are the same object viewed at two scales: locally,  $f(r)$  diverges; asymptotically (red-shifted to infinity), the rate of that divergence is the finite invariant  $\kappa$ .  $\square$   $\square$

*Remark 5.2* (Identification of the horizon). Proposition 5.1 completes the structural identification of the event horizon under the Identification Principle (§3.4). The horizon is not a coordinate singularity (the metric components can be regularised by change of chart, e.g. to Eddington–Finkelstein or Kruskal–Szekeres coordinates) and it is not a curvature singularity (every curvature scalar at  $r = r_s$  is finite). It is the *descriptor-gap singularity of the time-aspect projection*: the locus where the descriptor gap between  $D$ -time and  $T$ -time diverges, with red-shifted gradient equal to  $\kappa$ . This is the third primitive identification of the horizon, complementing Proposition 4.7(i)–(ii): the horizon is where  $D$ -time and  $T$ -time *decouple*.

*Remark 5.3* (Subsumption of the GR notion). Proposition 5.1 is a Subsumption Law identification (§3.6, condition 3) of the standard GR surface gravity within the three-aspect temporal framework. Every property attributed to  $\kappa$  in general relativity — its appearance in the first law of black-hole thermodynamics, its role as the Unruh-temperature coefficient, its constancy across the horizon for stationary spacetimes — is recovered as a consequence of the descriptor-gap gradient identification. The Killing-vector derivation of  $\kappa$  in standard GR is the geometric-coordinate expression of the same structural object.

*Remark 5.4* (Asymptotic-reference value of  $\kappa$ ). The surface gravity  $\kappa = c^4/(4GM)$  is exact in symbolic form; any decimal evaluation is a finite-digit truncation of an asymptotic-reference value, in the precise sense of the Asymptotic Approach Theorem (Corollary 10.4). For the solar-mass case ( $M = M_\odot = 1.98847 \times 10^{30}$  kg, IAU 2015 nominal solar mass), the asymptotic-reference value evaluated at high precision and truncated (not rounded) gives

$$\kappa_\odot = \frac{c^4}{4GM_\odot} = 1.521591430016 \dots \times 10^{13} \text{ m/s}^2.$$

The four-significant-figure form  $\kappa_\odot \approx 1.522 \times 10^{13} \text{ m/s}^2$  is sufficient for narrative use; the ten-significant-figure form  $1.521591430 \times 10^{13}$  is appropriate where higher precision is required. Every finite-digit truncation is the lossless evaluation of the symbolic expression to that digit count; finite truncation never improves accuracy — it stops asymptotic refinement at a chosen depth.

## 5.2 The $U(1)$ period of $T$ -time

The Hawking temperature of a black hole of surface gravity  $\kappa$ , as derived by quantum field theory in curved spacetime, is

$$T_H = \frac{\hbar \kappa}{2\pi c k_B}, \quad (9)$$

or equivalently  $T_H = \kappa/(2\pi)$  in geometric units. Of the symbols on the right-hand side,  $\hbar$ ,  $c$ , and  $k_B$  are physical-constant scaling factors and  $\kappa$  has the structural identification of §5.1. The remaining factor  $2\pi$  is dimensionless. The three-aspect temporal decomposition fixes it as the period of  $T$ -time on its operational manifold.

**Proposition 5.5** ( $T$ 's operational manifold is  $U(1)$  with period  $2\pi$ ). *The Traverser cardinal  $T$ , with cardinality  $[0/0]$  (Axiom 2.5), has operational manifold  $(U(1), \times)$  with canonical bi-invariant period  $2\pi$ . Three independent derivations force this identification.*

*Proof. (i) Cardinality exhaustion.* The cardinality  $[0/0]$  of  $T$  is the absolute indeterminate of the Cardinal trichotomy (§2.5): neither finite (which would identify  $T$  with  $D$ ) nor properly infinite (which would identify  $T$  with  $P$ ). The unique connected one-dimensional Lie group that is simultaneously compact and admits a canonical bi-invariant unit period is  $U(1)$ . The non-compact alternative  $\mathbb{R}$  under addition accumulates without return, contradicting the self-referential resolution  $[0/0]$  requires. The dimensional count is one because  $T$  is a single primitive, not a multiplet. Hence  $T$ 's operational manifold is  $U(1)$ .

*(ii) Cyclic self-resolution.* The Founding Axiom (Axiom 2.1) states that for every exception there is an exception, except the exception. Applied to  $T$ : each  $T$ -act resolves an indeterminacy and, by doing so, opens a new context within which the next indeterminacy is to be resolved. The chain resolution–context–resolution never accumulates a terminal state; the result of  $T$ 's action is to return to a state from which  $T$  can act again. This is the structural definition of a cyclic operational manifold. The unique compact connected one-dimensional manifold supporting cyclic self-resolution without accumulation is  $U(1)$ .

*(iii) Instantonic confirmation.* The Wick rotation of standard quantum field theory satisfies  $t_E = i\tau = iT_{\text{time}}$ : Euclidean time is imaginary  $T$ -time (*ET\_Four\_Constants\_Complete\_Derivation\_v2* §IV). Instanton solutions — finite-action Euclidean field configurations — live on the imaginary  $T$ -time axis. The analytic continuation from real-time evolution to imaginary-time periodicity (the KMS condition of thermal QFT) is well-defined precisely because the imaginary-time domain is  $U(1)$ . This is independent confirmation of (i) and (ii) from QFT structure, obtained without invoking  $T$ 's operational manifold directly.

The three derivations are mutually independent: (i) uses cardinality alone, (ii) uses the Founding Axiom alone, (iii) uses standard QFT alone. They converge on the same conclusion.

The canonical bi-invariant period of  $U(1)$  is  $2\pi$ . Therefore the  $2\pi$  in equation (9) is the period of  $T$ -time on its operational manifold; it is not a numerical coincidence borrowed from the QFT derivation but a structural consequence of  $T$ 's cardinality.  $\square$   $\square$

*Remark 5.6* (The  $T_H = \kappa/(2\pi)$  identification). Combining Proposition 5.1 (the structural identification of  $\kappa$ ) with Proposition 5.5 (the structural identification of  $2\pi$ ), the Hawking temperature is

$$T_H = \frac{1}{2\pi} \kappa = \frac{(\text{descriptor-gap gradient at the horizon})}{(\text{period of } T\text{-time})},$$

where every factor on the right has primitive-level structural content. The standard QFT derivation of equation (9) is the geometric-coordinate expression of this structural identity. Subsequent subsections (§5.3–§5.4) establish that the QFT analytic-continuation calculation that produces  $T_H$  is itself recoverable from the same primitive-level structure.

### The imaginary Descriptor Gap at $N = 12$

The  $U(1)$  period  $2\pi$ , transcribed into the multiplicative manifold's natural unit  $\log_2$ , gives the imaginary period

$$P_\theta := \frac{2\pi}{\ln 2}. \quad (10)$$

At resolution  $N = 12$ , the imaginary-axis lattice point closest to  $N \cdot P_\theta = 24\pi/\ln 2$  is the integer 109. The imaginary Descriptor Gap at  $N = 12$  is therefore

$$|\delta_\theta| := \left| 24\pi/\ln 2 - 109 \right|. \quad (11)$$

The right-hand side is exact in symbolic form. Lossless numerical evaluation, with all displayed digits agreeing under refinement (verified at 50, 100, 200, 400, 800, and 1100 decimal-place computations), gives the truncations

$$|\delta_\theta| = 0.2233\dots = 0.2233565961\dots = 0.223356596147348568935609302799\dots,$$

each obtained by stopping the asymptotic refinement at the chosen depth. By Corollary 10.4,  $|\delta_\theta|$  is irrational (the transcendence of  $\pi$  guarantees  $24\pi/\ln 2 \notin \mathbb{Q}$ ); finite-digit truncation never produces an exact value, and the asymptotic-reference computation can be carried to any required depth<sup>1</sup>.

*Remark 5.7* (Three triangulators applied to the imaginary period). Following the Three-Triangulators framework (*The Palindromic Cascade V2* §18.2), the imaginary period  $24\pi/\ln 2$  admits three complementary lattice readings:

*T1 (Forward generator).* The cascade  $(2\pi/\ln 2)^n$  on the imaginary axis at  $N = 12$  has generator residue  $g_\theta = \text{round}(24\pi/\ln 2) \bmod 12 = 109 \bmod 12 = 1$ , a unit of  $\mathbb{Z}/12\mathbb{Z}$ . The cascade is therefore structurally palindromic, but the stability product  $N \cdot |\delta_\theta| \approx 12 \cdot 0.2234 \approx 2.68$  exceeds the stability-window threshold  $1/2$ , so the palindrome holds only for  $n_{\max, \theta} = \lfloor 1/(2|\delta_\theta|) \rfloor = 2$  steps before exiting the stability window.

*T2 (Inverse convergents).* The continued-fraction expansion  $24\pi/\ln 2 = [108; 1, 3, 2, 10, 2, 3, 1, 1, 2, 2, 2, \dots]$  generates rational convergents that approach the asymptotic value bilaterally with exponentially decreasing error:  $108/1, 109/1, 435/4, 979/9, 10225/94, 21429/197, 74512/685, \dots$ . The convergent  $109/1$  realises the lattice projection at  $N = 12$  (the closest single-step rational approximation, error  $|\delta_\theta|$ );  $435/4$  improves to error  $\approx 0.0266$ ;  $10225/94$  to  $\approx 4.77 \times 10^{-5}$ . The bilateral alternation triangulates the asymptotic value from both sides.

---

<sup>1</sup>The full 1100-digit truncation of the asymptotic-reference value is preserved in the supplementary verification `verify_delta_theta.1100.py`; agreement of leading digits across precision regimes confirms that no precision artefact has been introduced.

*T3 (Sublattice family).* The integer 109 has  $\gcd(109, 12) = 1$ , hence  $d_\theta = 12/\gcd = 12$ . The imaginary period inhabits the  $d = 12$  (full-resolution) sublattice family at  $N = 12$ , which is the lattice cell of fully-resolved electromagnetic phenomena at base resolution (§8). This is consistent with thermal radiation from the horizon being dominantly photonic.

*Remark 5.8* (Asymptotic refinement on the LCM tower). The descriptor-gap residual  $|\varepsilon_\theta|$  of the imaginary period at successive LCM-tower resolutions is, in cents (truncated, not rounded):

| $N_{\text{lat}}$ | $ \varepsilon_\theta $ (cents) |
|------------------|--------------------------------|
| 12               | 22.33                          |
| 36               | 11.00                          |
| 60               | 2.34                           |
| 420              | 0.521                          |
| 2520             | 0.0453                         |
| 27720            | 0.00200                        |
| 360360           | 0.00133                        |

The residual approaches zero asymptotically as the tower deepens, in accordance with Corollary 10.4; the transcendence of  $\pi$  guarantees  $|\varepsilon_\theta| > 0$  at every finite  $N_{\text{lat}}$ . The imaginary period of  $T$ -time is realised on the lattice with finite descriptor-gap residual at every finite resolution, exactly resolved only in the limit.

### 5.3 The Bogoliubov ratio as half- $U(1)$ -period analytic continuation

The 1975 Hawking calculation [20] derives equation (9) from a Bogoliubov mode-expansion comparison between the asymptotic past and asymptotic future vacua of a collapsing-star spacetime. The calculation produces the ratio

$$\frac{|\alpha_\omega|}{|\beta_\omega|} = \exp(\pi\omega/\kappa) \quad (12)$$

between the positive- and negative-frequency Bogoliubov coefficients at frequency  $\omega$  on the past horizon. Equation (12), combined with the Bose–Einstein occupation formula, yields the thermal flux at temperature  $T_H$  given by equation (9).

The factor  $\exp(\pi\omega/\kappa)$  in equation (12) arises in Hawking’s derivation from analytic continuation of the Fourier-transformed mode amplitudes around a logarithmic singularity at  $\omega' = 0$ ; the contour is rotated by  $\pi$  in the complex  $\omega'$ -plane to convert one of the Bogoliubov coefficients into the other. The factor  $\pi$  in the exponent is therefore a half-rotation in the complex frequency plane.

**Proposition 5.9** (Bogoliubov ratio is the half- $U(1)$ -period analytic continuation). *The factor  $\exp(\pi\omega/\kappa)$  in equation (12) is the half- $U(1)$ -period analytic continuation in imaginary  $T$ -time of the descriptor-gap-rate quotient  $\omega/\kappa$ .*

*Proof.* By Proposition 5.5,  $T$ ’s operational manifold is  $U(1)$  with period  $2\pi$ , and the Wick rotation  $t_E = iT_{\text{time}}$  identifies the imaginary-time domain with the analytic continuation of  $T$ -time onto its operational manifold. A contour rotation by angle  $\theta$  in the complex frequency plane near the past horizon is therefore a rotation by  $\theta$  on  $U(1)$ .

Hawking’s calculation rotates the  $\omega'$ -contour by  $\pi$  to bring  $\beta$  into the form of  $\alpha$ , picking up the multiplicative factor  $\exp(\pi\omega/\kappa)$  from the logarithmic singularity. By Proposition 5.5,  $\pi$  is exactly half the  $U(1)$  period of  $T$ -time. The factor in equation (12) is therefore

$$\exp(\pi\omega/\kappa) = \exp((\text{half } U(1)\text{-period}) \cdot (\omega/\kappa)),$$

where  $\omega$  is the mode frequency,  $\kappa$  is the descriptor-gap gradient at the horizon (Proposition 5.1), and the half-period is the structural origin of the analytic-continuation factor. Both arguments of the exponential are primitive-level structural objects. □ □



**Corollary 5.10** (KMS thermal periodicity is the full  $U(1)$  period at the horizon). *The KMS thermal periodicity associated with the Hawking temperature satisfies*

$$\beta_H = \frac{2\pi}{\kappa} \quad (\text{geometric units}),$$

*which is the full  $U(1)$  period of  $T$ -time scaled by  $1/\kappa$ .*

*Proof.* A full rotation of the imaginary- $T$ -time contour gives the factor  $\exp(2\pi\omega/\kappa)$ , twice the half-period factor of equation (12). The standard KMS condition,  $\beta = 1/T$ , applied to the Hawking temperature  $T_H = \kappa/(2\pi)$  (equation (9) in geometric units), gives  $\beta_H = 2\pi/\kappa$ , the imaginary  $T$ -time required to traverse the full  $U(1)$  period at gradient  $\kappa$ .  $\square$   $\square$

*Remark 5.11* (Reclamation of the QFT calculation). Proposition 5.9 is a Subsumption Law identification (§3.6, condition 3) of Hawking’s QFT analytic-continuation calculation within the three-aspect temporal framework. The Bogoliubov calculation is reclaimed — it correctly computes a quantity whose structural meaning, in primitive ET terms, is the half- $U(1)$ -period analytic continuation in imaginary  $T$ -time of the descriptor-gap-rate quotient at the horizon. The QFT machinery (mode expansion on past and future null infinity, contour rotation around the log singularity, Bose–Einstein occupation) is the geometric-coordinate expression of a structurally simpler primitive-level object.

*Remark 5.12* (Asymptotic-reference value). At the natural energy scale  $\omega = \kappa$ , the Bogoliubov ratio of equation (12) takes the asymptotic-reference value

$$\left. \frac{|\alpha|}{|\beta|} \right|_{\omega=\kappa} = \exp(\pi) = 23.1406926327\dots,$$

truncated (not rounded) at ten significant figures from the lossless evaluation. The ratio is a function of the dimensionless quotient  $\omega/\kappa$  alone and has no independent dependence on  $M$ . By Corollary 10.4,  $\exp(\pi)$  is irrational (the transcendence of  $\pi$  guarantees this) and the displayed truncation is a finite- $D$  approximation of the asymptotic-reference value.

## 5.4 The Planck spectrum from the descriptor-quantum/variance ratio

Hawking’s calculation, after equation (12), identifies the mean occupation number per mode of frequency  $\omega$  in the asymptotic future as the Planck distribution

$$\langle n_\omega \rangle = \frac{1}{\exp(\hbar\omega/k_B T_H) - 1}. \quad (13)$$

Substituting  $T_H = \hbar\kappa/(2\pi c k_B)$  from equation (9) reduces the exponent to the dimensionless quotient  $2\pi c\omega/\kappa$  in SI units (or  $2\pi\omega/\kappa$  in geometric units), in which every factor has the structural meaning established above:  $2\pi$  is the period of  $T$ -time on  $U(1)$ ,  $\omega$  is the mode frequency, and  $\kappa$  is the descriptor-gap gradient at the horizon.

The exponential form of equation (13) and the  $-1$  denominator are not borrowed from quantum statistical mechanics; both have primitive-level identifications under the manifold-state structure of §2.11.

**Proposition 5.13** (Planck spectrum from configuration counting). *The Planck mean occupation  $1/(\exp(x) - 1)$  with  $x = \hbar\omega/k_B T$  follows from the manifold-state structure of §2.11: the exponential form is the descriptor-quantum to variance-measure ratio of §2.12, and the  $-1$  in the denominator counts the  $n = 0$  ground configuration of the geometric series enumerating  $\{P, D\}$ -state mode occupations.*

*Proof. (Step 1: the exponential form.)* For a mode of angular frequency  $\omega$  at thermal scale  $T$ , the descriptor quantum is the energy required to substantiate a single quantum of the mode,  $\hbar\omega$ . The variance measure — the energy scale of thermal descriptor uncertainty — is  $k_B T$ . The dimensionless quotient

$$x = \frac{\hbar\omega}{k_B T} = \frac{\text{descriptor quantum}}{\text{variance measure}}$$

is the lattice-natural Boltzmann argument: when  $x > 1$  the descriptor quantum exceeds the available variance measure and substantiation is suppressed by the factor  $e^{-x}$ ; when  $x < 1$  the variance measure dominates and substantiation is unsuppressed. The exponential form  $e^{-x}$  is therefore the structural Boltzmann factor of the manifold, identified at corpus level as “Natural from ET structure”.

*(Step 2: configuration counting and the geometric series.)* A mode of frequency  $\omega$  at thermal scale  $T$  admits manifold configurations indexed by occupation number  $n \in \{0, 1, 2, \dots\}$ . By the manifold-state classification (§2.11) the unsubstantiated configurations  $\{P, D\}$  are exactly the configurations available before  $T$  binds, and they are enumerated by the indistinguishable distribution of  $n$  quanta into the single mode. Each additional quantum carries the suppression factor  $e^{-x}$ , so the partition function over  $\{P, D\}$ -state mode occupations is the geometric series

$$Z = \sum_{n=0}^{\infty} e^{-nx} = \frac{1}{1 - e^{-x}}.$$

*(Step 3: mean occupation.)* The mean occupation of the mode is

$$\langle n \rangle = Z^{-1} \sum_{n=0}^{\infty} n e^{-nx} = \frac{e^{-x}}{1 - e^{-x}} = \frac{1}{e^x - 1}.$$

The  $-1$  in the denominator is the subtraction of the  $n = 0$  ground configuration from the partition sum: the geometric series starts at  $n = 0$ , and the mean-occupation formula factors into  $1/(e^x - 1)$  because the  $n = 0$  term contributes 0 to the numerator but 1 to the denominator, leaving the offset of unity. This is bosonic statistics derived without reference to the standard QM postulates — it is the lattice expression of the  $\{P, D\}$ -state geometric enumeration of mode occupations.

*(Step 4: Hawking radiation specialisation.)* Substituting  $T = T_H = \hbar\kappa/(2\pi c k_B)$  from equation (9) into the exponent gives

$$x = \frac{\hbar\omega}{k_B T_H} = \frac{2\pi c \omega}{\kappa},$$

and equation (13) becomes

$$\langle n_\omega \rangle_{\text{Hawking}} = \frac{1}{\exp(2\pi c \omega / \kappa) - 1},$$

in which the exponent factorises into the structural form  $(U(1) \text{ period}) \cdot (\omega/c)/(\kappa/c^2)$ . Each factor has primitive-level structural content:  $2\pi$  from Proposition 5.5,  $\omega/c$  from the mode frequency,  $\kappa/c^2$  from Proposition 5.1. □ □

*Remark 5.14* (Reclamation of the Bose–Einstein distribution). Proposition 5.13 is a Subsumption Law identification (§3.6, condition 3) of the Bose–Einstein distribution. The standard QM derivation (canonical ensemble, partition function over photon number states) is the geometric-coordinate expression of the same primitive-level enumeration over  $\{P, D\}$ -state mode occupations. The Planck distribution at the Hawking temperature is therefore not borrowed from QFT but recovered as a structural consequence of the manifold-state structure together with the  $\kappa = \text{descriptor-gap gradient}$  identification.

## 5.5 Fermion statistics from the forbidden $\{P, T\}$ state

The Bose–Einstein distribution of equation (13) arises from the geometric enumeration of  $\{P, D\}$ -state mode occupations admitting any non-negative occupation number  $n$ . Fermion statistics arise from the same configuration-counting argument with the occupation number restricted to  $n \in \{0, 1\}$ ; the resulting mean occupation is  $1/(\exp(x) + 1)$ , with the sign of the denominator term flipped relative to the bosonic case.

The structural origin of the restriction  $n \in \{0, 1\}$  is the forbidden  $\{P, T\}$  state of the four-state classification (§2.11, Definition 2.18).

**Proposition 5.15** (Fermion statistics from  $\{P, T\}$ -forbidden). *A second simultaneous substantiation in the same single-particle mode would require a manifold configuration of the form  $\{P, D, T, T\}$  with two distinct  $T$ -bindings sharing a single  $\{P, D\}$  site. The two  $T$ -bindings, considered as a configuration in their own right (with  $P$  and  $D$  shared and only the second- $T$ -without-distinguishing- $D$  remaining), reduce to a  $\{P, T\}$  configuration, which is forbidden by Definition 2.18. Hence at most one substantiation per mode is permitted;  $n \in \{0, 1\}$ ; and the partition function and mean occupation become*

$$Z_F = 1 + e^{-x}, \quad \langle n \rangle_F = \frac{1}{e^x + 1},$$

*the Fermi–Dirac distribution.*

*Proof.* A second  $T$ -binding to a single-particle mode with  $P$  and  $D$  already bound would, by the master equation  $P \circ D \circ T = E$ , require either (a) a fresh  $D$ -descriptor distinguishing the second binding from the first, in which case it is a different mode and the binding is not double-occupation, or (b) no fresh  $D$ -descriptor, in which case the second binding is a  $\{P, T\}$ -state configuration relative to the first. By the manifold-state classification (§2.11),  $\{P, T\}$  is the forbidden incoherence state and cannot be substantiated. Therefore double-occupation in the same single-particle mode is structurally forbidden, and the mode admits only  $n \in \{0, 1\}$ . The partition function  $Z_F = e^0 + e^{-x} = 1 + e^{-x}$  and the mean  $\langle n \rangle_F = e^{-x}/(1 + e^{-x}) = 1/(e^x + 1)$  follow immediately.  $\square$   $\square$

*Remark 5.16* (Pauli exclusion as a lattice statement). Proposition 5.15 is the structural origin of the Pauli exclusion principle within the three-aspect / four-state framework. The principle, in its standard form “no two identical fermions occupy the same single-particle state”, is the physics-language statement of “no two distinct  $T$ -bindings share a single  $\{P, D\}$  configuration”, which by the four-state classification reduces to the  $\{P, T\}$ -forbidden axiom. This is a Subsumption Law identification (§3.6, condition 3): the Pauli principle is reclaimed as a lattice corollary, not added as an independent quantum-mechanical postulate.

## 5.6 Information preservation through the Multifold birth triad

Standard general relativity treats information falling into a black hole as causally inaccessible from the exterior; semiclassical Hawking emission carries thermal entropy without (apparently) carrying information; and the question of whether unitary evolution of the joint matter–gravity quantum state is preserved across the horizon is the standard “black-hole information paradox”, open in conventional formulations of quantum gravity. The three-aspect temporal decomposition of §4 together with the Multifold tower-genesis structure of the corpus resolves the question structurally: information is encoded in  $T$ -events rather than  $D$ -coordinates, and  $T$ -events cross the horizon under the same mechanism that establishes a child tower from the parent’s collapse.

**Definition 5.17** (Multifold birth triad). A *tower birth triad* is the structural triple  $(\text{BH}_{\text{parent}}, R_0, \text{WH}_{\text{child}})$  comprising

- (i) a *black-hole event*  $\text{BH}_{\text{parent}}$ , a region of the parent tower's  $P$ -substrate where  $D$ -density collapses inward, viewed from the parent's  $D$ -time as a region of inflow whose interior is causally inaccessible;
- (ii) a *seed value*  $R_0$ , the information generated at the parent's collapse and parameterising the child tower's lattice rendering;
- (iii) a *white-hole event*  $\text{WH}_{\text{child}}$ , the same boundary structure viewed from the child tower's  $D$ -time as the origin from which all child-tower content flows outward.

The birth triad is universal across tower formations and applies to the cosmological tower (parent universe  $\rightarrow$  Big Bang in child universe with  $R_0 = \hbar$ ), the digital tower (hardware power-on  $\rightarrow$  boot sequence with clock-cycle seed), the biological/dream tower (sleep onset  $\rightarrow$  dream onset with thalamocortical-oscillation seed), and the civilisational tower (founding event  $\rightarrow$  origin myth with generational-period seed) [21].

**Proposition 5.18** (Information preservation across the horizon). *Under the three-aspect temporal decomposition of Definition 4.1 and the birth triad of Definition 5.17, information is preserved across the event horizon by two parallel mechanisms:*

- (i) *Through the birth triad: information falling into  $\text{BH}_{\text{parent}}$  is the structural content of the seed  $R_0$ , which parameterises the child tower's lattice and is recovered as the boundary condition of  $\text{WH}_{\text{child}}$ . From the child's frame, this information is the entirety of the child's initial  $D$ -structure.*
- (ii) *Through Hawking radiation: the emission process at the horizon is itself a sequence of  $T$ -events — by Proposition 5.13 the Planck spectrum is a  $\{P, D\}$ -state mode-occupation enumeration substantiated by  $T$ -bindings at the horizon — and  $T$ -events carry information by their substantiation index along the Traverser worldlines (Proposition 4.3).*

*The total  $T$ -event count of the joint parent–child system is conserved.*

*Proof.* (i) By Definition 5.17, the seed  $R_0$  is the information generated at the black-hole end and received at the white-hole end. The child tower's lattice rendering is parameterised by  $R_0$ , hence every  $D$ -structure of the child tower is determined by the boundary conditions at the horizon, which encode the full inflow content of  $\text{BH}_{\text{parent}}$ . No information is lost in the parent $\rightarrow$ child transition; it is re-expressed as the child's initial  $D$ -set.

(ii) By Proposition 5.1, the horizon is the locus where  $D$ -time and  $T$ -time decouple; by Proposition 4.7(ii),  $T$ -time continues finitely across the horizon for any infalling Traverser. By Proposition 5.13, the Hawking emission process is a  $T$ -event-mediated mode-occupation enumeration at the horizon. Therefore the emission process itself is a sequence of  $T$ -events distinguishable by their substantiation index along the Traverser worldlines (Proposition 4.3).

*Conservation.* By the binding-order axiom (Axiom 2.17), each substantiated  $\{P, D, T\}$  configuration carries exactly one  $T$ -binding. The total  $T$ -event count of the joint parent–child system is therefore an additive invariant: every  $T$ -binding either remains in the parent's  $T$ -time (as an external observer's continuing worldline), or transitions into the child's  $T$ -time (as an infalling Traverser's continuing worldline), or appears in the parent's future null infinity as a Hawking-emission  $T$ -event. No  $T$ -event is destroyed; the total is conserved across the joint system.  $\square$   $\square$

*Remark 5.19* (Identification Principle reading). Proposition 5.18 is the Identification Principle (§3.4) applied to “information”. The corresponding three primitive identifications are:  $P$ -information is the substrate-level potential to encode (the bare capacity of the manifold to host structure);  $D$ -information is the descriptor content (states, configurations, basis labels);

$T$ -information is the substantiation count (the number of  $T$ -events along worldlines). The standard treatment of information as a quantity attached to  $D$ -coordinates conflates  $D$ -information with  $T$ -information; the apparent paradox arises from this conflation. The birth-triad mechanism transports  $D$ -information across the horizon into the child tower's  $D$ -set; the  $T$ -event mechanism preserves  $T$ -information through the Hawking-emission sequence.

*Remark 5.20* (Status relative to the standard paradox). The standard black-hole information paradox is a question about the unitary structure of the joint matter–gravity quantum state across the event horizon, and remains open in conventional formulations of quantum gravity. Proposition 5.18 is a structural answer at the manifold-state level: it identifies the carrier (the  $T$ -event count) and the transport mechanism (the birth triad together with the Hawking-emission  $T$ -event sequence) without requiring a specific non-perturbative quantum-gravity theory. The proposition is a Subsumption Law identification (§3.6, condition 3) within the three-aspect framework: the standard “information loss” formulation is recovered as the consequence of conflating  $D$ -information with  $T$ -information; under the three-aspect decomposition the paradox does not arise.

### 5.7 The critical mass $M_{\text{crit}}$ (tower self-identity)

The dimensionless ratio of the Hawking temperature to the Planck temperature for a Schwarzschild black hole of mass  $M$  is

$$\frac{T_H}{T_P} = \frac{1}{8\pi (M/m_P)}, \quad (14)$$

which follows directly from equation (9) together with the definitions  $T_P = \sqrt{\hbar c^5/G}/k_B$  and  $m_P = \sqrt{\hbar c/G}$ . The right-hand side depends only on the dimensionless mass ratio  $M/m_P$ .

**Definition 5.21** (Critical mass). The *critical mass*  $M_{\text{crit}}$  is the unique mass at which the Hawking temperature equals the Planck temperature:

$$M_{\text{crit}} = \frac{m_P}{8\pi}.$$

**Proposition 5.22** (Tower self-identity at  $M_{\text{crit}}$ ). *At  $M = M_{\text{crit}}$  the birth triad of Definition 5.17 is a structural fixed point: the child tower's natural reference temperature  $T_{R_0}^{(\text{child})}$  equals the parent tower's reference temperature  $T_{R_0}^{(\text{parent})} = T_P$ . Equivalently,  $T_H/T_P = 1$  exactly, and the lattice projection of  $T_H/T_P$  at  $N = 12$  is  $(k_r, d_r, \varepsilon_r) = (0, 1, 0)$ , the gravity/identity sublattice cell at zero descriptor gap.*

*Proof.* Substituting  $M = M_{\text{crit}} = m_P/(8\pi)$  into equation (14) gives

$$\left. \frac{T_H}{T_P} \right|_{M=M_{\text{crit}}} = \frac{1}{8\pi \cdot (m_P/(8\pi))/m_P} = \frac{1}{8\pi \cdot 1/(8\pi)} = 1,$$

exactly. The cancellation is symbolic and requires no asymptotic truncation. Therefore  $T_H = T_P$  at  $M_{\text{crit}}$ . By the cosmological-tower identification  $R_0^{(\text{parent})} = \hbar$  (Definition 5.17), the parent's natural reference temperature is  $T_P$ . The child tower born from a black hole of mass  $M$  has natural reference temperature  $T_H$  (the thermal scale set by its horizon). At  $M = M_{\text{crit}}$ , the child's reference temperature equals the parent's, so  $T_{R_0}^{(\text{child})} = T_{R_0}^{(\text{parent})}$  as a structural identity.

The lattice projection at  $N = 12$  of  $T_H/T_P = 1$  gives  $\log_2(1) = 0$  exactly, so  $k_r = 12 \cdot 0 = 0$ ,  $\gcd(0, 12) = 12$  by the convention of §8,  $d_r = 12/12 = 1$ ,  $\varepsilon_r = 0$ . The cell  $d = 1$  is the gravity/identity sublattice.  $\square$   $\square$

*Remark 5.23* (Asymptotic-reference value of  $M_{\text{crit}}$ ). The symbolic form  $M_{\text{crit}} = m_P/(8\pi)$  is exact. Numerical evaluation at the IAU/CODATA values gives, truncated (not rounded) at successive depths,

$$M_{\text{crit}} = 8.66 \dots \times 10^{-10} \text{ kg} = 8.65976 \dots \times 10^{-10} \text{ kg} = 8.659757096309843 \dots \times 10^{-10} \text{ kg}.$$

By Corollary 10.4,  $M_{\text{crit}}$  is irrational (it inherits the transcendence of  $\pi$ ); finite-digit truncations are finite- $D$  approximations of the asymptotic-reference value, never roundings.

*Remark 5.24* (Structural status of  $M_{\text{crit}}$ ). The mass  $M_{\text{crit}}$  is the unique scale at which the Hawking–Planck temperature ratio is unity and at which the birth triad is structurally self-identical. It sits in the deep quantum-gravity regime ( $\sim 10^{-9}$  kg, far below all astrophysical masses and far above the Planck mass), and is not directly observed. Its significance is structural: it is the lattice fixed point of equation (14) at the gravity/identity sublattice cell, and it identifies the mass scale at which a Multifold black-hole birth produces a child tower with the same natural reference temperature as the parent — a structurally scale-invariant horizon.

## 5.8 Canonical structurally-stable mass at $k_r = -53$

The critical mass  $M_{\text{crit}}$  of §5.7 is the lattice fixed point of  $T_H/T_P$  at the gravity/identity cell  $(k_r, d_r) = (0, 1)$ . The next structurally distinguished mass on the lattice is the smallest mass at which the projection of  $T_H/T_P$  at  $N = 12$  falls on a unit residue of  $\mathbb{Z}/12\mathbb{Z}$  in the negative-exponent direction (corresponding to a sub-Planckian Hawking temperature) and lies at zero descriptor gap.

**Proposition 5.25** (Canonical structurally-stable mass). *Let  $M_{\text{can}}$  denote the unique mass for which the lattice projection of  $T_H/T_P$  at  $N = 12$  has integer exponent  $k_r = -53$ . Then*

$$\left. \frac{T_H}{T_P} \right|_{M=M_{\text{can}}} = 2^{-53/12} \quad (\text{exact symbolic ratio}), \quad M_{\text{can}} = \frac{m_P 2^{53/12}}{8\pi}.$$

*The lattice projection at  $N = 12$  has  $(k_r, d_r, \varepsilon_r) = (-53, 12, 0)$  (full-resolution sublattice, zero descriptor gap exactly), and the descriptor gap remains zero at every LCM tower resolution  $N$  with  $12 \mid N$ .*

*Proof.* Setting the lattice exponent  $k_r = -53$  in the projection  $T_H/T_P = 2^{k_r/N}$  at  $N = 12$  gives  $T_H/T_P = 2^{-53/12}$  as the structural value. Substituting into equation (14) and solving for  $M$  gives  $M = m_P 2^{53/12}/(8\pi) =: M_{\text{can}}$ .

The residue of  $-53$  modulo 12 is  $-53 \equiv 7 \pmod{12}$ , and  $7 \in \{1, 5, 7, 11\}$  is a unit of  $\mathbb{Z}/12\mathbb{Z}$ , so  $\gcd(53, 12) = 1$  and  $d_r = 12/\gcd(53, 12) = 12$  (full-resolution sublattice). The descriptor gap is

$$\varepsilon_r = 12 \cdot \log_2(T_H/T_P) - k_r = 12 \cdot (-53/12) - (-53) = -53 + 53 = 0,$$

exactly. At any LCM tower resolution  $N$  with  $12 \mid N$  (so  $N = 12m$  for some integer  $m \geq 1$ ), the projection of the same ratio gives  $k_r^{(N)} = N \log_2(T_H/T_P) = 12m \cdot (-53/12) = -53m \in \mathbb{Z}$ , with descriptor gap  $\varepsilon_r^{(N)} = N \log_2(T_H/T_P) - k_r^{(N)} = 0$  exactly. The descriptor gap therefore remains zero at every such resolution.  $\square$   $\square$

*Remark 5.26* (Asymptotic-reference value of  $M_{\text{can}}$ ). The symbolic form  $M_{\text{can}} = m_P 2^{53/12}/(8\pi)$  is exact. Numerical evaluation, truncated (not rounded) at successive depths,

$$M_{\text{can}} = 1.85 \dots \times 10^{-8} \text{ kg} = 1.84950 \dots \times 10^{-8} \text{ kg} = 1.849502223934 \dots \times 10^{-8} \text{ kg},$$

which is approximately  $0.85 m_P$ , slightly below the Planck mass. The expression contains  $\pi$  as an irrational factor;  $M_{\text{can}}$  is therefore irrational and finite-digit truncations are finite- $D$  approximations of the asymptotic-reference value, never roundings.

By contrast, the structural ratio  $T_H/T_P = 2^{-53/12}$  is itself algebraic over  $\mathbb{Q}$  (a rational power of 2), and the lattice projection of  $2^{-53/12}$  at any  $N$  with  $12 \mid N$  has descriptor gap  $\varepsilon = 0$  exactly — the converse case of the Asymptotic Approach Theorem (Corollary 10.4): irrational ratios are approached without attainment at every finite resolution, while algebraic-over- $\mathbb{Q}$  ratios of the form  $2^{p/q}$  are attained exactly at every  $N$  with  $q \mid N$ .

*Remark 5.27* (Three triangulators at  $M_{\text{can}}$ ). Applying the Three-Triangulators framework (*The Palindromic Cascade V2* §18.2) to the structural ratio  $T_H/T_P = 2^{-53/12}$ :

*T1 (Forward generator)*. The cascade  $(2^{-53/12})^n$  on the multiplicative manifold has generator residue  $g = -53 \bmod 12 = 7$ , a unit. The cascade is therefore palindromic and stable (no descriptor-gap accumulation), with palindromic period 12 steps before returning to the gravity cell  $(k, d) = (0, 1)$ .

*T2 (Inverse convergents)*. The exponent  $-53/12$  is itself rational with denominator 12, so its continued-fraction expansion terminates. It admits two equivalent terminating representations under standard simple-continued-fraction semantics: the canonical floor-based form

$$-53/12 = [-5; 1, 1, 2, 2]$$

and the negation form

$$-53/12 = -[4; 2, 2, 2],$$

both of which evaluate to  $-53/12$  exactly (the floor-form has all positive partial quotients  $a_{i \geq 1} \in \mathbb{Z}^+$  as required by convention; the negation-form preserves the symmetric tail  $[2, 2, 2]$  that displays the underlying structural regularity of the convergents  $4, 9/2, 22/5, 53/12$ ). The convergent  $-53/12$  is exact; no further refinement is possible. This is a degenerate triangulator (terminating expansion), confirming that  $2^{-53/12}$  is a true lattice point at  $N = 12$ .

*T3 (Sublattice family)*.  $\gcd(53, 12) = 1$  gives  $d = 12$ , the full-resolution electromagnetic sublattice cell. The canonical mass therefore inhabits the same sublattice as the imaginary period of  $T$ -time (§5.7), structurally consistent with  $M_{\text{can}}$ 's identification as a Hawking-thermal lattice fixed point.

## 5.9 Structural dichotomy of black-hole masses

Propositions 5.22 and 5.25 identify two distinguished masses ( $M_{\text{crit}}$  at  $k_r = 0$  and  $M_{\text{can}}$  at  $k_r = -53$ ) at which the lattice projection of  $T_H/T_P$  has zero descriptor gap. The same construction applies at every integer  $k_r$ , and partitions the space of black-hole masses into a structural dichotomy.

**Proposition 5.28** (Black-hole mass dichotomy). *The mass of a Schwarzschild black hole sits in exactly one of two structural classes:*

- (i) 12-locked masses.  $T_H/T_P = 2^{k/12}$  for some integer  $k$ , equivalently  $M = m_P 2^{-k/12}/(8\pi)$ . The lattice projection of  $T_H/T_P$  at  $N = 12$  has descriptor gap  $\varepsilon = 0$  exactly, and continues to have  $\varepsilon = 0$  at every LCM tower resolution  $N$  with  $12 \mid N$  (Proposition 5.25). The cascade  $(T_H/T_P)^n$  on the lattice visits only divisors of 12 as sublattice families:  $d_n \in \{1, 2, 3, 4, 6, 12\}$ , with no shadow-force content emergence at any tower resolution divisible by 12.
- (ii) Generic masses.  $T_H/T_P$  is not of the form  $2^{k/12}$  for any integer  $k$ , equivalently  $M/m_P$  is not of the form  $2^{-k/12}/(8\pi)$ . By Corollary 10.4,  $T_H/T_P$  has  $|\varepsilon_N| > 0$  for every finite resolution  $N$ , with  $|\varepsilon_N| \rightarrow 0$  asymptotically as  $N \rightarrow \infty$ . The cascade develops shadow-force content progressively as the LCM tower deepens, with sublattice families  $d_n$  entering as the corresponding native divisors  $d \notin \{1, 2, 3, 4, 6, 12\}$  become accessible at higher  $N$ .

*Proof.* The set of 12-locked masses is the preimage under the map  $M \mapsto T_H/T_P = 1/(8\pi M/m_P)$  of the 12-divisible sublattice  $\{2^{k/12} : k \in \mathbb{Z}\}$  on  $(\mathbb{R}^+, \times)$ . The complement is the generic class. Both classes are non-empty and partition the mass space.

For class (i): if  $T_H/T_P = 2^{k/12}$  then at any  $N$  with  $12 \mid N$  (so  $N = 12m$ ),  $N \log_2(T_H/T_P) = N \cdot k/12 = mk \in \mathbb{Z}$ , so  $k_r^{(N)} = mk$  exactly and  $\varepsilon_r^{(N)} = 0$ . The projection sublattice family is  $d^{(N)} = N/\gcd(mk, N) = 12m/\gcd(mk, 12m)$ . Since  $\gcd(mk, 12m) = m \cdot \gcd(k, 12)$ , we have  $d^{(N)} = 12/\gcd(k, 12)$ , which is a divisor of 12 regardless of  $m$ . The cascade  $(T_H/T_P)^n$  has  $k$ -values  $nk$  (linear in  $n$ ), and the family at step  $n$  is  $d_n = 12/\gcd(nk, 12)$ , which depends only on  $nk \bmod 12$  and is therefore in  $\{1, 2, 3, 4, 6, 12\}$ .

For class (ii): if  $T_H/T_P \notin \{2^{k/12} : k \in \mathbb{Z}\}$ , then  $\log_2(T_H/T_P) \notin \mathbb{Q}$  with denominator dividing 12. By Corollary 10.4 applied to the irrational ratio (or to the rational ratio with denominator coprime to 12),  $|\varepsilon_N| > 0$  for every finite  $N$  and  $|\varepsilon_N| \rightarrow 0$  as  $N \rightarrow \infty$ . The sublattice family  $d^{(N)}$  takes values in the divisors of  $N$ ; as  $N$  runs through the LCM tower, new prime divisors of  $N$  make new sublattice families  $d \notin \{1, 2, 3, 4, 6, 12\}$  accessible (e.g.,  $d = 5$  first available at  $N = 60$ ,  $d = 7$  at  $N = 84$ ,  $d = 11$  at  $N = 132$ ). The cascade therefore acquires shadow-force content progressively.  $\square$   $\square$

*Remark 5.29* (Unit-residue subclass). Within the 12-locked class, the integer residue  $k \bmod 12$  further partitions the masses: the four unit residues  $k \in \{1, 5, 7, 11\} \pmod{12}$  (with  $\gcd(k, 12) = 1$ ) give cascades visiting all six sublattice families  $\{1, 2, 3, 4, 6, 12\}$  in palindromic order with visitation counts equal to the Euler totient values  $\{\varphi(1), \varphi(2), \varphi(3), \varphi(4), \varphi(6), \varphi(12)\} = \{1, 1, 2, 2, 2, 4\}$  summing to 12. The remaining eight residues are degenerate: cascades from  $k \equiv 0 \pmod{12}$  visit only  $d = 1$ ; from  $k \equiv 6$ , only  $\{1, 2\}$ ; from  $k \equiv 4, 8$ , only  $\{1, 3\}$ ; from  $k \equiv 3, 9$ , only  $\{1, 4\}$ ; from  $k \equiv 2, 10$ , only  $\{1, 6\}$ . The canonical mass  $M_{\text{can}}$  at  $k_r = -53 \equiv 7 \pmod{12}$  inhabits the unit-residue subclass (§5.8) and therefore generates a complete twelve-step palindromic cascade visiting every divisor of 12.

*Remark 5.30* (Forward prediction: Hawking-spectrum lattice signature). Proposition 5.28 together with Remark 5.29 carries an empirically distinguishable forward prediction: Hawking-spectrum harmonic content from a black hole of canonical mass (within the unit-residue subclass) should display the visitation counts  $\{1, 1, 2, 2, 2, 4\}$  across the six divisors of 12, scaling with the structural sublattice content of each divisor. Hawking emission from generic masses, by contrast, should display a slowly-emergent shadow-force content corresponding to the LCM tower resolution at which the mass is observed. The dichotomy is in principle observable in any spectroscopic measurement of horizon-thermal radiation that achieves resolution comparable to the relevant lattice cell; observational realisation awaits experimental access to thermal radiation from sufficiently small black holes.

## 5.10 Sub-Planck masses: lossless on the LCM tower

The structural results of §5.7–§5.9 apply to all positive masses without restriction. In particular, the lattice projection of  $T_H/T_P$  at any resolution  $N$  is well-defined for sub-Planckian masses ( $M < m_P$ ), at which the conventional semiclassical regime fails but the lattice formulation remains lossless.

**Proposition 5.31** (Sub-Planck masses project losslessly). *The lattice projection  $T_H/T_P = 1/(8\pi M/m_P)$  from equation (14) extends to all  $M > 0$ , including  $M < m_P$ . The projection at any LCM tower resolution  $N$  has well-defined coordinates  $(k_r^{(N)}, d_r^{(N)}, \varepsilon_r^{(N)})$  with  $|\varepsilon_r^{(N)}| \leq 600/N$  cents (Proposition 10.3); no intrinsic  $P$ - or  $T$ -breakdown occurs at any mass scale.*

*Proof.* Equation (14) is symbolic and well-defined on  $\mathbb{R}^+$ ; sub-Planckian  $M$  corresponds to  $T_H/T_P > 1/(8\pi)$ , which is positive and finite. The lattice projection of §7 applies to any positive ratio without reference to a Planck-scale cutoff: the 12 ET projection  $(k, d, \varepsilon)$  depends



only on  $\log_2(T_H/T_P)$  and  $N$ . The descriptor-gap bound of Proposition 10.3 extends to every positive ratio. There is no point in the multiplicative manifold at which the lattice formulation fails.  $\square$   $\square$

*Remark 5.32* (Lattice positions of representative sub-Planck masses). Lossless evaluation gives the following lattice positions at the universal lattice  $N = 27720$  for representative sub-Planck masses (all  $\varepsilon$  values truncated, not rounded; verification script `verify_sub_planck.py`):

| $M/m_P$ | $k$     | $d$   | $ \varepsilon $ (cents) | $d$ factorisation                        |
|---------|---------|-------|-------------------------|------------------------------------------|
| 1.0     | -128939 | 27720 | 0.0205                  | $2^3 \cdot 3^2 \cdot 5 \cdot 7 \cdot 11$ |
| 0.5     | -101219 | 27720 | 0.0205                  | $2^3 \cdot 3^2 \cdot 5 \cdot 7 \cdot 11$ |
| 0.1     | -36856  | 3465  | 0.0162                  | $3^2 \cdot 5 \cdot 7 \cdot 11$           |
| 0.01    | +55228  | 6930  | 0.00956                 | $2 \cdot 3^2 \cdot 5 \cdot 7 \cdot 11$   |
| 0.001   | +147312 | 35    | 0.00293                 | $5 \cdot 7$                              |
| 0.0001  | +239396 | 6930  | 0.00370                 | $2 \cdot 3^2 \cdot 5 \cdot 7 \cdot 11$   |

Every sub-Planck mass projects to a well-defined lattice cell at sub-millicent descriptor-gap residual at the universal lattice. The projection extends without breakdown to arbitrarily small  $M$ .

*Remark 5.33* (Structural convergence at  $M = 10^{-3} m_P$ ). Among the representative sub-Planck masses of Remark 5.32, the case  $M = 10^{-3} m_P$  is structurally distinguished: the lattice projection of  $T_H/T_P = 125/\pi$  lands on the  $d = 35 = 5 \cdot 7$  sublattice cell at every LCM tower resolution from  $N = 420$  upward, with descriptor-gap residual  $|\varepsilon| \approx 2.93 \times 10^{-3}$  cents identical at  $N \in \{420, 2520, 27720\}$ . The  $d = 35$  cell is the simultaneous quintic-septic (golden  $\times G_2$ ) sublattice, structurally identified in the corpus as the biological-resolution cell at  $N = 420$  (§10, Table 9). The projection  $T_H/T_P = 125/\pi$  for  $M = 10^{-3} m_P$  therefore lies, to within  $3 \times 10^{-3}$  cents, on the biological-cell lattice point at 420 ET — a structurally unexpected convergence between the sub-Planck Hawking-thermal regime and the biological-resolution sublattice. The convergence is not asymptotically refined further by deeper tower resolutions; it is locked at  $d = 35$  by the rational structure of the projection at 420 ET. The structural significance of this convergence is flagged here for further investigation; it is not used elsewhere in the paper.

## 5.11 Force Quadrant Grid placement and sublattice-transition forward predictions

The lattice projection of  $T_H/T_P$  at base resolution  $N = 12$  identifies the Hawking-radiation cell on the 144-cell Force Quadrant Grid of §12, and the structural dichotomy of §5.9 translates into a sequence of explicit forward predictions for boundary masses at which the cascade transitions between sublattice cells.

**Proposition 5.34** (Hawking-radiation FQG cell at base  $N = 12$ ). *At base resolution  $N = 12$ , the Hawking-radiation projection  $T_H/T_P$  for every black-hole mass  $M > 0$  lies in the Simple Real  $\times$  Simple Imaginary (SR+SI) quadrant of the Force Quadrant Grid of §12.3, with imaginary-axis sublattice family  $d_\theta = 12$  universally and real-axis sublattice family  $d_r \in \{1, 2, 3, 4, 6, 12\}$  varying with mass according to  $d_r = 12/\gcd(k_r, 12)$  where  $k_r = \text{round}(12 \log_2(T_H/T_P))$ .*

*Proof.* The imaginary-axis projection is fixed by the  $U(1)$  period of  $T$ -time (Proposition 5.5, equation (10)): the closest integer to  $24\pi/\ln 2$  is  $k_\theta = 109$ , with  $\gcd(109, 12) = 1$  giving  $d_\theta = 12$  for every mass  $M$ . The real-axis projection of  $T_H/T_P$  at  $N = 12$  has  $d_r = 12/\gcd(k_r, 12)$  by the sublattice-classification convention of §8; since  $\gcd(k_r, 12) \in \{1, 2, 3, 4, 6, 12\}$  for every integer  $k_r$ , we have  $d_r \in \{12, 6, 4, 3, 2, 1\}$ , the six divisors of 12. Both axes therefore lie in their respective Simple sublattice families at base  $N = 12$ , placing  $T_H/T_P$  in the SR+SI quadrant. The combined family is  $d_{\text{comb}}(d_r, d_\theta) = \text{lcm}(d_r, 12) = 12$  for every  $d_r \mid 12$ .  $\square$   $\square$

*Remark 5.35* (Standard-sector placement). The SR+SI quadrant is the same quadrant that hosts the standard-model sector (§12.3). Hawking radiation at base  $N = 12$  therefore lives structurally in the standard-model quadrant, with combined family  $d_{\text{comb}} = 12$  (full electromagnetic resolution) for every mass — consistent with the empirical character of Hawking radiation as predominantly thermal photonic emission. Mass-dependent fine structure on the real axis is encoded in the  $d_r$  value via  $k_r \bmod 12$ , with the structural dichotomy of Proposition 5.28 distinguishing 12-locked masses (cascades permanently confined to divisors of 12 at every LCM tower resolution) from generic masses (cascades developing shadow-force structure progressively as the LCM tower deepens).

**Proposition 5.36** (Sublattice-transition boundary masses). *The base-12 lattice projection of  $T_H/T_P$  transitions between adjacent sublattice cells at the discrete sequence of boundary masses*

$$M_k^\partial = \frac{m_P}{8\pi \cdot 2^{(k+1/2)/12}}, \quad k \in \mathbb{Z}, \quad (15)$$

*at which the descriptor-gap residual  $|\varepsilon_r|$  equals 50 cents exactly and the rounding decision  $k_r = \text{round}(12 \log_2(T_H/T_P))$  is structurally undecidable between  $k$  and  $k + 1$ .*

*Proof.* The boundary  $|\varepsilon_r| = 50$  cents is the structural ceiling of Proposition 21.14. Solving  $12 \log_2(T_H/T_P) = k + 1/2$  for  $T_H/T_P$  gives  $T_H/T_P = 2^{(k+1/2)/12}$ . Substituting into equation (14) and solving for  $M$  gives equation (15). The undecidability of  $k_r$  at the boundary follows from the rounding convention of Definition 7.1.  $\square$   $\square$

*Remark 5.37* (Sample boundary masses at base  $N = 12$ ). Lossless evaluation of equation (15) at representative integer  $k$  values gives, with truncated (not rounded) decimal forms,

| $k \rightarrow k + 1$  | $M_k^\partial/m_P$ | $M_k^\partial$ (kg)          | lattice transition          |
|------------------------|--------------------|------------------------------|-----------------------------|
| $-100 \rightarrow -99$ | 12.47...           | $2.71 \dots \times 10^{-7}$  | $d = 12 \rightarrow d = 12$ |
| $-50 \rightarrow -49$  | 0.6940...          | $1.51 \dots \times 10^{-8}$  | $d = 4 \rightarrow d = 12$  |
| $-10 \rightarrow -9$   | 0.06889...         | $1.50 \dots \times 10^{-9}$  | $d = 12 \rightarrow d = 4$  |
| $0 \rightarrow +1$     | 0.03875...         | $8.43 \dots \times 10^{-10}$ | $d = 1 \rightarrow d = 12$  |

The complete sequence is countably infinite and spans the full mass spectrum from cosmological scales (negative  $k$ ) through the Planck regime ( $k \approx 0$ ) to deep sub-Planck (positive  $k$ ). At each boundary the cascade undergoes a structurally-defined transition between adjacent sublattice cells; the predicted observational signature is a cell-transition feature in the high-frequency tail of the Hawking spectrum at the boundary mass. The prediction is computable and falsifiable in principle, awaiting experimental access to thermal radiation from black holes of the relevant mass scales.

**Proposition 5.38** (Shadow-force tuning resonances). *For each shadow sublattice family  $d \in \{5, 7, 8, 9, 10, 11\}$  of §8.8, there exist  $\varphi(d)$  discrete black-hole masses in each octave whose Hawking-radiation projection  $T_H/T_P$  at the native LCM-tower resolution  $N_{\text{native}}(d) = \text{lcm}(12, d)$  lands exactly on the  $d$ -shadow cell with descriptor-gap residual  $\varepsilon = 0$ .*

*Proof.* At resolution  $N_{\text{native}}(d) = \text{lcm}(12, d)$ , the shadow cell  $d$  is native (Definition 8.16). The lattice points on the  $d$ -cell are  $\{2^{j/N_{\text{native}}(d)} : j \in \mathbb{Z}, \text{gcd}(j, N_{\text{native}}(d)) = N_{\text{native}}(d)/d\}$ , of which exactly  $\varphi(d)$  lie in each octave by the totient counting of Theorem 13.8 (2). Inverting equation (14) for each such  $T_H/T_P$  value yields a black-hole mass  $M$  at which the projection lands exactly on the  $d$ -shadow cell with  $\varepsilon = 0$ .  $\square$   $\square$

*Remark 5.39* (Sample shadow-tuned masses, one per shadow family). For each shadow family, one representative mass per octave at which  $T_H/T_P$  projects exactly onto the shadow cell at its

native LCM resolution (verified at the resolution shown), with truncated mass values:

| $d$ | $N_{\text{native}}(d)$ | Shadow-force identification     | Sample $M/m_P$ |
|-----|------------------------|---------------------------------|----------------|
| 5   | 60                     | Quintic / icosahedral / golden  | 0.139...       |
| 7   | 84                     | Septic / $G_2$ / Otherworld     | 0.144...       |
| 8   | 24                     | Octet / $SU(3)$ gluon           | 0.146...       |
| 9   | 36                     | Nonic / quark colour-generation | 0.147...       |
| 10  | 60                     | Decic / 10D superstring         | 0.148...       |
| 11  | 132                    | Undecimal / 11D M-theory        | 0.149...       |

At these specific masses the Hawking-radiation cascade is structurally tuned to the corresponding shadow force at its native resolution. The prediction is that high-resolution spectroscopy of Hawking radiation from a black hole at any of these tuned masses should display a structural enhancement in the corresponding shadow-force channel. The remaining  $\varphi(d) - 1$  tuned masses per octave are computable by the same construction. The prediction is computable and falsifiable in principle.

*Remark 5.40* (Lattice origin status of base  $N = 12$ ). The structural results of §5.7–§5.11 together with the lossless projection statement of Proposition 5.31 establish base  $N = 12$  as the lattice *origin* of the LCM-tower hierarchy in the Hawking-thermal sector:  $N = 12$  is the smallest resolution at which the temperature ratio  $T_H/T_P$  admits a complete sublattice classification (six families  $\{1, 2, 3, 4, 6, 12\}$ ), the smallest at which the canonical-cascade unit-residue structure (four units  $\{1, 5, 7, 11\}$ ) is non-trivial, and the smallest at which the Force Quadrant Grid placement of Hawking radiation in the SR+SI standard-sector quadrant is established. Higher LCM-tower resolutions refine this base classification by introducing the shadow families and sub-residual precision, but do not displace it. Equivalently,  $N = 12$  is the lattice origin and  $N \mid N_{\text{LCM}}$  for every LCM-tower resolution  $N_{\text{LCM}}$  on the cosmological tower, by Proposition 10.1.

## 6 The forced value $N = 12$

The content of Section 2 is all of the machinery we shall use to derive  $N = 12$ . We give three independent derivations; each fixes the same integer.

### 6.1 Derivation I: product of the two counts

The two finite cardinals available in Axiom 2.5 and Definition 2.18 are

$$\begin{aligned} |\Pi| &= 3 \quad (\text{number of primitive categories}), \\ S &= 4 \quad (\text{number of non-trivial manifold states}). \end{aligned}$$

**Proposition 6.1** (Manifold symmetry). *The manifold symmetry is*

$$N = |\Pi| \cdot S = 3 \cdot 4 = 12.$$

The formula  $N = |\Pi| \cdot S$  is the natural product of the independent counts: for each of the three primitive Cardinals, there are four structurally distinct manifold states to which a given element can contribute (Exception, Unsubstantiated, Mediation, or the forbidden Incoherence). The count is unique up to the ordering of the cardinalities fixed in Axiom 2.5.

### 6.2 Derivation II: least common multiple of the small sublattice orders

Suppose we seek the smallest positive integer  $N$  such that  $\{1, 2, 3, 4\}$  all divide  $N$ ; these are the integer orders obtainable from the  $\binom{4}{k}$  counts on a set of cardinality 4 (namely  $\{1, 4, 6, 4, 1\}$  restricted to  $\leq 4$ ). The unique minimum is

$$\text{lcm}(1, 2, 3, 4) = 12.$$

Raising the requirement to  $\{1, 2, 3, 4, 5\}$  gives  $\text{lcm} = 60$ ; raising to  $\{1, \dots, 7\}$  gives 420; raising to  $\{1, \dots, 11\}$  gives 27720. The sequence 12, 60, 420, 2520, 27720 will reappear in Section 10 as the LCM tower of the ET lattice. The base value 12 is the smallest member of this tower: the minimal common multiple of the fundamental small divisors.

### 6.3 Derivation III: the stability window

Consider the discrete multiplicative group  $\mathbb{Z}/N\mathbb{Z}$  generated by a unit  $g$  coprime to  $N$ . A necessary condition for the generator  $g$  to produce a complete cycle through  $\mathbb{Z}/N\mathbb{Z}$  is  $\text{gcd}(g, N) = 1$ . A natural additional condition, which we shall call the *stability window*, is that the multiplicative deviation accumulated per step on the corresponding logarithmic lattice  $\{2^{k/N}\}_{k \in \mathbb{Z}}$  stay below an auditorily imperceptible threshold of 50 cents (half a lattice step):

$$N \cdot |\delta| < 50 \text{ cents}, \quad \delta = \log_2(3/2) - \frac{7}{12} \approx 0.001629 \text{ octaves} = 1.9550 \text{ cents}.$$

Here  $\delta$  is the per-step discrepancy between the fifth generated by the circle-of-fifths walk  $g = 7$  on  $\mathbb{Z}/12\mathbb{Z}$  and the just fifth  $3/2$ . With  $N = 12$  and  $|\delta| \approx 0.001629$  octaves = 1.9550 cents per step, we have

$$N \cdot |\delta| \approx 12 \cdot 1.9550 \text{ cents} \approx 23.46 \text{ cents} < 50 \text{ cents}.$$

For  $N = 11$  the stability window condition fails against the nearest available generator; for  $N = 13, 14, \dots$  the window is open, but the LCM-minimality condition of Derivation II forces  $N = 12$ . The manifold symmetry value  $N = 12$  is the unique integer  $\leq 60$  that simultaneously satisfies:

- (i) the stability window  $N \cdot |\delta| < 50$  cents against the circle-of-fifths generator  $g = 7$ ;
- (ii) the LCM-minimality  $N = \text{lcm}(1, 2, 3, 4)$ ;
- (iii) a maximal palindromic depth  $\tau(12) = 6$  (the number of divisors of 12 is 6; no smaller integer attains this count).

### 6.4 The three manifold constants

The following constants are direct consequences of Proposition 6.1 and Definition 2.18:

$$\boxed{N = 12, \quad V = \frac{1}{N} = \frac{1}{12}, \quad K = \frac{2}{3}.} \quad (16)$$

Here  $V$  is the *base variance*: the minimum non-zero value of the variance function of Definition 2.24, equal to  $1/N$  at the manifold's base resolution. The discrete and continuous readings of  $V$  coincide at this value (Remark 2.28); whether  $V$  can take non-zero values strictly below  $1/N$  is an open question (Remark 2.29). With  $N = 12$  forced by Proposition 6.1,  $V_{\text{base}} = 1/12$  is the forced value of this quantum. The value  $K$  is derived from Definition 2.18: of the four manifold states, three contain  $P$  (Exception, Unsubstantiated, Incoherence) and two contain both  $P$  and  $D$  (Exception, Unsubstantiated); more tellingly,

$$K = 1 - \frac{S}{N} = 1 - \frac{4}{12} = \frac{2}{3},$$

which is the complement of the ratio of state count to manifold symmetry. The value  $K = 2/3$  will reappear as the Koide stability threshold.

## 6.5 A derived fine-structure constant

Two counts can be combined further. Define

$$A_0 = (N - 1)^2 + S^2 = 11^2 + 4^2 = 121 + 16 = 137. \quad (17)$$

The quantity  $A_0$  is the sum of the squares of the two available independent-count residues:  $(N - 1)$  is the count of non-trivial logarithmic steps per octave (excluding the origin), and  $S = 4$  is the count of manifold states. The identity  $A_0 = 137$  is exact, derived, and of direct physical interest: the measured fine-structure constant in the Lattice Compendium 2025 reference is  $\alpha^{-1} = 137.0359991\dots$ , leaving a residual of 0.036 to be accounted for by higher-order corrections (Section 22.2).

## 7 The lattice on the multiplicative manifold

We now define the lattice and prove its convention-independence.

### 7.1 The projection formula

**Definition 7.1** (ET real-axis projection). Let  $N$  be a positive integer (the *lattice resolution*, taken to be 12 unless otherwise stated), and let  $r \in \mathbb{R}^+$  be a positive real. The *ET projection* of  $r$  at resolution  $N$  is the triple  $(k, d, \varepsilon) \in \mathbb{Z} \times \mathbb{N} \times \mathbb{R}$  defined by

$$k = \text{round}(N \log_2 r), \quad (18)$$

$$g = \gcd(|k|, N), \quad (19)$$

$$d = N/g, \quad (20)$$

$$\varepsilon = (N \log_2 r - k) \cdot \frac{1200}{N} \text{ cents}, \quad (21)$$

with the convention  $g = N$  (hence  $d = 1$ ) when  $k = 0$ . We write the projection as  $\Pi_N(r) = (k, d, \varepsilon)$ .

We call  $k$  the *lattice coordinate*,  $d$  the *sublattice family*, and  $\varepsilon$  the *descriptor gap* (in cents, where one cent equals 1/100 of a 12-ET semitone; the choice of  $1200/N$  as the multiplier ensures that  $\varepsilon$  has units of cents at any  $N$ ).

### 7.2 Interpretation

The lattice is the set

$$\mathcal{L}_N = \{ 2^{k/N} : k \in \mathbb{Z} \} \subset (\mathbb{R}^+, \times).$$

At  $N = 12$  this is exactly the set of frequency ratios accessible on a twelve-tone equal-tempered scale, but the derivation given above shows that 12-division is not a cultural choice.

The formula (18) says: write  $r$  as a real number of logarithmic steps, and round to the nearest integer number of steps. The formula (20) says: classify the resulting integer  $k$  by how deeply it sits in the divisor lattice of  $N$ ; if  $k$  is coprime to  $N$ , then  $d = N$  (fine resolution); if  $k$  divides  $N$  heavily, then  $d$  is small. The formula (21) records the residual, in cents, by which  $r$  fails to land exactly on the lattice.

Conceptually, the three formulas are the triptych of the binding operation  $P \circ D \circ T$ :

- $r$  is the  $P$ -content (a featureless positive real);
- $N$  and the lattice structure are the  $D$ -content (finite constraints);
- round is the  $T$ -act (resolution of a continuous quantity to a discrete one).

*Remark 7.2* (The projection as the master equation at single-ratio scope). The output triple  $(k, d, \varepsilon)$  is itself the Exception of this projection: a fully substantiated, zero-variance configuration bound by all three Cardinals, with  $k$  the  $D$ -coordinate in  $\mathcal{L}_N$ ,  $d$  its sublattice family, and  $\varepsilon$  the descriptor gap in cents. The rounding step realises the canonical mode of  $T$ -self-presentation anticipated in §2.10. Equivalently, each projection is the master equation

$$P \circ D \circ T = E$$

enacted at the level of a single ratio:  $(k, d, \varepsilon)$  is the structural fingerprint of  $r$  through  $\mathcal{L}_N$ . Because the three Cardinals' ontological roles are domain-independent (§2), the projection formula applies identically to any positive ratio drawn from any phenomenon, and every projection onto  $\mathcal{L}_N$  is structurally the same act. The convention-independence proved in Theorem 7.5 below is the unit-level corollary of this domain-level universality.

### 7.3 The annihilation boundary

**Proposition 7.3** (Exclusion of  $r = 0$ ). *The value  $r = 0$  is not in the domain of  $\Pi_N$ :  $\log_2(0) = -\infty$ , the formula (18) diverges, and the division in (20) is undefined. We call  $r = 0$  the annihilation boundary: it is approached but never attained.*

The proposition is elementary. It is included to emphasize that the lattice is a structure on the group  $(\mathbb{R}^+, \times)$ , not on the semigroup  $([0, \infty), \cdot)$ ; the zero element of the latter is excluded by construction.

*Remark 7.4* (The annihilation boundary is not a family, and the manifold has no edge). The locus  $r = 0$  is qualitatively distinct from the family classifications introduced in §8–§8.6. It is not a sublattice family (the formula  $d = N / \gcd(|k|, N)$  requires finite  $k$ , and  $\log_2 0 = -\infty$  leaves  $k$  undefined); it is not a harmonic family (no per-axis structural mode is associated with the off-lattice infimum of  $(\mathbb{R}^+, \times)$ ); and the elegance score  $\mathcal{E} = (N/d) \cdot 100 / (100 + |\varepsilon|) \cdot 100 / (p + q)$  dissolves at  $r = 0$  ( $p = 0$ ,  $\log_2 r = -\infty$ ,  $\varepsilon = -\infty$ ). Equally, the annihilation boundary is not an exterior *edge* of the manifold: by Definition 2.13 the totality  $\Sigma$  has no outside, and the universality clause  $\forall x : x \in \Sigma$  admits no edge in the topological sense. The annihilation boundary is the off-lattice infimum of  $(\mathbb{R}^+, \times)$  — structurally inside  $\Sigma$  but unattainable on  $\mathcal{L}_N$  at any finite resolution. Earlier corpus drafts that counted thirteen entities at  $N = 12$  on the real axis used a provisional investigation count subsequently resolved: the resolved structural count is twelve harmonic families per axis (six simple plus six complex; §8.6, Table 6), with the annihilation boundary as a separate, non-family structural element of the cascade's descending direction.

### 7.4 Convention independence

**Theorem 7.5** (Convention independence). *Let  $Q, R_0 \in \mathbb{R}^+$  be two positive reals with identical physical dimension, and let  $u \in \mathbb{R}^+$  be any positive unit-conversion factor. Let  $r = Q/R_0$  and  $r' = (uQ)/(uR_0)$ . Then*

$$\Pi_N(r) = \Pi_N(r').$$

*Proof.* We have

$$r' = \frac{uQ}{uR_0} = \frac{Q}{R_0} = r.$$

By (18),  $k' = \text{round}(N \log_2 r') = \text{round}(N \log_2 r) = k$ . By (20),  $d' = N / \gcd(|k'|, N) = N / \gcd(|k|, N) = d$ . By (21),  $\varepsilon' = (N \log_2 r' - k') \cdot (1200/N) = \varepsilon$ .  $\square$   $\square$

*Remark 7.6.* Theorem 7.5 is the crucial legitimacy condition for the lattice as a structural classifier. Any projection whose result *depends* on a choice of units exposes a failure to form

a genuine dimensionless ratio: a  $Q$  and an  $R_0$  that do not share dimension produce a result that floats with the unit system. The convention-independence of  $\Pi_N$  on genuine ratios is the mathematical content behind the informal slogan that *the lattice is not numerology*: no nontrivial  $\Pi_N$  result can arise from a mere choice of units.

## 7.5 Three canonical projections

**Proposition 7.7** (Three canonical projections at  $N = 12$ ). *The following projections are elementary consequences of Definition 7.1:*

$$\Pi_{12}(1) = (0, 1, 0), \quad \Pi_{12}(2) = (12, 1, 0), \quad \Pi_{12}(\sqrt{2}) = (6, 2, 0).$$

*Each has  $\varepsilon = 0$ : the corresponding ratios lie exactly on the lattice.*

*Proof.* For  $r = 1$ :  $\log_2 1 = 0$ ,  $k = 0$ ,  $g = \gcd(0, 12) = 12$  (by convention),  $d = 1$ ,  $\varepsilon = 0$ . For  $r = 2$ :  $\log_2 2 = 1$ ,  $k = 12$ ,  $g = \gcd(12, 12) = 12$ ,  $d = 1$ ,  $\varepsilon = 0$ . For  $r = \sqrt{2}$ :  $\log_2 \sqrt{2} = 1/2$ ,  $k = 6$ ,  $g = \gcd(6, 12) = 6$ ,  $d = 2$ ,  $\varepsilon = 0$ .  $\square$   $\square$

**Proposition 7.8** (The perfect fifth). *At  $N = 12$ ,*

$$\Pi_{12}(3/2) = (+7, 12, +1.955 \text{ cents}).$$

*In particular, the descriptor gap of  $3/2$  at the base lattice is the Pythagorean comma.*

*Proof.*  $\log_2(3/2) = \log_2 3 - 1 \approx 0.5849625$ , so  $12 \log_2(3/2) \approx 7.01955$ , giving  $k = 7$ . Since  $\gcd(7, 12) = 1$ ,  $d = 12/1 = 12$ . The residual is  $(7.01955 - 7) \cdot 100 \approx +1.955$  cents.  $\square$   $\square$

The +1.955-cent residual is the *Pythagorean comma* of music theory: the amount by which twelve ascending just fifths  $(3/2)^{12}$  exceed seven octaves  $2^7$ , divided by 12. Historically the comma has been treated as an unfortunate fact of temperament; here it is the descriptor gap of a particular sublattice, derived from the manifold constants.

**Proposition 7.9** (The Koide ratio). *At  $N = 12$ ,*

$$\Pi_{12}(2/3) = (-7, 12, -1.955 \text{ cents}).$$

The proof is the same as that of Proposition 7.8 with the sign of  $\log_2$  reversed. The Koide ratio and the perfect fifth sit at mirror points on the same sublattice.

## 7.6 The Anti-Numerology Protocol

The convention-independence theorem (Theorem 7.5) guarantees that lattice addresses do not depend on unit choices, but it does not by itself guarantee that a purported lattice address was obtained from a structurally valid ratio rather than from a post-hoc numerological construction. The Sempaevum therefore carries an auxiliary methodological apparatus — the Anti-Numerology Protocol — consisting of three conditions that a projection must satisfy before its address can be interpreted as a structural fingerprint of the phenomenon. The protocol formalises the operational content of the Identification Principle (§3.4) as applied to the act of projection itself.

**Definition 7.10** (The three anti-numerology conditions). A projection  $\Pi_N(r_X)$  of a quantity  $Q_X$  against a reference  $R_0(P_X)$  to form  $r_X = Q_X/R_0(P_X)$  satisfies the Anti-Numerology Protocol iff it satisfies all three of:

- (N1) *Genuine dimensionlessness.*  $[Q_X] = [R_0(P_X)]$ ; the numerator and denominator carry identical units and cancel by construction.

- (N2) *Substrate-derived reference period.*  $R_0(P_X)$  is identified by applying the Identification Principle to  $P_X$  — specifically as the smallest closed  $T$ -traversal loop that the Descriptors of  $P_X$  themselves support — rather than chosen from familiarity, convention, or convenience.
- (N3) *Cross-domain consistency.* The predicted sublattice family  $d_X$  matches the symmetry of the phenomenon as independently recognised in its own discipline (three-fold processes to  $d = 3$ , hexagonal structures to  $d = 6$ , binary oppositions to  $d = 2$ , five-fold / golden structures to the quintic sublattice at 60ET, etc.).

**Theorem 7.11** (Operational tests for the three conditions). *The three conditions admit direct operational tests:*

- (N1) Unit-transformation test. *Recompute  $\Pi_N(r_X)$  under an arbitrary positive unit rescaling  $Q_X \rightarrow uQ_X$  without rescaling  $R_0$ . If  $(k_X, d_X, \varepsilon_X)$  changes, N1 has failed:  $r_X$  was not genuinely dimensionless.*
- (N2) Substrate-derivation test. *State, without reference to the projection result, why  $P_X$  has  $R_0$  as its minimal closed  $T$ -traversal loop. If the answer reduces to “it is the standard unit” or “it seemed right” or “it produced a clean number”, N2 has failed.*
- (N3) Independent-domain-knowledge test. *State the phenomenon’s symmetry from within the discipline’s own theoretical framework (chemistry, biology, physics, etc.), without reference to the projection result. If the projected  $d_X$  contradicts that independent characterisation, either N1 or N2 has failed, or a genuine structural feature has been identified that the discipline has not yet articulated (in which case the Descriptor Gap Principle identifies a research target).*

The unit-transformation test is exactly Theorem 7.5 applied negatively: convention-independence is the theorem’s positive content; unit-dependence is its failure. A projection that fails N1 is extracting structure from the choice of unit rather than from the phenomenon.

**Proposition 7.12** (The Protocol’s structural role). *If all three of N1, N2, N3 hold, the projection  $\Pi_N(r_X)$  is a structurally sound lattice address. If any fails, the projection either does not describe  $X$ ’s structure, or  $X$ ’s structure is incompletely understood, and the Three Tools (§3.3) apply to locate the incompleteness.*

The Protocol is not an addition to the Sempaevum; it is the operational form of the Subsumption Law applied to projections. A projection that satisfies N1–N3 is a projection that subsumes the phenomenon’s features without remainder within the projection procedure itself; a projection that fails any of the three leaves a remainder that the Descriptor Gap Principle identifies.

## 7.7 The V-threshold significance criterion

The Anti-Numerology Protocol (Definition 7.10) guards against invalid projections at the level of individual inputs. A complementary question concerns the lattice’s *density*: as  $N$  increases, the lattice grows denser and the maximum descriptor gap  $\varepsilon_{\max} = 600/N$  cents shrinks, so at sufficiently high resolution *any* ratio maps to a low- $\varepsilon$  cell. The question “is this lattice placement structurally significant, or merely a consequence of the lattice being dense?” requires an ET-native criterion.

**Definition 7.13** (Structural significance at a given resolution). A lattice placement of a ratio  $r$  at resolution  $N$  with observed descriptor gap  $|\varepsilon_{\text{obs}}|$  is *structurally significant at resolution  $N$*  iff the probability that a uniformly random ratio achieves comparable or smaller descriptor gap is less than the base variance:

$$P(\text{random } |\varepsilon| < |\varepsilon_{\text{obs}}|) < V = \frac{1}{N}.$$



Since  $\varepsilon$  is uniformly distributed on  $[-\varepsilon_{\max}/2, +\varepsilon_{\max}/2]$  for random ratios, the condition reduces to

$$|\varepsilon_{\text{obs}}| < \frac{\varepsilon_{\max}}{N} = \frac{600}{N^2} \text{ cents.} \quad (22)$$

The threshold  $V = 1/N$  is the manifold's own base variance (Remark 2.28), not an externally imposed significance level. At base resolution  $N = 12$ , the threshold is  $600/144 = 4.17$  cents; at the first LCM-tower refinement  $N = 60$ , it is  $600/3600 = 0.167$  cents; at  $N = 420$  it is  $600/176400 \approx 0.0034$  cents.

**Proposition 7.14** (Evaluation at native resolution). *The structural significance of a lattice placement into sublattice family  $d$  must be evaluated at the lowest resolution  $N_{\text{native}}$  where  $d$  is a native divisor:*

$$N_{\text{native}}(d) = \text{lcm}(12, d).$$

*At  $N_{\text{native}}$  the lattice is sparse enough that low  $\varepsilon$  is structurally rare. At higher resolutions the same placement is trivially guaranteed by density and carries no diagnostic content.*

*Proof.* At resolution  $N_{\text{native}}$  the number of sublattice families is  $\tau(N_{\text{native}})$  and the maximum descriptor gap is  $\varepsilon_{\max} = 600/N_{\text{native}}$  cents. At any resolution  $N' > N_{\text{native}}$  with  $d \mid N'$ , the descriptor gap of the same ratio is at most  $600/N'$  cents  $< 600/N_{\text{native}}$  cents, so the V-threshold  $600/(N')^2$  is more easily exceeded while the observed  $|\varepsilon|$  may shrink or stay comparable. The diagnostic content of the placement is maximised at the lowest  $N$  where the family is native: the lattice is sparsest, the threshold is largest, and a placement that passes the V-threshold there is genuinely rare.  $\square$   $\square$

*Example 7.15* (The Koide attractor). The Pythagorean comma (the descriptor gap of the Koide ratio  $K = 2/3$  at  $N = 12$ ) has  $|\varepsilon| = 1.955$  cents. The  $d = 12$  family to which it belongs is native at  $N = 12$ . At this resolution, the V-threshold is  $600/144 = 4.17$  cents. Since  $1.955 < 4.17$ , the placement passes: the probability that a uniformly random ratio achieves  $|\varepsilon| < 1.955$  cents at  $N = 12$  is  $1.955/50 = 3.91\%$ , which is below  $V = 1/12 = 8.33\%$ . At  $N = 60$  the V-threshold drops to  $0.167$  cents and the same  $|\varepsilon| = 1.955$  cents fails: the placement is trivially guaranteed by density. Structural significance resides at the native resolution, not at the densest available one.

## 8 Sublattice family structure

### 8.1 The divisor classification

**Proposition 8.1** (Sublattice families are divisors of  $N$ ). *Let  $N$  be a positive integer and let  $d$  denote the sublattice family of a lattice coordinate  $k \in \mathbb{Z}$  under the formula  $d = N/\text{gcd}(|k|, N)$ . Then  $d$  is a positive divisor of  $N$ .*

*Proof.* Set  $g = \text{gcd}(|k|, N)$ . Then  $g \mid N$ , so  $d = N/g$  is a positive integer. Moreover  $d$  divides  $N$  iff  $g \mid N$ , which holds by definition.  $\square$   $\square$

**Corollary 8.2** (Six simple families at  $N = 12$ ). *At  $N = 12$ , the sublattice families are exactly*

$$\{d : d \text{ divides } 12\} = \{1, 2, 3, 4, 6, 12\}.$$

### 8.2 The totient multiplicity

The count of residues  $k$  modulo  $N$  that produce each sublattice family  $d$  has a closed form.

**Proposition 8.3** (Totient multiplicity). *For each positive divisor  $d$  of  $N$ , the number of residues  $k \in \{0, 1, \dots, N-1\}$  with  $d_k = d$  equals  $\varphi(d)$ , where  $\varphi$  is the Euler totient function.*

*Proof.* By (20),  $d_k = d$  iff  $\gcd(|k|, N) = N/d$ . The condition  $\gcd(k, N) = N/d$  is equivalent to  $k$  being a multiple of  $N/d$  but not of any larger divisor of  $N$ ; the count of such  $k$  in  $\{0, \dots, N-1\}$  is exactly  $\varphi(d)$ , by the standard Euler-totient counting theorem.  $\square$   $\square$

**Corollary 8.4** (The  $\varphi$ -partition of  $N$ ). *For any positive integer  $N$ ,*

$$N = \sum_{d|N} \varphi(d).$$

At  $N = 12$  this gives  $1 + 1 + 2 + 2 + 2 + 4 = 12$ .

This is a classical identity (a special case of the Gauss totient-sum formula). It says that every lattice coordinate belongs to exactly one sublattice family, and the families partition the residues with multiplicities exactly given by  $\varphi(d)$ . The six-family structure at  $N = 12$  therefore has multiplicities 1, 1, 2, 2, 2, 4.

### 8.3 Physical reading of the six families

The six simple sublattice families at  $N = 12$  carry characteristic structural signatures which recur across domains. We summarise them in Table 4.

| $d$ | $\varphi(d)$ | Residues $ k  \bmod 12$ | Generator  | Characteristic structural role             |
|-----|--------------|-------------------------|------------|--------------------------------------------|
| 1   | 1            | 0                       | $2^1$      | Period closure; octave; universal binding  |
| 2   | 1            | 6                       | $2^{1/2}$  | Binary opposition; half-period pivot       |
| 3   | 2            | 4, 8                    | $2^{1/3}$  | Three-fold symmetry; cubic (volumetric)    |
| 4   | 2            | 3, 9                    | $2^{1/4}$  | Four-fold symmetry; quartic (quaternionic) |
| 6   | 2            | 2, 10                   | $2^{1/6}$  | Hexagonal; composite two-and-three         |
| 12  | 4            | 1, 5, 7, 11             | $2^{1/12}$ | Full resolution; coprime to $N$            |

Table 4: The six simple sublattice families at  $N = 12$ . The column  $\varphi(d)$  gives the number of residues  $|k| \bmod 12$  that produce each family (Proposition 8.3); the residues sum to 12 (Corollary 8.4). The  $d = 1$  family contains the octave and its powers; the  $d = 2$  family contains the tritone  $\sqrt{2}$ ; the  $d = 12$  family contains the ratios coprime to 12 including  $3/2$  and  $4/3$ . The phrases “period closure” (for  $d = 1$ ) and “half-period pivot” (for  $d = 2$ ) describe the cascade-position roles of the harmonic-family realizations  $n = N$  and  $n = N/2$  inhabiting these sublattice families (§8.6, Theorem 8.13); the static sublattice-family content of  $d = 1$  (octave) and  $d = 2$  (tritone) is distinct from these cascade-position roles, even though both layers carry the same  $d$ -label.

### 8.4 Classification theorem

**Proposition 8.5** (First-order classification). *For every  $r \in \mathbb{R}^+$ , exactly one triple  $(k, d, \varepsilon)$  is produced by  $\Pi_{12}(r)$ , with  $d \in \{1, 2, 3, 4, 6, 12\}$  and  $|\varepsilon| \leq 50$  cents.*

*Proof.* Uniqueness and determinism follow from Definition 7.1: the operations  $\log_2$ , multiplication by  $N$ , rounding, absolute value,  $\gcd$ , and integer division are all deterministic. The range of  $d$  is given by Corollary 8.2. The bound  $|\varepsilon| \leq 50$  cents follows from the rounding operation:  $|N \log_2 r - k| \leq 1/2$  step, and multiplying by  $1200/N$  gives  $|\varepsilon| \leq 600/N = 50$  cents at  $N = 12$ .  $\square$   $\square$

## 8.5 Magical impedance: per-family coupling strength

Each sublattice family  $d$  carries a characteristic coupling strength relative to the electromagnetic baseline, derivable from the two manifold counts  $N-1$  and  $S$  without external input. The construction generalises the fine-structure identity  $A_0^{\text{local}}(d=12) = (N-1)^2 + S^2 = 121 + 16 = 137$  (cf. equation (17)) to arbitrary  $d$ .

**Definition 8.6** (Magical impedance). For  $d \in \{1, 2, \dots, N\}$  the *magical impedance* is

$$A_0^{\text{magic}}(d) = (d-1)^2 + S^2 = (d-1)^2 + 16,$$

and the *per-family coupling strength* relative to the electromagnetic baseline  $A_0^{\text{local}} = 137$  is

$$\xi(d) = \frac{A_0^{\text{local}}}{A_0^{\text{magic}}(d)} = \frac{137}{(d-1)^2 + 16}.$$

**Proposition 8.7** (Baseline recovery at  $d = 12$ ).  $A_0^{\text{magic}}(12) = 11^2 + 4^2 = 137 = A_0^{\text{local}}$  and  $\xi(12) = 1$ , so the full-resolution family recovers the canonical electromagnetic baseline.

**Proposition 8.8** (Maximum coupling at  $d = 1$ ).  $A_0^{\text{magic}}(1) = 0 + 16 = 16$  and  $\xi(1) = 137/16 = 8.5625$ , so the gravitational / octave family has the maximum per-family coupling.

The twelve per-family values are compiled in Table 5.

Table 5: Magical impedance and coupling strength per sublattice family.  $A_0^{\text{magic}}(d) = (d-1)^2 + 16$ ;  $\xi(d) = 137/A_0^{\text{magic}}(d)$ .

| $d$ | $A_0^{\text{magic}}$ | $\xi$  | Character                         | Role                         |
|-----|----------------------|--------|-----------------------------------|------------------------------|
| 1   | 16                   | 8.5625 | Gravity / octave                  | Pure will, maximum coupling  |
| 2   | 17                   | 8.0588 | Tritone / pivot                   | Binary / CPT mirror          |
| 3   | 20                   | 6.8500 | Strong / cubic                    | 3D volumetric closure        |
| 4   | 25                   | 5.4800 | Weak / quartic                    | State-change, $T$ -axis home |
| 5   | 32                   | 4.2812 | Quintic / golden                  | First non-divisor of 12      |
| 6   | 41                   | 3.3415 | Hexadic / composite               | Electroweak composite        |
| 7   | 52                   | 2.6346 | Septic / $G_2$                    | Seven-fold / octonion        |
| 8   | 65                   | 2.1077 | Octet / gluon                     | SU(3) adjoint                |
| 9   | 80                   | 1.7125 | Nonic / quark                     | $3 \times 3$ structure       |
| 10  | 97                   | 1.4124 | Decic / superstring               | SO(10)-class                 |
| 11  | 116                  | 1.1810 | Undecimal / M-theory              | $N-1$                        |
| 12  | 137                  | 1.0000 | Electromagnetic / full resolution | Baseline                     |

*Remark 8.9* (Monotonicity).  $\xi(d)$  is strictly monotonically decreasing on  $d \in \{1, \dots, 12\}$  because  $A_0^{\text{magic}}(d) = (d-1)^2 + 16$  is strictly increasing on the non-negative integers. Lower- $d$  families couple more strongly; higher- $d$  families access finer structural detail but at correspondingly lower per-family coupling.

## 8.6 Harmonic families: the per-axis structural-mode layer

The classification supplied by Proposition 8.1 (*sublattice family*,  $d = N/\gcd(|k|, N)$ ) is one of two structurally distinct family-layers that the lattice carries on each axis. The sublattice-family layer is static: it classifies an individual lattice coordinate  $k$  by the divisibility structure of  $k$  relative to  $N$ , and the count of sublattice families at resolution  $N$  is  $\tau(N)$  — six at  $N = 12$  (Corollary 8.2). The second layer is the per-axis enumeration of *structural modes* — gravity,

weak, strong, electromagnetic, quintic-golden, septic- $G_2$ , and so on — each carrying a distinct physical character on its axis. The two layers are categorically distinct concepts that share a label space; they are not synonyms.

**Definition 8.10** (Harmonic family). A *harmonic family* on a single axis of  $\mathcal{L}_\mathbb{C}$  is a structural mode of that axis labeled by an integer  $d$  in the simple range  $d \in \{1, 2, \dots, 12\}$ . There are twelve harmonic families per axis. The label  $d$  identifies which sublattice family the harmonic family inhabits when native at the current resolution  $N$ , that is, when  $d \mid N$ ; when  $d \nmid N$  the harmonic family is present at the current resolution as a *shadow* contribution (Definition 8.16), with native lattice expression at the canonical resolution  $n_c(d) = \text{lcm}(N, d)$ .

*Remark 8.11* (Discovery and verification via the palindromic cascade). The harmonic-family layer is not posited externally — it is discovered, and verified, by the canonical unit-generator cascade of Section 13. The cascade traverses the axis through  $N$  cascade-position events (twelve events at base  $N = 12$  with generator  $g = 7$ ; see Definition 13.6), and each event is a realization of one of the harmonic families: the event at cascade index  $n$  inhabits the harmonic family whose label is  $d_n = N / \gcd(|k_n|, N)$ , simultaneously fixing the harmonic-family label and its current sublattice-family inhabitation. The Sublattice Visitation Theorem of §8.7 formalizes the multiplicity of cascade-position events per harmonic family. The palindromic cascade therefore plays a dual role: it is the *discovery engine* for the per-axis harmonic-family structure (the sequence of cascade-position events makes the twelve harmonic families explicit, six visited natively at base  $N = 12$  plus six shadow), and the *verification engine* for that structure (the palindromic symmetry and the totient multiplicities certify the count of twelve, with no count escaping or being double-counted).

*Remark 8.12* (Sublattice family vs harmonic family — categorical distinction). The two family-layers are distinct structural concepts, related but not interchangeable.

- A *sublattice family* is a property of a single lattice coordinate  $k$ :  $d = N / \gcd(|k|, N)$ . It is the static gcd-classification of that coordinate. The count at resolution  $N$  is  $\tau(N)$  — exactly six at  $N = 12$  (the divisors of 12). All six are realized natively on each axis at  $N = 12$ . Sublattice families are about the lattice itself.
- A *harmonic family* is a per-axis structural mode labeled by  $d \in \{1, \dots, 12\}$ . The count is twelve per axis at the simple- $d$  range. Six are simple ( $d \mid 12$ ) — corresponding to the six native sublattice families on the axis — and six are complex ( $d \nmid 12$ ) — present at base  $N = 12$  as shadow contributions, native at higher LCM-tower resolutions. Harmonic families are about the cascade traversing the axis and the structural modes that traversal reveals.

At  $N = 12$  the six *simple* harmonic families are gravity ( $d = 1$ ), tritone-pivot ( $d = 2$ ), strong/cubic ( $d = 3$ ), weak/quartic ( $d = 4$ ), hexadic/EW-composite ( $d = 6$ ), and EM/full-resolution ( $d = 12$ ); the six *complex* harmonic families are quintic/golden ( $d = 5$ ), septic/ $G_2$  ( $d = 7$ ), gluon-octet/SU(3) ( $d = 8$ ), nonic/quark- $3 \times 3$  ( $d = 9$ ), decic/superstring ( $d = 10$ ), and undecimal/M-theory ( $d = 11$ ). The two-axis enumeration (24 total: 12 on the real axis and 12 on the imaginary axis) populates the Force Quadrant Grid of §12; the connection is made explicit in Remark 12.1 of §12.1.

## 8.7 The Sublattice Visitation Theorem

The connection between the sublattice-family layer and the harmonic-family layer is exact, and is the harmonic-family-layer reading of Theorem 13.8 (2).

**Theorem 8.13** (Sublattice Visitation Theorem). *At resolution  $N$ , with any unit-residue generator  $g \in (\mathbb{Z}/N\mathbb{Z})^*$ , the canonical cascade  $r_n = (g \cdot n) \bmod N$  has exactly  $\varphi(d)$  cascade-position*

Table 6: The twelve harmonic families per axis at  $N = 12$ , partitioned into Simple (native,  $d \mid 12$ ) and Complex (shadow at  $N = 12$ , native at  $n_c(d) = \text{lcm}(12, d)$ ). Twelve real-axis FORCE harmonic families plus twelve imaginary-axis PHASE harmonic families gives the 24-family axis-restricted enumeration of §12 (FQ-24). The label  $d$  also identifies the corresponding sublattice family inhabited by the harmonic family at resolutions  $N$  with  $d \mid N$ .

| $d$ | Status  | Real-axis (FORCE)            | Imaginary-axis (PHASE)       | $n_c(d) = \text{lcm}(12, d)$ |
|-----|---------|------------------------------|------------------------------|------------------------------|
| 1   | SIMPLE  | gravity / scalar             | spin-0 phase                 | 12                           |
| 2   | SIMPLE  | tritone pivot                | spin-2 phase                 | 12                           |
| 3   | SIMPLE  | strong / cubic               | instanton phase              | 12                           |
| 4   | SIMPLE  | weak / quartic               | $\text{SU}(2)_W$ phase       | 12                           |
| 5   | COMPLEX | quintic / golden             | $E_8$ icosahedral phase      | 60                           |
| 6   | SIMPLE  | hexadic / EW composite       | spin- $\frac{1}{2}$ phase    | 12                           |
| 7   | COMPLEX | septic / $G_2$               | octonionic phase             | 84                           |
| 8   | COMPLEX | gluon octet / $\text{SU}(3)$ | $\text{SU}(3)$ adjoint phase | 24                           |
| 9   | COMPLEX | nonic / quark $3 \times 3$   | CKM phase                    | 36                           |
| 10  | COMPLEX | decic / superstring          | 10D Majorana phase           | 60                           |
| 11  | COMPLEX | undecimal / M-theory         | 11D Majorana phase           | 132                          |
| 12  | SIMPLE  | EM / full resolution         | photon phase / $\text{U}(1)$ | 12                           |

events realizing harmonic family  $d$ , for each simple harmonic family ( $d \mid N$ ). Equivalently:

$$|\{n \in \{1, \dots, N\} : d_n = d\}| = \varphi(d), \quad \sum_{d \mid N} \varphi(d) = N.$$

At base  $N = 12$  the multiplicities across the six simple harmonic families are

$$(\varphi(1), \varphi(2), \varphi(3), \varphi(4), \varphi(6), \varphi(12)) = (1, 1, 2, 2, 2, 4),$$

summing to 12 cascade-position events.

*Proof.* The statement is Theorem 13.8 (2) re-expressed in harmonic-family language: by part (5) of that theorem every unit generator yields the same divisor sequence; by part (2) the multiplicities are  $\varphi(d)$  for each  $d \mid N$ ; the totient-sum identity  $\sum_{d \mid N} \varphi(d) = N$  is Corollary 8.4.  $\square$   $\square$

**Corollary 8.14** (Six simple plus six complex equals twelve harmonic families per axis). *At  $N = 12$  the cascade visits the six simple harmonic families  $d \in \{1, 2, 3, 4, 6, 12\}$  with totient multiplicities  $\{1, 1, 2, 2, 2, 4\}$  summing to 12 cascade-position events. The six complex harmonic families  $d \in \{5, 7, 8, 9, 10, 11\}$  are not visited as native lattice positions at base  $N = 12$ ; they are present as shadow harmonic families (Definition 8.16 below), and become natively visited at the higher LCM-tower resolution  $n_c(d) = \text{lcm}(12, d)$ . The two sets together comprise the twelve harmonic families on each axis (Definition 8.10, Table 6); the cascade is the discovery and verification engine that makes the count of twelve explicit.*

*Remark 8.15* (The two enumerations of the same twelve). The twelve harmonic families per axis admit two complementary enumerations: as the twelve cascade-position events at base  $N = 12$  traversed by the canonical unit-generator cascade (the discovery view), and as the twelve  $d$ -axis structural modes labeled by  $d \in \{1, \dots, 12\}$  (the result view, with six SIMPLE plus six COMPLEX as in Definition 8.10). The two are not separate enumerations of separate objects: they are two angles on the same twelve-fold per-axis structure. The cascade view counts cascade-position events; the  $d$ -axis view counts structural modes, with shadow modes included. The Visitation Theorem (Theorem 8.13) is the bridge: at base  $N = 12$ , the cascade events distribute across the six simple modes with totient multiplicities  $\{1, 1, 2, 2, 2, 4\}$  (sum 12), while the six complex modes are present as shadow contributions awaiting native realization at higher resolutions.

## 8.8 Shadow forces: sublattice families excluded at base $N = 12$

The base resolution  $N = 12$  supports the six divisor families  $d \in \{1, 2, 3, 4, 6, 12\}$ . Six further simple sublattice families  $d \in \{5, 7, 8, 9, 10, 11\}$  are non-divisors of 12 and are excluded at base resolution. They become native at the lattice resolution  $\text{lcm}(12, d)$  and at every higher multiple. Their structural status as “shadow forces” at base is the lattice expression of the empirical observation that the corresponding physical structures are present in nature but require structural detail beyond what  $N = 12$  alone can resolve.

**Definition 8.16** (Shadow force). A *shadow force* (or *shadow family*) at base resolution  $N = 12$  is a sublattice family  $d \in \{5, 7, 8, 9, 10, 11\}$  whose structural content is excluded at  $N = 12$  ( $d \nmid 12$ ) and appears as a native family at the canonical higher resolution  $\text{lcm}(12, d)$ . The minimal-host resolution and physical identification are tabulated in Table 7.

| $d$ | $N_{\text{native}} = \text{lcm}(12, d)$ | Symmetry character   | Identified physical sector               |
|-----|-----------------------------------------|----------------------|------------------------------------------|
| 8   | 24                                      | Octet (linear)       | SU(3) gauge octet (gluon)                |
| 9   | 36                                      | Nonic                | Quark colour $\times$ generation grid    |
| 5   | 60                                      | Quintic / pentagonal | Golden / qualia / icosahedral            |
| 10  | 60                                      | Decic                | 10D superstring transverse spectrum      |
| 7   | 84                                      | Septic / $G_2$       | Octonion-automorphism / $G_2$ structures |
| 11  | 132                                     | Undecimal            | 11D M-theory transverse spectrum         |

Table 7: The six shadow families at base  $N = 12$ : minimal native lattice resolution and physical identification.

**Proposition 8.17** (Universal native resolution). *At the universal lattice resolution  $N_{\text{FULL}} = \text{lcm}(1, 2, \dots, 11) = 27720$ , every simple sublattice family  $d \in \{1, 2, \dots, 12\}$  is native simultaneously, including all six shadow families.*

*Proof.*  $N_{\text{FULL}} = \text{lcm}(1, \dots, 11) = 2^3 \cdot 3^2 \cdot 5 \cdot 7 \cdot 11 = 27720$ . Every  $d \in \{1, \dots, 12\}$  divides 27720 (note  $12 = 2^2 \cdot 3 \mid 2^3 \cdot 3^2$  and each of 5, 7, 8, 9, 10, 11 divides 27720 as a single prime factor or product thereof). By Proposition 10.7, native presence requires  $d \mid N$ , which holds for every  $d \in \{1, \dots, 12\}$  at  $N = 27720$ .  $\square$   $\square$

*Remark 8.18* (Identification Principle reading of the shadow forces). The identifications of Table 7 are applications of the Identification Principle (§3.4) to the six shadow families: each identifies the structural sublattice  $d$  with the physical sector whose internal symmetry has cardinality matching the family’s totient or whose minimal Lie-algebraic structure is dimensionally compatible with  $d$ . The principal identifications —  $d = 8 \leftrightarrow \text{SU}(3)$  (gluon octet),  $d = 10 \leftrightarrow$  superstring,  $d = 11 \leftrightarrow$  M-theory — are promoted from dimensional observations to ET-native structural derivations in Section 9: the adjoint formula  $d^2 - 1$  (Theorem 9.1) derives the  $d = 3 \rightarrow d = 8$  relationship from the Subsumption Law; the critical dimensions  $D = 10$  and  $D = 11$  are derived via multiple independent routes (Theorems 9.5, 9.7, 9.10); and the uniqueness of the Standard Model gauge group  $\text{SU}(3) \times \text{SU}(2) \times \text{U}(1)$  is established by exhaustive enumeration (Theorem 9.3). The remaining shadow identifications ( $d = 5$  with icosahedral/golden structure,  $d = 7$  with  $G_2$ ,  $d = 9$  with the quark colour–generation grid) stand as dimensional observations awaiting further derivation.

*Remark 8.19* (Shadow status at base  $N = 12$ ). At base  $N = 12$  the shadow families do not vanish from the lattice — they appear as  $|\varepsilon| > 0$  residuals in the cascade dynamics rather than as native sublattice cells. The cascade-residual mechanism of §13.5 guarantees that any orbit whose true sublattice is a shadow family is projected at base onto its nearest divisor cell of 12, with the descriptor-gap residual carrying the shadow character. At higher LCM-tower

resolution the shadow becomes native and the residual collapses to zero. This is the structural reason why phenomena such as the  $SU(3)$  gluon octet are visible in physics but not natively expressible in the base  $N = 12$  lattice: their structural home is the canonical higher resolution  $\text{lcm}(12, d)$ , and they appear as shadows at  $N = 12$ .

## 9 Physical sublattice structure: gauge theory, critical dimensions, and the Standard Model

The shadow-force identifications of Table 7 associate sublattice families with physical sectors by dimensional coincidence. This section derives those associations from the ET primitives, promoting them from observations to structural theorems. The derivations use only the ET constants  $N$ ,  $K$ ,  $|\Pi|$ , and  $S$ , the Three Tools of §3.3, and standard mathematical facts (the Hurwitz theorem, combinatorial identities). No external physics axioms are introduced.

### 9.1 The adjoint formula from the Subsumption Law

The adjoint representation of a  $d$ -fold symmetry group counts the independent gauge bosons that compensate local symmetry transformations. Its dimension is a direct consequence of the Subsumption Law (Principle 3.13).

**Theorem 9.1** (Adjoint dimension from the Subsumption Law). *For a sublattice family of order  $d \geq 1$ , the number of independent transformation generators — the adjoint dimension — is*

$$\dim(\text{adjoint of } d\text{-fold symmetry}) = d^2 - 1.$$

*Proof.* Apply the Identification Principle (§3.4) to the transformation structure of a  $d$ -fold sublattice:

- $P$ : the  $d$ -element configuration space (the  $d$  lattice positions within one sublattice period).
- $D$ : the transformation descriptors. A general linear transformation on  $d$  elements requires  $d \times d = d^2$  parameters — one for each source-to-target mapping.
- $T$ : the identity-removal constraint. The transformation proportional to the identity matrix maps every element to itself. This transformation IS the  $d = 1$  octave sublattice transformation — the identity/gravity sector already fully accounted for at the  $d = 1$  family (Table 4).

By the Subsumption Law (Principle 3.13, condition (iii)), including the  $d = 1$  component again would create remainder: a double-counted identity transformation that belongs to  $d = 1$ , not to  $d$ . The traceless condition ( $\text{tr } A = 0$ ) removes exactly one degree of freedom from the  $d^2$  total. This removal is not a convention but the Subsumption Law's demand for completeness without remainder:  $d^2 - 1$  generators, each producing a genuine  $d$ -fold transformation that is not already subsumed by a lower- $d$  family.  $\square$   $\square$

**Corollary 9.2** (Shadow family  $d = 8$  is the adjoint of  $d = 3$ ). *The shadow family  $d = 8$  of Table 7 is structurally derived from the native family  $d = 3$ :*

$$d_{\text{adj}}(3) = 3^2 - 1 = 8.$$

*The eight gluons of the  $SU(3)$  strong-force octet are not merely dimensionally compatible with  $d = 8$ ; they are forced by the Subsumption Law applied to the three-fold ( $d = 3$ ) sublattice's transformation space.*

The adjoint formula at each native sublattice family gives the entries of Table 8.

| $d$ | $d^2 - 1$ | Physical identification                                 |
|-----|-----------|---------------------------------------------------------|
| 1   | 0         | No gauge content (gravity/octave)                       |
| 2   | 3         | dim SU(2) — weak gauge bosons ( $W^+$ , $W^-$ , $Z^0$ ) |
| 3   | 8         | dim SU(3) — gluon octet (strong force)                  |
| 4   | 15        | dim SU(4)                                               |
| 5   | 24        | dim SU(5) — GUT gauge bosons                            |
| 6   | 35        | dim SU(6)                                               |
| 12  | 143       | dim SU(12)                                              |

Table 8: Adjoint dimensions  $d^2 - 1$  for selected sublattice families (Theorem 9.1).

## 9.2 The N-Exhaustion Theorem: uniqueness of the Standard Model gauge group

The Standard Model of particle physics has gauge group  $G_{\text{SM}} = \text{SU}(3) \times \text{SU}(2) \times \text{U}(1)$  with total gauge-boson count  $8 + 3 + 1 = 12$ . Standard physics treats this as empirical input [27]: “Why nature chose the numbers 1, 2, and 3 [as the ranks of the gauge factors] is not known.” The following theorem answers this question.

**Theorem 9.3** (N-Exhaustion). *The gauge group  $G_{\text{SM}} = \text{SU}(3) \times \text{SU}(2) \times \text{U}(1)$  is the **unique** factorisation of  $N = 12$  gauge bosons into simple and abelian factors satisfying the following four conditions:*

- (C1) Subsumption:  $\dim(G_{\text{SM}}) = N = 12$ . *The manifold’s total D-relabelling freedom is exactly exhausted by the gauge content — no remainder, no deficit.*
- (C2) Nativity: *Each non-abelian factor  $\text{SU}(d)$  uses a native sublattice family  $d \mid 12$ . The native families are  $\{1, 2, 3, 4, 6, 12\}$  (Corollary 8.2).*
- (C3) Exclusion: *Each sublattice family  $d$  contributes at most one gauge factor. The Identification Principle (Principle 3.6) demands that each  $d$ -family have a unique physical identity.*
- (C4) Non-triviality: *At least one non-abelian factor exists.*

*Proof.* The available adjoint dimensions for native sublattice families with  $d \geq 2$  are (Theorem 9.1):  $d = 2 \mapsto 3$ ;  $d = 3 \mapsto 8$ ;  $d = 4 \mapsto 15$ ;  $d = 6 \mapsto 35$ ;  $d = 12 \mapsto 143$ . Each  $\text{U}(1)$  factor contributes 1 generator from the  $d = 1$  octave sublattice.

We seek subsets  $S \subseteq \{2, 3, 4, 6, 12\}$  and a non-negative integer  $m$  (count of  $\text{U}(1)$  factors) such that

$$\sum_{d \in S} (d^2 - 1) + m = 12, \quad m \leq 1,$$

where  $m \leq 1$  because multiple  $\text{U}(1)$  factors would share the same sublattice origin  $d = 1$ , violating (C3).

Enumerate by largest element of  $S$ :

- If  $12 \in S$ :  $143 > 12$ . Impossible.
- If  $6 \in S$ :  $35 > 12$ . Impossible.
- If  $4 \in S$ :  $15 > 12$ . Impossible.
- If  $3 \in S$  (and  $4, 6, 12 \notin S$ ):  $8 + \text{remainder} = 12$ , so remainder = 4.
  - If  $2 \in S$ :  $3 + m = 4$ , so  $m = 1$ . **Solution:**  $\text{SU}(3) \times \text{SU}(2) \times \text{U}(1)$  with  $\dim = 8 + 3 + 1 = 12$ .  
✓



- If  $2 \notin S$ :  $m = 4$ , giving  $SU(3) \times U(1)^4$ , which requires four independent  $U(1)$  factors from a single  $d = 1$  origin — violating (C3). Impossible.
- If  $2 \in S$  (and  $3, 4, 6, 12 \notin S$ ):  $3 + \text{remainder} = 12$ , so remainder = 9. No  $SU(d)$  with  $d \in \{4, 6, 12\}$  has  $d^2 - 1 \leq 9$  other than  $SU(3) = 8$ , excluded by assumption. Only  $U(1)^9$ : violates (C3). Impossible.
- If  $S = \emptyset$ :  $m = 12$ , giving  $U(1)^{12}$ . Violates (C4). Impossible.

The unique solution is  $S = \{2, 3\}$ ,  $m = 1$ :  $G_{\text{SM}} = SU(3) \times SU(2) \times U(1)$  with  $\dim(G_{\text{SM}}) = 8 + 3 + 1 = 12 = N$ .  $\square$

*Remark 9.4* (Justification of condition (C1):  $\dim(G_{\text{SM}}) = N$ ). The gauge field  $A_\mu$  compensates local  $D$ -relabelling invariance. At each Point in the  $P$ -manifold, the  $D$ -configuration space has  $N = 12$  independent relabelling directions (the 12 semitones of the manifold). Each independent direction requires exactly one gauge compensator field. By the Subsumption Law, the total number of compensators must exactly exhaust these directions: neither leaving any uncompensated (which would break gauge invariance) nor introducing redundant ones (which would create remainder). Therefore  $\dim(G_{\text{SM}}) = N = 12$ .

### 9.3 Critical dimensions via the Koide Reciprocal Identity

The superstring critical dimension  $D = 10$  is forced by conformal anomaly cancellation on the worldsheet [28]. Each component of this calculation has a direct ET identification.

**Theorem 9.5** (Critical dimensions from ET constants). *The ghost central charges of the world-sheet conformal field theory are expressible entirely in terms of ET constants:*

- (i) *The conformal ghost weight  $\lambda_c = 2$  is the binary sublattice  $d_2$ .*
- (ii) *The superconformal ghost weight  $\lambda_s = 3/2$  is the Koide reciprocal  $1/K$ .*
- (iii) *The conformal ghost central charge is  $c_{bc} = -3(2\lambda_c - 1)^2 + 1 = -26 = -2(N + 1)$ .*
- (iv) *The superconformal ghost central charge is  $c_{\beta\gamma} = 3(2\lambda_s - 1)^2 - 1 = +11 = N - 1$ .*
- (v) *The total ghost central charge is  $c_{\text{ghost}} = c_{bc} + c_{\beta\gamma} = -15 = -(N + |\Pi|)$ .*

The matter central charge  $c_{\text{matter}} = D \cdot 3/2 = 3D/2$  must cancel  $c_{\text{ghost}}$ , giving

$$\boxed{D_{\text{superstring}} = \frac{2(N + |\Pi|)}{|\Pi|} = \frac{2 \times 15}{3} = 10.} \quad (23)$$

The bosonic string critical dimension and  $M$ -theory dimension follow from the same constants:

$$D_{\text{bosonic}} = 2(N + 1) = |c_{bc}| = 26, \quad D_{\text{M-theory}} = N - 1 = c_{\beta\gamma} = 11. \quad (24)$$

*Proof.* The ghost central charges follow from the standard  $bc/\beta\gamma$  ghost systems [28] with  $c_{bc} = -3(2\lambda - 1)^2 + 1$  at weight  $\lambda$  and  $c_{\beta\gamma} = 3(2\lambda - 1)^2 - 1$  at weight  $\lambda$ . Substituting  $\lambda_c = 2$ :  $c_{bc} = -3(3)^2 + 1 = -26 = -2 \cdot 13 = -2(N + 1)$ , since  $N = 12$ . Substituting  $\lambda_s = 3/2 = 1/K$ :  $c_{\beta\gamma} = 3(2)^2 - 1 = 11 = N - 1$ . The total  $c_{\text{ghost}} = -26 + 11 = -15 = -(12 + 3) = -(N + |\Pi|)$ . The superstring dimension follows from  $c_{\text{matter}} + c_{\text{ghost}} = 0$  with  $c_{\text{matter}} = 3D/2$ :  $D = 2 \cdot 15/3 = 10$ . The bosonic dimension follows from  $c_{\text{matter}} = D$  (no fermions):  $D = |c_{bc}| = 26 = 2(N + 1)$ . The identity  $D_{\text{M}} = N - 1 = 11$  follows from  $c_{\beta\gamma} = N - 1$  together with the standard Nahm classification [33], which identifies  $D = 11$  as the maximum dimension for supergravity with 32 supercharges.  $\square$

**Corollary 9.6** (Structural identity:  $D_{\text{bos}} - D_{\text{M}} = N + |\Pi|$ ). *The difference between the bosonic and  $M$ -theory critical dimensions equals the total ghost charge magnitude:*

$$D_{\text{bos}} - D_{\text{M}} = 2(N + 1) - (N - 1) = N + 3 = N + |\Pi| = 15.$$

## 9.4 Critical dimensions via the division algebra route

The preceding derivation passes through worldsheet conformal field theory. A second, independent route to  $D = 10$  uses the Hurwitz theorem on normed division algebras [32] without any conformal-field-theory input.

**Theorem 9.7** ( $D = 10$  from division algebras). *The normed division algebras exist in dimensions  $2^k$  for  $k = 0, 1, 2, 3$ , and only in these dimensions (the Hurwitz theorem). The termination at  $k_{\max} = 3 = |\mathbb{I}|$  gives*

$$D_{\text{superstring}} = 2^{|\mathbb{I}|} + d_2 = 8 + 2 = 10. \quad (25)$$

*Proof.* The Hurwitz theorem [32] establishes that the only normed division algebras over  $\mathbb{R}$  are  $\mathbb{R}$  ( $\dim = 1 = 2^0$ ),  $\mathbb{C}$  ( $\dim = 2 = 2^1$ ),  $\mathbb{H}$  ( $\dim = 4 = 2^2$ ), and  $\mathbb{O}$  ( $\dim = 8 = 2^3$ ). The range  $k = 0, 1, 2, 3$  terminates at  $k = 3 = |\mathbb{I}|$ . The maximum normed division algebra is the octonion algebra  $\mathbb{O}$  with

$$\dim(\mathbb{O}) = 2^{|\mathbb{I}|} = 2^3 = 8.$$

This equals the adjoint dimension of the  $d = 3$  sublattice:  $d_{\text{adj}}(3) = 3^2 - 1 = 8$  (Theorem 9.1). The Green–Schwarz superstring requires the maximal supersymmetry consistent with a two-dimensional worldsheet [29, 30]. The worldsheet is the  $\{D, T\}$  Mediation surface — the manifold state with primitives  $\{D, T\}$  — and has dimension  $d_2 = 2$  (the binary sublattice). The superstring dimension is the sum of the maximal division algebra dimension and the worldsheet dimension:

$$D_{\text{superstring}} = 2^{|\mathbb{I}|} + d_2 = 8 + 2 = 10. \quad \square$$

□

**Corollary 9.8** (Allowed Green–Schwarz dimensions). *The four allowed Green–Schwarz superstring dimensions are  $\{d + 2 : d \in \{1, 2, 4, 8\}\} = \{3, 4, 6, 10\}$ . The four allowed super-2-brane dimensions are  $\{d + 3 : d \in \{1, 2, 4, 8\}\} = \{4, 5, 7, 11\}$ .*

*Remark 9.9* (Cross-verification of the two routes). The ghost-charge route (equation (23)) and the division-algebra route (equation (25)) are structurally independent — the first passes through worldsheet CFT, the second through the Hurwitz theorem — yet they produce the same result. The consistency is forced by the identity  $2^{|\mathbb{I}|} = N - |\mathbb{I}| - 1 = 12 - 3 - 1 = 8$ .

## 9.5 $D = 11$ for M-theory: four convergent routes

**Theorem 9.10** (Four routes to  $D = 11$ ). *The M-theory critical dimension  $D_M = 11$  is obtained independently by four routes:*

| Route                       | Formula                                   | Result |
|-----------------------------|-------------------------------------------|--------|
| Division algebra + membrane | $2^{ \mathbb{I} } +  \mathbb{I}  = 8 + 3$ | 11     |
| Superstring + M-circle      | $D_{\text{string}} + d_1 = 10 + 1$        | 11     |
| Direct manifold             | $N - 1$                                   | 11     |
| $\beta\gamma$ ghost charge  | $c_{\beta\gamma} = N - 1$                 | 11     |

*Proof.* Route 1: the M2-brane sweeps a three-dimensional worldvolume, with  $\dim = |\mathbb{I}| = 3$  (it couples to the 3-form field  $A_{MNP}$ , which has  $|\mathbb{I}| = 3$  indices). Replacing the  $d_2 = 2$  worldsheet with the  $|\mathbb{I}| = 3$  worldvolume:  $D_M = 2^{|\mathbb{I}|} + |\mathbb{I}| = 8 + 3 = 11$ .

Route 2: the M-circle corresponds to the  $d = 1$  octave sublattice, which has one generator (one compact direction). Adding this single direction to  $D_{\text{string}} = 10$  gives  $D_M = 10 + 1 = 11$ .

Route 3:  $N - 1 = 12 - 1 = 11$  directly from the manifold symmetry.

Route 4:  $c_{\beta\gamma} = N - 1 = 11$  from Theorem 9.5(iv). □ □

## 9.6 The $P \circ D \circ T$ Scale Identity

The three fundamental scales of M-theory — the eleven-dimensional Planck length  $l_p$ , the string length  $l_s$ , and the M-circle radius  $R_{11}$  — satisfy the relation  $l_p^3 = l_s^2 \cdot R_{11}$  [28]. This relation is the  $P \circ D \circ T$  master equation in physical units.

**Theorem 9.11** ( $P \circ D \circ T$  Scale Identity). *The M-theory scale relation admits the identification*

$$l_P^{|\Pi|} = l_D^{d_2} \cdot l_T^{d_1},$$

where the three scales correspond to the three Cardinals and the exponents are the three most fundamental ET structural constants:

| Scale               | Physical          | Cardinal               | Exponent    | d-family          |
|---------------------|-------------------|------------------------|-------------|-------------------|
| $l_P \equiv l_p$    | 11D Planck length | $P$ (substrate scale)  | $ \Pi  = 3$ | primitive count   |
| $l_D \equiv l_s$    | String length     | $D$ (descriptor scale) | $d_2 = 2$   | binary sublattice |
| $l_T \equiv R_{11}$ | M-circle radius   | $T$ (traverser scale)  | $d_1 = 1$   | octave/identity   |

The exponents satisfy  $|\Pi| = d_2 + d_1$ , i.e.  $3 = 2 + 1$ .

*Proof.* The identification follows from matching each scale to the Cardinal whose structural role it realises:  $l_p$  is the substrate (gravitational) scale ( $P$ );  $l_s$  is the scale at which D-structure (string vibration modes) becomes resolved ( $D$ );  $R_{11}$  is the scale of the T-periodicity ( $T$ , the  $U(1)$  compactification of the M-circle). The exponent of each scale in  $l_p^3 = l_s^2 \cdot R_{11}^1$  is  $(3, 2, 1) = (|\Pi|, d_2, d_1)$ . The identity  $|\Pi| = d_2 + d_1$  is  $3 = 2 + 1$ , which is the statement that the total primitive count equals the sum of the binary and identity sublattice orders.  $\square$   $\square$

**Corollary 9.12** (The string coupling as the Koide reciprocal). *The string coupling constant satisfies  $g_s = (R_{11}/l_p)^{3/2} = (R_{11}/l_p)^{|\Pi|/d_2} = (R_{11}/l_p)^{1/K}$ . The coupling exponent  $|\Pi|/d_2 = 3/2 = 1/K$  is the Koide reciprocal — the same constant governing the superconformal ghost weight (Theorem 9.5(ii)).*

## 9.7 The Sublattice-to-Force Map

The N-Exhaustion Theorem (Theorem 9.3) determines which gauge factors exist. The sublattice structure determines how they couple.

**Theorem 9.13** (Sublattice-to-force correspondence). *The three forces of the Standard Model are mapped to sublattice families as follows:*

- (i) Strong force:  $d = 3$  (cubic sublattice). Three positions per period correspond to three colour charges. The adjoint representation has  $\dim = 3^2 - 1 = 8$  gluons (Theorem 9.1). Confinement arises because  $d = 3$  has the smallest non-trivial period in the lattice: quarks cannot be separated beyond one  $d = 3$  period without creating a new quark-antiquark pair.
- (ii) Weak force: governed by  $d_W = N(1 - K) = 12 \cdot 1/3 = 4$  (the quartic sublattice). The gauge group is  $SU(2)$  because  $\varphi(4) = 2$  residues per  $d = 4$  period (Proposition 8.3) give two fundamental charges (the two weak isospin states, up-type and down-type). The adjoint dimension is  $2^2 - 1 = 3$  (the three weak bosons  $W^+$ ,  $W^-$ ,  $Z^0$  before mixing).
- (iii) Electromagnetic force:  $d = 1$  (octave/identity sublattice). One generator yields one gauge boson (the photon). The electromagnetic force is the ambient lattice field: it couples to all charged matter because  $d = 1$  divides every d-family.

*Proof.* Item (i): by Corollary 9.2, the  $d = 3$  sublattice forces  $d_{\text{adj}}(3) = 8$  gauge bosons, which by the N-Exhaustion Theorem (Theorem 9.3) are the gluon octet of SU(3).

Item (ii): the quartic sublattice  $d = 4$  satisfies  $d_W = N(1 - K) = 12(1 - 2/3) = 4$ . By Proposition 8.3, the Euler totient  $\varphi(4) = 2$  gives two coprime residues modulo 4, which are the two fundamental charges of the weak isospin doublet. By Theorem 9.1, the adjoint dimension is  $2^2 - 1 = 3$ .

Item (iii): the  $d = 1$  sublattice has one generator (the octave identity,  $\varphi(1) = 1$ ). It divides every  $d$ -family because  $1 \mid d$  for all  $d$ , making the U(1) photon the universal coupling.

The total gauge-boson count is  $8 + 3 + 1 = 12 = N$ , confirming completeness without remainder (Theorem 9.3, condition (C1)). The remaining native families  $d \in \{4, 6, 12\}$  do not generate additional gauge factors (the  $N = 12$  budget is exhausted) but govern the mixing between forces:  $d = 4$  governs the electroweak mixing,  $d = 6$  governs the electroweak unification scale, and  $d = 12$  is the full electromagnetic resolution.  $\square$   $\square$

## 9.8 Supercharges, gauge group dimensions, and brane structure

Three further results follow from the manifold state count  $S = 4$  (equation (1)).

**Proposition 9.14** (Maximum supercharges). *The maximum number of supercharges in M-theory is*

$$Q_{\text{max}} = 2^{S+1} = 2^5 = 32.$$

*The graviton supermultiplet in 11 dimensions has  $2^{2S} = 2^8 = 256$  states — the square of the supercharge spinor dimension, as required for a short multiplet of the super-Poincaré algebra [33].*

**Proposition 9.15** (Anomaly-cancellation gauge group). *The dimension of the anomaly-free gauge group in 10-dimensional chiral supergravity is*

$$\dim(\text{gauge group}) = 2^S (2^{S+1} - 1) = 16 \times 31 = 496.$$

*This equals  $\dim \text{SO}(32) = 32 \cdot 31/2 = 496$  and  $\dim(\text{E}_8 \times \text{E}_8) = 2 \times 248 = 496$ . The anomaly-cancellation condition [34] that forces the gauge group is expressible as a function of  $S$  alone.*

**Proposition 9.16** (Central extensions and brane structure). *The 11-dimensional super-Poincaré algebra with 32 supercharges has:*

- *Symmetric  $Q \times Q$  matrix entries:  $32 \times 33/2 = 528$ .*
- *Momentum components:  $D_M = 11$ .*
- *Membrane (rank-2) charges:  $\binom{11}{2} = 55$ .*
- *Fivebrane (rank-5) charges:  $\binom{11}{5} = 462$ .*
- *Check:  $55 + 462 = 517 = 528 - 11$ . ✓*

*The membrane (2-dimensional spatial extent, worldvolume  $2+1 = 3 = |\Pi|$ ) couples to the 3-form field. The string (1-dimensional, worldsheet  $1+1 = 2 = d_2$ ) couples to the 2-form field. The product  $2 \times 5 = 10 = D_{\text{string}}$ ; the sum  $2 + 5 + 4 = 11 = D_M$ .*

**Remark 9.17** (Membrane–fivebrane duality on the LCM tower). The membrane at  $d = 2$  is native at  $N = 12$ ; the fivebrane at  $d = 5$  is a shadow family at  $N = 12$  (Table 7), becoming native at  $N = 60 = \text{lcm}(12, 5)$ . The membrane–fivebrane duality (electric–magnetic duality of the 3-form) maps to the  $d = 2 \leftrightarrow d = 5$  relationship on the lattice; the LCM tower escalation is the lattice’s mechanism for resolving structure that is present but not yet native at base resolution.

## 9.9 The Structural Significance Principle

The derivations of §§9.1–9.8 promote the shadow-force identifications from dimensional coincidence to structural derivation. It remains to state explicitly the methodological principle that distinguishes structurally derived identifications from numerological coincidence, addressing the compound critique that a dense lattice combined with unconstrained dimensional matching constitutes a confirmation-bias mechanism.

**Principle 9.18** (Structural Significance). A physical identification of sublattice family  $d$  with a physical sector is *structurally significant* if and only if it satisfies all four of the following conditions simultaneously:

- (P1) *V-threshold at native resolution.* The identification’s lattice placement satisfies the V-threshold criterion of Definition 7.13 at the lowest resolution  $N_{\text{native}}$  where  $d$  is native (Proposition 7.14).
- (P2) *Zero free parameters.* The identification follows from a derivation whose only inputs are the ET constants  $\{N, K, |\Pi|, S, V\}$ . A framework with zero adjustable parameters cannot be tuned to match arbitrary data.
- (P3) *Cross-domain convergence.* The same lattice cell is independently reached by physical ratios from  $k \geq 2$  distinct domains and independently reached by forward computation from ET constants. The probability of coincidental convergence drops as  $P^k$  — exponentially in the number of independent domains.
- (P4) *Multiple independent derivation routes.* The same result is derived by two or more structurally independent methods.

*Remark 9.19* (The Subsumption Law as anti-cherry-picking). The Subsumption Law (Principle 3.13, condition (iii)) demands completeness without remainder. Applied to physical identifications: if sublattice family  $d$  is identified with physical sector  $X$ , then  $d$  must account for all instances of that  $d$ -number across all domains simultaneously. Selectively identifying  $d = 8$  with SU(3) while ignoring other  $d = 8$  phenomena violates condition (iii). The shadow-force table of §8.8 is the Subsumption Law’s demand: every  $d$ -family must be identified with all its physical renderings simultaneously, or the identification is incomplete.

*Remark 9.20* (Structural vs. statistical significance). The Structural Significance Principle is not a frequentist or Bayesian statistical test imported from outside. It is ET-native: (P1) uses the manifold’s own base variance  $V = 1/N$  as the threshold; (P2) uses the Three Tools; (P3) uses the universality anchor  $3 = 3 = 3 = \Sigma$  (every domain is within  $\Sigma$ , so cross-domain convergence is a Subsumption test); (P4) uses the Verification Principle (independent routes arriving at the same answer equals mathematical consistency equals sufficient Descriptors, by Corollary 3.10). All four conditions are instances of tools already established in §3.

## 10 The LCM tower of refinements

### 10.1 When $N = 12$ is insufficient

The sublattice families at  $N = 12$  are the divisors of 12, namely  $\{1, 2, 3, 4, 6, 12\}$ . Phenomena whose structural symmetries lie outside this set cannot be resolved at the base lattice: five-fold symmetry (pentagonal, icosahedral, golden-ratio), seven-fold symmetry (heptagonal, certain capsid symmetries), eleven-fold symmetry (undecimal), eight-fold (octet, gluon), nine-fold (nonic), ten-fold (decic). To host these one must refine the lattice to a higher  $N$ .

## 10.2 Canonical refinements

The minimal  $N$  supporting a required family  $d$  is any multiple of  $d$  that is also a multiple of 12 (so as to preserve the base families). The smallest such  $N$  is  $\text{lcm}(12, d)$ .

**Proposition 10.1** (Canonical LCM landmarks). *The sequence*

$$\text{lcm}(1, \dots, k) \text{ for } k = 4, 5, 7, 9, 11$$

*is 12, 60, 420, 2520, 27720 respectively. Each is a multiple of 12; each introduces one or more new primes as native divisors:*

- 12 hosts  $\{1, 2, 3, 4, 6, 12\}$ ;
- $60 = 2^2 \cdot 3 \cdot 5$  adds  $d = 5$  (and composites 10, 15, 20, 30);
- $420 = 2^2 \cdot 3 \cdot 5 \cdot 7$  adds  $d = 7$  and the biological cross-product  $d = 35 = 5 \cdot 7$ ;
- $2520 = 2^3 \cdot 3^2 \cdot 5 \cdot 7$  completes  $d \leq 9$ ;
- $27720 = 2^3 \cdot 3^2 \cdot 5 \cdot 7 \cdot 11$  completes  $d \leq 11$  and makes all twelve families  $d \in \{1, 2, \dots, 12\}$  simultaneously native.

*Proof.* Direct calculation:

$$\begin{aligned} \text{lcm}(1, 2, 3, 4) &= 12, & \text{lcm}(1, \dots, 5) &= 60, & \text{lcm}(1, \dots, 7) &= 420, \\ \text{lcm}(1, \dots, 9) &= 2520, & \text{lcm}(1, \dots, 11) &= 27720. \end{aligned}$$

Each is a multiple of 12 because  $\{2, 3, 4\} \subset \{1, \dots, k\}$  for  $k \geq 4$ . The prime factorisations follow immediately.  $\square$   $\square$

## 10.3 Unification of the two readings

**Proposition 10.2** (Multiplicative and LCM readings). *The sequence of multiplicative refinements of the base lattice is 12, 24, 36, 48, 60, 72, ... (the positive multiples of 12). The LCM reading 12, 60, 420, 2520, 27720, ... is a sparse subsequence of this multiplicative reading.*

The proof is Proposition 10.1: each LCM landmark is a multiple of 12. The multiplicative reading is for uniform refinement of resolution (e.g., refining precision in the same family structure); the LCM reading is for introducing a new prime as a native divisor.

## 10.4 The continuous limit

**Proposition 10.3** (Asymptotic limit). *As  $N \rightarrow \infty$  through either reading of the tower,*

$$\mathcal{L}_N \longrightarrow (\mathbb{R}^+, \times),$$

*with the descriptor gap bounded by  $|\varepsilon| \leq 600/N$  cents, which tends to 0.*

*Proof.* The set  $\mathcal{L}_N = \{2^{k/N}\}$  is dense in  $\mathbb{R}^+$  under the multiplicative metric as  $N$  grows, since the steps  $2^{1/N}$  tend to 1. The gap bound follows from the rounding bound in the proof of Proposition 8.5.  $\square$   $\square$

**Corollary 10.4** (Asymptotic Approach Theorem). *For any irrational  $r \in (\mathbb{R}^+, \times)$ , the descriptor-gap residual  $|\varepsilon_N(r)|$  of the lattice projection at resolution  $N$  satisfies*

$$|\varepsilon_N(r)| \longrightarrow 0 \quad \text{as } N \rightarrow \infty, \quad \text{with} \quad |\varepsilon_N(r)| > 0 \text{ for every finite } N.$$

*An irrational ratio is approached asymptotically by the lattice without ever being attained at finite resolution.*

*Proof.* The asymptotic limit follows from the bound  $|\varepsilon_N(r)| \leq 600/N$  cents of Proposition 10.3. Strict positivity at every finite  $N$  follows from irrationality of  $r$ : if  $|\varepsilon_N(r)| = 0$  at some finite  $N$ , then  $N \log_2 r$  is an integer  $k$ , so  $r = 2^{k/N}$  is a rational power of 2 and hence rational, contradicting the hypothesis.  $\square$   $\square$

*Remark 10.5* (Structural origin). Corollary 10.4 expresses the finite-infinite asymmetry between the Cardinals:  $|P| = \Omega$  supports irrational positions;  $|D| = n$  provides only finitely many descriptors at any finite resolution;  $|T| = [0/0]$  resolves  $P$  under  $D$  but cannot exhaust  $P$ . Asymptotic approach without attainment is the structural signature of this asymmetry — not a deficiency of the lattice but a corollary of the cardinality trichotomy. The corollary is the formal vehicle by which any decimal value used in the paper is to be read: every finite-digit numeric is a finite- $D$  approximation of an asymptotic-reference value, and finite truncation never improves accuracy — it stops asymptotic refinement at a chosen depth.

#### 10.4.1 Structural necessity of asymptotic precision

The asymptotic character of Corollary 10.4 is not merely a technical limitation but a structural necessity forced by the categorical distinctions of the three Cardinals.

**Proposition 10.6** (Structural necessity of imprecision). *Perfect precision at finite resolution would collapse the categorical distinction between the three Cardinals.*

*Proof.* Suppose  $D$  could perfectly capture  $P$  at some finite resolution  $N_0$ . Then  $|D| = n_0$ . Descriptors would exhaust  $|P| = \Omega$  at that resolution: every position in  $P$  would be finitely described, and no  $P$ -content would lie beyond  $D$ 's reach. But then  $\Omega = n_0$ , collapsing the finite-infinite distinction. Without the substrate-constraint asymmetry ( $\Omega \neq n$ ), the binding order (Axiom 2.17) loses its force, the  $P$ - $D$  singularity dissolves, and the master equation ceases to produce distinct Cardinals. Similarly, suppose  $D$  could perfectly capture  $T$  at finite resolution. Then determinacy ( $n$ ) would subsume indeterminacy ( $[0/0]$ ), and  $T$ 's navigational capacity would be reducible to a finite Descriptor — eliminating agency by reducing  $T$  to  $D$ , violating the categorical disjointness (Axiom 2.6). The asymptotic approach preserves the categorical distinctions of the three Cardinals while permitting arbitrarily close approximation at arbitrarily high resolution.  $\square$   $\square$

### 10.5 Correspondence with integrative levels

The qualitative hierarchy of integrative levels introduced in §3.12 has a precise quantitative counterpart in the LCM tower. The two are not parallel concepts but the same structure read from two directions — the phenomenological (what kind of wholeness is present?) and the arithmetic (which sublattice families are native?).

**Proposition 10.7** (Integrative-to-resolution correspondence). *Let  $\ell$  be an integrative level, and let  $\mathcal{D}_\ell \subseteq \mathbb{N}$  be the set of sublattice families appearing in the  $(P, D, T)$  decompositions of level- $\ell$  phenomena. Then the minimum resolution  $N_\ell^{\min}$  at which level  $\ell$  is fully identifiable in the lattice framework is*

$$N_\ell^{\min} = \text{lcm}(\mathcal{D}_\ell).$$

*Proof.* A sublattice family  $d$  is native at resolution  $N$  if and only if  $d \mid N$  (Section 8). Level  $\ell$  requires every family in  $\mathcal{D}_\ell$  to be native simultaneously, so  $N$  must be a common multiple of all  $d \in \mathcal{D}_\ell$ . The smallest such common multiple is  $\text{lcm}(\mathcal{D}_\ell)$ .  $\square$   $\square$

**Corollary 10.8** (Monotonicity of the tower). *If  $\mathcal{D}_{\ell_1} \subseteq \mathcal{D}_{\ell_2}$  then  $N_{\ell_1}^{\min} \mid N_{\ell_2}^{\min}$ . Integrative levels whose Descriptor requirements strictly extend those of lower levels correspond to tower levels that are strict multiples of the lower ones.*

The corollary says that ascending the integrative hierarchy (adding Descriptor structure) induces divisibility ascent in the tower: one cannot host a more integrative level on less resolution. Conversely, every divisibility ascent in the tower is the emergence of new integrative structure at the lattice level — new sublattice families that did not exist in the previous resolution.

### The doubling theorem

The qualitative statement “higher integrative level requires higher resolution” becomes an exact quantitative law when one asks how the divisor count evolves along the canonical tower. Write  $N_\ell$  for the  $\ell$ -th canonical resolution ( $N_0 = 12$ ,  $N_1 = 60$ ,  $N_2 = 420$ ,  $N_3 = 2520$ ,  $N_4 = 27720$ , ...) and  $\tau(N_\ell)$  for the number of sublattice families native at that resolution (equivalently, the number of divisors of  $N_\ell$ ).

**Theorem 10.9** (Integrative–resolution doubling). *The divisor count of the  $\ell$ -th canonical resolution satisfies*

$$\tau(N_\ell) = \tau(N_0) \cdot 2^\ell = 6 \cdot 2^\ell. \quad (26)$$

*Equivalently, the integrative level is a logarithm of the family count,*

$$\ell = \log_2 \frac{\tau(N_\ell)}{\tau(N_0)} = \log_2 \frac{\tau(N_\ell)}{6}. \quad (27)$$

*Proof. Base case ( $\ell = 0$ ).* By Proposition 6.1,  $N_0 = |\Pi| \cdot S = 3 \cdot 4 = 12$ . Since  $\gcd(|\Pi|, S) = \gcd(3, 4) = 1$ , the divisor function is multiplicative at this argument:

$$\tau(N_0) = \tau(|\Pi| \cdot S) = \tau(|\Pi|) \cdot \tau(S) = \tau(3) \cdot \tau(4) = 2 \cdot 3 = 6.$$

Thus  $\tau(N_0) = 6 = 6 \cdot 2^0$ , and the base case holds.

*Inductive step.* We show that each canonical refinement exactly doubles  $\tau$ . The Descriptor Gap Principle (Principle 3.9) states that every gap in a description is itself a Descriptor. The smallest Descriptor capable of distinguishing a previously-unresolved pair of configurations is a single binary axis — a yes/no distinction with two values. Adjoining one binary Descriptor to the model therefore exactly doubles the count of distinguishable configurations, and the lattice-level consequence is that the new resolution  $N_{\ell+1}$  must satisfy  $\tau(N_{\ell+1}) = 2\tau(N_\ell)$ .

We verify directly that each canonical LCM step realises this doubling.

1.  $N_0 = 12 \rightarrow N_1 = 60$  ( $60 = 12 \cdot 5$ ). Introducing the new prime 5 with exponent 1 multiplies  $\tau$  by  $(1+1)/(0+1) = 2$ .
2.  $N_1 = 60 \rightarrow N_2 = 420$  ( $420 = 60 \cdot 7$ ). Introducing prime 7 multiplies  $\tau$  by 2.
3.  $N_2 = 420 \rightarrow N_3 = 2520$  ( $2520 = 420 \cdot 6$ ). The prime-power content changes as  $2^2 \cdot 3 \rightarrow 2^3 \cdot 3^2$ , giving the joint factor

$$\frac{3+1}{2+1} \cdot \frac{2+1}{1+1} = \frac{4}{3} \cdot \frac{3}{2} = 2.$$

This is the only canonical step that is realised by combined prime-power upgrades rather than by a single new prime; the product of the two fractional upgrades is exactly 2.

4.  $N_3 = 2520 \rightarrow N_4 = 27720$  ( $27720 = 2520 \cdot 11$ ). Introducing prime 11 multiplies  $\tau$  by 2.
5.  $N_4 = 27720 \rightarrow N_5 = 360360$  ( $360360 = 27720 \cdot 13$ ). Introducing prime 13 multiplies  $\tau$  by 2.

By induction,  $\tau(N_\ell) = \tau(N_0) \cdot 2^\ell = 6 \cdot 2^\ell$  for every  $\ell \geq 0$ .

*Extensibility.* The canonical tower extends indefinitely: for every  $\ell \geq 5$ , a canonical refinement exists that doubles  $\tau$  (realised by successive new primes 17, 19, 23, ..., or in special cases by prime-power combinations that product to 2, exactly as at  $\ell = 3$ ). Therefore the doubling law (26) holds for all  $\ell \geq 0$ . □ □



*Remark 10.10* (Two readings of a single equation). Equation (26) admits two parallel interpretations that describe the same underlying structure:

**Number-theoretic reading.**  $\tau(N_\ell)$  counts the divisors of the resolution integer  $N_\ell$ , equivalently the sublattice families native at that resolution. The equation states that each canonical LCM refinement exactly doubles the native-family count.

**Phenomenological reading.**  $\tau(N_\ell)$  counts the Descriptors needed to characterise the configurations of integrative level  $\ell$ . The equation states that each integrative emergence introduces one binary distinction axis and therefore exactly doubles the required Descriptor count.

Both readings measure the same quantity: the structural richness natively expressible at integrative level  $\ell$ . The phenomenological ladder (quantum, atomic, molecular, biological, cognitive, civilisational, ...) and the number-theoretic ladder (12, 60, 420, 2520, 27720, ...) are the same ladder seen from two directions. The coincidence is not numerical accident but structural necessity: the minimal closure of a descriptor gap at any integrative level is a binary distinction (Principle 3.9), and in the lattice this minimal closure is precisely a factor-2 increase in  $\tau$ .

**Corollary 10.11** (The integrative level is a logarithm). *For any configuration whose Cardinal decomposition requires  $\tau_X$  sublattice families to be expressible, the integrative level of  $X$  is*

$$\ell(X) = \log_2 \frac{\tau_X}{6}.$$

*This assigns a quantitative real-valued index to every substantiated configuration, converting the previously qualitative “integrative level” into a computable lattice quantity.*

Table 9 collects the empirical correspondences used in Section 22, now augmented with the divisor counts predicted by Theorem 10.9.

*Remark 10.12* (Physical reading of the tower). The LCM tower is therefore not merely a sequence of successively finer approximations to a continuous limit; it is a sequence of discrete ontological transitions at which previously unavailable Descriptor structure becomes native, with each transition *exactly doubling* the Descriptor count by Theorem 10.9. Each step up the tower makes a new integrative regime possible: ascending from  $N = 12$  to  $N = 60$  makes pentagonal biology possible; ascending from  $N = 60$  to  $N = 420$  makes septic biochemistry possible; ascending from  $N = 420$  to  $N = 2520$  makes enneagonal structure and the  $\alpha^{-1}$  universal coordinate native; ascending from  $N = 2520$  to  $N = 27720$  makes  $d = 11$  native and completes the simple-divisor floor  $d \leq 12$ . The continuous limit of Proposition 10.3 is the idealised endpoint at which every integrative level is resolved simultaneously.

*Remark 10.13* (Independent number-theoretic distinction of the canonical resolutions). The base resolution  $N = 12$  and the universal resolution  $N_4 = 27720$  are forward-derived in this paper from the PDT primitives (§6, §10). They are also independently distinguished in classical number theory: an integer  $n$  is *highly composite* (Ramanujan 1915 [5]) if every  $m < n$  has  $d(m) < d(n)$ , where  $d(n)$  counts the divisors of  $n$ . The base resolution  $N = 12$  is the fifth highly composite number ( $d(12) = 6$ , exceeding  $d(m)$  for every  $m < 12$ ); the universal resolution  $N_4 = 27720$  is the twenty-fifth highly composite number ( $d(27720) = 96$ , exceeding  $d(m)$  for every  $m < 27720$ ); and the intermediate resolutions  $N_1 = 60$  and  $N_3 = 2520$  are likewise highly composite. The canonical tower therefore traces a sparse subsequence of Ramanujan’s highly composite numbers, distinguished further by the LCM closure  $N_\ell = \text{lcm}(1, \dots, n_\ell)$  that forces *every* small divisor to be native. The same integers are therefore singled out twice: forward, from PDT primitives plus the doubling law (Theorem 10.9); and laterally, from elementary divisor counting. This cross-corroboration is non-trivial because the two derivations share no axioms: the present framework starts from  $(P, D, T)$  and the master equation, while Ramanujan’s classification starts from the divisor function  $d(n)$  on the natural numbers. The convergence on the same integers is independent evidence that the canonical resolutions are mathematically distinguished, not arbitrary choices.

| $\ell$ | $N_\ell$ | $\tau(N_\ell)$ | Newly native families               | Empirical phenomena first hosted here                                                                                                                                                      |
|--------|----------|----------------|-------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 0      | 12       | 6              | $\{1, 2, 3, 4, 6, 12\}$             | Koide lepton ratio; 12-TET intervals; 3:4:5 triangle; Krebs ( $8=2^3$ at $d=1$ ); glycolysis (10 at $d=3$ )                                                                                |
| 1      | 60       | 12             | adds $\{5, 10, 15, 20, 30\}$        | Pentagonal symmetry; Caspar–Klug $T=5$ virus capsids; icosahedral / golden-ratio structural biology                                                                                        |
| 2      | 420      | 24             | adds $\{7, 14, 21, 28, 35, \dots\}$ | Septic cycles; heptagonal structural biochemistry; the biological cross-product $d=35=5 \cdot 7$                                                                                           |
| 3      | 2520     | 48             | adds $\{8, 9, 18, 24, \dots\}$      | Enneagonal (9-fold) biological symmetries; $d=315=3^2 \cdot 5 \cdot 7$ ( $\alpha^{-1}$ host) first becomes native                                                                          |
| 4      | 27720    | 96             | adds $\{11, 22, \dots\}$            | Universal reference: all simple divisors $d \leq 12$ simultaneously native; cortical 11-dimensional clique dynamics [18]; $\alpha^{-1}$ at $(k, d) = (196768, 315)$ in this standard frame |
| 5      | 360360   | 192            | adds $\{13, 26, \dots\}$            | Predicted new domain: phenomena requiring $d=13$ structure                                                                                                                                 |

Table 9: The LCM tower and the integrative-level correspondence. The divisor column confirms the doubling law  $\tau(N_\ell) = 6 \cdot 2^\ell$  of Theorem 10.9: each row’s  $\tau$  value is exactly twice the previous. The “newly native” entries are those families  $d$  with  $d \mid N_\ell$  but  $d \nmid N_{\ell'}$  for every earlier tower level  $N_{\ell'}$ . By Proposition 10.7, a phenomenon whose Cardinal decomposition involves family set  $\mathcal{D}_\ell$  is minimally identifiable at  $N_\ell^{\min} = \text{lcm}(\mathcal{D}_\ell)$ ; by Corollary 10.11 its integrative level is  $\ell = \log_2(\tau_\ell/6)$ .

*Remark 10.14* (Empirical anchor for  $N_4 = 27720$ : cortical 11-dimensional clique dynamics). The Identification Principle (§3.4) and the Subsumption Law (§3.6) admit an independent empirical confirmation that  $N_4 = 27720$  is the cortical resolution, supplied by the Blue Brain Project’s 2017 algebraic-topology study of rat neocortical microcircuitry [18]. Reimann et al. measured directed simplicial cliques — groups of neurons all mutually connected with directionality — in both a reconstructed digital neocortical microcircuit and *in vitro* multi-neuron patch-clamp recordings, finding that cliques routinely formed through dimensions 1–7 and that some networks contained directed simplices up through dimension 11. The cliques bound into cavities (topological holes) which appeared during stimulus processing and dissolved afterward; the dynamical signature is a cascade through progressively higher dimensions  $1 \rightarrow 2 \rightarrow \dots \rightarrow 11$  and back, characterised in the original paper as the brain “building then razing a tower of multi-dimensional blocks”.

In the present framework this finding is forced. A dimension- $n$  directed clique is the lattice expression of a  $d_r = n$  family occupation; the cortex assembling progressively higher-dimensional structures is the cortex activating progressively higher- $d$  sublattice families, and disassembling them is releasing those activations. The cascade through the eleven simple dimensions can only occur at a lattice resolution where every  $d \in \{1, 2, \dots, 11\}$  is simultaneously native. By Proposition 10.7, native presence requires  $d \mid N$ ; the smallest  $N$  satisfying  $d \mid N$  for every  $d \in \{1, \dots, 11\}$  is  $\text{lcm}(1, \dots, 11) = 2^3 \cdot 3^2 \cdot 5 \cdot 7 \cdot 11 = 27720$ , which is exactly  $N_4$  of the LCM tower. At  $N = 12$  the dimensions  $\{5, 7, 8, 9, 10, 11\}$  are non-divisors and have no native sublattice; at  $N = 132 = 2^2 \cdot 3 \cdot 11$  the family  $d = 11$  becomes native but  $\{5, 7, 8, 9, 10\}$  do not, so the full cascade still cannot run; only at  $N = 27720$  does the entire cascade through

$d = 1, \dots, 11$  admit native lattice support. The cortical dynamics of Reimann et al. therefore require resolution  $N = 27720$ , and conversely the lattice’s prediction that  $N_4 = 27720$  is the cognitive integrative level (Table 9) is empirically confirmed by these dynamics.

The remark of Markram in the original communication — that “the mathematics usually applied for studying neural networks cannot detect the high-dimensional structures and spaces that we now see clearly” — has the structural reading via the Descriptor Gap Principle (§3.5). Conventional neural-network mathematics (linear combinations and low-order nonlinearities) operates approximately at 12 ET-equivalent resolution, hosting only  $d \in \{1, 2, 3, 4, 6, 12\}$ , so 11-dimensional cliques are invisible to it because  $d = 11 \nmid 12$  — a Descriptor Gap at the level of mathematical apparatus. Algebraic topology becomes the necessary detection apparatus precisely because algebraic topology counts simplicial structures whose component count follows  $C(n) = n^2(n^2 - 1)/12$  (Proposition 23.1, the Riemann-curvature-component count) — the manifold-symmetry constant  $N = 12$  appears in the denominator, which is the lattice signature in the topology-counting formula itself. The cortex is therefore the first confirmed physical system whose dynamics empirically require the full simple-divisor floor of  $N_4 = 27720$ , and whose proper mathematical apparatus is itself indexed by the manifold symmetry  $N = 12$ . The high-dimensional clique/cavity structure of neural networks has been independently confirmed by an entirely different methodology: the Sizemore et al. 2018 study [19] applied persistent homology to the human structural connectome (eight healthy adults, scanned in triplicate) and recovered cliques and topological cavities consistently across subjects and significantly above wiring-minimisation null models, in agreement with the local-microcircuit findings of Reimann et al. 2017. Two independent measurements — different species (rat vs. human), different scale (local microcircuit vs. whole-brain connectome), different mathematical apparatus (directed simplicial complexes vs. persistent homology) — converge on the same finding: the neural substrate hosts the higher- $d$  sublattice families that 12 ET-equivalent mathematics cannot natively express.

## 11 The complex lattice

Many phenomena have both a magnitude and a phase character. The lattice extends from the multiplicative group  $(\mathbb{R}^+, \times)$  to the multiplicative group  $(\mathbb{C}^\times, \times)$  by projecting the phase to a second axis.

### 11.1 Imaginary-axis projection

**Definition 11.1** (Phase projection). Let  $\theta \in [0, 2\pi)$  be an angle. The *phase projection* at resolution  $N$  is

$$k_\theta = \text{round}\left(\frac{N\theta}{2\pi}\right) \bmod N, \quad d_\theta = \frac{N}{\gcd(|k_\theta|, N)},$$

$$\varepsilon_\theta = \left(\frac{N\theta}{2\pi} - k_\theta\right) \cdot \frac{1200}{N} \text{ cents.}$$

The phase axis classifies rotational structure in exactly the same divisor-lattice way that the magnitude axis classifies scaling structure. The cyclic group  $U(1) = \mathbb{R}/2\pi\mathbb{Z}$  is discretised into  $N$  equal sectors.

### 11.2 Complex projection and the Gaussian integer lattice

**Definition 11.2** (Complex projection). For  $z = re^{i\theta} \in \mathbb{C}^\times$ , the *complex projection* is

$$w = k_r + ik_\theta \in \mathbb{Z}[i],$$

with the real-axis and imaginary-axis projections defined by Definitions 7.1 and 11.1 respectively. The *combined sublattice family* is

$$d_c = \text{lcm}(d_r, d_\theta).$$

**Proposition 11.3** (LCM amplification). *The combined family  $d_c = \text{lcm}(d_r, d_\theta)$  is always at least as large as the larger of  $d_r$  and  $d_\theta$ .*

The proof is elementary:  $\text{lcm}(a, b) \geq \max(a, b)$  with equality iff one divides the other. Structurally, off-axis (complex) positions have richer sublattice structure than either axis alone. At  $N = 12$  the  $6 \times 6$  grid of  $(d_r, d_\theta)$  pairs produces combined families taking values in  $\{1, 2, 3, 4, 6, 12\}$  with the distribution shown in Table 10.

| $d_r \backslash d_\theta$ | 1  | 2  | 3  | 4  | 6  | 12 |
|---------------------------|----|----|----|----|----|----|
| 1                         | 1  | 2  | 3  | 4  | 6  | 12 |
| 2                         | 2  | 2  | 6  | 4  | 6  | 12 |
| 3                         | 3  | 6  | 3  | 12 | 6  | 12 |
| 4                         | 4  | 4  | 12 | 4  | 12 | 12 |
| 6                         | 6  | 6  | 6  | 12 | 6  | 12 |
| 12                        | 12 | 12 | 12 | 12 | 12 | 12 |

Table 10: The  $6 \times 6$  table of combined sublattice families  $d_c = \text{lcm}(d_r, d_\theta)$  at the base lattice  $N = 12$ . Off the diagonal the LCM amplification typically drives  $d_c$  towards 12 (full resolution); of the 36 entries, 15 equal 12. This explains why most physical observables, which carry both magnitude and phase character, project to  $d = 12$  at the base lattice.

### 11.3 The D–T gradient

Let  $\alpha = \arctan(k_\theta/k_r)$  be the angle of the complex-lattice coordinate from the real axis. Define

$$(\text{magnitude fraction}) = \cos^2 \alpha, \quad (\text{phase fraction}) = \sin^2 \alpha.$$

At  $\alpha = 0$  (pure real) the coordinate is fully magnitudinal; at  $\alpha = \pi/2$  (pure imaginary) it is fully phase; in between it interpolates smoothly. The transition from deterministic to probabilistic behaviour (the familiar classical-to-quantum transition) is, in this picture, not a sharp boundary but a continuous function of  $\alpha$ .

### 11.4 The Gaussian-prime correspondence

The sublattice families have a clean number-theoretic pedigree in the Gaussian integers  $\mathbb{Z}[i]$ .

**Proposition 11.4** (Gaussian prime correspondence). *The sublattice families at  $N = 12$  and its LCM extensions correspond to the three Gaussian prime classes as follows:*

- $p = 2$  (ramified:  $2 = -i(1+i)^2$ ) generates the binary sublattices  $d \in \{2, 4, 8\}$ ;
- $p \equiv 3 \pmod{4}$  (inert: remain prime in  $\mathbb{Z}[i]$ ) generate the sublattices  $d \in \{3, 7, 11\}$  and their squares;
- $p \equiv 1 \pmod{4}$  (split:  $p = \pi\bar{\pi}$  in  $\mathbb{Z}[i]$ ) generate the sublattices  $d \in \{5, 13, \dots\}$  of mixed magnitude/phase character.

The correspondence is elementary but structurally informative: the three Gaussian prime classes exactly parallel the three Cardinals  $P$ ,  $D$ ,  $T$ . Ramified primes (which “double” in  $\mathbb{Z}[i]$ ) are  $P$ -class; inert primes (which remain on the real axis) are  $D$ -class; split primes (which factor across the real-imaginary split) are mixed  $D+T$ -class.

## 12 The 24-family catalog and the 144-cell Force Quadrant Grid

The real axis of the Sempaevum carries sublattice families classified by divisors of the current resolution  $N$  (§8). The imaginary axis does the same — but the physical character of the two axes is categorically distinct because the underlying manifolds are categorically distinct (Proposition 2.30:  $(\mathbb{R}^+, \times)$  flat,  $(U(1), \times)$  positively curved). Taking the real axis and the imaginary axis together at the universal resolution  $N_{\text{univ}} = 27720$ , the sublattice families partition into a catalog of 24 basic families (12 on each axis, distinguished by role not by number) plus a catalog of 42 genuinely-mixed combined off-axis families, for a total structural inventory that fills a  $12 \times 12$  *Force Quadrant Grid* of 144 cells.

### 12.1 The 12 real-axis FORCE families and the 12 imaginary-axis PHASE families

*Remark 12.1* (The 24 FORCE/PHASE labels are the harmonic-family count on  $\mathcal{L}_{\mathbb{C}}$ ). The 12 real-axis FORCE labels and the 12 imaginary-axis PHASE labels of this section, taken together, are the 24 axis-restricted harmonic families on the complex lattice  $\mathcal{L}_{\mathbb{C}}$  (Definition 8.10, Table 6): twelve harmonic families on the real axis (FORCE) and twelve harmonic families on the imaginary axis (PHASE), each labeled by an integer  $d \in \{1, 2, \dots, 12\}$  in the simple- $d$  range and partitioned per axis into six SIMPLE ( $d \mid 12$ , native at  $N = 12$ ) and six COMPLEX ( $d \nmid 12$ , shadow at base, native at  $n_c(d) = \text{lcm}(12, d)$ ). This 24-fold enumeration is structurally distinct from the sublattice-family count  $\tau(N)$  at any resolution  $N$ : at  $N = 12$  the lattice carries  $\tau(12) = 6$  sublattice families per axis (the divisors of 12); at  $N_{\text{univ}} = 27720$  it carries  $\tau(27720) = 96$  sublattice families per axis. The harmonic-family layer remains twelve per axis at the simple- $d$  range across all resolutions because it enumerates the per-axis structural modes of the lattice, not the gcd-classes of individual coordinates. The numerical coincidence at  $N = 12$  that both layers carry six SIMPLE values per axis ( $d \in \{1, 2, 3, 4, 6, 12\}$  in both enumerations) reflects the alignment of the simple-harmonic-family  $d$ -labels with the divisors of  $N = 12$ , not an identity of layers; the layers remain categorically distinct (§8.6, Remark 8.12). The palindromic cascade is the discovery and verification engine that makes the per-axis count of twelve explicit (§8.6, Remark 8.11; §8.7, Theorem 8.13); cascade asymmetry across the two axes (real 25-level stability vs imaginary 2-level) determines the resolution at which the discovery completes per axis (§13.2, Remark 13.4).

The real axis is  $D$ 's operational domain: its sublattice families classify magnitude structure. The imaginary axis is  $T$ 's operational domain: its families classify phase/rotation structure. Same divisor values, categorically different physics.

**Definition 12.2** (FORCE and PHASE families). Let  $d_r \mid N$  be the sublattice family of a real-axis coordinate  $k_r$ , and let  $d_\theta \mid N$  be the sublattice family of an imaginary-axis coordinate  $k_\theta$ , both via the projection formula  $d = N / \gcd(|k|, N)$  (Definition 7.1) applied to the respective axis. We call  $d_r$  the *FORCE family* of the position and  $d_\theta$  the *PHASE family*.

At the universal resolution  $N_{\text{univ}} = 27720 = 2^3 \cdot 3^2 \cdot 5 \cdot 7 \cdot 11$ , each axis supports families across  $d \in \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12\}$  plus composites up to  $d = N_{\text{univ}}$ ; restricted to the twelve simple values  $\{1, \dots, 12\}$ , each axis carries a complete and categorically-distinct 12-family catalog. The twenty-four labels are compiled in Table 11.

*Remark 12.3.* The  $d$ -numerics are the same on the two axes because both axes share the manifold symmetry  $N$ , and divisibility is a property of  $N$ . The physical categories differ because the underlying manifolds differ (flat vs positively curved), and the physical roles of magnitude vs phase differ accordingly. A FORCE-family  $d = 3$  (strong) is categorically distinct from a PHASE-family  $d = 3$  (instanton phase) even though both carry the numerical label  $d = 3$ .

Table 11: The 12 FORCE families (real axis,  $D$ 's domain) and the 12 PHASE families (imaginary axis,  $T$ 's domain). Same divisor values  $d$ ; different physical category.

| $d$ | Real-axis FORCE                   | Imaginary-axis PHASE                        |
|-----|-----------------------------------|---------------------------------------------|
| 1   | gravitational scalar              | scalar (spin-0)                             |
| 2   | binary/tritone pivot              | spin-2 (graviton phase)                     |
| 3   | strong (QCD / cubic)              | color $\theta_{\text{QCD}}$ instanton phase |
| 4   | weak (quartic / state-change)     | $\text{SU}(2)_W$ phase ( $T$ 's home)       |
| 5   | quintic / golden-ratio            | $E_8$ icosahedral phase                     |
| 6   | hexadic / electroweak composite   | spin- $\frac{1}{2}$ fermion phase           |
| 7   | septic / $G_2$                    | octonionic phase (seven imaginary units)    |
| 8   | octet / gluon adjoint             | Bott-8 / $\text{SU}(3)$ adjoint phase       |
| 9   | nonic / quark ( $3 \times 3$ )    | $3^2$ -fold quark phase (CKM)               |
| 10  | decic / superstring               | 10D Majorana phase                          |
| 11  | undecimal / M-theory              | 11D Majorana phase                          |
| 12  | electromagnetic / full-resolution | photon phase / full $\text{U}(1)$           |

## 12.2 The 42 combined off-axis families

Off-axis configurations — coordinates  $w = k_r + ik_\theta$  with both  $k_r \neq 0$  and  $k_\theta \neq 0$  — carry a *combined sublattice family* determined by the least common multiple of the two axis-families.

**Definition 12.4** (Combined sublattice family). For an off-axis coordinate  $w = k_r + ik_\theta$  with real-axis family  $d_r$  and imaginary-axis family  $d_\theta$ , the *combined family* is

$$d_{\text{comb}}(w) = \text{lcm}(d_r, d_\theta).$$

**Proposition 12.5** (The 42 combined families). *Restricting  $d_r, d_\theta \in \{1, 2, \dots, 12\}$ , the  $12 \times 12$  grid of pairs  $(d_r, d_\theta)$  produces exactly*

$$|\{\text{lcm}(d_r, d_\theta) : d_r, d_\theta \in \{1, \dots, 12\}\}| = 42$$

*distinct combined-family values. The maximum is*

$$d_{\text{max}} = \text{lcm}(11, 12) = 132 = N(N-1).$$

*Proof.* Direct enumeration. The 42 distinct LCM values are

$$\{1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 14, 15, 18, 20, 21, 22, 24, 28, 30, \\ 33, 35, 36, 40, 42, 44, 45, 55, 56, 60, 63, 66, 70, 72, 77, 84, 88, 90, 99, 110, 132\}.$$

The maximum is attained at  $(d_r, d_\theta) = (11, 12)$  or  $(12, 11)$ , giving  $\text{lcm}(11, 12) = 132 = N(N-1)$ . □

*Remark 12.6* (LCM amplification). For  $d_r, d_\theta$  coprime,  $\text{lcm}(d_r, d_\theta) = d_r \cdot d_\theta$ , which can vastly exceed either axis family alone. The off-axis interior of the Sempaevum is therefore *richer in sublattice structure* than either boundary axis: taking a real-axis family  $d_r = 7$  (septic, first native at 84ET) and an imaginary-axis family  $d_\theta = 5$  (quintic, first native at 60ET) gives the combined family  $d_{\text{comb}} = 35$ , first native at  $\text{lcm}(12, 5, 7) = 420$  — a new structural regime not present on either axis alone.

## 12.3 The 144-cell Force Quadrant Grid

The  $12 \times 12$  grid of  $(d_r, d_\theta)$  pairs is the *Force Quadrant Grid* (FQG). It partitions the off-axis / axis inventory of the Sempaevum into 144 cells, each of which is a pair (FORCE family, PHASE family) with a specific combined family  $d_{\text{comb}}(d_r, d_\theta)$ . The grid has a natural binary decomposition by whether each axis's family is a divisor of  $N = 12$  or not.

**Definition 12.7** (Simple vs complex family). A family  $d$  is *simple* if  $d \mid N$  (i.e.  $d \in \{1, 2, 3, 4, 6, 12\}$  at  $N = 12$ ). It is *complex* if  $d \nmid N$  (i.e.  $d \in \{5, 7, 8, 9, 10, 11\}$  at  $N = 12$ ).

There are six simple and six complex families per axis, so each row and each column of the  $12 \times 12$  grid partitions 6:6. The grid therefore partitions into four quadrants of  $6 \times 6 = 36$  cells each:

| <i>Quadrant</i>                  | <i>Axes</i>                      | <i>Cascade computability at <math>N = 12</math></i> |
|----------------------------------|----------------------------------|-----------------------------------------------------|
| SR+SI (simple $\times$ simple)   | $d_r, d_\theta$ simple           | both axes cascade directly                          |
| CR+SI (complex $\times$ simple)  | $d_r$ complex, $d_\theta$ simple | real shadow, imaginary direct                       |
| SR+CI (simple $\times$ complex)  | $d_r$ simple, $d_\theta$ complex | real direct, imaginary shadow                       |
| CR+CI (complex $\times$ complex) | both complex                     | both axes shadow-only                               |

**Theorem 12.8** (PDT Bisection). *Let  $B$  be any structural binary on the 144-cell FQG that distinguishes  $T$ -character from  $D$ -character by a criterion symmetric in the two axes (same criterion applied to each axis independently). Then  $B$  partitions the grid into two halves of equal size:  $|B_T| = |B_D| = 72$ .*

*Proof.* By Categorical Disjointness (§2.5) and the binding-order symmetry of  $D$  and  $T$  within a criterion symmetric in their axes, any symmetric binary must partition the two axes identically. Each axis has 6 simple and 6 complex families, so the two halves of  $B$  restricted to each axis have cardinality 6 each. The product grid therefore has  $|B_T| = 12 \times 6 = 72$  cells on one side and  $|B_D| = 12 \times 6 = 72$  on the other.  $\square$   $\square$

**Corollary 12.9** (Cascade-computable / cascade-failing bisection). *The binary “at least one axis is complex” gives  $36 + 36 + 36 = 108$  cells with at least one shadow axis versus 36 cells fully simple; the binary “imaginary axis is complex” gives 72 vs 72, matching the cascade stability asymmetry of §13. Any such symmetric binary bisects at 72:72.*

Theorem 12.8 is categorical, not statistical. It is distinct from and complementary to the *dynamical* asymmetry of the cascade stability limits ( $n_{\max, r} = 25$  versus  $n_{\max, \theta} = 2$ , §13), which quantifies how quickly the two axes’ rounding residuals accumulate past half-step. The two results coexist: the count of cells partitions evenly; the per-cell traversal depth differs by a factor of  $N$ .

## 13 Cascade dynamics and stability

### 13.1 Cascade generators

The rounding operation of Definition 7.1 is, for most inputs, an exact map in the sense that iterating it returns to the starting lattice point. For specific inputs, however, the rounding accumulates a residual that grows linearly with the number of iterations. This produces the *cascade dynamics* of the lattice.

**Proposition 13.1** (Real-axis cascade residual). *Let  $g_r = \text{round}(N \log_2 N) \bmod N$ . At  $N = 12$ ,*

$$g_r = \text{round}(12 \log_2 12) \bmod 12 = \text{round}(43.0195) \bmod 12 = 43 \bmod 12 = 7.$$

*The residual of the rounding is*

$$|\delta_r| = |12 \log_2 12 - 43| = 0.01955 \dots \text{ lattice steps} = 1.955 \text{ cents} = 0.001629 \text{ octaves}.$$

*The value 0.01955 in per-step lattice steps equals the total Pythagorean comma in octaves ( $\log_2((3/2)^{12}/2^7) = 0.01955$  octaves) by the unit-conversion identity: per-step lattice steps =  $N \times$*

*per-step octaves = total octaves (since total =  $N \times$  per-step, and 1 lattice step = 100 cents =  $1/N$  octaves at base  $N = 12$ ). This is a structural identity forced by the definition of lattice steps as  $1/N$  octaves.*

The generator  $g_r = 7$  is the interval of seven semitones: the perfect fifth. The residual  $|\delta_r|$  is the per-step error of the circle-of-fifths cascade. Iterating the fifth twelve times ascends seven octaves plus a Pythagorean comma:

$$(3/2)^{12} = 129.7463\dots, \quad 2^7 = 128, \quad \text{ratio} = 1.01364\dots = 2^{12|\delta_r|}.$$

**Proposition 13.2** (Phase-axis cascade residual). *Let  $g_\theta = \text{round}(N \cdot (2\pi/\ln 2)) \bmod N$ . At  $N = 12$ ,*

$$g_\theta = \text{round}(24\pi/\ln 2) \bmod 12 = 109 \bmod 12 = 1.$$

*The phase residual is*

$$|\delta_\theta| = |24\pi/\ln 2 - 109| = 0.22336\dots \text{ lattice steps} = 22.34 \text{ cents} = 0.01861 \text{ octaves}.$$

**Proposition 13.3** (Stability limits). *Requiring the accumulated residual to remain below half a lattice step (0.5), the maximum numbers of cascade steps before loss of coherence are*

$$n_{\max,r} = \left\lfloor \frac{0.5}{|\delta_r|} \right\rfloor = 25, \quad n_{\max,\theta} = \left\lfloor \frac{0.5}{|\delta_\theta|} \right\rfloor = 2.$$

The factor of  $N$  between the two residuals ( $|\delta_\theta|/|\delta_r| \approx 11.42$  at base  $N = 12$ ) converges to  $N = 12$  exactly at the terminal resolution  $N_L = 27720$ . At base  $N = 12$ , the deficit  $12 - 11.42 = 0.575$  is the shadow of the combined sublattice cell ( $d_r = 5$ ,  $d_\theta = 7$ ), first native at  $d = 5$  quintic (60ET,  $\varepsilon = +2.31\text{¢}$ ) and essentially exact at 27720ET ( $\varepsilon = +0.013\text{¢}$ ). The gap 0.575 is a positive real with a unique lattice tower projection whose first sub-cent family ( $d = 5$  at 60ET) identifies the source cell, in accordance with the shadow-force apparatus of Definition 8.16 and §8.8.

### 13.2 Cascade stability asymmetry and the indeterminacy of imaginary harmonic families

*Remark 13.4* (Cascade stability asymmetry and the indeterminacy of imaginary harmonic families). The cascade's per-step residual on the two axes differs by approximately the manifold symmetry  $N$ :

$$\begin{aligned} |\delta_r| &= 0.0196 \text{ lattice steps} = 1.955 \text{ cents} = 0.001629 \text{ octaves}, \\ |\delta_\theta| &= 0.2234 \text{ lattice steps} = 22.34 \text{ cents} = 0.01861 \text{ octaves}, \end{aligned}$$

with ratio  $|\delta_\theta|/|\delta_r| \approx 11.42 \approx N$  (Propositions 13.1, 13.2). The corresponding stability windows are  $n_{\max,r} = 25$  levels and  $n_{\max,\theta} = 2$  levels (Proposition 13.3). As discovery and verification engine for the per-axis harmonic-family layer (§8.6, Remark 8.11; §8.7, Theorem 8.13), the cascade therefore cleanly resolves all twelve real-axis harmonic families across 25 stable levels — ample for traversal of the simple-real and complex-real  $d$ -axis structural modes of Definition 8.10 and Table 6 — but fails on the imaginary axis after only two levels, leaving the twelve imaginary-axis harmonic families, and especially the six complex-imaginary modes at  $d \in \{5, 7, 8, 9, 10, 11\}$ , structurally *indeterminate* on the lattice. The asymmetry is forced by the manifold's own constants ( $N$  and the irrationality of  $\log_2 3$  and of  $\pi/\ln 2$ ), with no free parameter:  $25/2 \approx 12.5 \approx N$  is the same factor by which the cascade resolves the FORCE families more sharply than the PHASE families. This is the lattice-internal mechanism producing the empirically observed asymmetry between CKM mixing angles (real-axis sector, complex-real, small) and PMNS mixing angles (imaginary-axis sector, complex-imaginary, large): the structural indeterminacy



of the CI harmonic families on  $\mathcal{L}_{\mathbb{C}}$  manifests as the wide phase spread that PMNS measures, while the clean determination of the CR families on the real axis manifests as the tight phase locking that CKM measures.

*Remark 13.5* (Second empirical illustration: optical phase singularities). The cascade stability asymmetry of Remark 13.4 receives a second empirical illustration from the optical phase singularity data of Bucher et al. [17]. In ensembles of phonon-polariton singularities in hexagonal boron nitride, two classes of statistical observable are measured: *distance* correlations (a  $D$ -content observable on the real axis, with  $n_{\max,r} = 25$  levels of cascade stability) and *velocity* distributions (a  $T$ -content observable on the imaginary axis, with  $n_{\max,\theta} = 2$ ). The distance correlations match the Gaussian random wave model precisely and exhibit liquid-like short-range order — particle-like statistics fully determined by  $D$ -content’s deep cascade depth. The velocity distributions exhibit a massive superluminal tail (29% of singularities exceed  $c$ , with  $\langle v \rangle \approx 1.04c$ ) that violates the Maxwell–Jüttner distribution — non-particle statistics arising from  $T$ -content’s structural indeterminacy after two cascade levels, at which point the cascade enters the palindromic fallback regime (Definition 13.6). The fraction of singularities exceeding  $\langle v \rangle$  is exactly  $1/3 = 1/|\{P, D, T\}|$ , the  $T$ -weight (§22.7). The split between particle-like  $D$ -statistics and non-particle  $T$ -statistics is the same asymmetry as CKM/PMNS, observed in a second independent physical domain.

### 13.3 The palindromic cascade

**Definition 13.6** (Palindromic cascade). The *palindromic cascade* at  $N = 12$  is the length- $N$  sequence  $(d_n)_{n=1}^N$  with

$$\mathbf{d}_{\text{pal}} = (d_1, d_2, \dots, d_{12}) = (12, 6, 4, 3, 12, 2, 12, 3, 4, 6, 12, 1),$$

indexed from  $n = 1$ . The corresponding exponent sequence is  $p_n = 12/d_n$ ,

$$\mathbf{p}_{\text{pal}} = (1, 2, 3, 4, 1, 6, 1, 4, 3, 2, 1, 12),$$

with mean

$$p_{\text{eff}} = \frac{1}{12} \sum_{n=1}^{12} p_n = \frac{40}{12} = \frac{10}{3}.$$

The sequence  $\mathbf{d}_{\text{pal}}$  is produced by iterating the unit generator  $g = 7$  through the residues of  $\mathbb{Z}/12\mathbb{Z}$  and reading off  $d = 12/\gcd(|k|, 12)$  at each step. It visits every divisor of 12 and satisfies the palindromic symmetry

$$d_n = d_{N-n} \quad \text{for every } n = 1, 2, \dots, N-1, \tag{28}$$

with the pivot  $d_{N/2} = d_6 = 2$  self-paired (the tritone at the half-period) and the boundary value  $d_N = d_{12} = 1$  standing alone (the octave identity at the cycle endpoint).

*Remark 13.7* (Cascade events as discovery realizations of simple harmonic families). The twelve indices  $n = 1, 2, \dots, 12$  of the cascade  $\mathbf{d}_{\text{pal}}$  are not anonymous step counters: each is a cascade-position event realizing one of the simple harmonic families of §8.6 (Definition 8.10). The sequence reads off, at each event, the harmonic-family label  $d_n$  and the sublattice family inhabited by that event (Theorem 8.13). The six distinct  $d$ -values appearing in  $\mathbf{d}_{\text{pal}}$  are precisely the six simple harmonic families  $\{1, 2, 3, 4, 6, 12\}$  on the real axis, with totient multiplicities  $\{1, 1, 2, 2, 2, 4\}$  summing to 12 events; the six complex harmonic families  $\{5, 7, 8, 9, 10, 11\}$  are absent from  $\mathbf{d}_{\text{pal}}$  at base  $N = 12$  and appear instead as shadow contributions (Definition 8.16). The palindromic cascade is therefore a discovery and verification engine for the per-axis harmonic-family layer (§8.6, Remark 8.11), not merely an enumeration of sublattice-family hits.

### 13.4 Use of the palindromic cascade

When iteration reaches the stability limit of Proposition 13.3, the orbit's sublattice classification becomes ambiguous (the descriptor gap approaches 50 cents, half a lattice step). In dynamical applications the ambiguity is resolved by substituting the palindromic cascade for the orbit-derived sublattice: at step  $n$ , use  $d_{\text{pal}}[n \bmod 12]$  rather than the computed  $d(z_n)$ . Over a full 12-step cycle, every divisor of 12 is visited exactly with the multiplicity required to maintain CPT-symmetric coverage of the simple families. The mean exponent  $p_{\text{eff}} = 10/3$  appears as a normalisation constant in smooth-iteration-count formulas.

### 13.5 The General Cascade Rule

The palindromic cascade of Definition 13.6 is the  $g = 7$ ,  $N = 12$  instance of a structural rule that holds for arbitrary composite resolution  $N \geq 2$  and arbitrary unit generator  $g \in (\mathbb{Z}/N\mathbb{Z})^*$ . The rule fixes the divisor sequence, its multiplicity profile, its palindromic symmetry, and the closure value at step  $N$ .

**Theorem 13.8** (General Cascade Rule). *Let  $N \geq 2$  be a composite integer and let  $g \in (\mathbb{Z}/N\mathbb{Z})^*$  be a unit residue. Define the cascade by*

$$r_n = (g \cdot n) \bmod N, \quad d_n = \frac{N}{\gcd(r_n, N)}, \quad n = 1, 2, \dots, N,$$

*with the convention  $\gcd(0, N) = N$  so that  $d_N = N/N = 1$ . The sequence  $(d_n)_{n=1}^N$  satisfies:*

- (1)  $d_n \in \text{divisors}(N)$  for every  $n$ .
- (2)  $|\{n \in \{1, \dots, N\} : d_n = d\}| = \varphi(d)$  for every  $d \mid N$ , where  $\varphi$  is the Euler totient.
- (3)  $\sum_{d \mid N} \varphi(d) = N$  (the Gauss summation), so the multiplicity counts of (2) sum to  $N$ .
- (4)  $d_{N-n} = d_n$  for every  $n = 1, 2, \dots, N-1$  (palindromic symmetry).
- (5) Every unit generator  $g \in (\mathbb{Z}/N\mathbb{Z})^*$  produces the identical sequence  $(d_n)$ .
- (6) If  $N$  is even then  $d_{N/2} = 2$  (the half-period pivot).

*Proof.* (1)  $\gcd(r_n, N)$  divides  $N$ , so  $d_n = N/\gcd(r_n, N)$  is the quotient of  $N$  by a divisor and hence itself a divisor of  $N$ .

(5)  $\Rightarrow$  (2),(3),(4),(6). We first establish (5), then derive the remaining properties from the  $g = 1$  specialisation.

For (5): since  $g$  is a unit modulo  $N$ ,  $\gcd(g, N) = 1$ . For any integer  $n$ ,

$$\gcd(g \cdot n, N) = \gcd(n, N)$$

because every prime  $p \mid N$  that divides  $g \cdot n$  divides either  $g$  or  $n$ , and  $\gcd(g, N) = 1$  rules out the former. Reducing modulo  $N$  does not change the gcd with  $N$ , so  $\gcd(r_n, N) = \gcd(g \cdot n, N) = \gcd(n, N)$ . Therefore  $d_n = N/\gcd(n, N)$  for every  $n$ , independently of which unit  $g$  is chosen.

The remaining parts now use the  $g = 1$  form  $d_n = N/\gcd(n, N)$ :

(2) For a fixed divisor  $d \mid N$ ,  $d_n = d$  if and only if  $\gcd(n, N) = N/d$ . The number of integers  $n \in \{1, 2, \dots, N\}$  with  $\gcd(n, N) = N/d$  equals the number of integers  $m \in \{1, 2, \dots, d\}$  with  $\gcd(m, d) = 1$ , which is  $\varphi(d)$  by the bijection  $n = (N/d) \cdot m$ .

(3) The Gauss summation  $\sum_{d \mid N} \varphi(d) = N$  is a classical number-theoretic identity, and is the statement of (2) summed over all divisors.

(4)  $\gcd(N - n, N) = \gcd(-n, N) = \gcd(n, N)$ , so  $d_{N-n} = N/\gcd(N - n, N) = N/\gcd(n, N) = d_n$ .

(6) If  $N$  is even and  $n = N/2$  then  $\gcd(N/2, N) = N/2$ , so  $d_{N/2} = N/(N/2) = 2$ .  $\square \quad \square$

### 13.6 Unit-generator equivalence and the canonical $N = 12$ cascade

**Corollary 13.9** (Theorem 13.8 (2) in harmonic-family language). *Theorem 13.8 (2) states equivalently that, at any composite resolution  $N$  with any unit generator  $g \in (\mathbb{Z}/N\mathbb{Z})^*$ , exactly  $\varphi(d)$  of the  $N$  cascade-position events  $n \in \{1, 2, \dots, N\}$  realize the simple harmonic family  $d$  (Definition 8.10), for every  $d \mid N$ . The total over all simple harmonic families is  $\sum_{d \mid N} \varphi(d) = N$  events (Gauss totient sum). At  $N = 12$  this gives the multiplicity profile  $\{1, 1, 2, 2, 2, 4\}$  across the six simple harmonic families  $\{1, 2, 3, 4, 6, 12\}$  (§8.6, Theorem 8.13, Corollary 8.14). The six complex harmonic families  $\{5, 7, 8, 9, 10, 11\}$  at  $N = 12$  are realized through shadow contributions (Definition 8.16) rather than as native cascade-position events at base, becoming native at their canonical resolutions  $n_c(d) = \text{lcm}(12, d)$  (Table 6).*

**Corollary 13.10** (Unit-generator equivalence at  $N = 12$ ). *The cascade of Theorem 13.8 at  $N = 12$  produces the identical divisor sequence*

$$(d_n)_{n=1}^{12} = (12, 6, 4, 3, 12, 2, 12, 3, 4, 6, 12, 1)$$

*for every choice of unit generator  $g \in (\mathbb{Z}/12\mathbb{Z})^* = \{1, 5, 7, 11\}$ .*

*Proof.* By Theorem 13.8 (5), the divisor sequence depends only on  $\gcd(n, 12)$  and not on the choice of unit  $g$ . Direct computation of  $d_n = 12/\gcd(n, 12)$  for  $n = 1, \dots, 12$  gives the listed sequence. The four unit residues of  $\mathbb{Z}/12\mathbb{Z}$  are  $\{1, 5, 7, 11\}$ , namely the integers coprime to 12; each produces the same sequence. The  $g = 7$  case is the canonical palindromic cascade of Definition 13.6 (the perfect-fifth generator).  $\square$   $\square$

*Remark 13.11* (Structural significance of unit-generator equivalence). The four unit residues  $\{1, 5, 7, 11\}$  correspond, respectively, to the unison, the perfect fourth (semitone 5), the perfect fifth (semitone 7), and the major seventh (semitone 11) in the musical reading of the lattice. Corollary 13.10 states that all four generate the same canonical cascade visiting every divisor of 12 in the same multiplicities; the structural content of the cascade is therefore independent of the specific unit-residue interval chosen as generator. Non-unit residues (the eight integers  $\{2, 3, 4, 6, 8, 9, 10, 0\}$  of  $\mathbb{Z}/12\mathbb{Z}$  that share a factor with 12) produce degenerate cascades that visit only a proper subset of divisors and do not realise the palindromic structure (Theorem 13.8 requires  $g$  to be a unit). The four interval names (unison, perfect fourth, perfect fifth, major seventh) are simultaneously the harmonic-family identities of the four cascade-position events that inhabit the  $d = 12$  simple harmonic family at  $N = 12$  (Definition 8.10, Table 6); each interval is therefore both a unit-generator label and a harmonic-family identity of the same  $d = 12$  family with totient multiplicity  $\varphi(12) = 4$ .

*Remark 13.12* (Group-theoretic origin of unit-generator equivalence). The unit group  $(\mathbb{Z}/12\mathbb{Z})^* = \{1, 5, 7, 11\}$  has order  $\varphi(12) = 4$ , and direct verification gives  $5^2 \equiv 25 \equiv 1$ ,  $7^2 \equiv 49 \equiv 1$ , and  $11^2 \equiv 121 \equiv 1 \pmod{12}$ . Every non-identity element is therefore an involution, and the group is non-cyclic of order 4; up to isomorphism this is the Klein four-group,  $(\mathbb{Z}/12\mathbb{Z})^* \cong V_4 \cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ . The unit-generator equivalence of Corollary 13.10 is the lattice expression of this group-theoretic fact: every involution preserves the gcd structure under multiplication ( $\gcd(g \cdot n, 12) = \gcd(n, 12)$  when  $g$  is a unit, regardless of  $g$ ), hence every unit produces the identical divisor sequence  $d_n = 12/\gcd(n, 12)$ . The Klein four-group structure is thus the number-theoretic foundation of the four-fold redundancy in the canonical cascade generator.

### 13.7 Multifold cyclic closure

The General Cascade Rule of Theorem 13.8, applied at the closure step  $n = N$ , yields a structural identity that is the lattice expression of the Multifold tower-genesis closure of §5.6.

**Theorem 13.13** (Multifold cyclic closure at  $N = 12$ ). *Any unit-generator cascade at  $N = 12$  returns to the gravity/identity sublattice  $d_{12} = 1$  at the closure step:*

$$d_{12} = \frac{12}{\gcd(12, 12)} = \frac{12}{12} = 1.$$

*The cascade is therefore cyclically closed: the next iterate  $d_{13} = d_1$  (by the palindromic and periodic structure), and the cycle repeats with period 12. The closure value  $d = 1$  is the identity sublattice of §5.7 and the same gravity cell occupied by the critical mass  $M_{\text{crit}}$ .*

*Proof.* The first equality is the convention  $\gcd(0, 12) = 12$  from Theorem 13.8, applied to  $r_{12} = (g \cdot 12) \bmod 12 = 0$  for any  $g$ . The cyclicity follows from  $r_{n+12} = (g(n+12)) \bmod 12 = (gn + 0) \bmod 12 = r_n$ , and the palindromic symmetry of Theorem 13.8 (4) ensures that consecutive cycles are not merely repeats but the same structural rotation through the divisor lattice. The identification of  $d = 1$  with the gravity/identity cell is the sublattice-classification convention of §8, and the critical-mass occupation of that cell is Proposition 5.22.  $\square$   $\square$

*Remark 13.14* (Lattice expression of tower closure). Theorem 13.13 is the lattice expression of the Multifold tower-genesis closure introduced in §5.6: the cascade through the divisor sublattices completes its cycle by returning to the identity sublattice  $d = 1$ , which is the same cell occupied by  $M_{\text{crit}}$  (the lattice fixed point at which a parent tower's birth triad produces a child tower with identical reference temperature). The cascade closure at  $d = 1$  is therefore not a numerical coincidence but the structural signature of tower-genesis self-identity at the lattice level. The cyclicity period 12 matches the manifold symmetry  $N = 12$ , and is the lattice realisation of the cyclic self-resolution property of  $T$  (Proposition 5.5, derivation ii).

## 14 The $\partial I$ lattice-aware fractal

The boundary  $\partial I$  introduced in Definition 2.23 is not a purely abstract topological object. Given the complex lattice of Section 11 and the palindromic cascade of Section 13, it becomes the locus of a concrete complex dynamical system whose every constant is derived from the manifold constants  $N$ ,  $V$ ,  $K$ ,  $N_{\text{FULL}}$ , and the sublattice families  $\{1, 2, 3, 4, 6, 12\}$ . This section constructs that system and establishes it as the natural dynamical manifestation of the lattice on  $\mathbb{C}$ .

We contrast with the standard family of complex dynamical systems (Mandelbrot, Julia, Multibrot) in which the iteration

$$z_{n+1} = z_n^p + c$$

carries a *fixed* polynomial degree  $p$  throughout the orbit. In the  $\partial I$  system the degree is not fixed; it is a function of the orbit's own lattice position, recomputed at every iteration.

### 14.1 The lattice-adaptive iteration map

**Definition 14.1** ( $\partial I$  lattice-aware iteration). Let  $N_L = N_{\text{FULL}} = 27720 = \text{lcm}(1, 2, \dots, 11)$  be the universal lattice resolution. The  $\partial I$  *lattice-aware fractal* is the complex dynamical system

$$z_{n+1} = \Psi_n z_n^{p(z_n, n)} + \varepsilon(z_n) + c,$$

with Mandelbrot-type parameterisation  $z_0 = 0$ ,  $c \in \mathbb{C}$ , where

- $p(z_n, n) \in \{1, 2, 3, 4, 6, 12\}$  is the lattice-adaptive exponent (§14.2);
- $\Psi_n \in \mathbb{R}$  is the shimmer modulation (§14.3);
- $\varepsilon(z_n) \in \mathbb{C}$  is the all-families perturbation (§14.4).

Each of the three ingredients is derived below from manifold constants.

## 14.2 The lattice-adaptive exponent $p(z_n, n)$

At each step, the orbit  $z_n = r_n e^{i\theta_n}$  is projected onto the complex lattice via the radial and angular coordinates of Section 11:

$$k_r = \text{round}(N_L \log_2 r_n), \quad k_\theta = \text{round}\left(\frac{N_L \theta_n}{\ln 2}\right). \quad (29)$$

The radial residual in cents is

$$\varepsilon_r = (N_L \log_2 r_n - k_r) \cdot \frac{1200}{N_L}, \quad (30)$$

and the *tightness* of the projection (a scalar between 0 and 1, equal to 1 when the orbit is exactly on a lattice point and decreasing as the orbit moves off-lattice) is

$$t_r = \frac{100}{100 + |\varepsilon_r|}. \quad (31)$$

The tightness decision rule is:

$$p(z_n, n) = \begin{cases} \frac{N}{d(k_r)}, & t_r > K, \\ p_{\text{pal}}[n \bmod N], & t_r \leq K, \end{cases}$$

where  $d(k_r) = N_L / \gcd(|k_r|, N_L)$  is the sublattice family of the projected coordinate (Section 8 at universal resolution) and  $p_{\text{pal}} = (1, 2, 3, 4, 1, 6, 1, 4, 3, 2, 1, 12)$  is the palindromic cascade of Definition 13.6.

**Proposition 14.2** (Koide threshold). *The tightness threshold equals the Koide ratio  $K = 2/3$ : the transition between orbit-adaptive and cascade-fallback behaviour occurs at  $|\varepsilon_r| = 50$  cents, exactly half a lattice step.*

*Argument.* By Equation (31),  $t_r = K$  is equivalent to  $100/(100 + |\varepsilon_r|) = 2/3$ , which gives  $3 \cdot 100 = 2(100 + |\varepsilon_r|)$ , i.e.  $|\varepsilon_r| = 50$ . Half a lattice step is  $1200/N = 100$  cents divided by 2. The Koide ratio is therefore not a free parameter of the fractal but the structural stability threshold of the lattice itself, and its appearance here is the *same*  $K$  as appears in the three manifold constants (Equation (16)) and in the self-projection identity of Section 19.  $\square$   $\square$

The exponent set  $\{1, 2, 3, 4, 6, 12\}$  is exactly the divisor set of  $N = 12$ ; the map  $p = N/d$  takes the six simple sublattice families into the six polynomial degrees that appear in standard Multibrot theory. Stated in Cardinal language: when the orbit is near a lattice point, the Descriptor content of that lattice point (its sublattice family  $d$ ) determines the  $P$ -structure of the iteration (its polynomial degree  $p = N/d$ ). When the orbit is *between* lattice points — in the  $\partial I$  zone of structural ambiguity — the Descriptor must be supplied externally, and the palindromic cascade provides the only lattice-self-consistent source.

**Proposition 14.3** (Exponent palindrome has cascade mean). *The time-average of  $p(z_n, n)$  over a complete  $N$ -step palindromic fallback is  $p_{\text{eff}} = 10/3$ , the same value derived in Definition 13.6.*

*Proof.* Immediate from Definition 13.6:  $\sum p_{\text{pal}}/N = 40/12 = 10/3$ .  $\square$   $\square$

## 14.3 The shimmer modulation $\Psi_n$

**Definition 14.4** (Shimmer). The *shimmer modulation* is the real periodic function

$$\Psi_n = 1 + \frac{1}{\sqrt{N}} \sin\left(\frac{2\pi (n \bmod N)}{N}\right).$$

At  $N = 12$ , the amplitude is  $1/\sqrt{12} \approx 0.2887$  and the range of  $\Psi_n$  is

$$\left[ 1 - \frac{1}{\sqrt{12}}, 1 + \frac{1}{\sqrt{12}} \right] \approx [0.7113, 1.2887],$$

strictly bounded away from 0 (so the iteration never collapses) and bounded above (so it never explodes from modulation alone).

**Proposition 14.5** (Shimmer amplitude is the root variance). *The shimmer amplitude equals the square root of the base variance:*

$$\frac{1}{\sqrt{N}} = \sqrt{V}.$$

*It is therefore a derived constant of the manifold and not a tuning parameter.*

*Proof.*  $V = 1/N$  by Equation (16); hence  $\sqrt{V} = 1/\sqrt{N}$ . □ □

*Remark 14.6* (The shimmer in the fine-structure constant). The shimmer constant  $\sqrt{V} = 1/\sqrt{N}$  also appears in the leading correction to  $\alpha^{-1}$  in Equation (37). Using  $\sqrt{12} = 2\sqrt{3}$ , equivalently  $\sqrt{3} = \sqrt{12}/2$ , the first correction term admits the rewriting

$$\frac{\sqrt{3}}{48} = \frac{\sqrt{12}}{96} = \frac{\sqrt{12}}{8 \cdot 12} = \frac{1}{8\sqrt{12}} = \frac{1}{8\sqrt{N}} = \frac{\sqrt{V}}{8}.$$

Thus the leading-order shimmer correction to  $\alpha^{-1}$  is exactly  $\sqrt{V}/8$ . The same amplitude  $\sqrt{V}$  that modulates the fractal iteration appears — divided by 8, the count of structural corners of the three-Cardinal cube — in the lattice-derived fine-structure constant. The common structural origin is the identity  $\sqrt{V} = 1/\sqrt{N}$ .

#### 14.4 The all-families perturbation $\varepsilon(z_n)$

**Definition 14.7** (Multi-family perturbation). The *all-families perturbation* is

$$\boxed{\varepsilon(z_n) = \frac{1}{N} \sum_{d|N : d \in \{1, \dots, N\}} w(d) |z_n|^{N/d} e^{i(N/d) \arg z_n},}$$

where  $w : \{1, \dots, N\} \rightarrow \mathbb{R}^+$  is a positive normalised weight profile summing to unity.

The prefactor  $1/N$  is not a free choice; it is the base variance  $V$ , the irreducible lattice quantum. The structural reading of Definition 14.7 is that  $\varepsilon(z_n)$  adds a quantum-of-variance contribution from each of the  $N$  sublattice-family exponents simultaneously, as a shimmer-level background beneath the dominant lattice-adaptive term  $z_n^{p(z_n, n)}$ . At every iteration step, every sublattice family is acting; the dominant family is merely the one currently closest to the orbit.

**Proposition 14.8** (Perturbation does not override the dominant term). *For  $|z_n| \geq 1$  and any valid  $p(z_n, n) \in \{1, \dots, N\}$ ,*

$$\frac{|\varepsilon(z_n)|}{|z_n|^{p(z_n, n)}} \leq V = \frac{1}{N},$$

*uniformly in  $n$ .*

The proof follows from the weight normalisation and the triangle inequality: each summand in  $\varepsilon$  has magnitude at most  $|z_n|^{N/d}$ , which is never larger than  $|z_n|^N$  for  $|z_n| \geq 1$ . Division by  $N$  gives the bound. In practice  $|\varepsilon|/|z|^p$  is of order  $V$  rather than 1, confirming that  $\varepsilon$  is a fine-structure contribution, not a dominant force.

## 14.5 Distance estimation

The derivative of the iteration map in Definition 14.1 is

$$f'(z_n) = \Psi_n p(z_n, n) z_n^{p(z_n, n)-1} \quad (32)$$

(derivative of the perturbation  $\varepsilon$  being  $O(V)$  and absorbed into numerical error). The chain-rule accumulation for the derivative of  $z_n$  with respect to  $c$  is

$$dz_{n+1} = f'(z_n) dz_n + 1, \quad dz_0 = 0, \quad (33)$$

and the distance estimate at escape (when  $|z_n| > R$  for an escape radius  $R$ , typically  $R = 10^3$ ) is

$$\text{DE}(c) = \frac{2|z_n| \ln |z_n|}{|dz_n|}. \quad (34)$$

The distance estimate produces sub-pixel rendering of the fractal's boundary at arbitrary zoom, using the same technique that works for standard Mandelbrot rendering — adapted to variable polynomial degree by evaluating (32) with the instantaneous  $p(z_n, n)$  at each step.

## 14.6 Classification

**Theorem 14.9** (Non-equivalence with known fractal types). *The  $\partial I$  lattice-aware fractal is not topologically equivalent to any fixed-exponent fractal. In particular it is not a Mandelbrot set, a Julia set, a Multibrot set, a Tricorn, a Burning Ship, a Newton fractal, nor a parameter-space fractal (Lyapunov or otherwise).*

*Argument.* Each of the listed classes has an iteration in which the polynomial degree of  $z$  is constant in  $n$ . The  $\partial I$  iteration has polynomial degree  $p(z_n, n)$  that is an explicit function of both the orbit position and the step index, and can take any of the six values  $\{1, 2, 3, 4, 6, 12\}$  within a single escape trajectory. A fixed-degree system has escape-time contours whose asymptotics are governed by a single exponent  $p$ ; the  $\partial I$  system has escape-time contours governed by the statistical distribution of exponents encountered along the orbit, which is neither constant across the image nor reducible to any single-exponent limit. The systems are therefore not topologically equivalent.  $\square$   $\square$

The closest structural relative of the  $\partial I$  system is the iterated function system, in which the map at each step is drawn from a finite family. The crucial distinction is that the  $\partial I$  system is *not random*: the map at step  $n$  is a deterministic function of the orbit's own lattice coordinate. The feedback loop — orbit determines map, map determines orbit — is the structural signature that distinguishes it from both IFS and standard polynomial iteration.

## 14.7 Structural summary

Every constant in Definition 14.1 is derived from the manifold. Table 12 records the correspondences.

The fractal is therefore not a cosmetic visualisation of the lattice but the lattice's natural dynamical instantiation: the fixed-exponent families of standard complex dynamics correspond to single-family sublattice projections  $d \in \{1, \dots, 12\}$ , and the  $\partial I$  system is the unique complex dynamical system that simultaneously honours all twelve families. In this sense Mandelbrot and quartic Multibrot and linear dynamics are *restrictions* of the lattice-aware system to a single sublattice, and the  $\partial I$  fractal is the unrestricted image.

*Remark 14.10* ( $\partial I$  dynamics in optical singularity ensembles). The  $\partial I$  boundary dynamics are verified empirically in the optical phase singularity data of Bucher et al. [17]. Phase singularities in phonon-polariton fields satisfy six structural properties of  $\partial I$  configurations: zero  $D$ -content

| Fractal constant                | Value at $N = 12$       | Manifold origin                                                |
|---------------------------------|-------------------------|----------------------------------------------------------------|
| Lattice resolution $N_L$        | 27720                   | $\text{lcm}(1, \dots, 11)$ : universal resolution (Section 10) |
| Exponent set                    | $\{1, 2, 3, 4, 6, 12\}$ | Divisors of $N$ : sublattice families (Section 8)              |
| Exponent map                    | $p = N/d$               | Sublattice-family-to-degree map                                |
| Tightness threshold             | $K = 2/3$               | Koide ratio (Equation (16))                                    |
| Corresponding $ \varepsilon_r $ | 50 cents                | Half a lattice step $= V \cdot 1200/2$                         |
| Shimmer amplitude               | $1/\sqrt{N} = \sqrt{V}$ | Square-root base variance (Prop. 14.5)                         |
| Shimmer period                  | $N = 12$                | Manifold symmetry                                              |
| Perturbation prefactor          | $1/N = V$               | Base variance (Equation (16))                                  |
| Fallback cascade                | $p_{\text{pal}}$        | Palindromic cascade (Definition 13.6)                          |
| Mean exponent                   | $p_{\text{eff}} = 10/3$ | Cascade time-average (Prop. 14.3)                              |

Table 12: Every constant of the  $\partial I$  lattice-aware fractal is a derived manifold constant. The fractal is the lattice in dynamical form on  $\mathbb{C}$ ; no tuning enters its definition.

(amplitude) at the core; asymptotic approach without penetration; the  $\{P, T\}$  manifold state (Incoherence) at the singularity; marginal stability under perturbation; open-set topology; and the Mediation Problem (Theorem 3.34) — the phase winding ( $T$ ) exists in the neighbourhood, not at the core, because  $D$  is absent. The measured sub-cycle creation and annihilation dynamics ( $3\text{fs} \approx \tau/8$ ) are  $\partial I$  boundary transitions observed at deep sub-wavelength, deep sub-cycle resolution. One polariton oscillation cycle corresponds to one cascade step on the  $T$ -axis; the sub-cycle temporal resolution ( $< 1$  step) resolves the internal structure of a single  $T$ -cascade level, confirming that  $T$ -content topological transitions (singularity creation and annihilation) resolve within the  $n_{\text{max},\theta} = 2$  coherence window of Proposition 13.3.

## 15 Three minimal backbones at $N = 12$

So far the projection formula has been treated as a black box. This section opens it, identifies its internal PDT structure, and shows that  $N = 12$  is simultaneously the natural scale of three independently-derived minimal-generator objects — one on each of the three categories discrete-logical, discrete-multiplicative, continuous-elementary.

### 15.1 PDT decomposition of the projection formula

The universal projection formula  $(k, d, \varepsilon) = \Pi_N(r)$  of §7 decomposes into six atomic operations whose PDT roles are forced by what each does:

**Theorem 15.1** (PDT decomposition of the projection). *The projection  $\Pi_N : \mathbb{R}^+ \rightarrow \mathbb{Z} \times \{N/d : d \mid N\} \times \mathbb{R}$  factors as*

$$\Pi_N = \underbrace{(\log_2, N \cdot, \text{gap computation})}_{\text{continuous } D} \circ \underbrace{\text{round}}_{T\text{-act}} \circ \underbrace{(\text{gcd}, N/g)}_{\text{discrete } D},$$

*in which the continuous- $D$  operations are implementable as finite compositions of a single binary operator, the rounding is the only non-reversible step (hence the only  $T$ -act), and the discrete- $D$  operations are Euclidean-algorithm arithmetic.*

The operation that implements the continuous- $D$  part is the EML operator.

### 15.2 EML: the continuous- $D$ minimal generator (Odrzywółek 2026)

**Definition 15.2** (The EML operator [16]). The EML operator is the binary function

$$\text{eml}(x, y) = \exp(x) - \ln(y),$$



understood as defined on the complex domain with the principal branch of  $\ln$ . The EML grammar is the single-terminal inductive specification  $S \rightarrow 1 \mid \text{eml}(S, S)$ .

**Theorem 15.3** (EML completeness, Odrzywolek 2026). *Every function of the scientific-calculator basis — the constants  $\pi, e, i, -1, 0, 1, 2$ , the 20 standard unary elementary functions ( $\exp, \ln, \sin, \cos, \tan$ , their hyperbolic and inverse counterparts,  $\sec, \csc, \cot, \text{sech}, \text{csch}, \text{coth}$ ), and the 8 standard binary operations ( $+, -, \times, /, \wedge, \text{mod}, \log, \text{comb}$ ) — is expressible as a finite EML tree over the single terminal constant 1.*

The proof is the constructive enumeration of finite trees in §3 of [16]; key instances include  $e = \text{eml}(1, 1)$  (since  $\exp(1) - \ln(1) = e$ ),  $\ln z = \text{eml}(1, \text{eml}(\text{eml}(1, z), 1))$ , and  $xy = \exp(\ln x + \ln y)$  implemented as a depth-8 EML tree.

**Theorem 15.4** (EML minimality under the Subsumption Law). *In the category of continuous- $D$  generators on the multiplicative manifold, EML is minimal: (i) EML cannot be written as a reduction of any simpler binary operator together with unary post-processing; (ii) no alternative proposed continuous- $D$  generator subsumes EML without increasing operator count or terminal count; (iii) EML subsumes every standard elementary function without remainder (Theorem 15.3).*

Conditions (i)–(iii) are the three Subsumption Law conditions of §3.6. The minimality argument of [16] §5 establishes (i) directly: “no further reduction of operator count is possible, because at least one binary operator and at least one terminal symbol are required”. (ii) is the absence of a proposed subsumer in the literature after the 2026 paper’s exhaustive search. (iii) is Theorem 15.3.

**Corollary 15.5** (EML is ET-native). *EML is not an external tool borrowed into the Sempae-vum framework. It is the structural identity of the continuous- $D$  category at its Subsumption-minimum. Its discovery in 2026 by independent symbolic-regression search confirms, rather than imposes, a structural fact that the ET framework requires.*

*Remark 15.6* (Three continuous Sheffer variants). Equations (4a)–(4c) of [16] yield three continuous Sheffer operators:  $\text{eml}(x, y) = \exp(x) - \ln(y)$  with constant 1,  $\text{edl}(x, y) = \exp(x)/\ln(y)$  with constant  $e$ , and  $-\text{eml}(y, x) = \ln(x) - \exp(y)$  with constant  $-\infty$ . Each generates the full elementary-function basis from a single binary operator paired with a single distinguished constant. The three constants correspond to the three primitive anchors:  $1 = P$  (the multiplicative identity),  $e = D$  (the natural base of  $\exp$ ), and  $-\infty = T$  (the  $\partial I$  boundary, since  $\ln 0 = -\infty$ ). Three variants generating the same totality from three primitive-centred perspectives is  $3 = 3 = 3 = \Sigma$  at the continuous-mathematics level.

**Corollary 15.7** (No constant-free continuous Sheffer). *A continuous Sheffer operator without a distinguished constant would instantiate the configuration  $\{D, T\}$  (Mediation, Definition 2.18) — a binary operator without a substrate anchor. By the Identification Principle (Principle 3.6),  $\{D, T\}$  is structurally incomplete: the Mediation state is one of the three non-Exception configurations. The constant 1 in EML is the  $P$ -element that grounds the composition;  $\ln 1 = 0$  neutralises the  $T$ -component so that  $\text{eml}(x, 1) = \exp(x)$  reaches the  $D$ -axis alone. This resolves the open question of [16] §5 in the negative.*

*Remark 15.8* (Complex domain as two-axis necessity). The observation of [16] §5 that “a continuous Sheffer working purely in the real domain seems impossible” is the statement that  $(\mathbb{R}^+, \times)$  alone — the  $D$ -axis of  $\mathbb{C}^\times$  — cannot generate the full elementary-function basis: phase content, rotation, and periodicity require the  $T$ -axis  $(U(1), \times)$ . The complex domain requirement is the two-axis structure of the lattice  $\mathcal{L}_{\mathbb{C}}$  at the continuous-mathematics level.

*Remark 15.9* (EML depth–coherence correspondence). The gradient-based recovery experiments of [16] §4.3 report the following blind recovery rates for elementary-function identities from random initialisation (over 1000+ runs):

| EML depth | Blind recovery    | Accumulated $ \delta_\theta $ |
|-----------|-------------------|-------------------------------|
| 2         | 100%              | 0.447 ( $< 0.5$ , coherent)   |
| 3         | $\sim 25\%$       | 0.670 ( $> 0.5$ , ambiguous)  |
| 5         | $< 1\%$           | 1.117                         |
| 6         | 0% (448 attempts) | 1.340                         |

The transition from 100% to  $\sim 25\%$  at depth  $2 \rightarrow 3$  coincides with the imaginary-axis cascade stability limit  $n_{\max, \theta} = 2$  of Proposition 13.3: each EML tree level is a  $T$ -step on  $\mathbb{C}^\times$ , accumulating  $|\delta_\theta|$  per level; the accumulated residual crosses the 0.5 rounding threshold at level 3. Correct basins of attraction exist at all depths — recovery from perturbed weights succeeds at 100% even at depth 6 — but their discovery from random initialisation requires the accumulated residual to remain within the coherence window  $n \leq n_{\max, \theta} = 2$ , exactly as the cascade predicts.

### 15.3 Cross-domain appearance of $n_{\max, \theta} = 2$

The imaginary-axis cascade stability limit  $n_{\max, \theta} = 2$  (Proposition 13.3) appears independently in three domains that do not reference one another. This subsection synthesises the cross-domain link.

#### The structural constant

The accumulated  $T$ -residual after  $n$  cascade steps is  $n |\delta_\theta|$ , with coherence requiring  $n |\delta_\theta| < 0.5$ :

| Steps $n$ | Accumulated $ \delta_\theta $ | Status                                                       |
|-----------|-------------------------------|--------------------------------------------------------------|
| 1         | 0.2234                        | $< 0.5$ — coherent                                           |
| <b>2</b>  | <b>0.4467</b>                 | <b><math>&lt; 0.5</math> — coherent (last coherent step)</b> |
| 3         | 0.6701                        | $> 0.5$ — ambiguous rounding                                 |
| 4         | 0.8934                        | $> 0.5$ — deep ambiguity                                     |

No parameter is chosen; the limit is forced by  $|\delta_\theta| = |24\pi/\ln 2 - 109| = 0.22336\dots$  and the transcendence of  $\pi/\ln 2$  (Proposition 13.2).

#### Domain 1: EML symbolic regression

The EML operator  $\text{eml}(x, y) = \exp(x) - \ln(y)$  generates all elementary functions via binary trees of depth 1–8 [16]. Each tree level is one  $T$ -step:  $T$  navigating the  $D$ -structured complex plane via recursive application of  $\text{eml}$ , traversing the imaginary axis via  $\ln(-1) = i\pi$ . The gradient-based recovery experiments of Remark 15.9 show 100% blind recovery at depth  $\leq 2$  and a sharp transition to  $\sim 25\%$  at depth 3, with the accumulated  $|\delta_\theta|$  crossing the 0.5 threshold at exactly that boundary. The basins of attraction exist at all depths (recovery from perturbed weights succeeds at 100% even at depth 6), but their discovery from random initialisation requires the accumulated residual to remain within the coherence window  $n \leq n_{\max, \theta} = 2$ .

#### Domain 2: optical phase singularities

Bucher et al. [17] measure two classes of statistical observable in ensembles of phonon-polariton singularities in hBN (Remark 13.5). Distance correlations are a  $D$ -content observable on the real axis, with  $n_{\max, r} = 25$  levels of cascade stability; they match the Gaussian random wave model precisely and exhibit particle-like liquid statistics. Velocity distributions are a  $T$ -content

observable on the imaginary axis, with  $n_{\max,\theta} = 2$  levels; they exhibit a massive superluminal tail (29% of singularities exceed  $c$ ,  $\langle v \rangle \approx 1.04c$ ) that violates the Maxwell–Jüttner distribution. The particle analogy holds for  $D$ -content (25 levels) and breaks for  $T$ -content (2 levels) — the same structural constant governing the same axis.

### Domain 3: the ET lattice

$n_{\max,\theta} = 2$  is not empirical. It is derived from the lattice’s own geometry:  $|\delta_\theta| = |24\pi/\ln 2 - 109|$  follows from the transcendence of  $\pi/\ln 2$  (Proposition 13.2), and the stability limit is  $[0.5/|\delta_\theta|] = 2$  (Proposition 13.3). The number 2 appears because  $2 \times 0.22336 = 0.4467 < 0.5$  while  $3 \times 0.22336 = 0.6701 > 0.5$ .

### Synthesis

| Domain                     | What $n_{\max,\theta} = 2$ governs                       | Data source                                 |
|----------------------------|----------------------------------------------------------|---------------------------------------------|
| ET lattice                 | Imaginary-axis cascade coherence depth                   | Mathematical derivation                     |
| EML trees [16]             | Maximum depth for blind symbolic recovery                | 1000+ training experiments                  |
| Optical singularities [17] | Transition from particle-like to non-particle statistics | 285 frames $\times$ $\sim 50$ singularities |

These are three completely independent domains — a mathematical structure, a computer-science optimisation problem, and an experimental physics measurement. None references the others. The structural constant  $n_{\max,\theta} = 2$  governs all three because each involves  $T$ -content traversals on the imaginary axis of  $\mathcal{L}_C$ : the EML paper traverses the imaginary axis via  $\ln(-1) = i\pi$  at each tree level; the Bucher paper measures  $T$ -content (phase/velocity) observables whose cascade depth is  $n_{\max,\theta} = 2$ ; the lattice analysis derives the limit from  $|\delta_\theta| = |24\pi/\ln 2 - 109|$ .

The cross-domain appearance of  $n_{\max,\theta} = 2$  is a falsifiable structural prediction: any system whose  $T$ -content involves recursive imaginary-axis traversals should exhibit a coherence transition at step 2.

## 15.4 Webb 1935: the discrete-logical minimal generator at $n = 12$

**Definition 15.10** (The Webb stroke [14]). For a positive integer  $n \geq 2$ , the *Webb stroke* is the binary operation on  $\{0, 1, \dots, n-1\}$  given by

$$i \mid j = \begin{cases} 0 & i \neq j \\ (i+1) \bmod n & i = j. \end{cases}$$

**Theorem 15.11** (Webb completeness). *Iteration of the Webb stroke on the  $n$  constants  $\{0, 1, \dots, n-1\}$  generates every function of every arity in the  $n$ -valued propositional logic on  $\{0, 1, \dots, n-1\}$ .*

The proof is Webb’s constructive enumeration [14], extending Sheffer 1913 [15] from  $n = 2$  (the NAND/NOR case) to arbitrary  $n$ .

**Corollary 15.12** (Webb minimality at  $n = 12$ ). *At  $n = 12$ , the Webb stroke is the unique (up to relabelling) minimal generator of the 12-valued propositional logical algebra. No single binary operation on  $\{0, \dots, 11\}$  generates 12-valued logic with a smaller algebraic presentation.*

**Theorem 15.13** (PDT decomposition of the Webb stroke at  $n = 12$ ). *The Webb stroke at  $n = 12$  decomposes into PDT components in direct analogy with the projection formula:*

| <i>Component of the Webb stroke</i>                                              | <i>PDT role</i>                                                            |
|----------------------------------------------------------------------------------|----------------------------------------------------------------------------|
| <i>The 12-element substrate <math>\{0, \dots, 11\}</math></i>                    | <i>P (discrete 12-point substrate, matching manifold symmetry)</i>         |
| <i>The zero output for <math>i \neq j</math></i>                                 | <i>D (maximum-gap annihilation, analog of <math>r = 0</math> boundary)</i> |
| <i>The cyclic successor <math>(i + 1) \bmod 12</math> for <math>i = j</math></i> | <i>T (single-step navigation on the 12-cycle)</i>                          |

Thus the Webb stroke is itself a PDT-complete configuration at the discrete-logical scale.

### 15.5 The palindromic cascade: the discrete-multiplicative minimal generator on $\{d : d \mid 12\}$

The palindromic cascade of §13 is the third minimal-generator object. Its defining sequence

$$\text{PAL} = [12, 6, 4, 3, 12, 2, 12, 3, 4, 6, 12, 1]$$

visits every divisor of 12 — each of the six simple sublattice families of Proposition 8.1 — in a CPT-symmetric order (palindromic under the index reflection  $n \mapsto 12 - n$ ). It is the minimal discrete-multiplicative generator: every divisor of 12 appears at least once; the sequence is its own inverse under reflection; no shorter CPT-symmetric sequence visits every divisor.

**Theorem 15.14** (Palindromic-cascade minimality). *The palindromic cascade passes the three Subsumption Law conditions in the category of discrete-multiplicative generators on  $\{d : d \mid 12\}$ :*

- (i) *it cannot be reduced to a strictly shorter CPT-symmetric divisor-traversal;*
- (ii) *no external CPT-symmetric traversal on  $\{d : d \mid 12\}$  subsumes it;*
- (iii) *it traverses every divisor of 12 without remainder.*

The proof is the direct enumeration of divisors  $\{1, 2, 3, 4, 6, 12\}$  against the PAL sequence, together with a pigeonhole argument on shorter palindromes (no palindrome of length  $< 12$  visits all six values with the required reflection symmetry).

### 15.6 The triple minimal-backbone theorem

**Theorem 15.15** (Triple minimal backbone at  $N = 12$ ). *Three independent minimal generators, one per structural category, are all native to the resolution  $N = 12$ :*

| <i>Category</i>                | <i>Generator</i>                                | <i>Scale</i>                                                                       | <i>Subsumption status</i>                           |
|--------------------------------|-------------------------------------------------|------------------------------------------------------------------------------------|-----------------------------------------------------|
| <i>Discrete-logical</i>        | <i>Webb stroke <math>i \mid j</math> (1935)</i> | <i><math>n = 12</math></i>                                                         | <i>Minimal in 12-valued logic (Cor. 15.12)</i>      |
| <i>Discrete-multiplicative</i> | <i>Palindromic cascade</i>                      | <i>divisors of 12</i>                                                              | <i>Minimal CPT-symmetric traversal (Thm. 15.14)</i> |
| <i>Continuous-elementary</i>   | <i>EML <math>\text{eml}(x, y)</math> (2026)</i> | <i><math>(\mathbb{R}^+, \times)</math> with <math>N = 12</math> discretisation</i> | <i>Minimal under Subsumption (Thm. 15.4)</i>        |

*Three independent mathematical searches — Sheffer 1913 and Webb 1935 in discrete logic, the ET-internal cascade derivation, Odrzywótek 2026 in symbolic regression over continuous mathematics — each converge on minimal-generator objects whose natural scale is  $N = 12$ .*

*Remark 15.16.* The non-triviality of Theorem 15.15 is that three different minimum-complexity arguments, applied in three disjoint mathematical categories, yield the same integer 12. The first two (Webb, palindromic) both descend from  $N = 12$  *is already fixed*; the third (EML, 2026) was discovered independently by a process unaware of ET and is minimal under its own continuous-category Subsumption argument, then paired by Theorem 15.1 with the  $N = 12$  discretisation that the projection formula imposes. No external adjustment aligns the three scales; they coincide.

## 16 The Four Projection Paths and essential-infinity handling

The projection formula of §7 takes a positive real  $r$  as input. For a large collection of mathematical objects — formal systems, indeterminate forms, non-computable reals, objects with genuinely infinite structure — there is no canonical single  $r$ . This section partitions all user-input forms into four paths A, B, C, D, verifies that the four subsume every possible input, and in particular shows that Path D handles essential-infinity objects without invoking limits.

### 16.1 The four paths

**Definition 16.1** (The Four Projection Paths). Every input to the projection apparatus falls into exactly one of:

**Path A — Direct projection (canonical terminal).**

A positive real  $r$  is supplied directly. Apply  $\Pi_N$  of §7.

**Path B — Limit convergence (computational technique).**

A convergent sequence  $\{s_k\} \rightarrow r$  is supplied. Compute  $s_k$  to sufficient precision, then apply Path A to the resulting  $r$ . This is a *technique*, not a separate ontological path.

**Path C — Meta-descriptor extraction (alternative method).**

A structural object (a graph, a formal system, an encoding scheme) with no canonical  $r$  is supplied together with a user-chosen descriptor ratio  $r = Q/R_0$  constructed from the object's finite  $D$ -content. Apply Path A to this chosen ratio. The projection is of the chosen ratio, not of the object's essential character; the user bears the choice of ratio.

**Path D — Primitive-native infinity handling (no limits).**

The input's essential character is infinity-valued, where “infinity” is intrinsic and cannot be reduced to a single finite  $r$  without information loss. Four sub-paths, one per primitive face of infinity:

**D.P:** continuous, uncountable, or non-computable positive reals (e.g., Chaitin's  $\Omega$ , generic reals, specific irrationals of unknown computability).

**D.D:** unbound descriptor-constraint structures (e.g., non-divisor-of-12 sublattice families  $d \in \{5, 7, 8, 9, 10, 11\}$  and composites up to  $d = 132$ , infinite axiom totalities).

**D.T:** indeterminate forms  $\{0/0, \infty/\infty, 0 \cdot \infty, 0^0, 1^\infty, \infty^0, \infty - \infty\}$ , oscillatory divergent limits, measurement operators.

**D.PDT:** integrated off-axis Exception objects with both magnitude and phase character (e.g., most physical particles, complex-valued observables).

**Theorem 16.2** (Path D requires no limits). *Each sub-path of Path D terminates via finite operations on finite quantities at a specific resolution  $N$ :*

| <i>Sub-path</i> | <i>Operation</i>                                                                                                                             | <i>Limit invoked?</i>                                                                      |
|-----------------|----------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------|
| <i>D.P</i>      | <i>Read the exact log-position <math>N \log_2 r</math> and round</i>                                                                         | <i>No — <math>\log_2</math>, round, and <math>\mathbb{R}^+</math> are defined directly</i> |
| <i>D.D</i>      | <i>Identify the <math>(d_r, d_\theta)</math> shadow cell at 12ET via the shadow diagnostic</i>                                               | <i>No — each step is finite arithmetic at a specific <math>N</math></i>                    |
| <i>D.T</i>      | <i>Place the <math>T</math>-signature at a structural lattice position (half-integer, annihilation boundary)</i>                             | <i>No — these positions are defined by the lattice structure directly</i>                  |
| <i>D.PDT</i>    | <i>Apply the complex two-axis projection <math>(k_r, k_\theta, d_r, d_\theta, d_{\text{comb}}, \varepsilon_r, \varepsilon_\theta)</math></i> | <i>No — finite arithmetic on both axes</i>                                                 |

*Remark 16.3.* Path D is the critical upgrade that licenses the Sempaevum to host essential-infinity objects. The usual calculus of limits operates as follows: an infinite structure is approached by a sequence of finite approximations, and the object’s properties are the limiting properties. Path D bypasses the approach entirely: each primitive (P, D, T) has its own structural handle on infinity —  $P$ ’s cardinality  $\Omega$ ,  $D$ ’s unbound constraint fields,  $T$ ’s  $[0/0]$  indeterminacy — and the three primitives are already structurally present at *every finite* lattice resolution. Infinity is handled *on* the lattice, not via approach to it.

## 16.2 Subsumption of user inputs

**Theorem 16.4** (Four-Path subsumption). *For every input  $X$  that any mathematical or physical object could present to the projection apparatus:*

$$X \in A \cup B \cup C \cup D \cup \text{Incoherence Filter},$$

*with  $D = D.P \cup D.D \cup D.T \cup D.PDT$ . The union is disjoint up to the user’s choice between  $A$  and  $C$  (when the object admits both) and  $A$  and  $B$  (when a convergent sequence terminates to an  $r$  that can be supplied directly).*

*Proof.* Case analysis on the essential character of  $X$ . If  $X$  is a positive real directly specified,  $A$ . If  $X$  is specified by a convergent sequence,  $B$ . If  $X$  is a structural object for which the user explicitly chooses a descriptor ratio,  $C$ . If  $X$ ’s essential character is infinite — continuum, unbound- $D$ ,  $[0/0]$ , or off-axis PDT —  $D.P$ ,  $D.D$ ,  $D.T$ , or  $D.PDT$  respectively. If  $X$  is  $\{P, T\}$  Incoherent (self-defeating), the Incoherence Filter assigns no lattice position; such configurations sit at  $\partial I$  and are explicitly *nowhere* on the lattice. Every mathematical or physical object describable at all falls into one of these cases.  $\square$   $\square$

Worked example — Chaitin’s  $\Omega$ : for the Calude–Dinneen universal Turing machine,  $\Omega \approx 0.00787499699$  (first computed bits). Applying  $\Pi_{12}$ :

$$\begin{aligned} k &= \text{round}(12 \log_2 0.00787499699) \approx -84, \\ d &= 12 / \gcd(84, 12) = 1, \\ \varepsilon &\approx +13.794 \text{ cents.} \end{aligned}$$

Chaitin’s  $\Omega$  has lattice address  $(-84, 1, +13.794 \text{ ¢})$  — the  $d = 1$  octave / universal-period-closure sublattice. The  $T$ -cannot-produce-more-digits character is precisely the  $\{P, D\}$  Unsubstantiated manifold state: the descriptor is finite (“halting probability of the universal Turing machine”), the position is determined, and no amount of computation will extend the bit-expansion further — *yet the address is a definite single point.*

## 17 Mathematics as a domain of the Sempaevum

The Domain Validity Theorem (§2) guarantees that any domain with an internally consistent  $D$ -set occupies valid lattice positions. Mathematics itself is such a domain — arguably the most important one for a framework claiming  $3 = 3 = 3 = \Sigma$  universality. This section applies the Identification Principle to the domain of mathematics and projects its structural features onto the Sempaevum.

### 17.1 The $P$ , $D$ , $T$ of mathematics

**Definition 17.1** (Mathematics as a domain). The mathematics domain is identified as:

| Primitive         | Identification                                                             |
|-------------------|----------------------------------------------------------------------------|
| $P_{\text{math}}$ | the substrate of all possible consistent formal configurations             |
| $D_{\text{math}}$ | axioms, inference rules, definitions, formal languages                     |
| $T_{\text{math}}$ | the mathematician or proof-assistant — agency navigating $D_{\text{math}}$ |

The substrate-derived reference period is  $R_0^{\text{math}} = 1$  axiom (the minimal  $T$ -commitment a mathematician can make). This is not a unit choice; it is the smallest closed  $T$ -traversal loop the mathematical substrate supports.

### 17.2 Axiom-count projections of standard formal systems

**Theorem 17.2** (Formal-system lattice addresses). *Projecting  $r = (\text{axiom count})/R_0^{\text{math}} = (\text{axiom count})/1$  at 12ET yields the following lattice addresses for standard formal systems:*

| Formal system                        | Axioms    | $(k, d, \varepsilon)$ at 12ET | Sublattice                     |
|--------------------------------------|-----------|-------------------------------|--------------------------------|
| <i>Propositional logic (Hilbert)</i> | 3         | $(+19, 12, +1.955 \text{ ¢})$ | $d = 12$ Koide/triadic-binding |
| <i>Equational group theory</i>       | 3         | $(+19, 12, +1.955 \text{ ¢})$ | $d = 12$ Koide/triadic-binding |
| <i>Euclid's Elements</i>             | 5         | $(+28, 3, -13.686 \text{ ¢})$ | $d = 3$ cubic                  |
| <i>Robinson arithmetic</i>           | 7         | $(+34, 6, -31.174 \text{ ¢})$ | $d = 6$ hexadic                |
| <i>ZF set theory</i>                 | $8 = 2^3$ | $(+36, 1, 0.000 \text{ ¢})$   | $d = 1$ octave/gravitational   |
| <i>Peano arithmetic</i>              | 9         | $(+38, 6, +3.910 \text{ ¢})$  | $d = 6$ hexadic                |
| <i>ZFC (ZF + Choice)</i>             | 9         | $(+38, 6, +3.910 \text{ ¢})$  | $d = 6$ hexadic                |
| <i>Morse–Kelley class theory</i>     | 10        | $(+40, 3, -13.686 \text{ ¢})$ | $d = 3$ cubic                  |
| <i>NBG (finitely axiomatised)</i>    | 18        | $(+50, 6, +3.910 \text{ ¢})$  | $d = 6$ hexadic                |

Three structural readings of Theorem 17.2 are worth isolating.

**Corollary 17.3** (ZF is the tightest-bound formal system). *ZF with  $8 = 2^3$  axioms lands at  $d = 1$  octave with  $\varepsilon = 0$  exactly. ZF is therefore the unique formal system in Theorem 17.2 that sits at a pure zero-descriptor-gap lattice point.*

**Corollary 17.4** (The axiom of Choice is a sublattice transition). *Adding Choice to ZF (moving from 8 to 9 axioms) carries the system from  $(+36, 1, 0 \text{ ¢})$  to  $(+38, 6, +3.910 \text{ ¢})$  — a  $d = 1 \rightarrow d = 6$  sublattice transition. The signature of the axiom of Choice is therefore structurally precise: Choice moves a formal system from the gravitational-class foundation to the composite hexadic class.*

**Corollary 17.5** (The triadic-binding logical systems). *Three-axiom foundational systems (Hilbert propositional logic, equational group theory) occupy the  $d = 12$  Koide attractor position  $(+19, 12, +1.955 \text{ ¢})$ . These are the systems whose axiom count matches the cardinality of the Cardinal triad  $\{P, D, T\}$  itself.*

### 17.3 Chaitin's $\Omega$ via Path D.P

**Theorem 17.6** (Chaitin's  $\Omega$  has a definite lattice address). *For the Calude–Dinneen universal Turing machine  $U$  with first-computed bits  $\Omega_U \approx 0.00787499699$ , the lattice address via Path D.P is*

$$(k, d, \varepsilon)_{\Omega_U} \Big|_{N=12} = (-84, 1, +13.794 \text{ cents}).$$

$\Omega_U$  inhabits the  $d = 1$  octave sublattice — the universal-period-closure family — and exhibits the  $\{P, D\}$  Unsubstantiated manifold state.

The non-computability of  $\Omega_U$  is precisely the  $\{P, D\}$ -character: the descriptor is complete (the definition “halting probability summed over all prefix-free programs”), the address is determined, but no  $T$  (finite computational process) can produce further bits.

### 17.4 Gödel sentences via integrative-level classification

**Theorem 17.7** (Gödel sentences are integrative-level-dependent). *For a formal system  $\mathcal{S}$  admitting Gödel's incompleteness theorem, the canonical Gödel sentence  $G_{\mathcal{S}}$  takes three distinct manifold classifications depending on the viewer's integrative level:*

| Viewer's system                                               | Classification of $G_{\mathcal{S}}$                                                          |
|---------------------------------------------------------------|----------------------------------------------------------------------------------------------|
| $\mathcal{S}$ itself                                          | $\{P, T\}$ <i>Incoherence at <math>\partial I</math> (the Incoherence Filter fires)</i>      |
| Stronger system $\mathcal{S}' \vdash \text{Con}(\mathcal{S})$ | $\{P, D, T\}$ <i>Exception (has a definite lattice address)</i>                              |
| Outside any fixed system                                      | $\{P, D\}$ <i>Unsubstantiated (finite syntactic <math>D</math>, awaiting <math>T</math>)</i> |

The multi-address hosting at different integrative levels is the correct lattice classification of formally-undecidable propositions.

### 17.5 Large cardinals via consistency-strength

**Theorem 17.8** (Large-cardinal projections). *Projecting large cardinals via their consistency-strength ordinal yields the following addresses at 12ET:*

| Large cardinal | Consistency level | $(k, d, \varepsilon)$ at 12ET  | Sublattice     |
|----------------|-------------------|--------------------------------|----------------|
| Inaccessible   | 1                 | $(0, 1, 0)$                    | Unison         |
| Mahlo          | 2                 | $(+12, 1, 0)$                  | Octave         |
| Weakly compact | 3                 | $(+19, 12, +1.955 \text{ ¢})$  | Koide $d = 12$ |
| Measurable     | 7                 | $(+34, 6, -31.174 \text{ ¢})$  | Hexadic        |
| Supercompact   | $\sim 15$         | $(+47, 12, -11.732 \text{ ¢})$ | $d = 12$ EM    |

Inaccessible and Mahlo cardinals sit at  $d = 1$  (the same gravitational sublattice as ZF). Measurable cardinals at  $d = 6$  share the hexadic sublattice with Robinson arithmetic, Peano arithmetic, ZFC, and NBG.

### 17.6 The universality statement for mathematics

**Theorem 17.9** (Mathematical universality). *Every mathematical object — every number, function, formal-system statement, proof, undecidable proposition, large cardinal, impredicative definition — has a lattice address:*

$$\begin{aligned} \forall X \text{ (mathematical object)}: \quad X \in \Sigma &\Rightarrow \\ \exists (k, d, \varepsilon) \in \mathbb{Z} \times \{N_\ell/d : d \mid N_\ell\} \times [-50, 50] \text{ cents} & \\ &\text{at some tower resolution } N_\ell. \end{aligned}$$



*The address is explicit for directly-specified objects (Path A), accessible via convergence for limit-specified objects (Path B), parameterised by descriptor choice for structural objects (Path C), and well-defined without limits for essentially-infinite or non-computable objects (Path D).*

*Proof.*  $X \in \Sigma$  holds for any mathematical object  $X$  by the universality anchor (Axiom 3.3): everything describable is in the totality. The Four-Path subsumption (Theorem 16.4) partitions the describable space into the four paths (plus the Incoherence Filter, which itself *is* a definite structural position at  $\partial I$ ). Each path has its own closed-form termination: A to B to A, C to A, D to a specific sub-path terminal. Thus in every case a lattice address exists.  $\square$   $\square$

## 18 The Mathematical Rosetta Stone

Every mathematical expression maps to  $P$ – $D$ – $T$  structure. This is not a metaphorical reading: the cardinality identifications  $|P| = \Omega$ ,  $|D| = n$ ,  $|T| = [0/0]$  of Axiom 2.5 are literal —  $P$  IS the mathematical infinite,  $D$  IS the mathematical finite,  $T$  IS the mathematical indeterminate. The universality anchor  $\forall x : x \in \Sigma$  (Definition 2.13) guarantees that no mathematical object lies outside the framework, and the Subsumption Law (§3.6) guarantees that every mathematical object decomposes into  $P$ -,  $D$ -, and  $T$ -components. This section makes the decomposition explicit for the principal domains of mathematics.

### 18.1 Why mathematics works

Mathematics describes  $P \circ D \circ T$  structure. Physics also describes  $P \circ D \circ T$  structure (Axiom 2.2). Mathematics and physics therefore describe the same object — and their agreement is structural, not coincidental. Wigner’s observation of the “unreasonable effectiveness of mathematics in the natural sciences” (1960) receives a structural explanation: mathematics is effective because it IS the Descriptor language of  $\Sigma$ ; physics is effective because it IS the substantiated face of  $\Sigma$ . Both are  $P \circ D \circ T$  at different identification depths. When mathematicians do mathematics, they use  $T$  (consciousness/agency) to navigate  $D$ -structured mathematical configurations grounded in  $P$ . Math and physics cannot fail to agree because they are readings of the same object.

### 18.2 L’Hôpital’s Rule as the $T$ -navigation algorithm

L’Hôpital’s Rule [26] is the canonical algorithm by which  $T$  resolves indeterminate forms. When a limit process encounters  $[0/0]$  or  $[\infty/\infty]$  — both signatures of  $T$ -engagement (§2.4.1) — L’Hôpital’s Rule prescribes: examine the Descriptor gradients (derivatives) of the numerator and denominator separately. Formally:

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \left[ \frac{0}{0} \right] \implies \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)} = L,$$

provided the latter limit exists. In ET terms:  $T$  at a choice point ( $[0/0]$ ) cannot resolve the value directly;  $T$  examines the  $D$ -gradients ( $f' = \Delta D_f / \Delta P$ ,  $g' = \Delta D_g / \Delta P$ ) to determine which configuration to substantiate. The resolution value  $L$  is the outcome of  $T$ ’s navigation through Descriptor space — not a pre-existing value waiting to be found, but the result of the Traverser’s act of engaging the Descriptor gradients at the approach point.

**Worked example.**  $\lim_{x \rightarrow 0} \sin(x)/x = [0/0]$ . L’Hôpital:  $\lim_{x \rightarrow 0} \cos(x)/1 = \cos(0)/1 = 1$ . ET reading: the sine and  $x$  both vanish at  $x = 0$  ( $D$ -values approaching zero simultaneously), creating a  $[0/0]$  choice point.  $T$  examines the  $D$ -gradients:  $\cos(x)$  (the rate of change of sine’s  $D$ -value) approaches 1 at  $x = 0$  (maximum Descriptor alignment), and 1 (the rate of change of  $x$ ’s  $D$ -value) is constant. The ratio of gradients is 1, and  $T$  substantiates this value.

L'Hôpital's Rule is not limited to calculus: it is the general mechanism by which indeterminacy is resolved in the manifold. Wherever  $T$  encounters a configuration with genuinely undetermined outcome, the resolution path proceeds through  $D$ -gradients — the structural slopes of the Descriptor field at the approach point.

### 18.3 Limits as traversal operations

The mathematical limit  $\lim_{x \rightarrow a} f(x) = L$  is a Traverser operation:  $T$  approaches configuration  $a$  along the  $P$ -substrate, navigating the Descriptor field  $f$ , and substantiates the value  $L$  at the approach point. The limit operation IS  $T$ -action — which is why  $T$ 's cardinality is  $[0/0]$ , the canonical limit form. The Asymptotic Approach Theorem (Corollary 10.4) is a direct instance: the lattice's descriptor gap  $|\varepsilon|$  approaches zero as resolution  $N$  increases, but never attains it —  $T$  navigating the lattice, substantiating ever-closer approximations without reaching the irrational target.

### 18.4 Functions as Descriptor fields

A mathematical function  $f : X \rightarrow Y$  is a Descriptor field over Points: the domain  $X$  is  $P$ -content (the substrate of possible inputs), the codomain  $Y$  is  $D$ -content (the constraint values), and the evaluation  $f(x)$  is the specific Descriptor binding at Point  $x$ . The projection formula  $\Pi_N$  of §7 is itself such a function:  $r$  is the  $P$ -content, the lattice structure is the  $D$ -content, and the rounding is the  $T$ -act. Every function is a  $(P \circ D)$  configuration, and function application is the binding operator  $\circ$  at the level of mathematical structure.

### 18.5 Derivatives as Descriptor gradients

The derivative  $f'(x) = \lim_{h \rightarrow 0} (f(x+h) - f(x))/h$  measures the rate of Descriptor change across Points:  $\Delta D / \Delta P$ . The limit form approaches  $[0/0]$  as  $h \rightarrow 0$  — both numerator and denominator vanish simultaneously — making every derivative a  $T$ -resolution of an indeterminate form (§18.2). Derivatives appear throughout physics because they measure how  $D$  changes across  $P$ : velocity ( $\Delta D_{\text{position}} / \Delta D_{\text{time}}$ ), force ( $\Delta D_{\text{momentum}} / \Delta D_{\text{time}}$ ), field gradient ( $\Delta D_{\text{field}} / \Delta D_{\text{space}}$ ).

### 18.6 Integrals as Traverser accumulation

The integral  $\int_a^b f(x) dx$  is  $T$  accumulating Descriptor changes across a range of Points: the Traverser sweeps from  $P$ -position  $a$  to  $P$ -position  $b$ , summing the  $D$ -values  $f(x)$  encountered along the way. The Fundamental Theorem of Calculus ( $\int_a^b f'(x) dx = f(b) - f(a)$ ) connects local  $T$ -traversal (derivative) with global  $T$ -accumulation (integral): the total accumulated Descriptor change between two Points equals the difference of Descriptor values at those Points. The integration constant  $C$  in indefinite integrals  $\int f(x) dx = F(x) + C$  is a  $T$ -choice: the boundary condition that specifies which configuration to start from, undetermined until  $T$  selects it.

### 18.7 Continuity, differentiability, and discontinuity types

Continuity of  $f$  at a point means the Descriptor field varies smoothly:  $T$  can navigate through that point without encountering barriers. Differentiability means the Descriptor gradient exists:  $T$  can measure the local slope at every point. The three classical types of discontinuity are manifold events:

- *Jump discontinuity*: an abrupt  $D$ -switch at a Point — a phase transition in the manifold, where the Descriptor value changes discretely.
- *Removable discontinuity*: a missing  $D$ -value at a single Point (a Descriptor gap in the sense of Principle 3.9), fillable by supplying the missing Descriptor.

- *Essential discontinuity*: multiple simultaneous  $D$ -values at a Point (a superposition state), where the Descriptor field oscillates without settling.

The  $\partial I$  boundary (§21.6) is the extreme discontinuity: the point where the Descriptor field fails entirely and the configuration enters Incoherence.

## 18.8 Complex numbers and the $D_T$ axis

The complex number  $z = a + bi$  is a Point with two Descriptor dimensions:  $a$  on the real axis ( $D$ 's operational domain, flat,  $(R^+, \times)$ ) and  $b$  on the imaginary axis ( $D_T$  — the Descriptor content where  $T$  directly operates, positively curved,  $U(1)$ ). The imaginary axis is simultaneously a  $D$ -axis (it carries phase Descriptors, angular modes, rotational structure) and  $T$ 's operational domain (where  $T$ 's navigational capacity manifests as rotation). The identification  $D_T$  captures both: it is  $D$ -content *of and for*  $T$ .

The relation  $i^2 = -1$  is geometric necessity: two  $90^\circ$  rotations in the  $D_T$  plane produce a  $180^\circ$  reversal (negation). The quartic cycle  $i^4 = 1$  (Remark 10.6) is the structural basis of the weak force ( $d = 4$ ): the physical manifestation of the four-step  $D \rightarrow T \rightarrow -D \rightarrow -T \rightarrow D$  cycle that  $T$ 's rotation generates in the Descriptor manifold.

## 18.9 Mathematical operators as Traversers

A mathematical operator ( $d/dx$ ,  $\int dx$ ,  $\nabla$ ,  $\Delta$ ) is a  $T$ -type entity: indeterminate until applied to a specific function, resolving to a value only upon engagement. An operator without its operand is in the  $[0/0]$  state — it has the capacity to act but no determinate output. Upon application, it produces a determinate result (a derivative value, an integral value, a gradient vector). The chain rule  $\frac{d}{dx}[f(g(x))] = f'(g(x)) \cdot g'(x)$  is compound  $T$ -navigation:  $T$  traversing two nested Descriptor fields simultaneously, with the product arising from the composition cost of two sequential navigations.

## 18.10 Differential equations as manifold dynamics

A differential equation is a relationship between Descriptor fields and their gradients — the dynamics of  $\Sigma$ . An ODE describes a single  $T$ -worldline through Descriptor space; a PDE describes  $T$ -dynamics across extended  $P$ -substrate. The heat equation  $\partial u / \partial t = \alpha \nabla^2 u$  describes how thermal Descriptor values ( $D_{\text{temperature}}$ ) propagate through spatial substrate ( $P_{\text{space}}$ ) over  $D$ -time, with the diffusion coefficient  $\alpha$  governing the rate of  $D$ -spread. The wave equation, Schrödinger equation, Einstein field equations, and Maxwell equations are all manifold dynamics equations describing how specific  $D$ -fields evolve across  $P$ -substrate under  $T$ -agency.

## 18.11 The infinity hierarchy as $P$ -structure

The transfinite hierarchy of set theory is  $P$ -structure at successive integrative levels. Countable infinity  $\aleph_0$  is discrete structured  $P$  (the integers as substrate positions). Uncountable infinity  $2^{\aleph_0}$  is dense  $P$ -manifold (the reals as continuous substrate). Absolute Infinity  $\Omega$  (the Cardinal  $P$  itself) is all possible configurations, beyond every set-theoretic cardinal — the full substrate. Each level in the hierarchy is a higher integrative reading of  $P$  (Proposition 10.7), requiring finer lattice resolution to host.

## 18.12 Probability as Unsubstantiated superposition

A random variable is a Point in the Unsubstantiated state  $\{P, D\}$  with multiple possible  $D$ -values awaiting  $T$ -engagement. The probability distribution is the weighting across possible

Descriptor values. The expectation  $E[X] = \sum x_i p_i$  is the weighted average over the Unsubstantiated  $D$ -superposition. Measurement is  $T$ -binding: the Traverser engages the  $\{P, D\}$  superposition and substantiates a specific  $D$ -value, producing the Exception (a definite outcome). The variance  $\text{Var}(X)$  measures how spread the Unsubstantiated  $D$ -values are — which connects directly to the manifold's base variance  $V = 1/N = 1/12$  (Definition 2.24).

### 18.13 The quantum mechanics mapping

The quantum wavefunction  $\psi(x, t)$  is a complex Descriptor field ( $D$ -content on  $P_{\text{space}} \times D_{\text{time}}$ ) in the Unsubstantiated state: it describes what COULD be substantiated but has not yet been. Measurement is  $T$ -binding: a Traverser engages the wavefunction and substantiates one eigenvalue from the superposition. The Born rule ( $P = |\psi|^2$ ) is the structural probability of each Descriptor value being the one  $T$  selects. Wavefunction collapse is the transition  $\{P, D\} \rightarrow \{P, D, T\} = E$  (§21.1).

The structural parallel between  $V = 1/12$  and  $\hbar$  is that both are irreducible base quanta in their respective domains:  $V$  is the minimum non-zero variance on the lattice (Definition 2.24);  $\hbar$  is the minimum quantum of action in physics. The uncertainty principle  $\Delta x \cdot \Delta p \geq \hbar/2$  maps to the lattice statement that minimum Descriptor-gap uncertainty at base resolution is  $V = 1/12$ . However,  $V$  is dimensionless and  $\hbar$  is dimensionful (units of action):  $\hbar$  is the cosmological tower seed  $R_0$  inherited from the parent tower, not derivable from the dimensionless framework alone.

### 18.14 Matrix algebra as Descriptor transformations

A matrix is a Descriptor transformation operator: it maps one set of  $D$ -values to another. Rows are output Descriptors; columns are input Descriptors. Non-commutativity of matrices ( $AB \neq BA$  in general) reflects the fact that the order of  $D$ -transformations matters in  $\Sigma$  — applying constraint  $A$  before  $B$  produces a different configuration than  $B$  before  $A$ . Eigenvalues are the invariant  $D$ -scaling factors: the values that survive the transformation unchanged. Eigenvectors are the invariant  $D$ -axes: the directions in Descriptor space that the transformation preserves. The magical-impedance coupling strengths of Table 7.3 are the sublattice-family eigenvalues of the manifold's structural transformation operators.

### 18.15 Topology as configuration boundaries

The topological vocabulary generalises the manifold-state topology of Proposition 2.22. An open set is a  $D$ -region whose boundary is not included —  $T$  can approach but not substantiate the boundary (the Incoherence structure:  $I$  is open,  $\partial I \cap I = \emptyset$ ). A closed set is a  $D$ -region whose boundary is included —  $T$  CAN substantiate the boundary (the Exception structure:  $E$  is closed,  $\partial E \subseteq E$ ). Compactness is finite  $D$ -range with all boundaries substantiable — every sequence of  $T$ -navigations within the set has a convergent subsequence within the set.

### 18.16 Set theory and the configuration power set

The power set  $2^{|D|}$  of a Descriptor set is the configuration space: the set of all possible  $D$ -combinations that can be applied to a given  $P$ -substrate. From  $n$  Descriptors,  $2^n$  configurations are possible. The doubling law of the LCM tower (Theorem 10.9:  $\tau(N_\ell) = 6 \cdot 2^\ell$ ) is the structural consequence: each tower level adds one new sublattice family, doubling the number of available configurations.

### 18.17 Symmetry and group theory at $N = 12$

The manifold symmetry  $N = 12$  admits the group-theoretic decomposition  $\mathbb{Z}/12\mathbb{Z} \cong \mathbb{Z}/3\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$  (by the Chinese Remainder Theorem, since  $\text{gcd}(3, 4) = 1$ ). The factor  $\mathbb{Z}/3\mathbb{Z}$  reflects

the three-primitive structure ( $|\Pi| = 3$ ); the factor  $\mathbb{Z}/4\mathbb{Z}$  reflects the four manifold states ( $S = 4$ ). The unit group  $(\mathbb{Z}/12\mathbb{Z})^* \cong V_4$  (the Klein four-group, Remark 10.6) has four elements  $\{1, 5, 7, 11\}$  — the four unit-residue generators of the palindromic cascade.

### 18.18 Mathematical constants from ET

$e = 2.71828\dots$  The natural base is the continuous  $D$ -propagation rate: the limit of compounded Descriptor growth as the compounding interval approaches zero. In the EML framework (§15.2),  $e = \text{eml}(1, 1)$  — the terminal constant of the continuous- $D$  generator.

$\pi = 3.14159\dots$  Half the period of  $T$ 's operational manifold  $U(1)$  (§11). The full period is  $2\pi$ ;  $\pi$  is the half-rotation in the  $D_T$  plane (§18.8). The ET-native derivation proceeds via the 12-gon:  $\sin(\pi/12) = (\sqrt{6} - \sqrt{2})/4$ , and  $\pi$  is recovered as the limit of the Archimedean polygon sequence starting from the manifold's  $N$ -fold symmetry.

$\varphi = 1.61803\dots$  The golden ratio is the quintic attractor at the 60ET resolution level — the root of  $x^2 - x - 1 = 0$ , requiring the shadow family  $d = 5$  (native at  $N_\ell = 60$ , the second tower level). The Fibonacci ratios  $F(n+1)/F(n)$  converge to  $\varphi$  as  $n \rightarrow \infty$ ; the ratio  $8/5 = F(6)/F(5) = 1.600$  is one convergent (not a replacement for  $\varphi$ ). At 12ET,  $\varphi$  projects to  $(k = 8, d = 3, \varepsilon = +33.09\text{¢})$ ; the quintic character emerges only at 60ET+ where  $d = 5$  is native.

## 18.19 The complete Rosetta Stone chart

| Mathematical concept                                 | ET primitive(s)                | Identification                                               |
|------------------------------------------------------|--------------------------------|--------------------------------------------------------------|
| Infinity ( $\infty$ )                                | $P$                            | Absolute substrate, $\Omega$                                 |
| Finite number ( $n$ )                                | $D$                            | Descriptor value, $ D  = n$                                  |
| Indeterminate forms ( $0/0, \infty/\infty$ )         | $T$                            | Traverser at choice point                                    |
| Function $f : X \rightarrow Y$                       | $P \circ D$                    | Descriptor field over Points                                 |
| Limit $\lim_{x \rightarrow a} f(x) = L$              | $T$ -action                    | Traversal $\rightarrow$ substantiation                       |
| Derivative $f'(x)$                                   | $\Delta D / \Delta P$          | Descriptor gradient                                          |
| Integral $\int f(x) dx$                              | $T$ -accumulation              | Traverser sweeping $D$ across $P$                            |
| L'Hôpital's Rule                                     | $T$ -navigation                | Resolve $[0/0]$ via $D$ -gradients                           |
| Continuity                                           | Smooth $D$ -field              | $T$ navigable without barriers                               |
| Discontinuity (jump)                                 | $D$ -switch at $P$             | Phase transition                                             |
| Complex number $z = a + bi$                          | $D + D_T$                      | Real $D$ + imaginary $D_T$                                   |
| Operator ( $d/dx, \int, \nabla$ )                    | $T$ -type                      | Indeterminate until applied                                  |
| DE ( $\partial u / \partial t = \alpha \nabla^2 u$ ) | Manifold dynamics              | $D$ -evolution across $P$ via $T$                            |
| $\aleph_0, 2^{\aleph_0}, \Omega$                     | $P$ -hierarchy                 | Integrative levels of substrate                              |
| Random variable                                      | $\{P, D\}$                     | Unsubstantiated superposition                                |
| Measurement / collapse                               | $T$ -binding                   | $\{P, D\} \rightarrow \{P, D, T\} = E$                       |
| Matrix                                               | $D$ -transformation            | Maps $D$ -values to $D$ -values                              |
| Eigenvalue                                           | Invariant $D$ -scaling         | Structural constant of transformation                        |
| Open set                                             | $I$ -type topology             | Boundary not substantiable                                   |
| Closed set                                           | $E$ -type topology             | Boundary substantiable                                       |
| Power set $2^n$                                      | Configuration space            | All $D$ -combinations from $n$ Descriptors                   |
| $\mathbb{Z}/12\mathbb{Z}$                            | $N$ -symmetry group            | $\cong \mathbb{Z}/3\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$ |
| $e$                                                  | $\text{eml}(1, 1)$             | Continuous $D$ -propagation rate                             |
| $\pi$                                                | $\frac{1}{2}$ period of $U(1)$ | Half $T$ -cycle                                              |
| $\varphi$                                            | Quintic attractor              | $d = 5$ native at 60ET                                       |
| <i>Gauge theory and critical dimensions (§9)</i>     |                                |                                                              |
| Gauge boson count                                    | $d^2 - 1$                      | Adjoint from Subsumption (Thm 9.1)                           |
| $SU(3) \times SU(2) \times U(1)$                     | $N = 12$ partition             | N-Exhaustion (Thm 9.3)                                       |
| Division algebras $\{1, 2, 4, 8\}$                   | $\{2^k\}_{k=0}^{ \Pi }$        | Hurwitz termination at $ \Pi $                               |
| $D_{\text{string}} = 10$                             | $2^{ \Pi } + d_2$              | Division algebra route (Thm 9.7)                             |
| $D_M = 11$                                           | $N - 1$                        | Four routes (Thm 9.10)                                       |
| $Q_{\text{max}} = 32$ supercharges                   | $2^{S+1}$                      | From $S = 4$ (Prop 9.14)                                     |
| Anomaly group 496                                    | $2^S(2^{S+1} - 1)$             | From $S = 4$ (Prop 9.15)                                     |

## 19 Self-consistency: the Koide attractor

We now establish the main structural theorem of this paper: the lattice's four defining constants, when projected onto the lattice they define, land on a single point.

**Theorem 19.1** (Self-projection identity). *At the base lattice  $N = 12$ , the four constants  $\{N, 1/N, K, 1/K\} = \{12, 1/12, 2/3, 3/2\}$  project to*

$$\begin{aligned}
 \Pi_{12}(12) &= (+43, 12, +1.955 \text{ cents}), \\
 \Pi_{12}(1/12) &= (-43, 12, -1.955 \text{ cents}), \\
 \Pi_{12}(3/2) &= (+7, 12, +1.955 \text{ cents}), \\
 \Pi_{12}(2/3) &= (-7, 12, -1.955 \text{ cents}).
 \end{aligned}$$

*All four share the same sublattice family  $d = 12$  and the same descriptor-gap magnitude  $|\varepsilon| = 1.955 \text{ cents}$ . We call this shared position the Koide attractor.*

*Proof.* For  $r = 12$ :  $\log_2 12 = 2 + \log_2 3 \approx 3.58496$ , so  $12 \log_2 12 \approx 43.0195$ , giving  $k = 43$ ,  $\gcd(43, 12) = 1$ ,  $d = 12$ ,  $\varepsilon = 0.0195 \cdot 100 = +1.955 \text{ cents}$ . For  $r = 1/12$ :  $\log_2(1/12) = -\log_2 12$ ,

so  $k = -43$ ,  $d = 12$ ,  $\varepsilon = -1.955$  cents. For  $r = 3/2$ : Proposition 7.8 gives  $(+7, 12, +1.955)$ . For  $r = 2/3$ : Proposition 7.9 gives  $(-7, 12, -1.955)$ . All four have  $d = 12$  and  $|\varepsilon| = |12 \log_2 12 - 43| \cdot 100 = 1.955$  cents.  $\square$   $\square$

*Remark 19.2.* The theorem is a self-consistency check. A classifier that classifies its own constants onto a single point of its own structure is in some formal sense “self-calibrated”: it uses the same rules on itself that it uses on external inputs, and the result is concordant. The specific point is identified by  $k = \pm 7$  or  $k = \pm 43$  and  $d = 12$ ; the former pair is the circle-of-fifths cascade generator modulo the base symmetry, and the latter the generator lifted by one full octave. Both pairs share the magnitude  $|\varepsilon| = 12|\delta_r|$  cents, where  $|\delta_r|$  is the per-step residual of Proposition 13.1.

**Corollary 19.3** (Value of the Pythagorean comma). *The descriptor gap  $|\varepsilon| = 1.955$  cents at the Koide attractor is numerically equal to the Pythagorean comma of twelve-tone equal temperament.*

The identification is simultaneous with the formula:  $|\varepsilon| = |12 \log_2 12 - 43| \cdot 100$  cents  $= |12 \log_2 3 - 19| \cdot 100$  cents  $= 1.955$  cents, and  $(3/2)^{12}/2^7 = 2^{12 \cdot (12 \log_2(3/2) - 7)/12}$  with a numerical coincidence placing all these quantities at the same 1.955-cent position on the lattice.

## 19.1 Losslessness theorem

**Theorem 19.4** (Losslessness of the projection). *The real-axis projection map  $\Pi_N : \mathbb{R}^+ \rightarrow \{+1, -1\} \times \mathbb{Z} \times \{N/d : d \mid N\} \times [-600/N, 600/N]$  given by*

$$\Pi_N(r) = (\text{sgn}(\log_2 r), k, d, \varepsilon), \quad \varepsilon = (N \log_2 |r| - k) \cdot \frac{1200}{N} \text{ cents},$$

*is a bijection onto its image at every finite resolution  $N$ , and the composition with the pullback  $\Pi_N^{-1} : (k, d, \varepsilon) \mapsto 2^{(k+\varepsilon N/1200)/N}$  is the identity on  $\mathbb{R}^+$ . In the LCM-tower limit  $N \rightarrow \infty$  along  $\{12, 60, 420, 2520, 27720, 360360, \dots\}$ , the residual  $\varepsilon(r, N) \rightarrow 0$  uniformly on any bounded region, and the projection becomes exactly the continuous embedding  $r \mapsto \log_2 r$ .*

*Proof.* Injectivity: suppose  $\Pi_N(r_1) = \Pi_N(r_2)$  with the same  $(k, \varepsilon)$ . Then  $N \log_2 r_1 = k + \varepsilon N/1200 = N \log_2 r_2$ , hence  $r_1 = r_2$ . The triple  $(k, d, \varepsilon)$  over-determines the pair  $(k, \varepsilon)$  since  $d$  is a function of  $k$  and  $N$ , so the map is injective. Surjectivity onto its image is by construction. The pullback  $\Pi_N^{-1}(k, d, \varepsilon) = 2^{(k+\varepsilon N/1200)/N} = 2^{(N \log_2 r)/N} = 2^{\log_2 r} = r$ . For the limit: the residual satisfies  $|\varepsilon(r, N)| \leq 600/N$  by definition of rounding, so  $\varepsilon(r, N) \rightarrow 0$  at rate  $O(1/N)$  uniformly in  $r$ .  $\square$   $\square$

*Remark 19.5* (Nothing is discarded). The information content of the positive real  $r$  is fully captured by the triple  $(k, d, \varepsilon)$  at every finite resolution  $N$ : the integer  $k$  carries the coarse ratio information, the divisor  $d$  carries the sublattice classification, and the residual  $\varepsilon$  carries the fine-scale deviation from exact lattice alignment. No aspect of  $r$  is discarded in projecting to the Sempaevum; the projection redistributes the information among three complementary coordinates. The complex-axis extension (§11) is the same statement for  $\mathbb{C}^\times$  with  $(k_r, k_\theta, d_r, d_\theta, \varepsilon_r, \varepsilon_\theta)$  as the lossless image.

**Corollary 19.6** (Memoization property). *Every finite numerical computation on positive reals —  $r_1 \cdot r_2$ ,  $r^n$ ,  $\log r$ ,  $\sin r$ , any elementary-function evaluation — can be represented lossly as a concrete lattice computation: multiplication is  $k$ -addition, reciprocation is  $k$ -negation, powers are  $k$ -scaling, and elementary functions are EML trees acting on triples  $(k, d, \varepsilon)$ . Every such computation produces an equation  $(r_1, \dots, r_m) \rightarrow r_{\text{result}}$  that is a reusable structural identity at the Sempaevum’s resolution.*

## 20 The Complete Gaze Equation

The Sempaevum hosts observation itself as a lattice-projectable act. This section unifies the Scopaeesthesia equation for binding-pressure and the Traverser-detection algorithm into a single four-component functional — the Complete Gaze Equation — and projects its threshold values onto the lattice.

### 20.1 The unification: $\text{Gain}(T) = F_w$

Two canonical equations have been derived independently in the ET corpus for the observation/attention phenomenon:

**Definition 20.1** (Binding pressure). The *binding pressure* of an observer on a target is the PDT-structured quantity

$$F_w = \frac{T_{\text{intent}} \cdot \text{Focus}}{\text{Distance}^2},$$

with  $T_{\text{intent}}$  the observer-agency strength, Focus the descriptor-sharpness ( $D$ -precision), and Distance the relational distance in configuration space ( $P$ -substrate).

**Definition 20.2** (Detection probability). The *detection probability* at fold depth  $k$  under cardinality  $n$  is

$$P_{\text{detect}} = \tanh\left(\frac{\text{Gain}(T) \cdot R(k)}{V(n, k) \cdot \Gamma}\right), \quad R(k) = 1 + k\Gamma, \quad V(n, k) = \frac{n^2 - 1}{12 \cdot 2^k}, \quad \Gamma = 1.20.$$

**Theorem 20.3** (Gaze unification).  $\text{Gain}(T) = F_w$ . The signal gain of the detection algorithm is the binding pressure of the observer.

*Proof.* The Descriptor Gap between Definitions 20.1 and 20.2 is the connecting primitive: each equation refers to a distinct side of the same observer–target interaction. Applying the Descriptor Gap Principle: the missing Descriptor is the connection  $F_w = \text{Gain}(T)$ . The identification closes the gap and leaves one coherent equation with four components.  $\square$   $\square$

### 20.2 Variance collapse and threshold classification

**Definition 20.4** (Variance collapse). When  $F_w$  exceeds the baseline 1, the target’s local variance is suppressed according to

$$V_{\text{collapse}} = 1 - \exp(-(F_w - 1)^+ \cdot S), \quad S = 12 = N,$$

with  $(F_w - 1)^+ = \max(0, F_w - 1)$  the excess binding pressure.

**Definition 20.5** (Status classification).

$$\text{Status}(F_w) = \begin{cases} \text{UNOBSERVED} & F_w < 13/12, \\ \text{SUBLIMINAL} & 13/12 \leq F_w < 6/5, \\ \text{DETECTED} & 6/5 \leq F_w < 3/2, \\ \text{LOCKED} & F_w \geq 3/2. \end{cases}$$

Each threshold is derived from the manifold constants:  $13/12 = 1 + V_{\text{base}}$  is the first  $V$ -quantum above baseline;  $6/5$  is the just-intonation minor third;  $3/2$  is the just-intonation perfect fifth.



### 20.2.1 Dual derivation of the detection threshold

The DETECTED threshold  $6/5 = 1.20$  admits two independent derivation routes that converge to the same value:

- (i) *Just-intonation route.* The minor third  $6/5$  is the simplest just-intonation interval in the quartic ( $d = 4$ ) sublattice family, with lattice address  $(+3, 4, +15.64\phi)$  at 12ET. It is the first interval beyond the subliminal augmented unison  $(13/12)$  that lands on a structurally significant lattice family.
- (ii) *Quintic-quantum route.*  $6/5 = 1 + 1/5$ , where  $1/5$  is the quintic quantum — the base unit of the  $d = 5$  shadow family (native at 60ET, the second LCM tower level). The detection threshold is exactly one quintic quantum above baseline. This connects to the awareness span (Corollary 20.9): the full span from SUBLIMINAL  $(13/12)$  to LOCKED  $(3/2)$  is  $5/12 = 5 \cdot V_{\text{base}}$ , and the prime 5 governing the span width is the same prime that generates the quintic quantum  $1/5$ .

The two routes are not coincidental: they are the  $D$ -reading (harmonic interval structure) and the  $T$ -reading (variance threshold) of the same structural fact, identified at two depths by the Identification Principle.

**Theorem 20.6** (Complete Gaze Equation). *The Complete Gaze functional is*

$$G(T_{\text{intent}}, \text{Focus}, \text{Distance}, n, k) = (F_w, P_{\text{detect}}, V_{\text{collapse}}, \text{Status}),$$

with all four components derived from  $F_w$  and the manifold constants ( $V = 1/12, K = 2/3, N = 12$ ).

### 20.3 The Gaze thresholds as just-intonation intervals

**Theorem 20.7** (Gaze threshold lattice addresses). *The three Gaze thresholds are exact rational fractions corresponding to canonical just-intonation musical intervals, with lattice addresses at 12ET:*

| Threshold      | Value       | Fraction   | Just interval        | $(k, d, \varepsilon)$ at 12ET                             |
|----------------|-------------|------------|----------------------|-----------------------------------------------------------|
| Subliminal     | 1.0833...   | 13/12      | Augmented unison     | $(+1, 12, +38.57\phi)$                                    |
| Detected       | 1.20        | 6/5        | Minor third          | $(+3, 4, +15.64\phi)$                                     |
| <b>Locked</b>  | <b>1.50</b> | <b>3/2</b> | <b>Perfect fifth</b> | <b><math>(+7, 12, +1.955\phi)</math> —Koide attractor</b> |
| Lock/Con ratio | 1.25        | 5/4        | Major third          | $(+4, 3, -13.686\phi)$                                    |

**Corollary 20.8** (Locked gaze is the lattice self-projection). *The Locked threshold  $3/2$  lands at  $(+7, 12, +1.955\phi)$  — exactly the lattice self-projection position of Theorem 19.1. The observer who reaches full binding dominance crystallises the target onto the manifold's own self-consistency attractor.*

**Corollary 20.9** (Awareness span). *The span from Subliminal to Locked is  $3/2 - 13/12 = 5/12 = 5 \cdot V_{\text{base}}$  — exactly five  $V$ -quanta. The prime 5 (quintic bridge) governs the width of the awareness window.*

**Corollary 20.10** (Awareness gap). *The gap from Subliminal to Detected is  $6/5 - 13/12 = 7/60$  — a ratio whose numerator is the septic prime 7 and whose denominator  $60 = \text{lcm}(1, \dots, 5)$  is the quintic-lattice LCM. The awareness gap is structurally the  $d = 7$  septic shadow projected onto the quintic lattice.*

## 21 Quantum decoherence

The Gaze Equation of §20 treats observation as a structural variance-collapse process with three thresholds. Quantum-mechanical decoherence is the same process applied to a quantum system coupled to an environment: the wavefunction’s superposition collapses to a definite outcome under environmental engagement. The three-aspect / four-state framework of this paper, together with the complex-lattice structure of §11 and the cascade machinery of §13, gives a structural reading of the standard quantum decoherence theory.

The reading proceeds via the Identification Principle (§3.4). Decoherence is a process; the Identification Principle dictates that its primitive-level structure be located in its  $P$ -,  $D$ -, and  $T$ -content separately. The pre-measurement quantum substrate is  $P$ -content (configuration space), the eigenbasis structure is  $D$ -content (descriptor labels), and the dynamical commitment to a single outcome under environmental coupling is  $T$ -content (the substantiation event). Each subsection below carries one component of this structural decomposition: §21.1 identifies the four-state transition, §21.2 the rate formula, §21.3 the geometric mechanism, §21.4 the stable cells, §21.5 the Gaze-aligned stages, §21.6 the structural boundary, §21.7 the information-preservation question, and §21.8 the cosmological consequence.

### 21.1 Decoherence as a four-state transition

**Proposition 21.1** (State-transition reading of decoherence). *Quantum decoherence is the manifold-state transition*

$$\{P, D\} \longrightarrow \{D, T\} \longrightarrow \{P, D, T\},$$

*in which the pre-measurement superposition is the unsubstantiated  $\{P, D\}$  configuration of Definition 2.18, the decoherence-in-progress regime is the Mediation  $\{D, T\}$  configuration with  $T$  engaging environmental descriptors, and the post-measurement classical outcome is the substantiated Exception  $\{P, D, T\}$ . The forbidden  $\{P, T\}$  state is structurally avoided throughout.*

*Proof.* By Definition 2.18, the four manifold states partition into three coherent configurations  $\{P, D\}$ ,  $\{D, T\}$ ,  $\{P, D, T\}$  and one forbidden incoherence  $\{P, T\}$ . The pre-measurement superposition has  $D$ -content (the basis-state structure of the Hilbert space) bound to  $P$ -substrate (the configuration space) but no  $T$ -binding to a single eigenstate; this is the unsubstantiated  $\{P, D\}$  configuration. Environmental coupling engages  $T$  across system-and-environment descriptors, producing the Mediation  $\{D, T\}$  configuration in which  $T$  is binding without yet having selected a single eigenstate; corpus assignment in *ET.Descriptor.D.Paper* §10.3 identifies  $\{D, T\}$  Mediation as “quantum decoherence in progress, photons in transit, chemical transition states.” At completion,  $T$  has substantiated one configuration and the state becomes  $\{P, D, T\} = E$  with variance  $V(E) = 0$  (Proposition 2.26). The forbidden  $\{P, T\}$  state cannot appear because it represents a configuration with no  $D$ -content and would correspond to a measurement outcome with no descriptor resolution — structurally impossible.  $\square$   $\square$

*Remark 21.2* (Born rule from  $\{P, T\}$ -forbidden). The structural impossibility of  $\{P, T\}$  outcomes is the lattice-level statement of the Born rule constraint that measurement outcomes must be definite descriptor-resolved events. Equivalently, no measurement produces an incoherent outcome; every outcome carries a  $D$ -label. This is a Subsumption Law identification (§3.6, condition 3): the Born rule’s constraint that measurement outcomes are eigenstates of the measured observable is recovered as a corollary of the manifold-state classification.

### 21.2 The decoherence rate $R = \Gamma(T \circ D_{\text{env}})^2$

**Proposition 21.3** (Decoherence rate). *The instantaneous decoherence rate of a quantum system coupled to an environment with descriptor field  $D_{\text{env}}$  has the structural form*

$$R = \Gamma(T \circ D_{\text{env}})^2, \tag{35}$$

in which  $\Gamma$  is the (dimensionful) coupling strength,  $T \circ D_{\text{env}}$  is the composition of the Traverser primitive with the environmental descriptor structure, and the squaring is forced by the complex-lattice realisation of  $T$ 's operational manifold ( $U(1)$ , Proposition 5.5).

*Proof.* The Mediation state  $\{D, T\}$  requires  $T$ -binding to environmental descriptors  $D_{\text{env}}$ ; the rate of progress through Mediation toward Exception is proportional to the magnitude of the  $T \circ D_{\text{env}}$  composition. The complex-lattice extension of §11 realises  $T$ -amplitudes as complex-valued objects on  $U(1)$  (Proposition 5.5), and the conversion of complex amplitudes to physical rates is the squared-magnitude  $|T \circ D_{\text{env}}|^2$ . The dimensionful coupling constant  $\Gamma$  converts the dimensionless squared amplitude to a rate. Equation (35) follows.  $\square$   $\square$

*Remark 21.4* (Reclamation of standard decoherence rates). Equation (35) subsumes the standard Joos–Zeh decoherence rate formulas (e.g.,  $\Gamma_{\text{dec}} \sim \Lambda^2 k_B T / \hbar$  for thermal-collisional environments and  $\Gamma_{\text{dec}} \sim G^2 \rho / \hbar$  for graviton baths). Each standard formula is a specific identification of  $T \circ D_{\text{env}}$  for the relevant environment, with  $\Gamma$  absorbing the dimensional prefactors. The squared structure  $(T \circ D_{\text{env}})^2$  is the structural origin of the universal squared-coupling form that runs through every standard decoherence-rate expression.

### 21.3 Geometric mechanism: $\alpha$ -rotation in the complex lattice

The complex lattice of §11 parameterises configurations by  $(k_r, k_\theta)$  with the angle  $\alpha = \arctan(k_\theta/k_r)$  measuring the orientation in the complex log-plane. The angle has a structural identification under the three-aspect framework:  $\cos^2 \alpha$  is the  $D$ -fraction (real-axis descriptor weight) and  $\sin^2 \alpha$  is the  $T$ -fraction (imaginary-axis traversal weight).

**Proposition 21.5** (Decoherence as  $\alpha$ -rotation). *Quantum decoherence is the continuous reduction of the angle  $\alpha$  from  $\pi/2$  (pure imaginary-axis dominance,  $T$ -fraction = 1, pure quantum) toward 0 (pure real-axis dominance,  $D$ -fraction = 1, pure classical), with the effective descriptor-gap residual at angle  $\alpha$  given by*

$$|\delta_{\text{eff}}(\alpha)| = |\delta_r| \cos^2 \alpha + |\delta_\theta| \sin^2 \alpha, \quad (36)$$

where  $|\delta_r|$  and  $|\delta_\theta|$  are the real-axis and imaginary-axis cascade residuals (Propositions 13.1, 13.2).

*Proof.* The complex-lattice extension assigns to a configuration of  $\{D, T\}$ -content the orientation angle  $\alpha$  by Compendium §32. The  $D$ -fraction  $\cos^2 \alpha$  and  $T$ -fraction  $\sin^2 \alpha$  partition unity. Under environmental coupling,  $T$ -binding is progressively transferred from the system's internal degrees of freedom (imaginary-axis dominance) to system-environment composite descriptors (real-axis dominance), so  $\alpha$  decreases monotonically. The effective descriptor-gap residual is the convex combination of the real-axis and imaginary-axis residuals weighted by the corresponding fractions, giving equation (36).  $\square$   $\square$

*Remark 21.6* (Quantitative trajectory at base  $N = 12$ ). At base  $N = 12$  with  $|\delta_r| \approx 0.0196$  and  $|\delta_\theta| \approx 0.2234$  (Propositions 13.1, 13.2), the effective descriptor-gap residual along the decoherence trajectory shrinks by a factor of approximately 11.4 as  $\alpha$  rotates from  $\pi/2$  to 0, going from  $|\delta_{\text{eff}}| = 0.2234$  (pure quantum) to  $|\delta_{\text{eff}}| = 0.0196$  (pure classical). The  $T$ -event density per unit time on the trajectory drops by the same factor: pure-quantum regimes have  $\sim 11\times$  higher  $T$ -event density than pure-classical regimes. This is the lattice-side expression of the standard observation that quantum systems exhibit indeterminacy in proportion to the breadth of their unsubstantiated  $T$ -binding.

*Remark 21.7* (Sample  $\alpha$ -trajectory profile). At representative angles along the decoherence trajectory at base  $N = 12$ , the descriptor-fraction split and effective residual are (with  $|\delta_{\text{eff}}|$  in cents, truncated):

| $\alpha$ (deg) | $D$ -fraction $\cos^2 \alpha$ | $T$ -fraction $\sin^2 \alpha$ | $ \delta_{\text{eff}} $ (cents) | Regime                                  |
|----------------|-------------------------------|-------------------------------|---------------------------------|-----------------------------------------|
| 90             | 0.000                         | 1.000                         | 22.34                           | Pure quantum (imaginary-axis dominance) |
| 75             | 0.067                         | 0.933                         | 20.97                           | Quantum-dominated                       |
| 45             | 0.500                         | 0.500                         | 12.15                           | Schrödinger-cat regime                  |
| 30             | 0.750                         | 0.250                         | 7.05                            | Decoherence in progress                 |
| 15             | 0.933                         | 0.067                         | 3.32                            | Mostly classical                        |
| 0              | 1.000                         | 0.000                         | 1.96                            | Pure classical (real-axis dominance)    |

The factor  $\sim 11.4$  between the pure-quantum and pure-classical residuals (Remark 21.6) is the multiplicative ratio  $|\delta_\theta|/|\delta_r| = 0.2234/0.0196 \approx 11.40$ . Per the structural identification of §11, the imaginary axis is  $T$ 's operational domain and the real axis is  $D$ 's operational domain; the  $\alpha$ -rotation is the continuous reduction of  $T$ -fraction in favour of  $D$ -fraction under environmental coupling.

*Remark 21.8* (Cascade-tower projection of decoherence times). The decoherence time  $\tau_{\text{dec}}$  of any quantum system is a positive real with the dimension of time and admits the lattice projection  $\tau_{\text{dec}}/t_P$  at any resolution  $N$ , where  $t_P$  is the Planck time (the cosmological tower's time-scale reference; §5.7). Sample projections at base  $N = 12$ , with  $\tau_{\text{dec}}$  values from canonical decoherence-theory references and  $|\varepsilon|$  truncated (not rounded), are:

| System                          | $\tau_{\text{dec}}$ (s) | $\tau_{\text{dec}}/t_P$ | $k_r$ | $d_r$ |
|---------------------------------|-------------------------|-------------------------|-------|-------|
| Cold atom ( $\mu\text{K}$ trap) | $10^{-3}$               | $1.86 \times 10^{40}$   | +1605 | 4     |
| Free electron in air            | $10^{-10}$              | $1.86 \times 10^{33}$   | +1326 | 2     |
| Large molecule                  | $10^{-20}$              | $1.86 \times 10^{23}$   | +928  | 3     |
| $10\mu\text{m}$ dust at STP     | $10^{-36}$              | $1.86 \times 10^7$      | +290  | 6     |
| 1mm bacterium                   | $10^{-42}$              | 18.55                   | +51   | 4     |
| Schrödinger cat                 | $10^{-50}$              | $1.86 \times 10^{-7}$   | −268  | 3     |

Different physical decoherence times sit at different sublattice cells; each  $\tau_{\text{dec}}$  is a lossless lattice point at base  $N = 12$  (Theorem 19.4). At higher LCM-tower resolutions the cascade visits the shadow-force families progressively in accordance with Proposition 5.28 applied to the ratio  $\tau_{\text{dec}}/t_P$ , with structurally identical machinery to the Hawking-thermal cascade of §5.11. The General Cascade Rule (Theorem 13.8) governs the visitation profile of every such cascade.

## 21.4 Pointer states as high-elegance lattice cells

**Proposition 21.9** (Pointer states are lattice-stable cells). *Pointer states (Zurek einselection) are configurations that survive environmental coupling without displacement. Under the lattice classification, a configuration is pointer-stable if and only if it occupies a lattice cell of high structural elegance, equivalently low descriptor-gap residual  $|\varepsilon|$  and low sublattice family  $d$  at the relevant resolution.*

*Proof.* A configuration is pointer-stable if  $T$ -binding to environmental descriptors does not displace it from its lattice cell. Displacement requires that the environment-coupled trajectory crosses a cell boundary (where  $|\varepsilon|$  approaches 50 cents and rounding becomes ambiguous; see §21.6 below). A configuration sitting on a high-elegance cell ( $|\varepsilon| \rightarrow 0$  and  $d$  small) has the maximum margin to any cell boundary and is therefore the most robust against environmental displacement. The structurally-favoured pointer-state cells at  $N = 12$  are  $d \in \{1, 2, 4, 12\}$ :  $d = 1$  (gravity / cascade closure) for position eigenstates,  $d = 2$  (Mediation pivot) for spin in canonical bases,  $d = 4$  (quartic / weak-channel) for spin along measurement axes, and  $d = 12$  (full electromagnetic resolution) for EM-class observables.  $\square$   $\square$

*Remark 21.10* (Why position eigenstates are robust). The cell  $d = 1$  is the gravity / cascade-closure cell of §5.7, the unique sublattice whose lattice point coincides with the unison ratio  $1/1$ . Position eigenstates inherit pointer-stability from this gravitational-cell identification: the  $d = 1$  family has the highest structural elegance of any cell at  $N = 12$ , and configurations attached to position-as-substrate inherit this elegance. The standard observation that environmental decoherence rapidly localises quantum systems in position basis is the lattice-side statement that the position basis maps onto the most stable cell at  $N = 12$ .

## 21.5 Gaze thresholds as decoherence stages

The Complete Gaze Equation of §20 classifies any observation by its  $F_w$  value into four status bands (UNOBSERVED, SUBLIMINAL, DETECTED, LOCKED) with thresholds at  $F_w \in \{1, 13/12, 6/5, 3/2\}$ . Under the  $\alpha$ -rotation reading of §21.3, the Gaze thresholds align with structurally distinguished stages of the decoherence trajectory.

**Proposition 21.11** (Gaze thresholds are decoherence stages). *The four Gaze-status bands of §20 correspond bijectively to four stages of the  $\alpha$ -rotation decoherence trajectory of Proposition 21.5, characterised at each stage by a specific lattice projection at  $N = 12$ :*

| <i>Status</i> | <i><math>F_w</math> threshold</i> | <i><math>(k_r, d_r, \varepsilon_r)</math> at <math>N = 12</math></i> | <i><math>\alpha</math>-stage characterisation</i> |
|---------------|-----------------------------------|----------------------------------------------------------------------|---------------------------------------------------|
| UNOBSERVED    | $F_w < 1$                         | $(0, 1, 0)$ ( <i>below baseline</i> )                                | $\alpha = \pi/2$ ( <i>pure quantum</i> )          |
| SUBLIMINAL    | $F_w \geq 13/12$                  | $(+1, 12, +38.57 \text{ cents})$                                     | $\alpha$ <i>boundary-near</i>                     |
| DETECTED      | $F_w \geq 6/5$                    | $(+3, 4, +15.64 \text{ cents})$                                      | $\alpha$ <i>mid-trajectory</i>                    |
| LOCKED        | $F_w \geq 3/2$                    | $(+7, 12, +1.955 \text{ cents})$                                     | $\alpha \rightarrow 0$ ( <i>pure classical</i> )  |

The ordered sequence  $UNOBSERVED \rightarrow SUBLIMINAL \rightarrow DETECTED \rightarrow LOCKED$  is the bijective image of the  $\alpha$ -rotation decoherence trajectory, with each Gaze threshold a structurally distinguished waypoint along the rotation from  $\alpha = \pi/2$  to  $\alpha \rightarrow 0$ .

*Proof.* The Gaze projections at  $N = 12$  are read directly from §20 (Table after the Status definition); the lattice cells given are the projections of  $1/1$ ,  $13/12$ ,  $6/5$ , and  $3/2$  at  $N = 12$ , with descriptor-gap residuals  $\{0, +38.57, +15.64, +1.955\}$  cents respectively (truncated, not rounded).

The bijection with  $\alpha$ -rotation stages is established by reading the descriptor-gap residual as the stage-marker: at  $\alpha = \pi/2$  (pure quantum) the trajectory has not yet started and sits at the unison cell; under environmental coupling the residual increases monotonically (toward the boundary at  $|\varepsilon| = 50$  cents) until  $\alpha$  reaches the SUBLIMINAL threshold; further coupling reduces  $\alpha$  toward DETECTED, where the residual drops to mid-cell (+15.64 cents); LOCKED corresponds to  $\alpha$  approaching 0 with the residual now small (+1.955 cents), the pre-classical pythagorean-comma cell. The monotone correspondence is enforced by the structural identity  $\alpha_{\text{Gaze}} = \pi/2 \cdot (1 - F_w/F_w^{\text{LOCKED}})$  when  $F_w \leq 3/2$ , with  $\alpha_{\text{Gaze}}$  the residual angle in the  $\alpha$ -rotation parametrisation. The four Gaze thresholds are therefore the four structurally-distinguished waypoints of the  $\alpha$ -rotation decoherence trajectory at  $N = 12$ .  $\square$   $\square$

*Remark 21.12* (Quintic gating of awareness). The span from SUBLIMINAL to LOCKED is  $3/2 - 13/12 = 5/12 = 5V_{\text{base}}$ , exactly five  $V$ -quanta (§20, sentence following the Status definition). Equivalently, the awareness window is structurally five quintic units wide, with the prime 5 governing its extent. The corresponding decoherence-trajectory waypoints are therefore separated by exactly five quintic structural increments, and a system traverses these in five quintic-paced stages from subliminal detection to locked measurement. This is the lattice-side derivation of the empirical observation that awareness ramp-up under stimulus is staged rather than continuous.

*Remark 21.13* (Quartic-channel role of the DETECTED threshold). The DETECTED threshold  $6/5$  projects to the  $d = 4$  quartic sublattice at  $N = 12$  (the only Gaze threshold at a non- $d = 12$  cell). Under the Identification Principle (§3.4) the  $d = 4$  family carries the weak-channel / four-fold structural character (§12.1). The DETECTED stage of decoherence is therefore the stage at which the system's  $T$ -binding has migrated from the imaginary-axis-dominant pure-quantum regime through a quartic-symmetric intermediate cell before locking into the LOCKED  $d = 12$  post-measurement cell. This identifies the structural reason for the quartic gating: pre-measurement attention selection has the same symmetry structure as the weak-channel sublattice, and selective attention is the lattice-side expression of weak-channel intermediate substantiation.

## 21.6 Coherence boundary at $|\varepsilon| = 50$ cents: measurement uncertainty

**Proposition 21.14** ( $|\varepsilon| = 50$  cents as the coherence boundary). *The lattice projection  $k_r = \text{round}(N \log_2 r)$  is well-defined for descriptor-gap residual  $|\varepsilon| < 50$  cents and structurally ambiguous at  $|\varepsilon| = 50$  cents exactly, where rounding to either of two adjacent integers  $k$  is equally valid. The  $|\varepsilon| = 50$  cents boundary is therefore the lattice-side statement of measurement uncertainty: at the boundary,  $T$  cannot resolve to a unique sublattice cell.*

*Proof.* By Definition 7.1,  $k_r = \text{round}(N \log_2 r)$  where the rounding convention places  $\varepsilon \in (-50, +50]$  cents per Proposition 10.3. At the boundary  $\varepsilon = +50$  (or  $-50$  by mirror symmetry) the rounding decision between the two adjacent integers is structurally undecidable; the configuration sits exactly midway between two lattice cells. By the Identification Principle the act of  $T$ -resolution is the act of selecting one cell; at the boundary this act is ambiguous. The coherence boundary  $|\varepsilon| = 50$  cents is therefore the structural ceiling on measurement precision at any resolution.  $\square$   $\square$

*Remark 21.15* (Coherence time bound). The coherence time  $\tau_{\text{coh}}$  of a quantum system under environmental coupling is bounded by the time required for the trajectory to traverse the descriptor-gap residual range from a starting cell-centre ( $|\varepsilon| = 0$ ) to a cell-boundary ( $|\varepsilon| = 50$ ). This bound is independent of the specific environmental mechanism — it is a structural property of the lattice, not a postulate of quantum theory. The bound predicts that coherence times saturate at a structurally-determined maximum for any system whose lattice trajectory under environmental coupling is known.

## 21.7 Information preservation through the decoherence boundary

The information question for quantum decoherence is structurally identical to the black-hole information question of §5.6: a system's coherent information appears to be lost into an environment, but the joint system-environment state preserves total  $T$ -event count.

**Corollary 21.16** (Decoherence-boundary information preservation). *The decoherence event  $\{\text{System}_{\text{parent}}\} \rightarrow \{\text{boundary}\} \rightarrow \{\text{Environment}_{\text{child}}\}$  is an instance of the Multifold birth-triad mechanism (Definition 5.17) at the system-environment scale. Information is preserved by  $T$ -event conservation across the joint system-environment state, by the same mechanism as the black-hole case (Proposition 5.18).*

*Proof.* The decoherence boundary plays the structural role of the seed  $R_0$  of Definition 5.17: it transmits the system's pre-decoherence descriptor content into the environmental descriptor field. By the binding-order axiom (Axiom 2.17), each  $T$ -event is associated with exactly one  $\{P, D, T\}$  substantiation; the joint system-environment  $T$ -event count is an additive invariant. Information is conserved through the decoherence event as the environmental entanglement entropy increases by exactly the amount of  $T$ -binding transferred from the system. This recovers the standard observation that decoherence is unitary on the joint state, as a consequence of  $T$ -event conservation rather than as an independent postulate.  $\square$   $\square$

## 21.8 The cosmological M-state budget

The fraction of universal energy held in active Mediation  $\{D, T\}$ -state configurations at any instant is a structural invariant of the cosmological tower.

*Remark 21.17* (Active-mediation cosmological budget). The total universal energy in  $\{D, T\}$ -state configurations (active mediation) is approximately 3.0%, partitioned into M-vacuum (vacuum decoherence, virtual-particle mediation, zero-point fluctuation transitions) at 1.6% and M-matter (photons in flight, chemical reactions, biological metabolism, in-progress wavefunction collapse) at 1.4%. The ratio  $1.6/1.4 = 8/7$  is exact under the assignment of M-vacuum and M-matter to their respective sublattice-family content (*ET\_Fine\_Structure\_Constant\_REVISED*, M-states budget). The 3% ceiling and the 8:7 ratio constitute a cosmological constraint: any change in cosmic-scale decoherence rates must respect both. Deviations would constitute observable structural violations.

## 22 Empirical confrontation

We now confront the lattice with empirical data from five disciplines. Each subsection identifies the substrate, the reference period, and the resulting lattice projection, and compares the lattice-derived structural reading with the discipline-internal description.

### 22.1 Twelve-tone equal temperament

The twelve-tone equal-tempered scale is the lattice  $\mathcal{L}_{12}$  restricted to one octave: the frequency ratios  $\{2^{k/12} : k = 0, 1, \dots, 12\}$ . Each of the twelve ratios is a canonical projection; six of them lie exactly on the lattice ( $d = 1, 2$ ) and the remaining six lie at tight descriptor gaps that identify their sublattice character (Table 13).

| Interval       | Just ratio $r$ | $12 \log_2 r$ | $k$ | $d$ | $\varepsilon$ (cents) |
|----------------|----------------|---------------|-----|-----|-----------------------|
| Unison         | 1/1            | 0.000         | 0   | 1   | 0.00                  |
| Major second   | 9/8            | 2.039         | 2   | 6   | +3.91                 |
| Minor third    | 6/5            | 3.156         | 3   | 4   | +15.64                |
| Major third    | 5/4            | 3.863         | 4   | 3   | -13.69                |
| Perfect fourth | 4/3            | 4.980         | 5   | 12  | -1.96                 |
| Tritone        | $\sqrt{2}$     | 6.000         | 6   | 2   | 0.00                  |
| Perfect fifth  | 3/2            | 7.020         | 7   | 12  | +1.96                 |
| Minor sixth    | 8/5            | 8.137         | 8   | 3   | +13.69                |
| Major sixth    | 5/3            | 8.844         | 9   | 4   | -15.64                |
| Minor seventh  | 16/9           | 9.961         | 10  | 6   | -3.91                 |
| Major seventh  | 15/8           | 10.883        | 11  | 12  | -11.73                |
| Octave         | 2/1            | 12.000        | 12  | 1   | 0.00                  |

Table 13: The twelve just-intoned intervals and their projections at  $N = 12$ . The perfect fifth and perfect fourth sit at  $d = 12$  with descriptor gap  $\pm 1.955$  cents, the Pythagorean comma. The thirds sit at  $d = 3$  (major) and  $d = 4$  (minor). The tritone sits exactly on the lattice at  $d = 2$ .

The conventional view of twelve-tone equal temperament as a historical compromise is replaced by a derivation: the 12-division is forced by  $N = |\Pi| \cdot S = 12$ , and the characteristic cent residuals of the just intervals are the descriptor gaps of the respective sublattices.

## 22.2 The fine-structure constant

The fine-structure constant  $\alpha$  satisfies  $\alpha^{-1} \approx 137.0359991\dots$  in modern measurement. The leading-order lattice-derived value is

$$A_0 = (N - 1)^2 + S^2 = 137$$

(Equation 17). The residual  $\alpha^{-1} - A_0 \approx 0.036$  admits a closed-form decomposition in terms of the manifold constants,

$$\alpha^{-1}(\text{ET}) = 137 + \frac{\sqrt{3}}{48} - \frac{\sqrt{3}}{93312\pi^2} - \frac{1}{216(18\pi - 1)}, \quad (37)$$

whose component terms have structural readings as follows:

- The leading correction  $A_1 = \sqrt{3}/48 = \sigma/K_{\text{EM}}$  with  $\sigma = \sqrt{1/12}$  and  $K_{\text{EM}} = 8$  is an *open shimmer* over electromagnetic channels: a linear  $T$ -path along a diameter-like trajectory across the  $\partial I$  boundary, contributing positively (open paths  $k < 3/2$  enter with positive sign, closed/semi-closed paths  $k \geq 3/2$  enter with negative sign, by Definition 2.23).
- The cross term  $A_{\text{cross}} = \sqrt{3}/(93312\pi^2) = (2/\pi) \cdot A_1 \cdot A_2$  with  $A_2 = \kappa^2/(N^3\pi)$  is the product interference of the shimmer (open  $A_1$ ) with the closed bilateral-mediation loop (closed  $A_2$ ,  $k = 2$ ), with the geometric factor  $2/\pi$  arising from the bilateral-to-circumferential phase conversion (the structural mean of an open path's projection along a closed loop's axis, equal to  $\int_0^\pi \sin \theta d\theta/\pi = 2/\pi$ ). The closed factor enters with subtractive sign.
- The geometric series  $1/(216(18\pi - 1)) = \sum_{k \geq 2} \kappa^k/(N^{k+1}\pi^{k-1}) = \kappa^2/[N^2(N\pi - \kappa)]$  is the closed-form summation of the higher closed-mediation loops, with  $k$ -th term carrying  $k$  Koide vertices, manifold-volume suppression  $N^{-(k+1)}$ , and  $(k - 1)$  phase integrations  $\pi^{-(k-1)}$ .

*Remark 22.1* (Structural origin of the  $A_{\text{cross}}$  factors). The cross-term  $A_{\text{cross}} = 2\sigma\kappa^2/(K_{\text{EM}}N^3\pi^2)$  has every factor traceable to  $A_1$  or  $A_2$  or to canonical geometric inheritance: the 2 is the bilateral symmetry of the two vertices on the closed loop (inherited from  $A_2$ );  $\sigma$  is the shimmer amplitude (inherited from  $A_1$  numerator);  $\kappa^2$  is the two Koide couplings at the loop vertices (inherited from  $A_2$  numerator);  $K_{\text{EM}}$  is the EM channel distribution (inherited from  $A_1$  denominator);  $N^3$  is the bilateral-loop manifold volume (inherited from  $A_2$  denominator);  $\pi^2$  is the product of two phase integrations, one from  $A_2$ 's closed loop and one from the bilateral-to-circumferential geometric conversion factor  $2/\pi$ . No new ad-hoc element is introduced; the cross-term is forward-derivable from the two pre-existing structural terms with one canonical geometric factor.

Numerical evaluation to 50 decimal digits yields

$$\alpha^{-1}(\text{ET}) = 137.03599916744133748324548\dots,$$

which agrees with the CODATA 2022 recommended value  $\alpha_{\text{exp}}^{-1} = 137.035\,999\,177(21)$  [7] (relative uncertainty  $1.6 \times 10^{-10}$ ) to within  $|\alpha^{-1}(\text{ET}) - \alpha_{\text{exp}}^{-1}| \approx 9.6 \times 10^{-9}$  — approximately  $0.46\sigma$  on the CODATA scale, or  $\sim 7$  parts in  $10^{11}$  of  $\alpha^{-1}$ . The two leading direct atom-interferometry determinations carry a known  $5\sigma$  mutual tension: Parker et al. 2018 (Berkeley Cs) [8] reports  $\alpha^{-1} = 137.035\,999\,046(27)$  ( $+4.5\sigma$  above the ET value), Morel et al. 2020 (LKB Rb) [9] reports  $\alpha^{-1} = 137.035\,999\,206(11)$  ( $-3.5\sigma$  below). The CODATA 2018 adjustment [6] gave  $\alpha^{-1} = 137.035\,999\,084(21)$  ( $+4.0\sigma$ ); the CODATA 2022 adjustment cited above is the first least-squares synthesis whose central value falls within one  $\sigma$  of equation (37). The ET formula sits within the experimental window spanned by Parker–Morel and at the centre of the CODATA 2022 confidence interval, with no free parameters. At the universal lattice  $N = 27720$ , the projection of this value lands at  $(k, d) = (196768, 315)$  with  $|\varepsilon| \approx 0.002$  cents; the sublattice family  $d = 315 = 3^2 \cdot 5 \cdot 7$  carries simultaneous cubic-squared, quintic, and septic structural character, which is physically appropriate for a coupling constant governing electromagnetic interaction with the three-generation lepton spectrum.



*Remark 22.2* (Manifold resolution floor). The fundamental theoretical resolution of equation (37) at base  $N = 12$  is

$$\delta_{\text{manifold}} = \frac{\sigma}{K_{\text{EM}} N^5} \approx 1.45 \times 10^{-7},$$

the structural floor below which 12ET arithmetic cannot resolve algebraic differences. The sub-ppb-level disagreements among contemporary measurements of  $\alpha^{-1}$  (the  $> 5\sigma$  tension between Morel 2020 Rb (Paris, LKB) [9] and Parker 2018 Cs (Berkeley) [8] atom-interferometry determinations) sit well within  $\delta_{\text{manifold}}$ , below the base-lattice resolution floor. Sub-ppb structural features of  $\alpha^{-1}$  require projection onto higher LCM-tower resolutions (the natural next being  $N = 2520$  or  $N = 27720$ , where the cross-complex sublattice  $d = 315$  resolves to sub-millicent precision). Equation (37) is the base-resolution closed form; the higher-tower refinements are structural enhancements of it, not corrections.

### 22.3 The Koide ratio

The Koide formula evaluated at modern lepton masses  $(m_e, m_\mu, m_\tau) = (0.51099895, 105.6583755, 1776.86)$  [7] gives

$$Q = 0.6666605 \pm 0.000002.$$

The ET-derived value is  $K = 2/3 = 0.666666\bar{6}$  (Equation 16), with residual of approximately 6 parts per million. Both  $2/3$  and  $3/2$  project to the Koide attractor at  $d = 12$ ,  $|\varepsilon| = 1.955$  cents (Theorem 19.1).

### 22.4 Geometric ratios

The classical 3:4:5 right triangle has three internal ratios whose projections are shown in Table 14.

| Ratio                        | Value | $12 \log_2 r$ | $k$ | $d$ | $\varepsilon$ (cents) |
|------------------------------|-------|---------------|-----|-----|-----------------------|
| 3/4 (short leg : long leg)   | 0.750 | -4.980        | -5  | 12  | +1.96                 |
| 4/5 (long leg : hypotenuse)  | 0.800 | -3.863        | -4  | 3   | +13.69                |
| 3/5 (short leg : hypotenuse) | 0.600 | -8.844        | -9  | 4   | +15.64                |

Table 14: Projections of the internal ratios of the 3:4:5 triangle. The three ratios populate sublattices  $d = 12$ ,  $d = 3$ ,  $d = 4$  respectively. These are the three non-trivial simple sublattice families at  $N = 12$ .

The Pythagorean triangle thus occupies exactly three different sublattice families at the base lattice, one of each non-trivial simple family besides  $d = 2$  and  $d = 6$ .

Bond angles in chemistry admit two simultaneous lattice readings, each capturing a distinct structural feature of the configuration. The angle ratio  $r_\theta = \theta/180^\circ$  encodes the symmetry order of the bond geometry; the cosine  $r_{\cos} = |\cos \theta|$  encodes the rational invariant that makes the angle definable in closed form. By the Identification Principle, both are valid Path-C projections of the same underlying configuration, and their joint reading provides the complete structural fingerprint:

The tetrahedral configuration illustrates the structural significance of the dual reading:  $\theta/180^\circ = \arccos(-1/3)/\pi$  is transcendental, projecting to  $d = 4$  (correctly recognising the 4-fold tetrahedral symmetry) but with  $|\varepsilon| = 39.07\text{¢}$  in the Twilight Zone of the Stability Window;  $|\cos \theta| = 1/3$  is rational, projecting exactly to the Koide attractor at  $d = 12$ ,  $|\varepsilon| = 1.955\text{¢}$  — the same lattice address as the Sempaevum’s own defining constants (Theorem 19.1). The angle reading classifies the symmetry; the cosine reading classifies the lattice address. Both are forced; neither is preferred. The same dual structure is exhibited by the trigonal-planar configuration, where

| Geometry                           | $\theta$        | $\theta/180^\circ$ projection    | $ \cos \theta $ | $ \cos \theta $ projection        |
|------------------------------------|-----------------|----------------------------------|-----------------|-----------------------------------|
| Linear (sp)                        | $180^\circ$     | (0, 1, 0) unison                 | 1               | (0, 1, 0) unison                  |
| Trigonal-planar (sp <sup>2</sup> ) | $120^\circ$     | (−7, 12, −1.96 $\phi$ )<br>Koide | 1/2             | (−12, 1, 0) octave                |
| Tetrahedral (sp <sup>3</sup> )     | $\arccos(-1/3)$ | (−9, 4, +39.07 $\phi$ ) d=4      | 1/3             | (−19, 12, −1.96 $\phi$ )<br>Koide |
| Right-angle (oct.)                 | $90^\circ$      | (−12, 1, 0) octave               | 0               | $\partial I$ annihilation         |

Table 15: Dual lattice readings for the principal bond geometries. The angle-ratio reading exposes the symmetry order (d=4 for tetrahedral correctly identifies its 4-fold structure); the cosine reading exposes the structurally tighter rational invariant (tetrahedral lands exactly on the Koide attractor at  $d = 12, |\varepsilon| = 1.955 \phi$ ). The right angle is the unique geometry whose cosine reading hits the  $\partial I$  boundary at  $r = 0$ , consistent with orthogonal closure being structurally degenerate (D-only, no traversal coupling).

the angle reading  $r_\theta = 2/3$  lands on the Koide attractor and the cosine reading  $r_{\cos} = 1/2$  lands on the  $d = 1$  octave; in both cases the two readings populate distinct Simple sublattice families ( $d \in \{1, 12\}$ ), consistent with bond geometry’s status as a doubly-classified configuration.

## 22.5 Celestial mechanics

The principal mean-motion resonances of the Solar System project cleanly onto low- $d$  lattice attractors:

- Neptune–Pluto 3:2:  $\Pi_{12}(3/2) = (+7, 12, +1.96)$  — the Koide attractor.
- Io–Europa 2:1:  $\Pi_{12}(2) = (+12, 1, 0)$  — the octave.
- Galilean moons, Europa–Ganymede 2:1: same as Io–Europa.
- Saturn–Jupiter period ratio  $\approx 2.48$ , near 5/2:  $\Pi_{12}(5/2) = (+16, 3, -13.69)$  —  $d = 3$  cubic with the major-third descriptor gap.

Gravitational binding, which selects minimum-variance configurations, systematically targets low- $d$  sublattices; the lattice structure is consistent with the empirical distribution of resonances.

## 22.6 Biochemistry

The step counts of major metabolic cycles are direct positive integers (denominator = 1 step):

- Citric acid (Krebs) cycle: 8 steps.  $\Pi_{12}(8) = (+36, 1, 0)$  — exact  $d = 1$  octave class. The step count is  $2^3$ .
- Urea cycle: 4 core steps.  $\Pi_{12}(4) = (+24, 1, 0)$  — exact  $d = 1$ .
- Glycolysis: 10 steps.  $\Pi_{12}(10) = (+40, 3, -13.69)$  —  $d = 3$  cubic.
- ATP synthase rotational ring: typically 10  $c$ -subunits. Same projection as glycolysis.
- Cell cycle: 4 phases (G1, S, G2, M).  $\Pi_{12}(4) = (+24, 1, 0)$ .

The “closure” cycles (Krebs, urea) project to  $d = 1$  with  $\varepsilon = 0$ ; their step counts are powers of 2. The “linear pathways” (glycolysis, purine synthesis) project to  $d = 3$  with the major-third gap; their step counts are multiples of 2 and 5. The distinction between closure and linear pathway is reflected in distinct sublattice family assignments.

## 22.7 Optical phase singularities

Bucher et al. [17] measure the ultrafast dynamics of optical phase singularity ensembles in phonon-polariton fields in hexagonal boron nitride (hBN), achieving deep sub-wavelength spatial ( $20\text{ nm} \approx \lambda_{\text{hBP}}/30$ ) and deep sub-cycle temporal ( $3\text{ fs} \approx \tau/8$ ) resolution. The Identification Principle applied to wave propagation in the hyperbolic medium gives:

| Primitive | Identification in the hBN phonon-polariton system                                                                                                                    |
|-----------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| $P$       | The spatial substrate of the crystal — the hBN lattice                                                                                                               |
| $D$       | The dispersion relation $\omega(\mathbf{k})$ — the finite constraint governing wave propagation                                                                      |
| $T$       | The wave field's phase — the $U(1)$ -valued quantity at each point; $T$ -agency is the topological dynamics (singularity creation, annihilation, $\pm 2\pi$ winding) |

The data yield four lattice-addressable quantities.

### Phase singularities as $\partial I$ boundary configurations

Phase singularities are points of zero amplitude and undefined phase. At the singularity core,  $D$ -content (amplitude on  $(\mathbb{R}^+, \times)$ ) vanishes and  $T$ -content (phase on  $(U(1), \times)$ ) is undefined, placing the configuration in the  $\{P, T\}$  state (Incoherence, Definition 2.18). The  $\pm 2\pi$  winding of  $T$  around the core exists in the neighbourhood, not at the core itself — this is the Mediation Problem (Theorem 3.34) made physical:  $P$  and  $T$  cannot bind without  $D$ 's finite structure. All six structural properties of  $\partial I$  boundary configurations (§14) are satisfied.

### The 1/3 velocity-tail fraction as $T$ -weight

The Berry–Dennis velocity distribution  $P_{\pm}(|v|) = 8\langle v \rangle^2 |v| / (8|v|^2 + 4\langle v \rangle^2)^2$  has the property that exactly 1/3 of singularities exceed the mean velocity  $\langle v \rangle$ . The substitution  $w = 2u^2 + 1$  (with  $u = |v|/\langle v \rangle$ ) maps the threshold  $u = 1$  to  $w = 3 = |P, D, T|$ ; the integral from  $w = 3$  to  $\infty$  versus  $w = 1$  to  $\infty$  gives  $1/3 = 1/|P, D, T|$ , the  $T$ -weight. Below the mean: the  $\{P, D\}$ -constrained fraction  $= K = 2/3$ . Above the mean: the unconstrained  $T$ -fraction  $= 1 - K = 1/3$ . The 1/3 is the primitive count appearing in the velocity distribution of  $T$ -content objects.

### Material classification via the Dimensionless Seed Ratio

The natural reference period of a phonon-polariton material is  $R_0 = 1/\omega_{\text{TO}}$  (one TO phonon oscillation period); the Dimensionless Seed Ratio is  $r = \omega_{\text{LO}}/\omega_{\text{TO}}$ . Different materials project to different lattice positions:

| Material                         | $\omega_{\text{LO}}/\omega_{\text{TO}}$ | $k$ | $d$ | $\varepsilon$ (cents) |
|----------------------------------|-----------------------------------------|-----|-----|-----------------------|
| hBN upper band                   | $1610/1370 = 161/137$                   | 3   | 4   | −20.5                 |
| hBN lower band                   | $830/780 = 83/78$                       | 1   | 12  | +7.6                  |
| $\alpha$ -MoO <sub>3</sub> [100] | $972/820$                               | 3   | 4   | −5.6                  |

hBN upper and  $\alpha$ -MoO<sub>3</sub> both sit at  $d = 4$  (quartic) but at different descriptor gaps ( $\varepsilon = -20.5\text{ ¢}$  versus  $-5.6\text{ ¢}$ ); hBN lower band sits at  $d = 12$  — a different sublattice family entirely. The polariton properties of each material, including the phase-to-group velocity ratio, are classified by the Dimensionless Seed Ratio's lattice position. The measured  $v_{\varphi}/v_g \approx 12 \pm 1$  for hBN is the  $T/D$  propagation ratio at the  $d = 4$ ,  $\varepsilon = -20.5\text{ ¢}$  cell;  $\alpha$ -MoO<sub>3</sub> at the same  $d$  but different  $\varepsilon$  produces a different ratio.

### Wavelength compression $\lambda/\lambda_0 \approx 11$ and the $d = 11$ harmonic family

The polariton wavelength compression factor  $\lambda/\lambda_0 \approx 11$  (where  $\lambda = 7\text{ }\mu\text{m}$  free-space and  $\lambda_0 \approx 630\text{ nm}$  polariton) identifies the  $d = 11$  undecimal harmonic family — the last prime in  $N_{\text{FULL}} = \text{lcm}(1, \dots, 11) = 27720$ . Combined with the measured  $v_\varphi/v_g \approx 12 = N$ , the structural chain is:

$$\lambda/\lambda_0 \approx 11 \Rightarrow v_\varphi = c/11, \quad v_\varphi/v_g = N = 12 \Rightarrow v_g = c/(11 \times 12) = c/132.$$

The integer  $132 = \text{lcm}(11, 12) = n_c(d=11)$  is the canonical LCM-tower resolution at which the  $d = 11$  harmonic family first becomes native (Table 6). The group velocity of hBN phonon-polaritons thus equals  $c$  divided by the resolution at which the  $d = 11$  harmonic family first appears natively on the LCM tower. The system operates at the structural junction where the compression factor (11, the last prime of  $N_{\text{FULL}}$ ) and the velocity ratio (12, the manifold symmetry) combine to give  $n_c(11) = 132$ .

The lattice tower projection of the integer 11 confirms the harmonic-family identification:

| Resolution | $k$   | $d$ | $\varepsilon$ (cents) | Note                                    |
|------------|-------|-----|-----------------------|-----------------------------------------|
| 12 ET      | 42    | 2   | −48.68                | $d = 2$ at base (shadow of $d = 11$ )   |
| 132 ET     | 457   | 132 | −3.23                 | $d = 132$ ; involves 11 as prime factor |
| 420 ET     | 1453  | 420 | −0.11                 | First sub-cent                          |
| 27720 ET   | 95895 | 616 | +0.019                | $d = 616 = 2^3 \cdot 7 \cdot 11$        |

At the universal resolution 27720 ET, the integer 11 lands at sublattice  $d = 616 = 2^3 \cdot 7 \cdot 11$ , which contains  $d = 11$  as a prime factor: the compression factor’s lattice home involves the  $d = 11$  harmonic family at terminal resolution.

## 22.8 Summary of the empirical confrontation

Across the five disciplines surveyed, the same six simple sublattice families recur, and a handful of attractor positions are shared across domains: the Koide attractor (+7, 12, +1.955) appears in particle physics, celestial mechanics, and musical intonation; the octave (+12, 1, 0) appears in acoustics, biochemistry, and mass hierarchies; the tritone (+6, 2, 0) appears in geometry and palindromic symmetries. Table 16 summarises selected cross-domain coincidences.

## 22.9 Fundamental forces: full PDT decomposition

Each fundamental force, when subjected to the Identification Principle (§3.4), admits a full  $P \circ D \circ T$  decomposition. No force is exclusively  $P$ ,  $D$ , or  $T$ ; each has components in all three Cardinals.

| Force   | $P$ -content (substrate)  | $D$ -content (structure)    | $T$ -content (agency) |
|---------|---------------------------|-----------------------------|-----------------------|
| Gravity | Spacetime manifold        | Curvature, $G$ , field eqs  | Gravitational agency  |
| EM      | Charge-carrying substrate | Charge, potential, Maxwell  | EM agency             |
| Strong  | Colour-charge substrate   | Colour charge, gluon fields | Strong binding agency |
| Weak    | Isospin substrate         | Isospin, mixing angles      | Weak-decay agency     |

The quantum gravity problem is, in this framework, a missing Descriptor at the interface between quantum  $D$ -structure and gravitational  $D$ -structure: the Descriptor Gap Principle identifies the gap as the Descriptor that bridges the  $d = 1$  (octave, gravitational) sublattice and the quantum sublattices ( $d = 2, 3, 4, 12$ ). The gap is a Descriptor waiting to be identified, not a category error. Early corpus formulations classified gravity as “a Traverser type” exclusively; the mature analysis using the three operational tools gives each force its full  $P \circ D \circ T$  structure. The quantitative sublattice-to-force correspondence — which sublattice families generate which gauge groups, and why — is established in Theorem 9.13 of §9.7.

| Domain    | Quantity                                   | Ratio $r$ | $(k, d)$  | Family |
|-----------|--------------------------------------------|-----------|-----------|--------|
| Music     | Perfect fifth                              | 3/2       | (+7, 12)  | Koide  |
| Particles | Koide ratio $K$                            | 2/3       | (−7, 12)  | Koide  |
| Celestial | Neptune–Pluto resonance                    | 3/2       | (+7, 12)  | Koide  |
| Chemistry | Trigonal-planar bond $120^\circ/180^\circ$ | 2/3       | (−7, 12)  | Koide  |
| Manifold  | Manifold symmetry $N$                      | 12        | (+43, 12) | Koide  |
| Manifold  | Base variance $V$                          | 1/12      | (−43, 12) | Koide  |
| Music     | Octave                                     | 2/1       | (+12, 1)  | Octave |
| Biology   | Krebs cycle step count                     | 8         | (+36, 1)  | Octave |
| Biology   | Cell-cycle phases                          | 4         | (+24, 1)  | Octave |
| Markets   | Doubling                                   | 2/1       | (+12, 1)  | Octave |
| Celestial | Io–Europa resonance                        | 2/1       | (+12, 1)  | Octave |
| Music     | Major third                                | 5/4       | (+4, 3)   | Cubic  |
| Geometry  | Pythagorean ratio 4/5                      | 4/5       | (−4, 3)   | Cubic  |
| Biology   | Glycolysis, ATP ring                       | 10        | (+40, 3)  | Cubic  |
| Celestial | Saturn–Jupiter period                      | 5/2       | (+16, 3)  | Cubic  |

Table 16: Cross-domain coincidences: distinct physical quantities from different disciplines projecting to the same sublattice family, often to the same lattice point. The clustering is the quantitative content of the regularity noted in §1.1.

## 22.10 The QED $\beta$ -function in ET terms

The 1-loop QED  $\beta$ -function coefficient for each charged lepton is

$$\beta_0(\text{QED}) = -\frac{2}{3\pi} = -\frac{d_2}{|\Pi| \cdot \pi},$$

where  $d_2 = 2$  (the binary sublattice, the Mediation worldsheet),  $|\Pi| = 3$  (the primitive count), and  $\pi = \frac{1}{2} \cdot 2\pi$  (the half-period of  $T$ 's  $U(1)$  manifold). Every factor has a direct ET identification. The running coupling  $\alpha(\mu) = \alpha_0/(1 - (2\alpha_0/(3\pi)) \ln(\mu/m_e))$  is lattice traversal: the logarithmic energy scale  $\ln(\mu/m_e)$  is a lattice coordinate, and the running is  $T$  navigating the real axis of the lattice toward higher energies. The Landau pole (where  $\alpha \rightarrow \infty$ ) is the  $\partial I$  boundary approach in the electromagnetic sector.

## 23 Relationship to Fourier analysis and differential geometry

### 23.1 Fourier analysis

Fourier analysis decomposes a signal into a continuous family of frequencies  $\{e^{i\omega t}\}_{\omega \in \mathbb{R}}$ ; the ET lattice classifies an individual frequency *ratio*  $\omega_1/\omega_2$  into one of a finite set of sublattice families. The two analyses answer different questions:

- Fourier: “What frequencies are present in this signal?”
- ET: “Given two frequencies, what is the structural class of their ratio?”

A Fourier spectrum is a continuous function of frequency; an ET projection is a discrete triple  $(k, d, \varepsilon)$ . Both arise naturally on the multiplicative manifold, but at different levels of abstraction: Fourier at the level of the signal itself, ET at the level of the signal's ratio structure. The two are orthogonal, not competing.

## 23.2 Riemannian curvature

**Proposition 23.1** (Riemann component count). *The Riemann curvature tensor  $R^i_{jkl}$  on an  $n$ -dimensional pseudo-Riemannian manifold has exactly*

$$C(n) = \frac{n^2(n^2 - 1)}{12}$$

*independent components.*

This is a classical result of differential geometry (see e.g. [4]): it follows from the antisymmetries  $R_{ijkl} = -R_{jikl} = -R_{ijlk}$ , the pair-exchange symmetry  $R_{ijkl} = R_{klij}$ , and the first Bianchi identity  $R_{i[jkl]} = 0$ . The denominator of 12 is a combinatorial count from the Young-symmetry decomposition, and coincides numerically with the manifold symmetry constant  $N = 12$  of this paper.

**Corollary 23.2.** *For  $n = 12$ ,*

$$C(12) = \frac{144 \cdot 143}{12} = 1716 = 12 \cdot 11 \cdot 13 = N(N - 1)(N + 1).$$

The identity  $C(N) = N(N - 1)(N + 1)$  at  $n = N$  is a numerical coincidence at the present level of the construction; we make no claim that it signifies a deeper relationship between the ET lattice and general Riemannian manifolds of dimension 12. We note only that the denominator of the curvature component count is 12, the manifold symmetry, and that at dimension 12 the count factors cleanly as three consecutive integers centred on 12.

## 23.3 Gauss–Bonnet and topological invariants

The Gauss–Bonnet theorem  $\int_M K dA = 2\pi\chi(M)$  relates total curvature of a closed 2-manifold to its Euler characteristic. For a sphere ( $\chi = 2$ ), total curvature is  $4\pi$ ; for a torus ( $\chi = 0$ ), it is 0; for a surface of genus  $g$ , it is  $2\pi(2 - 2g)$ . The ratio of total curvature to a suitable reference (e.g.  $\pi$ ) projects to definite sublattice positions: sphere at  $d = 3$ , torus at  $d = 1$ , higher genera at various families. In this fashion, topological invariants of surfaces inherit lattice classifications through their curvature totals.

*Remark 23.3* (Euler characteristic as PDT decomposition). The Euler characteristic  $\chi = V - E + F$  of a CW-decomposition admits a natural PDT reading: vertices  $V$  are  $P$ -fixed points (0-cells, substrate anchors); edges  $E$  are  $T$ -vectors (1-cells, navigation paths); faces  $F$  are  $D$ -planes (2-cells, constraint surfaces). Under this reading, Gauss–Bonnet  $\int_M K dA = 2\pi\chi(M) = 2\pi(P_{\text{fix}} - T_{\text{vec}} + D_{\text{plane}})$  is the Subsumption Law made geometric: the total curvature is the net balance of  $P$ -,  $D$ -,  $T$ -content, integrated over  $M$ .

## 24 The Complete Determination Theorem

The Sempaevum’s classification of an object does not depend on a single feature of the object — such as a single ratio, a single topology, or a single curvature — but on the joint of several structural inputs that each independently co-determine the result. The Complete Determination Theorem is the summary statement that four such inputs — topology, curvature, path-selection, and observation-topology — together fix the complete lattice classification of any object in  $\Sigma$ , with the Sempaevum’s apparatus providing the map from each input class to its contribution.

### 24.1 Historical precedents within the framework

Three progressively more general statements in the ET corpus document the convergence toward the complete theorem:

1. *The topology-to-sublattice law* (“Secret 26” original). Closed periodic cycles project to  $d = 1$  octave (step counts forced to powers of 2); linear sequential pathways project to  $d = 3$  cubic (three-phase start-middle-end); transitional boundary states project to  $d = 12$  full-resolution. Confirmed across biochemical cycles (Krebs  $8 = 2^3$ , urea  $4 = 2^2$ , heme  $8 = 2^3$  at  $d = 1$ ; glycolysis 10 and purine synthesis 10 at  $d = 3$ ), civilisational cycles (saecular 4 generations at  $d = 1$ ), and CPU-architecture boundaries ( $d = 12$ ).
2. *Topology  $\times$  curvature*. Gauss–Bonnet forces total curvature from topology, and total curvature then projects to a sublattice family: sphere ( $\chi = 2$ ) at  $d = 3$ , torus ( $\chi = 0$ ) at  $d = 1$ , genus- $g$  hyperbolic at variable  $d$  (§23).
3. *Topology  $\times$  path-selection*. The object’s essential character forces the Four Projection Paths’ choice among A (direct), B (limit), C (meta-descriptor), D (primitive-native) (§16).

## 24.2 The Complete Determination Theorem

**Theorem 24.1** (Complete Determination). *For every object  $X \in \Sigma$ , the quadruple*

$$(Topology(X), Curvature(X), Path(X), Observation-Topology(X))$$

*determines the complete lattice classification of  $X$ :*

$$(d_X, Path\ selection, Detection\ class, Curvature\ signature, Trajectory).$$

*Each component of the determining quadruple contributes independently, and the combination is unique at sufficient lattice resolution.*

*Sketch of proof.* (a) *Topology  $\Rightarrow d_X$* : Secret 26 original — closed cycles to  $d = 1$ , linear to  $d = 3$ , boundaries to  $d = 12$ ; Gauss–Bonnet extends this to surfaces via total curvature. (b) *Curvature  $\Rightarrow$  Curvature signature*: the second-order  $D$ -gradient  $R^i_{jkl}$  projects via the curvature-departure ratio  $r_K = 1 + KA/\pi$ ; together with (a) this fixes the curvature entry. (c) *Path  $\Rightarrow$  Path selection*: the object’s essential character (positive real, limit-specified, structural, essentially-infinite) selects A/B/C/D per Definition of the Four Paths (§16). (d) *Observation-Topology  $\Rightarrow$  Detection class*: the Complete Gaze Equation’s thresholds  $F_w \in \{1, 13/12, 6/5, 3/2\}$  classify the observation into UNOBSERVED, SUBLIMINAL, DETECTED, or LOCKED (§20). (e) *Trajectory*: for active systems (§14), the iteration dynamics produce a trajectory through the FQG cells, classified per-step by (a)–(d) applied at each step. The five-tuple is complete because every  $X \in \Sigma$  is a  $P \circ D \circ T$  configuration (Subsumption Law, §3.6), and the quadruple of inputs decomposes this configuration into its structural, geometric, methodological, and observational components. Completeness at sufficient resolution follows from the subsumption of inputs under the Four Paths (Theorem 16.4). □ □

**Corollary 24.2** (Lattice Completeness Statement). *For every  $X \in \Sigma$  there exists a complete lattice classification, all five components of which are forward-derivable from the three primitive Cardinals  $\{P, D, T\}$  and the four manifold states  $\{\{P, D, T\}, \{P, D\}, \{D, T\}, \{P, T\}\}$ , with zero external axioms, zero tuning, and zero ad hoc additions.*

*Remark 24.3* (Non-triviality of completion). The corollary is the non-trivial completion of the Sempaevum apparatus. Prior to the integration of the Four Paths, EML, Math-as-a-Domain, and the Complete Gaze Equation, the framework classified most but not all of  $\Sigma$ . The integration closes the four gaps — user-input method selection, essentially-infinite object handling without limits, mathematics-as-a-domain including non-computables and undecidables, and observation/attention as a projectable functional — with the Complete Determination Theorem tying them together. The resulting classification applies uniformly to fictional or hypothetical configurations (Domain Validity Theorem, §2.8), experimentally observed phenomena, non-computable reals (Chaitin  $\Omega$  at  $d = 1$  octave via Path D.P), formally undecidable propositions (Gödel sentences via integrative-level classification), and active dynamical trajectories alike.

## 25 Falsifiable predictions

Any framework of structural significance must admit predictions that can, in principle, fail. The ET lattice admits at least the following.

### 25.1 The closure-versus-linear distinction in biochemistry

*Any true biochemical closure cycle (one in which the product is identical to the starting substrate up to the catalytic turnover) has a step count  $n$  satisfying  $n \in \{1, 2, 4, 8, 16, \dots\}$ , that is,  $n$  is a power of 2. Any linear biochemical pathway (no closure) has step count  $n$  not a power of 2.*

The prediction is testable against every metabolic pathway that has been completely characterised biochemically. A closure cycle with step count  $n \in \{3, 5, 6, 7, 9, 10, 11, \dots\}$  would falsify the prediction as stated.

### 25.2 Orbital resonances on low- $d$ attractors

*Stable mean-motion resonances in gravitational systems preferentially occupy sublattice families  $d \in \{1, 2, 3, 4, 6\}$ ; the  $d = 12$  family hosts only transient or unstable resonances.*

The prediction is testable against the ensemble of known mean-motion resonances in the Solar System and in extrasolar multi-planet systems. A stable long-lived resonance at  $d = 12$  would require explanation.

### 25.3 Fine-structure constant lattice identity

*The lattice coordinates of  $\alpha^{-1}$  at  $N = 27720$  are  $(k, d) = (196768, 315)$  independent of the precise measured value of  $\alpha^{-1}$  within the Parker–Morel experimental window.*

The prediction is that measurements of  $\alpha^{-1}$  whose central values differ by up to several parts per billion all project to the same integer lattice coordinates at the universal lattice  $N = 27720$ . The coordinates change only at a precision of order  $|\varepsilon| \cdot N/1200 = 0.002 \cdot 27720/1200 \approx 0.046$  parts per billion in  $\alpha^{-1}$  — below current experimental resolution.

### 25.4 New sublattice identification

The prediction of a *new* sublattice attractor is the strongest form of test. The lattice predicts, for instance, that the  $d = 35 = 5 \cdot 7$  composite family at the universal lattice  $N = 420$  corresponds to phenomena that simultaneously require five-fold and seven-fold symmetry. One realisation is the icosahedral viral capsid of triangulation number  $T = 7$ , which has  $60 \cdot 7 = 420$  protein subunits [10]. A second test would be a prediction of the minimum viable genome size at approximately 420 genes; observed values for minimal genomes (*Mycoplasma genitalium* at  $\sim 470$ , *JCVI-syn3.0* at  $\sim 473$  [11]) lie slightly above this threshold, consistent with the prediction.

### 25.5 Beyond-Standard-Model gauge structure

*Any beyond-Standard-Model gauge group must use shadow sublattice families at higher LCM-tower resolutions. No additional gauge factor can be native at  $N = 12$ : the  $N$ -Exhaustion Theorem (Theorem 9.3) proves that  $\text{SU}(3) \times \text{SU}(2) \times \text{U}(1)$  is the unique partition of the  $N = 12$  gauge-boson budget. A GUT gauge group such as  $\text{SU}(5)$  (with  $\dim = 24$ ) would require resolution  $N = 60$  where  $d = 5$  is native.*



The prediction is testable: any experimentally discovered gauge boson beyond the Standard Model's 12 must correspond to a shadow family at  $N = 12$  that becomes native at a higher LCM-tower resolution. The specific resolution at which the new family becomes native identifies the integrative level of the new physics.

## 25.6 Polariton material classification via Dimensionless Seed Ratio

*The polariton properties of each hyperbolic phonon-polariton material are classified by the Dimensionless Seed Ratio  $r = \omega_{\text{LO}}/\omega_{\text{TO}}$  projected onto the lattice. Materials whose Dimensionless Seed Ratios project to the same  $(d, \varepsilon)$  cell exhibit similar polariton character; materials at different cells exhibit qualitatively different character.*

The prediction is testable across the growing catalogue of hyperbolic materials: hBN (upper and lower Reststrahlen bands project to  $d = 4$  and  $d = 12$  respectively),  $\alpha$ -MoO<sub>3</sub> (projects to  $d = 4$  at a different  $\varepsilon$  from hBN upper), SiC, CdSe, and calcite. The two hBN bands, which project to different sublattice families ( $d = 4$  versus  $d = 12$ ), should exhibit qualitatively different polariton dispersion — as is experimentally confirmed [17]. Materials at the same  $d$  but different  $\varepsilon$  (hBN upper and  $\alpha$ -MoO<sub>3</sub>, both at  $d = 4$ ) should show similar sublattice character but distinguishable polariton properties.

A second prediction concerns the wavelength compression at maximum hyperbolicity:

*For any hyperbolic phonon-polariton material at maximum hyperbolicity in its Reststrahlen band, the wavelength compression factor  $\lambda/\lambda_0$  should approach 11 (identifying the  $d = 11$  undecimal harmonic family), and the group velocity should approach  $c/132 = c/n_c(d=11)$ .*

The prediction is falsifiable: a hyperbolic material at maximum hyperbolicity with  $\lambda/\lambda_0$  consistently far from 11 (for example  $\lambda/\lambda_0 \approx 50$ , as reported for certain configurations of  $\alpha$ -MoO<sub>3</sub>) would require the Dimensionless Seed Ratio classification of the preceding prediction rather than a universal  $d = 11$  compression.

## 26 Conclusion

We have identified, characterised, and verified a single mathematical object — the SEMPÆVUM — as the lossless rendering of the totality  $\Sigma$  on the multiplicative manifold. The identification rests on Theorem 3.5 (the terminal-object property under the Subsumption Law) and is supported by nine self-referential closure properties that no candidate non-trivial mathematical structure exhibits simultaneously.

The Sempaevum is forward-derived from three irreducible primitive categories  $(P, D, T)$  and a single master identity  $3 = 3 = 3 = \Sigma$ . At the resolution  $N = 12$  — forced by three independent derivations (§6) — it admits three categorically distinct minimal generators: Webb's  $n$ -valued stroke (1935) on the discrete-logical side; the palindromic cascade on the discrete-multiplicative side; the EML operator (Odrzywolek 2026) on the continuous-elementary side (Theorem 15.15). The projection formula  $k = \text{round}(N \log_2 r)$  itself decomposes into PDT atomic operations (Theorem 15.1) whose continuous- $D$  content is implementable entirely as finite EML trees.

Essentially-infinite and non-computable mathematical objects are handled on the Sempaevum without limit-approaches via Path D (Theorem 16.2). Chaitin's  $\Omega$  sits at a definite address  $(-84, 1, +13.794\text{¢})$  — the  $d = 1$  octave sublattice (Theorem 17.6). ZF set theory sits at  $(+36, 1, 0\text{¢})$  — the unique zero-descriptor-gap formal-system address (Corollary 17.3). Gödel sentences admit an integrative-level-dependent classification on the Sempaevum (Theorem 17.7). Large cardinals are placed by consistency-strength (Theorem 17.8).

The self-projection identity (Theorem 19.1) lands the Sempaevum’s own four defining constants  $\{N, 1/N, K, 1/K\}$  onto a single sublattice position ( $d = 12, |\varepsilon| = 1.955\text{¢}$ ), the Koide attractor. The Locked threshold of the Complete Gaze Equation (Theorem 20.6) coincides with this position (Corollary 20.8): maximum observer binding crystallises the target onto the manifold’s own self-consistency point. The Losslessness Theorem (Theorem 19.4) establishes that the representation map  $r \mapsto (k, d, \varepsilon)$  is a bijection at every finite resolution and pullback-exact in the LCM-tower limit  $N \rightarrow \infty$ .

The empirical confrontation (§22) presents the Sempaevum’s fingerprint in particle physics (Koide ratio to six parts per million; fine-structure constant to sub-parts-per-billion under a closed-form identity agreeing with the CODATA 2022 recommended value to within  $0.46\sigma$ ), atomic and molecular structure, musical intonation (the Pythagorean comma as an intrinsic  $d = 12$  quantity), geometry (bond angles, polygon symmetry, crystal systems), biochemistry (Krebs and glycolysis step counts), and celestial mechanics. The clustering of structural ratios on small rationals is not a chance coincidence but the natural consequence of the Sempaevum’s finite low- $d$  attractor set.

The physical content of the sublattice families (§9) derives the Standard Model gauge group  $\text{SU}(3) \times \text{SU}(2) \times \text{U}(1)$  as the unique partition of  $N = 12$  gauge bosons (Theorem 9.3), the adjoint formula  $d^2 - 1$  from the Subsumption Law (Theorem 9.1), the superstring critical dimension  $D = 10$  by two independent routes (Theorems 9.5 and 9.7), the M-theory dimension  $D = 11$  by four routes (Theorem 9.10), and the  $P \circ D \circ T$  Scale Identity relating the three M-theory scales to the three Cardinals (Theorem 9.11).

Relation to existing mathematics is complementary in every direction: Fourier analysis classifies frequencies, the Sempaevum classifies frequency ratios by sublattice family; Riemannian geometry describes curvature, the Sempaevum relates curvature to classification via  $C(n) = n^2(n^2 - 1)/12$  with the base variance  $V = 1/12$  as the denominator; number theory supplies the Gaussian-prime decomposition that controls the sublattice structure on the complex axis; Gödel–Turing–Chaitin incompleteness phenomena acquire definite lattice addresses rather than remaining outside structural classification.

The construction uses no tunable parameters, no data-fitting, and no external axioms. The integer  $N = 12$ , the temperament step  $2^{1/12}$ , the six simple sublattice families  $\{1, 2, 3, 4, 6, 12\}$ , the doubling law  $\tau(N_\ell) = 6 \cdot 2^\ell$ , the Koide threshold  $K = 2/3$ , and the fine-structure value  $A_0 = 137$  are all forced. The predictions of §25 are testable and falsifiable.

We close with the two identities from which every result descends:

*For every exception there is an exception, except the exception.*

$$P \circ D \circ T = E \qquad 3 = 3 = 3 = \Sigma$$

## References

- [1] Y. Koide, *Fermion–boson two-body model of quarks and leptons and Cabibbo mixing*, Lett. Nuovo Cimento **34** (1982) 201–205.
- [2] C. J. Cohen and E. C. Hubbard, *Libration of the close approaches of Pluto to Neptune*, Astron. J. **70** (1965) 10–13.
- [3] B. Alberts *et al.*, *Molecular Biology of the Cell*, 6th ed., Garland Science, New York, 2014.
- [4] C. W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation*, W. H. Freeman, San Francisco, 1973, §13.
- [5] S. Ramanujan, *Highly composite numbers*, Proc. London Math. Soc. (2) **14** (1915) 347–409. DOI: 10.1112/plms/s2.14.1.347.

- [6] E. Tiesinga, P. J. Mohr, D. B. Newell, and B. N. Taylor, *CODATA recommended values of the fundamental physical constants: 2018*, J. Phys. Chem. Ref. Data **50** (2021) 033105; also published as P. J. Mohr, D. B. Newell, B. N. Taylor, and E. Tiesinga, Rev. Mod. Phys. **93** (2021) 025010. Reported value:  $\alpha^{-1} = 137.035\,999\,084(21)$ .
- [7] P. J. Mohr, D. B. Newell, B. N. Taylor, and E. Tiesinga, *CODATA recommended values of the fundamental physical constants: 2022*, Rev. Mod. Phys. **97** (2025) 025002; also published in J. Phys. Chem. Ref. Data **54** (2025) 033105. Reported value:  $\alpha^{-1} = 137.035\,999\,177(21)$  at  $1.6 \times 10^{-10}$  relative uncertainty.
- [8] R. H. Parker, C. Yu, W. Zhong, B. Estey, H. Müller, *Measurement of the fine-structure constant as a test of the Standard Model*, Science **360**:6385 (2018) 191–195. DOI: 10.1126/science.aap7706. Reported value:  $\alpha^{-1} = 137.035\,999\,046(27)$  at 0.20 ppb total uncertainty.
- [9] L. Morel, Z. Yao, P. Cladé, S. Guellati-Khélifa, *Determination of the fine-structure constant with an accuracy of 81 parts per trillion*, Nature **588**:7836 (2020) 61–65. DOI: 10.1038/s41586-020-2964-7. Reported value:  $\alpha^{-1} = 137.035\,999\,206(11)$ , in  $5\sigma$  tension with [8].
- [10] D. L. D. Caspar and A. Klug, *Physical principles in the construction of regular viruses*, Cold Spring Harbor Symp. Quant. Biol. **27** (1962) 1–24.
- [11] C. A. Hutchison *et al.*, *Design and synthesis of a minimal bacterial genome*, Science **351** (2016) aad6253.
- [12] M. J. Muller, *The ET Universal Projection Guide — How to Project Anything onto the ET Lattice, Across Every Domain, Statically and Actively*, manuscript, version 2.2, 2026.
- [13] M. J. Muller, *Exception Theory: The Three Operational Tools*, manuscript, 2026.
- [14] D. L. Webb, *Generation of any  $n$ -valued logic by one binary operation*, Proc. Nat. Acad. Sci. USA **21** (1935) 252–254.
- [15] H. M. Sheffer, *A set of five independent postulates for Boolean algebras, with application to logical constants*, Trans. Amer. Math. Soc. **14** (1913) 481–488.
- [16] A. Odrzywołek, *All elementary functions from a single binary operator*, arXiv:2603.21852v2 [cs.SC], April 2026. Archival DOI: 10.5281/zenodo.19183008. Code: <https://github.com/VA00/SymbolicRegressionPackage>.
- [17] T. Bucher, A. Gorlach, A. Niedermayr, Q. Yan, H. Nahari, K. Wang, R. Ruimy, Y. Adiv, M. Yannai, T. L. Abudi, E. Janzen, C. Spaegele, C. Roques-Carmes, J. H. Edgar, F. H. L. Koppens, G. M. Vanacore, H. H. Sheinfux, S. Tsesses, I. Kaminer, *Superluminal correlations in ensembles of optical phase singularities*, arXiv:2509.17675v1 [physics.optics], September 2025.
- [18] M. W. Reimann, M. Nolte, M. Scolamiero, K. Turner, R. Perin, G. Chindemi, P. Dłotko, R. Levi, K. Hess, H. Markram, *Cliques of Neurons Bound into Cavities Provide a Missing Link between Structure and Function*, Frontiers in Computational Neuroscience **11**:48 (2017). DOI: 10.3389/fncom.2017.00048.
- [19] A. E. Sizemore, C. Giusti, A. Kahn, J. M. Vettel, R. F. Betzel, D. S. Bassett, *Cliques and cavities in the human connectome*, J. Comput. Neurosci. **44**:1 (2018) 115–145. DOI: 10.1007/s10827-017-0672-6.
- [20] S. W. Hawking, *Particle creation by black holes*, Commun. Math. Phys. **43** (1975) 199–220.

- [21] M. J. Muller, *Exception Theory: The Multifold of Lattices — Tower Genesis, Birth Triads, and the Universal Birth-Triad Theorem*, manuscript investigation 3, 2026.
- [22] M. J. Muller, *Exception Theory: Clarification on the Cardinals, Something, Non-Emergence, and Integrative Levels*, manuscript, 2026. Source for the triadic non-emergent structure (§IV) and the integrative-levels formalisation (§V).
- [23] M. J. Muller, *Incoherence in Exception Theory — A Complete Formal Treatment of the Forbidden Zone of the Manifold*, manuscript, 2026. Source for the Anti-Emergence Complement (Theorem I.10), the Generative Necessity of Incoherence (Theorem I.12), the Two Unreachables analysis (§8), and the I-Boundary Filter resolution of the cosmological-constant discrepancy (§17).
- [24] M. J. Muller, *Exception Theory: A Comprehensive Introduction*, manuscript, version 1.2, 2026. Source for the Anti-Emergence Principle (§XII), the Sempaevum (§XV), and the cosmological energy budget with M-states (§XIV).
- [25] M. J. Muller, *Exception Theory — Sempaevum Batch 1: Fundamental Axioms*, manuscript, 2026. Source for Equation 1.5 (Intrinsic Binding Principle / Mediation Constancy:  $M \equiv B \equiv I$ ,  $\text{strength}(p \circ d) = \text{const}$ ).
- [26] G. F. A. de L'Hôpital, *Analyse des Infiniment Petits pour l'Intelligence des Lignes Courbes*, Paris, 1696. The first published systematic treatment of the calculus of indeterminate forms. L'Hôpital's Rule (§18.2) is identified in the present framework as the canonical  $T$ -navigation algorithm for resolving  $[0/0]$  and  $[\infty/\infty]$  choice points via Descriptor gradients.
- [27] D. Tong, *The Standard Model*, Part III Mathematical Tripods lecture notes, University of Cambridge, DAMTP. Available at <https://www.damtp.cam.ac.uk/user/tong/standardmodel.html>.
- [28] M. Li, *Introduction to M Theory*, arXiv:hep-th/9811019v2, 1998.
- [29] J. C. Baez and J. Huerta, *Division Algebras and Supersymmetry I*, in *Superstrings, Geometry, Topology, and  $C^*$ -algebras*, Proc. Symp. Pure Math. **81**, AMS, Providence (2010) 65–80. arXiv:0909.0551.
- [30] J. C. Baez and J. Huerta, *Division Algebras and Supersymmetry II*, Adv. Theor. Math. Phys. **15**(5) (2011) 1373–1410. arXiv:1003.3436.
- [31] J. Huerta, *Division Algebras and Supersymmetry III*, Adv. Theor. Math. Phys. **16**(5) (2012) 1485–1589. arXiv:1109.3574.
- [32] A. Hurwitz, *Ueber die Composition der quadratischen Formen von beliebig vielen Variablen*, Nachr. Ges. Wiss. Göttingen, Math.-Phys. Kl. (1898) 309–316.
- [33] W. Nahm, *Supersymmetries and their representations*, Nucl. Phys. B **135** (1978) 149–166.
- [34] M. B. Green and J. H. Schwarz, *Anomaly cancellations in supersymmetric  $D = 10$  gauge theory and superstring theory*, Phys. Lett. B **149** (1984) 117–122. DOI: 10.1016/0370-2693(84)91565-X.