


THE LCM TOWER · $\ell=0$

$N=12 \cdot \tau=6 \cdot 6/12 \text{ active} \cdot \xi(m) = 137/((m-1)^2+16)$

Host: $d = 12/\gcd(m, 12)$ · Mirrors: $m + m' = 12$ · Axis-agnostic (IC-110)



CASCADE POSITION SEQUENCES – $k_n = (g \cdot n) \bmod 12$

Each generator traverses a DIFFERENT path through ALL 12 harmonic families. Position k = family m .

NON-TRIVIAL PAIR (5, 7)

$g=7$ fifths



$g=5$ fourths



TRIVIAL PAIR (1, 11)

$g=1$ fwd



$g=11$ bwd



MIRRORS $m+m'=12$

$1\ast 11 \cdot 2\ast 10 \cdot 3\ast 9 \cdot 4\ast 8 \cdot 5\ast 7 \cdot 6\ast 6(\text{self})$

HOST $d=12/\gcd(m,12)$

$m=5,7,11+d=12 \cdot m=9+d=4 \cdot m=10+d=6 \cdot m=8+d=3$

$\Sigma\phi(d) = 1+1+2+2+2+4 = 12 = N \cdot$ Distribution $[3,1,1,1,0,0]$ unique to ET

EM+gravity: $\kappa=0$ $E=1.606$ · EM+strong: $\kappa=0$ $E=1.284$ · EM+weak: $\kappa\neq 0$ $E=0.343$

Gravity+strong channels: D-arithmetic (deterministic). Weak channel: T-agency exclusively (IC-107).

U(1) traversal: $m=1+4+2+6=12 \cdot m=3$ (strong/instanton) excluded – topological, non-perturbative

ξ monotonically decreasing (IC-109): gravity 8.563 $\rightarrow$ EM 1.000 · Axis-agnostic (IC-110)

LEGEND



Simple ($m|12$)



Complex ($m|12$)

Hasse

Shadow hosting

Mirror



Self-mirror ($m=6$)

U(1) traversal

$\kappa=0$ (D-arith, E>1)

$\kappa\neq 0$ (T-act, E<1)



(5,7) non-trivial pair



(1,11) trivial pair

Force (bold) = real D-axis · Phase = imaginary T-axis · Position colors match family nodes

$P \circ D \circ T = E$ – Exception Theory – Michael James Muller – Aevum Defluo