Study ID

Title

Author

Year

Search **Go to Record**

Country (of cows)

General notes about the study (if needed)

>> Trials >>

ID

Study id expo

Trial expo

Group

Group order

Active ingredient ⌄ ☐ TS

order-active ingredient

AB info

Concentration per dose

Administration regime ⌄

Selection criteria ⌄

AB preparation ⌄

AB trade name

Frequency of administration

Interval (days)

Route ⌄

TS info

TS Conc (%)

TS prepration ⌄

TS active ingredient ⌄

TS trade Name

More drug info. (copy/paste)

Role in PICO*

PICO 1-2*

○ Control   ○ Treatment   ○ Not-used

PICO 3-4*

○ Control   ○ Treatment   ○ Not-used

PICO 5*

○ Control   ○ Treatment   ○ Not-used

Quality of administration technique

Teat preparation before DCT infusion*

○ Yes   ○ No   ○ Not-reported

Depth of insertion of DCT syringe cannula*

○ Full   ○ Partial   ○ Not-reported

Teat spray or dipping after DCT infusion*

○ Yes   ○ No   ○ Not-reported

>> Outcomes >>

ID

Study id expo ☐

Trial expo ☐

Group expo ☐

OUTCOME_ARM

Outcome info

Outcome	⌄
Def	
Detection Method	⌄
Stratifying variable	⌄
Total number of quarters at risk	

Pathogen-arm

Time info

Time order	☐
days post calving	from ☐ to ☐
COV (1000 cells/ml)	
Ounit	⌄

Change Time settings

Stratification levels	Pathogen spp.	Path. Def. (if others)	+Ve	Total

>> Contrast

ID

study id expo [   ]   T [   ]

trial expo [   ]   U [   ]

Group expo [   ]

OUTCOME_CONTRAST

Estimate/Coefficient characteristics

Estimate type [▾]   Estimate scale [▾]

CI level [▾]

Pathogen-Contrast

Stratification levels	Pathogen spp.	Path.Def. (if others)	Comparison group	Estimate	SE	LCI	UCI	P-value

Back to Arm