

Using Bioinformatics to Transform the Diagnosis and Treatment of Ovarian Cancer

Word count (excluding references): 1985

Introduction

Many diseases remain undetected or undiagnosed in their early stages and are only discovered after they have spread this is particularly true for ovarian cancer (Siegel *et al.*, 2017). This malignant tumor is among the most dangerous cancers affecting women, with an overall five year survival rate of only around 49% across all stages (Vincent Schilling *et al.*, 2023). Current screening tools, such as CA-125 blood tests or medical imaging, are not sufficiently reliable for early detection (Siegel *et al.*, 2017). In many cases, the disease progresses and develops resistance to chemotherapy due to late diagnosis, by which time the cancer has reached advanced stages (Vincent Schilling *et al.*, 2023). These challenges have motivated researchers in the field of bioinformatics to explore more effective approaches for diagnosing and managing ovarian cancer, especially in recent years.

In this context, bioinformatics has helped scientists identify genes that behave abnormally in cancerous cells compared to healthy ones. This is made possible through the integration of computational and statistical techniques that allow for the analysis of massive biological datasets. Such approaches contribute to more on patient-specific responses (Hibba Rashid *et al.*, 2023).

This Essay explores examples of how bioinformatics contributes to early diagnosis and treatment planning for ovarian cancer and discusses the challenges associated with its clinical implementation.

Early Diagnosis Using RNA Sequencing

Resistance to treatment in ovarian cancer is often mediated by mechanisms such as drug efflux transporters and PARP related pathways (Sun *et al.*, 2015). Because ovarian cancer symptoms are varied and similar to other conditions, the usual diagnostic methods are often not accurate or sufficient (Siegel *et al.*, 2017). This highlights a growing need for molecular based diagnostic strategies that leverage bioinformatics and gene expression profiling to identify novel therapeutic targets and improve clinical outcomes (Schilling *et al.*, 2023). One of the most widely used techniques in this context is RNA sequencing, which allows researchers to analyze gene expression patterns across thousands of genes simultaneously (Liu *et al.*, 2022). One area where bioinformatics has

made a difference is in analyzing RNA-seq data. This technique helps researchers see which genes are turned on or off in cancer cells. For example, Liu et al. analyzed RNA-seq data and found around 590 genes that were expressed differently between ovarian cancer and normal tissue (Liu *et al.*, 2022). Many of these genes were linked to how cells grow and how the immune system works.

They applied machine learning techniques like LASSO regression and SVM-RFE to narrow down important genes. This confirms the relevance of these genes in detecting and distinguishing cancer samples, such as FOLR1, BUB1, and PSAT1. Another study by Zheng et al. showed that some of these genes are also connected to poor patient outcomes, so they might help with prognosis as well. (Zheng *et al.*).

RNA-seq has also been useful in identifying subtypes of ovarian cancer, which could explain why some patients respond differently to the same treatment. This isn't fully applied in hospitals yet, but it's a direction that researchers are exploring. There's also been interest in combining RNA-seq data with other types of omics like DNA methylation or proteomics, to get a more layered understanding of what's going on inside the tumor. In a few newer studies, single-cell RNA-seq has allowed scientists to map which cells inside a tumor are contributing to disease progression. This is important because sometimes the bulk RNA data hides variations between cell types. Tools like Seurat or Scanpy are now being used more often in ovarian cancer datasets for this reason (Yue You and Qing Yang).

in addition to transcriptome-based diagnostics like RNA sequencing, recent studies have investigated the role of epigenetic markers, particularly DNA methylation, in the early detection of ovarian cancer. Lien et al. (2024) developed a methylation based, machine learning model, using cervical scraping samples, aiming to identify patients with ovarian cancer in a non-invasive manner (Lien *et al.*, 2025). This approach leverages abnormal DNA methylation patterns associated with tumor progression and highlights the potential of integrating epigenetic signals into bioinformatics pipelines to improve diagnostic accuracy (Lien *et al.*, 2025; Yuan Tian *et al.*, 2025).

Treatment Personalization and Response Prediction

Not only is bioinformatics having an impact on ovarian cancer diagnosis, but it's also contributing to your treatment. One of these is FOLR1, which has been investigated as a diagnostic marker, as well as a candidate target for therapy. Kim et al. (2024) demonstrated that targeting FOLR1 with an antibody-drug conjugate

impeded cell migration and demonstrated reduced tumor volume in ovarian cancer models (Kim *et al.*, 2020). Bioinformatics is also helping explain why certain tumors stop responding to chemotherapy. A common mechanism is the ABCB1 gene (also called MDR1), which is often overexpressed in drug-resistant ovarian tumors. Sun *et al.* (2022), microRNA miR-186 may ultimately contribute to restoring chemosensitivity via down-regulating ABCB1 (Sun *et al.*, 2015). Such discoveries could also inform new strategies to make existing treatments more effective and less prone to resistance.

A few predictive models were generated recently to predict the survival or potentially responders for immunotherapy in ovarian cancer. For example, Zhao X *et al.* developed a glycosylation-related gene signature that was associated with patient survival and immune cell infiltration (Zhao *et al.*, 2022). Similarly, Zhao C *et al.* proposed a glycosyltransferase gene-based gene signature to predict patients who are most likely to benefit from immunotherapy (Xu *et al.*, 2023).

On the immunological front, Liu *et al.* analyzed the relationship between specific genes and immune cell presence within the tumor microenvironment (Liu *et al.*, 2022).

In work aimed at investigating potential therapeutic approaches, Block *et al.* trialled a vaccine designed to stimulate Th17 cells, and found a positive immune response (Block *et al.*, 2020). Concomitantly, a small fraction of studies aimed at grouping tumors based on immune infiltration. This line of research has the potential to lead to more personalized immunotherapy strategies, but more validation is needed (Feng *et al.*, 2019).

Challenges and Where Things Stand

A recent study has shown that the hundreds of thousands of genes contained in The Cancer Genome Atlas also enable it to identify and classify molecular transcripts. By combining TCGA data with a new technique called factorial hierarchical clustering (FHC), researchers found it easier and faster to sort through this vast amount of information. These molecular profiles have shown promise in helping anticipate how patients might respond to treatment and in estimating overall outcomes. However, their application in clinical practice remains limited. Liquid biopsies have recently emerged as a promising tool for improving early cancer detection. (Roy *et al.*, 2016). Liquid biopsies have recently emerged as a promising tool for improving early cancer detection. By analyzing circulating tumor DNA (ctDNA) and other blood-based biomarkers, researchers can gain valuable insight into tumor location, mutation dynamics,

and underlying molecular changes without the need for invasive procedures. Although not yet commonly used in clinical settings, preliminary findings indicate that this approach could significantly improve early diagnostic accuracy and ongoing monitoring (Della Corte *et al.*, 2024).

A systematic review revealed that, even under current national guidelines, a considerable number of women at elevated risk of ovarian cancer are not being referred for genetic testing or receiving appropriate counseling. For instance, only 24% – 26% of Black patients were referred for counseling or completed testing, compared to 40% – 43% of White patients. Similar gaps were noted based on insurance status, with privately insured patients being significantly more likely to receive services than those on Medicaid or uninsured (Kolla and Parikh, 2024).

One of the biggest barriers and challenges is the difficulty of integrating next-generation sequencing (NGS) output into hospital systems. This problem is not only limited to technical information, but many factors lead to it. In addition, its seed lies in the lack of standardized clinical guidelines and a deficiency among medical professionals for interpreting biocomputing reports. Solving this problem is not a matter of creating sophisticated computing infrastructure but instead requires other important aspects which will largely make life-saving difference to patients' lives. This involves all disciplines working together with bioinformaticians, clinicians and health policy makers as well as initiating policies on believable clinical reporting. Carter and Pan Tanowitz (2016) describes issues such as lack of standards, slow data flows, and difficulty with variant discovery communications to patients or non-specialist physicians (Denisenko *et al.*, 2024). These problems must be addressed if genetic information is to become a routine part of medical decision making. Recently, a few researchers have tried to improve biological database systems in the form of a cloud service. They believe that this will be a forward step for global users due to its ease and flexibility of application. Restrictions cannot be placed on genetic information solely to prevent or deal with crime. There are real people who are patients themselves and their families. Their urgent needs should not be ignored (Gene 2015). (Thirunavukkarasu *et al.*, 2025) Another difficulty that must be addressed is the cancerous complexity of it all. The individual qualities of ovarian tumors even within the same patient can vary greatly, and environmental factors around them change over time (Bellazzi *et al.*, 2012).

Recent developments in AI have facilitated the creation of improved models for diagnosis, prognosis, and selection of treatment of cancer. Nevertheless, to this point, no single data modality is sufficiently informative to account for all the biological complexity of tumors in current models. In this respect, deep learning

(DL) approaches have shown potential in integrating multimodal and high-dimensional biological data to build deep, complex predictive models. One of the notable works in this field is the unified multi-task DL framework of OmiEmbed proposed by Zhang et al. (2021), that was developed for multi-omic data integration of gene expression, proteomics and metabolomics. OmiEmbed tackles several clinical tasks, tumor classification, phenotype reconstruction and survival prediction, within a single learning framework in which deep representations of biological features are learned using task-specific modules. Importantly, this method outperforms state of the art machine learning approaches in a comparison study in several benchmarks in classification accuracy and survival estimation (Zhang *et al.*, 2021).

The privacy of patient genetics is a concern (particularly after data sets are pooled together among research groups). Some researchers are working on developing techniques so that bioinformatics models are also transparent and interpretable, and thus we can demand that they will be able to be trusted in real world medical systems (Integrated genomic analyses of ovarian carcinoma). Still, there are some things to look for. Single cell sequencing and spatial transcriptomics may aid to more finely resolve (Denisenko *et al.*, 2024). profile tumors. There is suggestion that incorporating different types of measurement such as gene expression, protein concentrations, or metabolic values may provide a more complete view of the biology of the tumor. Nevertheless, the broad implementation of these integrative methods into routine clinical practice is impeded by variability in standardization and interpretation of the assays and by practical construction and deployment of these tools (Thirunavukkarasu *et al.*, 2025).

And as hopeful as deep learning models seem, they require a wider array of datasets before we can trust the predictions they're making. There's also a lot of hope around federated learning, where models are trained over decentralized data, without sharing sensitive information directly (Jenny Lin *et al.*, 2021).

Conclusion

The application of bioinformatics methods has significantly contributed to a breakthrough in the biological processes of ovarian cancer, particularly in identifying crucial genes and predicting individuals' responses to therapies. Techniques such as RNA-seq and integrative multi-omics analysis show some promise, but most of what has been achieved remains within the realm of scientific research. Most of these tools are not easy to use in actual medical practice, and so far, there are only a few well-organized and informative datasets.

If we truly want to move forward, stronger integration between research and the clinic is essential, and these tools must be designed in a way that is understandable for everyday use. Not every model will succeed, but if there are enough of them, and with some testing and refinement, it helps preserve patients lives.

References

- Bellazzi,R. *et al.* (2012) Clinical Bioinformatics: challenges and opportunities. *BMC Bioinformatics*, **13**, S1, 1471-2105-13-S14-S1.
- Block,M.S. *et al.* (2020) Th17-inducing autologous dendritic cell vaccination promotes antigen-specific cellular and humoral immunity in ovarian cancer patients. *Nat Commun*, **11**, 5173.
- Della Corte,L. *et al.* (2024) The role of liquid biopsy in epithelial ovarian cancer: State of the art. *Critical Reviews in Oncology/Hematology*, **194**, 104263.
- Denisenko,E. *et al.* (2024) Spatial transcriptomics reveals discrete tumour microenvironments and autocrine loops within ovarian cancer subclones. *Nat Commun*, **15**, 2860.
- Feng,H. *et al.* (2019) Identification of significant genes with poor prognosis in ovarian cancer via bioinformatical analysis. *J Ovarian Res*, **12**, 35.
- Hibba Rashid *et al.* (2023) Identification of Novel Genes and Pathways of Ovarian Cancer Using a Comprehensive Bioinformatic Framework. **169**, 3056–3075.
- Integrated genomic analyses of ovarian carcinoma 2011.
- Jenny Lin *et al.* (2021) Disparities in genetic testing for women with ovarian cancer. *Gynecologic Oncology*, **162**, S135–S136.
- Kim,O. *et al.* (2020) Targeting progesterone signaling prevents metastatic ovarian cancer. *Proc. Natl. Acad. Sci. U.S.A.*, **117**, 31993–32004.
- Kolla,L. and Parikh,R.B. (2024) Uses and limitations of artificial intelligence for oncology. *Cancer*, **130**, 2101–2107.
- Lien *et al.* (2025) Exploring potential methylation markers for ovarian cancer from cervical scraping samples.
- Liu,J. *et al.* (2022) Identification and Validation of the Diagnostic Characteristic Genes of Ovarian Cancer by Bioinformatics and Machine Learning. *Front. Genet.*, **13**, 858466.
- Roy,S. *et al.* (2016) Next-Generation Sequencing Informatics: Challenges and Strategies for Implementation in a Clinical Environment. *Archives of Pathology & Laboratory Medicine*, **140**, 958–975.
- Schilling,V. *et al.* (2023) A Bioinformatics Analysis of Ovarian Cancer Data Using Machine Learning. *Algorithms*, **16**, 330.
- Siegel *et al.* (2017) Cancer statistics 2017. *CA: A Cancer Journal for Clinicians*, 67.
- Sun,K.-X. *et al.* (2015) MicroRNA-186 induces sensitivity of ovarian cancer cells to paclitaxel and cisplatin by targeting ABCB1. *J Ovarian Res*, **8**, 80.
- Thirunavukkarasu *et al.* (2025) Transcriptome profiling and metabolic pathway analysis towards reliable biomarker discovery in early-stage lung cancer. **66**, 115–126.
- Vincent Schilling *et al.* (2023) A Bioinformatics Analysis of Ovarian Cancer Data Using Machine Learning. **16**, 20.
- Xu,X. *et al.* (2023) A signature based on glycosyltransferase genes provides a promising tool for the prediction of prognosis and immunotherapy responsiveness in ovarian cancer. *J Ovarian Res*, **16**, 5.

- Yuan Tian *et al.* (2025) The Chip Analysis Methylation Pipeline (ChAMP).
- Yue You and Qing Yang Glycosylation-related genes mediated prognostic signature contribute to prognostic prediction and treatment options in ovarian cancer: based on bulk and single-cell RNA sequencing data. *2024*, 16.
- Zhang,X. *et al.* (2021) OmiEmbed: A Unified Multi-Task Deep Learning Framework for Multi-Omics Data. *Cancers*, **13**, 3047.
- Zhao,C. *et al.* (2022) Glycosylation-Related Genes Predict the Prognosis and Immune Fraction of Ovarian Cancer Patients Based on Weighted Gene Coexpression Network Analysis (WGCNA) and Machine Learning. *Oxidative Medicine and Cellular Longevity*, **2022**, 1–23.
- Zheng,M.-J. *et al.* Identification of molecular marker associated with ovarian cancer prognosis using bioinformatics analysis and experiments. *11 January 2019*, **234**, 11023–11036.