

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/343940944>

A field and video-annotation guide for baited remote underwater stereo-video surveys of demersal fish assemblages

Article in *Methods in Ecology and Evolution* · August 2020

DOI: 10.1111/2041-210X.13470

CITATIONS

127

40 authors, including:



Tim J. Langlois

University of Western Australia

130 PUBLICATIONS 6,279 CITATIONS

[SEE PROFILE](#)



Todd Bond

University of Western Australia

46 PUBLICATIONS 1,027 CITATIONS

[SEE PROFILE](#)

READS

792



Jordan Scott Goetze

Curtin University

22 PUBLICATIONS 890 CITATIONS

[SEE PROFILE](#)



Jacquomo Monk

University of Tasmania

74 PUBLICATIONS 1,550 CITATIONS

[SEE PROFILE](#)

A field and video annotation guide for baited remote underwater stereo-video surveys of demersal fish assemblages

Tim Langlois¹  | Jordan Goetze^{2,3}  | Todd Bond¹  | Jacquomo Monk⁴  |
 Rene A. Abesamis⁵  | Jacob Asher^{6,7}  | Neville Barrett⁴  | Anthony T. F. Bernard^{8,9}  |
 Phil J. Bouchet¹⁰  | Matthew J. Birt¹¹  | Mike Cappo¹² | Leanne M. Currey-Randall¹²  |
 Damon Driessen³ | David V. Fairclough^{3,13}  | Laura A. F. Fullwood³ | Brooke A. Gibbons¹ |
 David Harasti¹⁴  | Michelle R. Heupel¹²  | Jamie Hicks¹⁵ | Thomas H. Holmes^{1,2} |
 Charlie Huveneers¹⁶  | Daniel Ierodiaconou¹⁷  | Alan Jordan⁴ | Nathan A. Knott¹⁸  |
 Steve Lindfield¹⁹  | Hamish A. Malcolm²⁰  | Dianne McLean^{1,11}  | Mark Meekan¹¹  |
 David Miller¹⁵ | Peter J. Mitchell²¹ | Stephen J. Newman^{3,13}  | Ben Radford¹¹ |
 Fernanda A. Rolim²²  | Benjamin J. Saunders³  | Marcus Stowar¹² |
 Adam N. H. Smith²³  | Michael J. Travers^{3,13}  | Corey B. Wakefield^{3,13} |
 Sasha K. Whitmarsh¹⁶  | Joel Williams¹⁴  | Euan S. Harvey³ 

¹The UWA Oceans Institute and School of Biological Sciences, The University of Western Australia, Perth, WA, Australia; ²Marine Science Program, Biodiversity and Conservation Science, Department of Biodiversity, Conservation and Attractions, Kensington, WA, Australia; ³School of Molecular and Life Sciences, Curtin University, Perth, WA, Australia; ⁴Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, Tas, Australia; ⁵Angelo King Center for Research and Environmental Management, Silliman University, Dumaguete City, Philippines; ⁶NOAA Fisheries, Pacific Islands Fisheries Science Center, Science Operations Division, NOAA Inouye Regional Center, Honolulu, HI, USA; ⁷Joint Institute for Marine and Atmospheric Research, University of Hawai'i at Mānoa, Honolulu, HI, USA; ⁸South African Institute for Aquatic Biodiversity, Grahamstown, South Africa; ⁹Department of Zoology and Entomology, Rhodes University, Grahamstown, South Africa; ¹⁰Centre for Research into Ecological & Environmental Modelling, School of Mathematics and Statistics, University of St Andrews, St Andrews, UK; ¹¹Australian Institute of Marine Science, Indian Ocean Marine Research Centre, Perth, WA, Australia; ¹²Australian Institute of Marine Science, Townsville, Qld, Australia; ¹³Western Australian Fisheries and Marine Research Laboratories, Department of Primary Industries and Regional Development, Government of Western Australia, North Beach, WA, Australia; ¹⁴Fisheries Research, NSW Department of Primary Industries, Taylors Beach, NSW, Australia; ¹⁵Marine Science Program, Science & Corporate Services Division, Department for Environment and Water, Adelaide, SA, Australia; ¹⁶Southern Shark Ecology Group, College of Science and Engineering, Flinders University, Bedford Park, SA, Australia; ¹⁷School of Life and Environmental Sciences, Centre for Integrative Ecology, Deakin University, Warrnambool, Vic., Australia; ¹⁸Fisheries Research, NSW Department of Primary Industries, Huskisson, NSW, Australia; ¹⁹Coral Reef Research Foundation, Koror, Palau; ²⁰Fisheries Research, NSW Department of Primary Industries, Coffs Harbour, NSW, Australia; ²¹Centre for Environment, Fisheries and Aquaculture Science, Lowestoft, UK; ²²Elasmobranch Research Laboratory, Institute of Biosciences, São Paulo State University, São Vicente, SP, Brazil and ²³School of Natural and Computational Sciences, Massey University, Auckland, New Zealand

Correspondence

Jordan Goetze
Email: gertzaj@gmail.com

Funding information

Australian Government's National Environmental Science Program; Australian Research Data Commons; Gorgon-Barrow Island Gorgon Barrow Island Net Conservation Benefits Fund

Handling Editor: Edward Codling

Abstract

1. Baited remote underwater stereo-video systems (stereo-BRUVs) are a popular tool to sample demersal fish assemblages and gather data on their relative abundance and body size structure in a robust, cost-effective and non-invasive manner. Given the rapid uptake of the method, subtle differences have emerged in the way stereo-BRUVs are deployed and how the resulting imagery is annotated. These disparities limit the interoperability of datasets obtained across studies, preventing broadscale insights into the dynamics of ecological systems.

Tim Langlois and Jordan Goetze are joint lead authors.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. *Methods in Ecology and Evolution* published by John Wiley & Sons Ltd on behalf of British Ecological Society

2. We provide the first globally accepted guide for using stereo-BRUVs to survey demersal fish assemblages and associated benthic habitats.
3. Information on stereo-BRUVs design, camera settings, field operations and image annotation are outlined. Additionally, we provide links to protocols for data validation, archiving and sharing.
4. Globally, the use of stereo-BRUVs is spreading rapidly. We provide a standardized protocol that will reduce methodological variation among researchers and encourage the use of Findable, Accessible, Interoperable and Reusable workflows to increase the ability to synthesize global datasets and answer a broad suite of ecological questions.

KEYWORDS

monitoring (population ecology), population ecology, sampling

1 | INTRODUCTION

Our understanding of fish ecology and ability to manage populations appropriately require accurate data on species occurrence, abundance, body size, distribution and behaviour. Remote video-based sampling methods are increasingly being adopted due to: (a) their non-destructive nature, (b) ability to sample rare species (Goetze et al., 2019; Harvey, Santana-Garcon, Goetze, Saunders, & Cappel, 2018), over broad depth ranges (Heyns-Veale et al., 2016; Wellington et al., 2018), (c) provision of a permanent record that can be reviewed to reduce interobserver variability (Cappel, De'ath, Stowar, Johansson, & Doherty, 2009), (d) ability to collect concomitant data on habitat (Bennett, Wilson, Shedrawi, McLean, & Langlois, 2016; e.g. epibenthic cover and substrate, Collins et al., 2017) and (e) provision of images for science communication. Remote underwater video sampling methods are not subject to diver safety restrictions, nor do they suffer from the behavioural biases resulting from diver presence (Gray et al., 2016; Lindfield, Harvey, McIlwain, & Halford, 2014). Multiple remote systems can be deployed in the field consecutively to make efficient use of field time and enable spatially extensive sampling (Langlois, Radford, et al., 2012).

The use of bait with remote underwater video (BRUV) systems increases the relative abundance and diversity of fishes observed, particularly species targeted by fisheries, without precluding the sampling of fishes not attracted to bait (Coghlan, McLean, Harvey, & Langlois, 2017; Harvey, Cappel, Butler, Hall, & Kendrick, 2007; Speed, Rees, Cure, Vaughan, & Meekan, 2019). Biases associated with bait use have been discussed in various studies (Coghlan et al., 2017; Dorman, Harvey, & Newman, 2012; Goetze et al., 2015; Hardinge, Harvey, Saunders, & Newman, 2013). Variation in bait plume dispersal and the sensitivity of different fish species to bait is unknown (Harvey et al., 2007), and species-specific (Bernard & Götz, 2012), with cryptic and sedentary species potentially under-represented (Stat et al., 2019; Watson, Harvey, Anderson, & Kendrick, 2005). Despite these limitations, BRUVs have been shown to provide relative measures of species richness and abundance for a range of species in a diverse array of conditions and habitats (Cappel, Harvey, & Shortis, 2006).

BRUV systems with stereo-video cameras (stereo-BRUVs) enable precise measurements of body size (Harvey, Fletcher, & Shortis, 2001), which surpass estimates made by divers (Harvey et al., 2001). Both length and biomass distribution data are recognized as essential metrics for biodiversity conservation and fisheries management reporting (Langlois, Harvey, & Meeuwig, 2012). Importantly, stereo-BRUVs provide comparable body size distribution data to fisheries-dependent methods such as trawls (Cappel, Speare, & De'ath, 2004), hook and line (Langlois, Fitzpatrick, et al., 2012) and trap fishing (Langlois et al., 2015). Despite being considered unsuitable for estimating density, stereo-BRUVs provide a cost-effective and statistically powerful method to detect spatio-temporal changes in the relative abundance, length and biomass distribution of fish assemblages (Born et al., 2015; Harvey, Cappel, Kendrick, & McLean, 2013; Malcolm, Schultz, Sachs, Johnstone, & Jordan, 2015). However, in over 275 studies using stereo-BRUVs for a range of objectives (Supporting Information 1), Whitmarsh, Fairweather, and Huveneers (2017) found widespread variation in methodology, which may prevent interoperability of the data.

We provide a widely accepted protocol for the use of benthic stereo-BRUVs including information on design, field operation, image annotation, data validation, archiving and synthesis. By providing a standardized protocol for stereo-BRUVs surveys, we aim to reduce variation in methodologies among researchers, and encourage the use of Findable, Accessible, Interoperable and Reusable (FAIR, Wilkinson et al., 2016) workflows to increase the ability to synthesize datasets and answer broadscale ecological questions.

2 | STEREO-BRUVs DESIGN

Stereo-BRUV systems consist of a frame (Figure 1a), protecting two convergent video cameras inside waterproof housings, attached to a base bar (Figure 1b), with some form of baited container fixed in front of the cameras (Figure 1e). Systems are generally tethered by rope to surface buoys to facilitate relocation and retrieval




FIGURE 1 Equipment required for baited remote underwater stereo-video system surveys, including (a) mild-steel galvanized frame and bridle, (b) stereo base-bar and camera housings, (c) rope with detachable float line and two floats, (d) storage container for equipment and bait, (e) PVC bait arm (reinforced with fiberglass rod) with mesh bait bag and supporting metal diode arm, (f) metal weights for deep-water or strong current, (g) long-armed glove for handling bait, and (h) dry kit including spare cameras, spare batteries, battery charger, micro-SD card reader, micro-SD cards, standard tools, cable ties to secure bait bags, silicone grease for o-rings and (i) calibrated cameras securely fixed to face plates

(Figure 1c). Ballast can be added to frames for use in deep water or areas of strong current (Figure 1f).

2.1 | Cameras and photogrammetry

We recommend cameras with full, high-definition resolution of at least $1,920 \times 1,080$ pixels (Harvey, Goetze, McLaren, Langlois, & Shortis, 2010) and a capture rate of at least 30 frames per second (note: some models of action cameras can overheat at high resolution e.g. 4K). Higher camera resolution will improve identification of fish and the pixel selection required for measurement. Higher frame rates reduce blur on fast-moving species. To maintain stereo-calibrations, cameras must have video stabilization disabled, and a fixed focal length can facilitate measurements both close to and far from the camera systems when correctly calibrated (Boutros, Shortis, & Harvey, 2015; Shortis, Harvey, & Abdo, 2009). The field of view should be standardized and chosen to limit distortion in the image (e.g. no more than a medium angle, $\sim 95^\circ$ H-FOV). When sampling demersal

fish assemblages at typical maximum range (8 m) from the cameras, Boutros et al. (2015) suggested a camera separation < 500 mm will result in a decrease in the accuracy of measurements, with measurement precision being a function of $1/(\text{camera separation})$. Cameras are fixed to a rigid base bar to preserve the stereo-calibration required to calculate accurate length and range measurements (Boutros et al., 2015; Harvey & Shortis, 1995, 1998; Shortis & Harvey, 1998; Shortis et al., 2009). The stereo system pictured in Figure 1 uses two GoPro Hero 5 Black cameras, with camera housings separated by 700 mm with 7° convergence angle on a steel base bar, although 500 mm with a 5° convergence angle is also common.

Stereo-calibrations must be made both prior to and following a field campaign. Given the required tolerances involved with stereo-BRUVs design, we recommend seeking manufacture and calibration advice from recognized providers or adhering to strict specifications. Any changes in camera positioning (e.g. if a camera is dismounted during battery replacement) will disrupt the stereo-calibration, resulting in measurement error. For this reason, most 'off-the-shelf' housings remain unsuitable for stereo-BRUVs. Figure 1i provides an example of a camera that is secured to the housing faceplate to ensure stability. Each housing and camera should be uniquely identified, ensuring the latter are only used on the system they are calibrated for.

2.2 | Bait

As a general rule, locally sourced, sardine-type oily bait is recommended (Dorman et al., 2012), as the oil disperses to attract fish. Sourcing sardine bait locally from factory discards (e.g. fish heads, tails and guts) will reduce the survey's ecological footprint, cost of sampling and potential for disease translocation. We recommend 0.8–1 kg of roughly crushed bait, positioned between 1.2 and 1.5 m in front of the cameras with the mesh bait bag as close to the benthos as possible. Positioning further than 1.5 m from the camera will reduce the ability to identify and measure individuals. Bait should be replaced after each deployment.

2.3 | Deployment duration

Benthic stereo-BRUVs should be deployed for a standard duration. We recommend deployments of 60 min (bottom time), to allow species detection (Currey-Randall, Cappo, Simpfendorfer, Farabaugh, & Heupel, 2020), and facilitate comparison with historical data. Deployments of 30 min have been demonstrated to be sufficient for sampling particular species of finfish on shallow temperate reefs (Bernard & Götz, 2012; Harasti et al., 2015).

2.4 | Sampling design

Sampling strategies should be designed to ensure valid inferences and interpretations of resulting data (Smith, Anderson, & Pawley, 2017). We

recommend spatially balanced statistical routines, such as R package MBHDESIGN (Foster et al., 2019), which can incorporate environmental information and legacy sites to create sampling designs with known inclusion probabilities (Foster et al., 2017, 2018). Due to the need to revisit each site to retrieve stereo-BRUVs after deployment, spatially balanced designs may be inefficient for sampling large regions (>10 min transit time between samples) and clustered sampling designs may be preferred (Hill et al., 2018).

Individual stereo-BRUVs samples should be separated when set simultaneously to reduce the likelihood of non-independence due to individuals being concurrently sampled by adjacent stereo-BRUVs. Separation distance will depend on the mobility of the species and the habitat being studied; for typical demersal fish assemblages, a minimum of 400 m for 1-hr deployments is recommended (Bond, Partridge, et al., 2018) or 250 m for 30-min deployments (Cappo et al., 2001).

2.5 | Field logistics

Vessels fitted with a swinging davit arm, or pot-tipper and winch are ideal for deploying and retrieving stereo-BRUVs in deeper waters (Figure 2); however, light-weight stereo-BRUVs (Supporting Information 2) can be retrieved by hand. Comparable trap fishing retrieval methods are generally the most efficient. Each retrieval design remains dependent on the type of vessel used, stereo-BRUVs weight and size and prevailing sea conditions. Local fishers familiar

with a study location can provide valuable advice on sampling logistics. Multiple stereo-BRUVs can be deployed concurrently, with ~10 stereo-BRUVs systems providing optimum logistical efficiency for 60-min deployment times. Crepuscular periods should be avoided (if not the purpose of the study) due to demonstrated changes in fish behaviour during these times (Bond, Langlois, et al., 2018; Myers, Harvey, Saunders, & Travers, 2016). When sampling in low light conditions, both blue (450–465 nm) and white (550–560 nm) lights can be used. White can provide the best imagery for identification (Birt, Stowar, Currey-Randall, McLean, & Miller, 2019), but blue has been found to avoid potential behavioural biases and reduce backscatter from plankton at night (Fitzpatrick, McLean, & Harvey, 2013). Field methodology checklists are provided in Supporting Information 3.

2.6 | Image annotations

2.6.1 | Software

Software specifically designed to annotate and measure fish from stereo-video will substantially increase the cost-efficiency and consistency of image annotation (Gomes-Pereira et al., 2016). For stereo-video, the challenge is not the annotation, but the calibration of imagery to provide accurate length and range measurement. Annotation software and packages with measurement capabilities include Vision Measurement System (Harman, Harvey, & Kendrick, 2003), NIH Image (Dunbrack, 2006), SEBASTES package in Python (Boldt, Williams,




FIGURE 2 Methods to safely deploy and retrieve baited remote underwater stereo-video systems (BRUVs) from different size vessels using different equipment. (a) Deploying a stereo-BRUVs using an A-frame and pulley at the vessel's stern; (b) deploying a stereo-BRUV with weights and a light from the side of a vessel; (c) deploying light-weight stereo-BRUV from a small rigid inflatable (see Supporting Information 2); (d) using a 'pot winch' and 'pot tipper' to quickly retrieve stereo-BRUVs in deep water; (e) retrieving a stereo-BRUVs using a davit arm from the side of a vessel; (f) retrieving stereo-BRUVs by hand using a repurposed anchor hauler in the Philippines

Rooper, Towler, & Gauthier, 2018), STEREO MORPH package in R (Olsen & Westneat, 2015) and EventMeasure plus CAL (calibration software) from SeaGIS (seagis.com.au). EventMeasure and CAL is the most widely used option due to its established workflow, ability to create 3D stereo-calibrations and active development, which enables cost-effective and consistent point and stereo annotation of video imagery. Manual image annotation and measurement can be time consuming, but the emerging field of automated image annotation provides promise of increased cost efficiency and collection of novel metrics (Marini et al., 2018).

2.6.2 | Annotation metadata

Field metadata (Supporting Information 4) should be used to populate a unique code for each sample and annotation set. Time on the seabed should be annotated to provide a start time for the stereo-BRUVs deployment period. It is important that the link between annotations and imagery is maintained.

2.6.3 | Abundance estimates

We recommend all fish be identified to the lowest taxonomic level possible. The standard metric of abundance is MaxN, the maximum number of individuals of a given species present in a single video frame (Priede, Bagley, Smith, Creasey, & Merrett, 1994). MaxN is widely used for BRUVs (Whitmarsh et al., 2017), as it is conservative and ensures that no individual is counted more than once (Schobernd, Bacheler, & Conn, 2013). While it has frequently been suggested that MaxN underestimates both small- and large-bodied individuals, the only study so far to evaluate this has found MaxN provides a representative sample of size distributions (Coghlan et al., 2017). Synchronized and calibrated left and right cameras allow the analyst to determine the range of fish in the field of view and ensure they are within a predefined distance from the cameras. Typically, fish are counted within a maximum distance of 8 m, beyond which length estimates are likely to be inaccurate unless specialist calibrations have been conducted. Annotations of the current MaxN may be updated when individual fish are more clearly visible, and therefore easier to measure, by taking photogrammetric measurements of individual body length at the last MaxN annotated.

2.6.4 | Body size measurements

Synchronized and calibrated stereo-video streams are used to accurately measure body size. All individuals of each species should be measured at their MaxN. We recommend measuring fork length rather than total length, as it is more easily definable across a range of species. Biomass estimates typically rely on total length, but fork length to total length conversions can be used to complete these calculations (Froese & Pauly, 2019). For species where total length can be unreliable

or there is no definable fork, body size is estimated using other measures (e.g. disk measurements for rays). Photogrammetric length measurements are typically made with some degree of error, which can be minimized by measuring individuals when they are as close to cameras as possible with both the nose and the tail-fork clearly visible, still or slowly moving, at an angle $<45^\circ$ perpendicular to the cameras and straight (not bent from turning). Defining cut-offs for measurement error across projects will help to maintain accurate and precise body size estimates, we provide recommended stereo-measurement length rules for EventMeasure in Supporting Information 5. If fish cannot be measured within these parameters, a '3D point' may be used for annotation, which records the 3D location of the fish to ensure it is within the sampling area (Harvey, Fletcher, Shortis, & Kendrick, 2004). To create a relative abundance metric standardized to a consistent sample area, abundance should be summed from the lengths and 3D points at the MaxN for each species. For biomass estimates, 3D points provide a basis for extrapolating a median length value to fish that could not be measured (Wilson, Graham, Holmes, MacNeil, & Ryan, 2018). When large tightly packed schools are encountered, fish that cannot be measured should have 3D points. When lengths or 3D points are not possible for every fish, multiple individuals can be assigned to a single length or 3D point, but care should be taken to represent the range of body sizes within a school.

2.6.5 | Behaviour

A range of behavioural observations, including time of first arrival, time to first feed and minimum approach distance, may also be calculated (Coghlan et al., 2017; Goetze et al., 2017).

2.6.6 | Interoperable and reproducible annotations

Video imagery enables annotators to work collaboratively to ensure identifications are consistent. A library of reference images, such as that supported by EventMeasure, will assist with identification and training. It is acknowledged that some genera cannot be consistently identified to species level from imagery, so individuals are recorded at genus-family levels (e.g. flathead: *Platycephalus* spp). For unidentified individuals, a common convention is that fish that are potentially identifiable later are annotated to *Genus* sp1–10, this permits a batch rename at a later stage if the species is successfully identified. Individuals that are clearly unidentifiable to species are annotated as *Genus* sp.

2.6.7 | Habitat classification

Information on relief, habitat types and benthic composition (e.g. percent cover of benthos types) should be recorded from each deployment (Bennett et al., 2016; Collins et al., 2017), to facilitate investigation of fish-habitat relationships and to enable the sampling field of view to be standardized or controlled for subsequent data

analysis (McLean et al., 2016). It is important that these data are annotated consistently and it is recommended that they are mapped to the CATAMI classification scheme (Althaus et al., 2015) and a 0–5 estimate of benthic relief (Polunin & Roberts, 1993; Wilson, Graham, & Polunin, 2007). An example of habitat composition and relief annotation schema is provided in a GitHub repository (Langlois, 2017). Forward facing imagery can be annotated in a range of software, including TransectMeasure from SeaGIS (seagis.com.au), Benthobox (https://benthobox.com), CoralNet (https://coralnet.ucsd.edu/) and Squidle+ (https://squidle.org), which all provide suitable workflows and comparable data outputs.

2.6.8 | Quality control and data curation

Quality control and data curation are vital to ensure FAIR data workflows (Wilkinson et al., 2016). All corrections should be made within the original annotation files to ensure data consistency over time. We recommend the following approaches to ensure quality control:

- Annotators should complete 'training' videos where species IDs and MaxN are known and can be used to assess competency.
- A different annotator should complete the MaxN and length measurement annotations to provide an independent check of the species identifications.
- Quality assurance should be carried out by a senior video analyst or researcher and involves a random review of 10% of annotated videos and data within a project. If accuracy is below 95% for all identifications and estimates of MaxN, reannotation should be undertaken.
- Unique identifiers of annotators and dates of when imagery was annotated should be maintained to provide a data checking trail (see Supporting Information 4).

R workflows and function packages are provided in a GitHub repository (Langlois, 2020) to enable validation with regional species lists and likely minimum and maximum sizes for each species.

2.6.9 | Data storage, discoverability and release

We encourage open data policies and recommend archiving and sharing stereo-BRUVs annotations on global biodiversity data repositories, such as Ocean Biogeographic Information System, Global Biodiversity Information Facility and the recently developed GlobalArchive (globalarchive.org). GlobalArchive is a centralized repository that allows open access and private sharing of fish image annotation data from stereo-BRUVs or similar imagery-based sampling techniques. GlobalArchive allows users to store data in a standardized and secure manner and makes meta-data discoverable, thus encouraging collaboration and synthesis of datasets within the community of practice. We recommend all quality-controlled annotation data and any associated calibration, taxa and

habitat data should be uploaded to GlobalArchive and we encourage that all data should be made publicly available via the public data option. As an example, the Australian standards for data management, discoverability and release are provided in Supporting Information 6.

3 | CONCLUSION

Globally, stereo-BRUVs usage is increasing rapidly. The standardization of stereo-BRUVs surveys and annotation will facilitate the synthesis of comparable data over continental and global scales and provide rich and interoperable data to inform natural resource management. Variation in methodology has constrained the interoperability of these data to date (Whitmarsh et al., 2017), we encourage researchers to standardize and share technical improvements and issues via an established on-line forum or working group (Supporting Information 7).

Achieving consistent field methodology and FAIR annotation, with data archiving and sharing protocols, provide the greatest barrier to the globally consistent uptake and impact of stereo-BRUVs. We provide a standardized protocol that will reduce methodological variation among researchers and encourage the use of FAIR workflows to increase the ability to synthesize datasets and answer a range of ecological questions.

ACKNOWLEDGEMENTS

The authors would like to thank James Seager (SeaGIS.com.au) for support with software and both James Seager and Ray Scott for stereo equipment and advice. Researchers T.L., B.A.G., J.W., N.B. and J.M. were supported by the Marine Biodiversity Hub through funding from the Australian Government's National Environmental Science Program. Data validation scripts and GlobalArchive.org were supported by the Australian Research Data Commons, the Gorgon-Barrow Island Gorgon Barrow Island Net Conservation Benefits Fund, administered by the Government of Western Australia and the BHP/UWA Biodiversity and Societal Benefits of Restricted Access Areas collaboration. This manual has been endorsed by GOOS Biology and Ecosystems Panel of Experts as a globally accepted best practice for conducting research with baited remote underwater stereo-video systems.

AUTHORS' CONTRIBUTIONS

All authors conceived the ideas and designed methodology; T.L. and J.G. led the writing of the manuscript. All authors contributed critically to the drafts and gave final approval for publication.

PEER REVIEW

The peer review history for this article is available at <https://publons.com/publon/10.1111/2041-210X.13470>.

DATA AVAILABILITY STATEMENT

No data were presented.

ORCID

Tim Langlois  <https://orcid.org/0000-0001-6404-4000>
 Jordan Goetze  <https://orcid.org/0000-0002-3090-9763>
 Todd Bond  <https://orcid.org/0000-0001-6064-7015>
 Jacquomo Monk  <https://orcid.org/0000-0002-1874-0619>
 Rene A. Abesamis  <https://orcid.org/0000-0001-7456-1415>
 Jacob Asher  <https://orcid.org/0000-0003-3997-8439>
 Neville Barrett  <https://orcid.org/0000-0002-6167-1356>
 Anthony T. F. Bernard  <https://orcid.org/0000-0003-0482-6283>
 Phil J. Bouchet  <https://orcid.org/0000-0002-2144-2049>
 Matthew J. Birt  <https://orcid.org/0000-0003-2233-0716>
 Leanne M. Currey-Randall  <https://orcid.org/0000-0002-3772-1288>
 David V. Fairclough  <https://orcid.org/0000-0002-9620-5064>
 David Harasti  <https://orcid.org/0000-0002-2851-9838>
 Michelle R. Heupel  <https://orcid.org/0000-0002-8245-7332>
 Charlie Huveneers  <https://orcid.org/0000-0001-8937-1358>
 Daniel Ierodiaconou <http://orcid.org/0000-0002-7832-4801>
 Nathan A. Knott  <https://orcid.org/0000-0002-7873-0412>
 Steve Lindfield  <https://orcid.org/0000-0002-7933-6206>
 Hamish A. Malcolm  <https://orcid.org/0000-0001-7315-1537>
 Dianne McLean  <https://orcid.org/0000-0002-0306-8348>
 Mark Meekan  <https://orcid.org/0000-0002-3067-9427>
 Stephen J. Newman  <https://orcid.org/0000-0002-5324-5568>
 Fernanda A. Rolim  <https://orcid.org/0000-0003-3761-3970>
 Benjamin J. Saunders  <https://orcid.org/0000-0003-1929-518X>
 Adam N. H. Smith  <https://orcid.org/0000-0003-0059-6206>
 Michael J. Travers  <https://orcid.org/0000-0002-3072-1699>
 Sasha K. Whitmarsh  <https://orcid.org/0000-0001-8934-2354>
 Joel Williams  <https://orcid.org/0000-0002-4173-3855>
 Euan S. Harvey  <https://orcid.org/0000-0002-9069-4581>

REFERENCES

- Althaus, F., Hill, N., Ferrari, R., Edwards, L., Przeslawski, R., Schönborg, C. H. L., ... Gowlett-Holmes, K. (2015). A standardised vocabulary for identifying benthic biota and substrata from underwater imagery: The CATAMI classification scheme. *PLoS ONE*, 10, e0141039. <https://doi.org/10.1371/journal.pone.0141039>
- Bennett, K., Wilson, S. K., Shedrawi, G., McLean, D. L., & Langlois, T. J. (2016). Can diver operated stereo-video surveys for fish be used to collect meaningful data on benthic coral reef communities? *Limnology and Oceanography: Methods*. <https://doi.org/10.1002/lom3.10141>
- Bernard, A. T. F., & Götz, A. (2012). Bait increases the precision in count data from remote underwater video for most subtidal reef fish in the warm-temperate Agulhas bioregion. *Marine Ecology Progress Series*, 471, 235–252. <https://doi.org/10.3354/meps10039>
- Birt, M. J., Stowar, M., Currey-Randall, L. M., McLean, D. L., & Miller, K. J. (2019). Comparing the effects of different coloured artificial illumination on diurnal fish assemblages in the lower mesophotic zone. *Marine Biology*, 166, 154. <https://doi.org/10.1007/s00227-019-3595-0>
- Boldt, J. L., Williams, K., Rooper, C. N., Towler, R. H., & Gauthier, S. (2018). Development of stereo camera methodologies to improve pelagic fish biomass estimates and inform ecosystem management in marine waters. *Fisheries Research*, 198, 66–77. <https://doi.org/10.1016/j.fishres.2017.10.013>
- Bond, T., Langlois, T. J., Partridge, J. C., Birt, M. J., Malseed, B. E., Smith, L., & McLean, D. L. (2018). Diel shifts and habitat associations of fish assemblages on a subsea pipeline. *Fisheries Research*, 206, 220–234. <https://doi.org/10.1016/j.fishres.2018.05.011>
- Bond, T., Partridge, J. C., Taylor, M. D., Langlois, T. J., Malseed, B. E., Smith, L. D., & McLean, D. L. (2018). Fish associated with a subsea pipeline and adjacent seafloor of the North West Shelf of Western Australia. *Marine Environmental Research*, 141, 53–65. <https://doi.org/10.1016/j.marenvres.2018.08.003>
- Bornt, K. R., McLean, D. L., Langlois, T. J., Harvey, E. S., Bellchambers, L. M., Evans, S. N., & Newman, S. J. (2015). Targeted demersal fish species exhibit variable responses to long-term protection from fishing at the Houtman Abrolhos Islands. *Coral Reefs*, 34, 1297–1312. <https://doi.org/10.1007/s00338-015-1336-5>
- Boutros, N., Shortis, M. R., & Harvey, E. S. (2015). A comparison of calibration methods and system configurations of underwater stereo-video systems for applications in marine ecology. *Limnology and Oceanography: Methods*, 13, 224–236. <https://doi.org/10.1002/lom3.10020>
- Cappo, M., De'ath, G., Stowar, M., Johansson, C., & Doherty, P. (2009). *The influence of zoning (closure to fishing) on fish communities of the deep shoals and reef bases of the southern Great Barrier Reef Marine Park*. Cairns, Australia: Reef and Rainforest Research Centre Limited.
- Cappo, M., Harvey, E., & Shortis, M. (2006). Counting and measuring fish with baited video techniques – An overview. In J. M. Lyle, D. M. Furlani, & C. D. Buxton (Eds.), *Proceedings of the 2006 Australian Society of Fish Biology Conference and Workshop Cuttingedge Technologies in Fish and Fisheries Science*. Hobart, August 2006. (pp. 101–114)
- Cappo, M., Speare, P., & De'ath, G. (2004). Comparison of baited remote underwater video stations (BRUVS) and prawn (shrimp) trawls for assessments of fish biodiversity in inter-reefal areas of the Great Barrier Reef Marine Park. *Journal of Experimental Marine Biology and Ecology*, 302, 123–152. <https://doi.org/10.1016/j.jembe.2003.10.006>
- Cappo, M., Speare, P., Wassenberg, T. J., Rees, M., Heyward, A., & Pitcher, R. (2001). Use of baited remote underwater video stations (BRUVS) to survey demersal fish - How deep and meaningful. In E. S. Harvey, & M. Cappo (Eds.), *Direct sensing of the size frequency and abundance of target and non-target fauna in Australian fisheries - A national workshop* (pp. 63–71). Rottneest Island: Fisheries Research Development Corporation.
- Coghlan, A. R., McLean, D. L., Harvey, E. S., & Langlois, T. J. (2017). Does fish behaviour bias abundance and length information collected by baited underwater video? *Journal of Experimental Marine Biology and Ecology*, 497, 143–151. <https://doi.org/10.1016/j.jembe.2017.09.005>
- Collins, D. L., Langlois, T. J., Bond, T., Holmes, T. H., Harvey, E. S., Fisher, R., & McLean, D. L. (2017). A novel stereo-video method to investigate fish-habitat relationships. *Methods in Ecology and Evolution/British Ecological Society*, 8, 116–125. <https://doi.org/10.1111/2041-210X.12650>
- Currey-Randall, L. M., Cappo, M., Simpfendorfer, C. A., Farabaugh, N. F., & Heupel, M. R. (2020). Optimal soak times for baited remote underwater video station surveys of reef-associated elasmobranchs. *PLoS ONE*, 15, e0231688. <https://doi.org/10.1371/journal.pone.0231688>
- Dorman, S. R., Harvey, E. S., & Newman, S. J. (2012). Bait effects in sampling coral reef fish assemblages with stereo-BRUVs. *PLoS ONE*, 7, e41538. <https://doi.org/10.1371/journal.pone.0041538>
- Dunbrack, R. L. (2006). In situ measurement of fish body length using perspective-based remote stereo-video. *Fisheries Research*, 82, 327–331. <https://doi.org/10.1016/j.fishres.2006.08.017>
- Fitzpatrick, C., McLean, D., & Harvey, E. S. (2013). Using artificial illumination to survey nocturnal reef fish. *Fisheries Research*, 146, 41–50. <https://doi.org/10.1016/j.fishres.2013.03.016>
- Foster, S. D., Hosack, G. R., Lawrence, E., Przeslawski, R., Hedge, P., Caley, M. J., ... Hayes, K. R. (2017). Spatially balanced designs that incorporate legacy sites. *Methods in Ecology and*

- Evolution/British Ecological Society*, 8, 1433–1442. <https://doi.org/10.1111/2041-210X.12782>
- Foster, S. D., Hosack, G. R., Monk, J., Lawrence, E., Barrett, N. S., Williams, A., & Przeslawski, R. (2019). Spatially balanced designs for transect-based surveys. *Methods in Ecology and Evolution*, 11(1), 95–105. <https://doi.org/10.1111/2041-210X.13321>
- Foster, S. D., Monk, J., Lawrence, E., Hayes, K. R., Hosack, G. R., & Przeslawski, R. (2018). Statistical considerations for monitoring and sampling. In R. Przeslawski & S. Foster (Eds.), *Field manuals for marine sampling to monitor Australian waters* (pp. 23–41). National Environmental Science Programme (NESP). Retrieved from <https://survey-design-field-manual.github.io/>
- Frøese, R., & Pauly, D. (2019). *FishBase*. www.fishbase.org
- Goetze, J. S., Bond, T., McLean, D. L., Saunders, B. J., Langlois, T. J., Lindfield, S., ... Harvey, E. S. (2019). A field and video analysis guide for diver operated stereo-video. *Methods in Ecology and Evolution*, 10, 1083–1090. <https://doi.org/10.1111/2041-210X.13189>
- Goetze, J. S., Januchowski-Hartley, F. A., Claudet, J., Langlois, T. J., Wilson, S. K., & Jupiter, S. D. (2017). Fish wariness is a more sensitive indicator to changes in fishing pressure than abundance, length or biomass. *Ecological Applications*, 27, 1178–1189. <https://doi.org/10.1002/eap.1511>
- Goetze, J. S., Jupiter, S. D., Langlois, T. J., Wilson, S. K., Harvey, E. S., Bond, T., & Naisilisili, W. (2015). Diver operated video most accurately detects the impacts of fishing within periodically harvested closures. *Journal of Experimental Marine Biology and Ecology*, 462, 74–82. <https://doi.org/10.1016/j.jembe.2014.10.004>
- Gomes-Pereira, J. N., Auger, V., Beisiegel, K., Benjamin, R., Bergmann, M., Bowden, D., ... Santos, R. S. (2016). Current and future trends in marine image annotation software. *Progress in Oceanography*, 149, 106–120. <https://doi.org/10.1016/j.pocean.2016.07.005>
- Gray, A. E., Williams, I. D., Stamoulis, K. A., Boland, R. C., Lino, K. C., Hauk, B. B., ... Kosaki, R. K. (2016). Comparison of reef fish survey data gathered by open and closed circuit SCUBA divers reveals differences in areas with higher fishing pressure. *PLoS ONE*, 11, e0167724. <https://doi.org/10.1371/journal.pone.0167724>
- Harasti, D., Malcolm, H., Gallen, C., Coleman, M. A., Jordan, A., & Knott, N. A. (2015). Appropriate set times to represent patterns of rocky reef fishes using baited video. *Journal of Experimental Marine Biology and Ecology*, 463, 173–180. <https://doi.org/10.1016/j.jembe.2014.12.003>
- Hardinge, J., Harvey, E. S., Saunders, B. J., & Newman, S. J. (2013). A little bait goes a long way: The influence of bait quantity on a temperate fish assemblage sampled using stereo-BRUVs. *Journal of Experimental Marine Biology and Ecology*, 449, 250–260. <https://doi.org/10.1016/j.jembe.2013.09.018>
- Harman, N., Harvey, E. S., & Kendrick, G. A. (2003). Differences in fish assemblages from different reef habitats at Hamelin Bay, south-western Australia. *Marine and Freshwater Research*, 54, 177–184. <https://doi.org/10.1071/MF02040>
- Harvey, E. S., Cappel, M., Butler, J., Hall, N., & Kendrick, G. (2007). Bait attraction affects the performance of remote underwater video stations in assessment of demersal fish community structure. *Marine Ecology Progress Series*, 350, 245–254. <https://doi.org/10.3354/meps07192>
- Harvey, E. S., Cappel, M., Kendrick, G. A., & McLean, D. L. (2013). Coastal fish assemblages reflect geological and oceanographic gradients within an Australian zootone. *PLoS ONE*, 8, e80955. <https://doi.org/10.1371/journal.pone.0080955>
- Harvey, E., Fletcher, D., & Shortis, M. (2001). Improving the statistical power of length estimates of reef fish: A comparison of estimates determined visually by divers with estimates produced by a stereo-video system. *Fishery Bulletin-National Oceanic and Atmospheric Administration*, 99, 72–80.
- Harvey, E., Fletcher, D., Shortis, M. R., & Kendrick, G. A. (2004). A comparison of underwater visual distance estimates made by scuba divers and a stereo-video system: Implications for underwater visual census of reef fish abundance. *Marine and Freshwater Research*, 55, 573–580. <https://doi.org/10.1071/MF03130>
- Harvey, E. S., Goetze, J. S., McLaren, B., Langlois, T., & Shortis, M. R. (2010). Influence of range, angle of view, image resolution and image compression on underwater stereo-video measurements: High-definition and broadcast-resolution video cameras compared. *Marine Technology Society Journal*, 44, 75–85. <https://doi.org/10.4031/MTSJ.44.1.3>
- Harvey, E. S., Santana-Garcon, J. S., Goetze, J. S., Saunders, B. J., & Cappel, M. (2018). The use of stationary underwater video for sampling sharks. In J. C. Carrier, M. R. Heithaus, & C. A. Simpfendorfer (Eds.), *Shark research: Emerging technologies and applications for the field and laboratory* (pp. 111–132). Boca Raton, FL: CRC Press.
- Harvey, E., & Shortis, M. (1995). A system for stereo-video measurement of sub-tidal organisms. *Marine Technology Society Journal*, 29, 10–22.
- Harvey, E. S., & Shortis, M. R. (1998). Calibration stability of an underwater stereo-video system: Implications for measurement accuracy and precision. *Marine Technology Society Journal*, 32, 3.
- Heyns-Veale, E. R., Bernard, A. T. F., Richoux, N. B., Parker, D., Langlois, T. J., Harvey, E. S., & Götz, A. (2016). Depth and habitat determine assemblage structure of South Africa's warm-temperate reef fish. *Marine Biology*, 163, 1–17. <https://doi.org/10.1007/s00227-016-2933-8>
- Hill, N. A., Barrett, N., Ford, J. H., Peel, D., Foster, S., Lawrence, E., ... Hayes, K. R. (2018). Developing indicators and a baseline for monitoring demersal fish in data-poor, offshore Marine Parks using probabilistic sampling. *Ecological Indicators*, 89, 610–621. <https://doi.org/10.1016/j.ecolind.2018.02.039>
- Langlois, T. J. (2017). Habitat annotation of forward facing benthic imagery. Retrieved from <https://github.com/TimLanglois/Habitat-annotation-of-forward-facing-benthic-imagery>
- Langlois, T. J. (2020). Stereo or mono video annotation workflows. Retrieved from <https://github.com/UWAMEGFisheries/Stereo-or-mono-video-annotation-workflows>
- Langlois, T. J., Fitzpatrick, B. R., Fairclough, D. V., Wakefield, C. B., Hesp, S. A., McLean, D. L., ... Meeuwij, J. J. (2012). Similarities between line fishing and baited stereo-video estimations of length-frequency: Novel application of kernel density estimates. *PLoS ONE*, 7, e45973. <https://doi.org/10.1371/journal.pone.0045973>
- Langlois, T. J., Harvey, E. S., & Meeuwij, J. J. (2012). Strong direct and inconsistent indirect effects of fishing found using stereo-video: Testing indicators from fisheries closures. *Ecological Indicators*, 23, 524–534. <https://doi.org/10.1016/j.ecolind.2012.04.030>
- Langlois, T. J., Newman, S. J., Cappel, M., Harvey, E. S., Rome, B. M., Skepper, C. L., & Wakefield, C. B. (2015). Length selectivity of commercial fish traps assessed from in situ comparisons with stereo-video: Is there evidence of sampling bias? *Fisheries Research*, 161, 145–155. <https://doi.org/10.1016/j.fishres.2014.06.008>
- Langlois, T. J., Radford, B. T., Van Niel, K. P., Meeuwij, J. J., Pearce, A. F., Rousseaux, C. S. G., ... Harvey, E. S. (2012). Consistent abundance distributions of marine fishes in an old, climatically buffered, infertile seascape: Abundance distributions of fishes in stable seascapes. *Global Ecology and Biogeography*, 21, 886–897. <https://doi.org/10.1111/j.1466-8238.2011.00734.x>
- Lindfield, S. J., Harvey, E. S., McIlwain, J. L., & Halford, A. R. (2014). Silent fish surveys: Bubble-free diving highlights inaccuracies associated with SCUBA-based surveys in heavily fished areas. *Methods in Ecology and Evolution*, 5, 1061–1069. <https://doi.org/10.1111/2041-210X.12262>
- Malcolm, H. A., Schultz, A. L., Sachs, P., Johnstone, N., & Jordan, A. (2015). Decadal changes in the abundance and length of snapper (*Chrysophrys auratus*) in subtropical marine sanctuaries. *PLoS ONE*, 10, e0127616. <https://doi.org/10.1371/journal.pone.0127616>
- Marini, S., Fanelli, E., Sbragaglia, V., Azzurro, E., Del Rio Fernandez, J., & Aguzzi, J. (2018). Tracking fish abundance by underwater image

- recognition. *Scientific Reports*, 8, 13748. <https://doi.org/10.1038/s41598-018-32089-8>
- McLean, D. L., Langlois, T. J., Newman, S. J., Holmes, T. H., Birt, M. J., Bornt, K. R., ... Fisher, R. (2016). Distribution, abundance, diversity and habitat associations of fishes across a bioregion experiencing rapid coastal development. *Estuarine, Coastal and Shelf Science*, 178, 36–47. <https://doi.org/10.1016/j.ecss.2016.05.026>
- Myers, E. M. V., Harvey, E. S., Saunders, B. J., & Travers, M. J. (2016). Fine-scale patterns in the day, night and crepuscular composition of a temperate reef fish assemblage. *Marine Ecology*, 37(3), 668–678. <https://doi.org/10.1111/maec.12336>
- Olsen, A. M., & Westneat, M. W. (2015). StereoMorph: An R package for the collection of 3D landmarks and curves using a stereo camera set-up. *Methods in Ecology and Evolution*, 6(3), 351–356. <https://doi.org/10.1111/2041-210X.12326>
- Polunin, N. V. C., & Roberts, C. M. (1993). Greater biomass and value of target coral-reef fishes in two small Caribbean marine reserves. *Marine Ecology Progress Series*, 100, 167–176. <https://doi.org/10.3354/meps100167>
- Priede, I. G., Bagley, P. M., Smith, A., Creasey, S., & Merrett, N. R. (1994). Scavenging deep demersal fishes of the Porcupine Seabight, north-east Atlantic: Observations by baited camera, trap and trawl. *Journal of the Marine Biological Association of the United Kingdom*, 74, 481–498. <https://doi.org/10.1017/S0025315400047615>
- Schobernd, Z. H., Bacheler, N. M., & Conn, P. B. (2013). Examining the utility of alternative video monitoring metrics for indexing reef fish abundance. *Canadian Journal of Fisheries and Aquatic Sciences*, 71, 464–471. <https://doi.org/10.1139/cjfas-2013-0086>
- Shortis, M. R., & Harvey, E. S. (1998). Design and calibration of an underwater stereo-video system for the monitoring of marine fauna populations. *International Archives of Photogrammetry and Remote Sensing*, 32, 792–799.
- Shortis, M., Harvey, E., & Abdo, D. (2009). A review of underwater stereo-image measurement for marine biology and ecology applications: An annual review. *Oceanography and Marine Biology Annual Review*, 47, 257–292.
- Smith, A. N. H., Anderson, M. J., & Pawley, M. D. M. (2017). Could ecologists be more random? Straightforward alternatives to haphazard spatial sampling. *Ecography*, 40, 1251–1255. <https://doi.org/10.1111/ecog.02821>
- Speed, C. W., Rees, M. J., Cure, K., Vaughan, B., & Meekan, M. G. (2019). Protection from illegal fishing and shark recovery restructures meso-predatory fish communities on a coral reef. *Ecology and Evolution*, 9, 10553–10566. <https://doi.org/10.1002/ece3.5575>
- Stat, M., John, J., DiBattista, J. D., Newman, S. J., Bunce, M., & Harvey, E. S. (2019). Combined use of eDNA metabarcoding and video surveillance for the assessment of fish biodiversity. *Conservation Biology*, 33, 196–205. <https://doi.org/10.1111/cobi.13183>
- Watson, D. L., Harvey, E. S., Anderson, M. J., & Kendrick, G. A. (2005). A comparison of temperate reef fish assemblages recorded by three underwater stereo-video techniques. *Marine Biology*, 148, 415–425. <https://doi.org/10.1007/s00227-005-0090-6>
- Wellington, C. M., Harvey, E. S., Wakefield, C. B., Langlois, T. J., Williams, A., White, W. T., & Newman, S. J. (2018). Peak in biomass driven by larger-bodied meso-predators in demersal fish communities between shelf and slope habitats at the head of a submarine canyon in the south-eastern Indian Ocean. *Continental Shelf Research*, 167, 55–64. <https://doi.org/10.1016/j.csr.2018.08.005>
- Whitmarsh, S. K., Fairweather, P. G., & Huveneers, C. (2017). What is Big BRUVver up to? Methods and uses of baited underwater video. *Reviews in Fish Biology and Fisheries*, 27, 53–73. <https://doi.org/10.1007/s11160-016-9450-1>
- Wilkinson, M. D., Dumontier, M., Aalbersberg, I. J. J., Appleton, G., Axton, M., Baak, A., ... Mons, B. (2016). The FAIR guiding principles for scientific data management and stewardship. *Scientific Data*, 3, 160018. <https://doi.org/10.1038/sdata.2016.18>
- Wilson, S. K., Graham, N. A. J., Holmes, T. H., MacNeil, M. A., & Ryan, N. M. (2018). Visual versus video methods for estimating reef fish biomass. *Ecological Indicators*, 85, 146–152. <https://doi.org/10.1016/j.ecolind.2017.10.038>
- Wilson, S. K., Graham, N. A. J., & Polunin, N. V. C. (2007). Appraisal of visual assessments of habitat complexity and benthic composition on coral reefs. *Marine Biology*, 151, 1069–1076. <https://doi.org/10.1007/s00227-006-0538-3>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: Langlois T, Goetze J, Bond T, et al.

A field and video annotation guide for baited remote underwater stereo-video surveys of demersal fish

assemblages. *Methods Ecol Evol.* 2020;00:1–9. <https://doi.org/10.1111/2041-210X.13470>