

MICT-5101: Probability and Stochastic Process¹

Dr. Md. Rezaul Karim

PhD(KULeuven & UHasselt), MS(Biostatistics), MS(Statistics)

Professor, Department of Statistics and Data Science

Jahangirnagar University (JU), Savar, Dhaka - 1342, Bangladesh

MS-2024



¹These course slides should not be reproduced nor used by others (without permission).

Lecture Outline I

1 Chapter 7. Renewal Process and Branching Process

1.1 Classification of Renewal Process

1.2 Distribution of Renewal Process

1.3 Finding the Renewal Function

1.4 Branching Processes

1.5 Key Features of Branching Process

1.6 Mathematical Model of a Branching Process

1.7 Mean of Branching Process

1.8 Variance of Branching Process



Chapter 7. Renewal Process and Branching Process



1 Chapter 7. Renewal Process and Branching Process

1.1 Classification of Renewal Process

1.2 Distribution of Renewal Process

1.3 Finding the Renewal Function

1.4 Branching Processes

1.5 Key Features of Branching Process

1.6 Mathematical Model of a Branching Process

1.7 Mean of Branching Process

1.8 Variance of Branching Process




Renewal Process

We have seen that a Poisson process is a counting process for which the times between successive events are independent and identically distributed **exponential** random variables. One possible generalization is to consider a counting process for which the times between successive events are independent and identically distributed with an arbitrary distribution. Such a counting process is called a **renewal process**.

Definition 1.1

Let $\{N(t), t \geq 0\}$ be a counting process and let X_n denote the time between the $(n-1)$ st and the n th event of this process, $n \geq 1$. If the sequence of nonnegative random variables $\{X_1, X_2, X_3, \dots\}$ is independent and identically distributed, then the counting process $\{N(t), t \geq 0\}$ is said to be a **renewal process**. It was introduced as a generalization of **Poisson process**. It was considered as a process in continuous time.





Let X_1, X_2, \dots be i.i.d random variables with $\mathbb{E}[X_i] < \infty$, and $P(X_i > 0) = 1$. Let

$$S_0 = 0, \quad S_n = X_1 + \dots + X_n, \quad n \geq 1.$$

Define

$$N(t) = \max\{n : S_n \leq t\}.$$

Then $\{N(t), t \geq 0\}$ is called a **renewal process**.

- Events are called “**renewals**”. The interarrival times between events X_1, X_2, \dots are also called “renewals”
- A more general definition allows the first renewal X_1 to be of a different distribution, called a **delayed renewal process**.



Renewal Process

Example:

Suppose, we have an infinite number of electric light bulbs whose life times are iid random variables. Suppose we use a single light bulb at a time and we immediately replace it by a new one if it fail. Let $N(t)$ represents the number of light bulbs that have failed by time t , then $\{N(t); t \geq 0\}$ is a Renewal Process.



Renewal process used in a variety of situations such as

- demography
- manpower studies
- reliability
- replacement and maintenance
- inventory control
- queuing process
- simulation and so on.



Classification of Renewal Process

1 Delayed/General/Modified renewal process

A renewal process $\{N(t); t \geq 0\}$ is said to be delayed/general/modified renewal process if the first arrival time X_1 has a different distribution than the other X_2, X_3, \dots

2 Equilibrium renewal process

A renewal process $\{N(t); t \geq 0\}$ is said to be equilibrium renewal process if the distribution of the 1st interarrival time is specified which is

$$F_e(t) = \int_0^t \frac{1 - F(x)}{\mu} dx$$

where μ = mean of F . and it different from the iid distributed other.



Difference between Renewal Process and Poisson Process

(i) Interarrival times of the renewal process $\{N(t); g \geq 0\}$ are iid with arbitrary distribution.

(ii) The mean function is

$$m(t) = \sum_{n=1}^{\infty} F_n(t)$$

(iii) Renewal process is the generalization of Poisson process.

(iv) mean function may be linear.

(i) Interarrival times have independently identical exponentially distributed.

(ii) The mean function is

$$m(t) = \lambda t$$

(iii) Renewal process is a particular case of renewal process.

(iv) Mean function always linear.



General Renewal Process: Distribution of $N(t)$

Let X_1, X_2, X_3, \dots be i.i.d. random variables representing the inter-arrival times in a renewal process. Define the total time of the n -th renewal by:

$$S_n = X_1 + X_2 + \dots + X_n$$

The number of renewals $N(t)$ by time t is:

$$N(t) = \max\{n : S_n \leq t\}$$

The probability that exactly n renewals occur by time t is:

$$P(N(t) = n) = P(S_n \leq t \text{ and } S_{n+1} > t)$$

This can be written as the difference of cumulative distribution functions (CDFs) of S_n :

$$P(N(t) = n) = P(S_n \leq t) - P(S_{n+1} \leq t) = F_n(t) - F_{n+1}(t)$$



Example 7.1 Suppose that $P\{X_n = i\} = p(1-p)^{i-1}, i \geq 1$. That is, suppose that the interarrival distribution is geometric. Now $S_1 = X_1$ may be interpreted as the number of trials necessary to get a single success when each trial is independent and has a probability p of being a success. Similarly, S_n may be interpreted as the number of trials necessary to attain n successes, and hence has the negative binomial distribution

$$P\{S_n = k\} = \begin{cases} \binom{k-1}{n-1} p^n (1-p)^{k-n}, & k \geq n \\ 0, & k < n \end{cases}$$

Thus, from Equation (7.3) we have that

$$\begin{aligned} P\{N(t) = n\} &= \sum_{k=n}^{[t]} \binom{k-1}{n-1} p^n (1-p)^{k-n} \\ &\quad - \sum_{k=n+1}^{[t]} \binom{k-1}{n} p^{n+1} (1-p)^{k-n-1} \end{aligned}$$



Equivalently, since an event independently occurs with probability p at each of the times $1, 2, \dots$

$$P\{N(t) = n\} = \binom{[t]}{n} p^n (1-p)^{[t]-n} \quad \blacksquare$$

Another expression for $P(N(t) = n)$ can be obtained by conditioning on S_n . This yields

$$P(N(t) = n) = \int_0^\infty P(N(t) = n | S_n = y) f_{S_n}(y) dy$$

Now, if the n th event occurred at time $y > t$, then there would have been less than n events by time t . On the other hand, if it occurred at a time $y \leq t$, then there would be exactly n events by time t if the next interarrival exceeds $t - y$. Consequently,

$$\begin{aligned} P(N(t) = n) &= \int_0^t P(X_{n+1} > t - y | S_n = y) f_{S_n}(y) dy \\ &= \int_0^t \bar{F}(t - y) f_{S_n}(y) dy \end{aligned}$$

where $\bar{F} = 1 - F$.



Example 7.2 If $F(x) = 1 - e^{-\lambda x}$ then S_n , being the sum of n independent exponentials with rate λ , will have a gamma (n, λ) distribution. Consequently, the preceding identity gives

$$\begin{aligned}P(N(t) = n) &= \int_0^t e^{-\lambda(t-y)} \frac{\lambda e^{-\lambda y} (\lambda y)^{n-1}}{(n-1)!} dy \\&= \frac{\lambda^n e^{-\lambda t}}{(n-1)!} \int_0^t y^{n-1} dy \\&= e^{-\lambda t} \frac{(\lambda t)^n}{n!}\end{aligned}$$



Finding the Renewal Function

The mean value of a renewal process $\{N(t); t \geq 0\}$ is known as the renewal function. Let $P_n(t) = P(N(t) = n)$ is a probability mass function of $\{N(t); t \geq 0\}$. Then the expression of the renewal function is

$$\begin{aligned}m(t) &= E[N(t)] \\&= \sum_{n=1}^{\infty} nP_n(t) \\&= P_1(t) + 2P_2(t) + 3P_3(t) + 4P_4(t) + \cdots \\&= P_1(t) + P_2(t) + P_3(t) + P_4(t) + \cdots \\&\quad + P_2(t) + P_3(t) + P_4(t) + \cdots \\&\quad + P_3(t) + P_4(t) + \cdots \\&= \Pr(N(t) \geq 1) + \Pr(N(t) \geq 2) + \Pr(N(t) \geq 3) + \cdots \\&= \sum_{n=1}^{\infty} \Pr(N(t) \geq n) = \sum_{n=1}^{\infty} \Pr(S_n \leq t) = \sum_{n=1}^{\infty} F_n(t)\end{aligned}$$



As $t \rightarrow \infty$, if the inter-arrival times have mean μ , the renewal process exhibits the following asymptotic behavior:

$$\lim_{t \rightarrow \infty} \frac{m(t)}{t} = \frac{1}{\mu}$$



Introduction to Branching Processes

- A **branching process** is a stochastic model that describes the evolution of populations where each individual in the population can produce offspring.
- Individuals reproduce and generate offspring, and the population size evolves over time.
- The key feature of a branching process is its probabilistic nature.
- The **Galton-Watson branching process** is a common model for such processes.



Key Features

- **Generations:** The population is divided into generations.
- **Offspring Distribution:** The number of offspring each individual has is determined by a random variable with a specific distribution.
- **Extinction:** The process may eventually result in the extinction of the population (i.e., no individuals are left).



Mathematical Model of a Branching Process

Consider a **Galton-Watson branching process**:

- Let Z_n be the population size at generation n .
- The population starts with Z_0 individuals.
- Each individual produces offspring independently according to some probability distribution p_k , where p_k is the probability that an individual produces k offspring.

The size of the population in the next generation is given by:

$$Z_n = \sum_{i=1}^{Z_{n-1}} X_i$$

where X_i is the number of offspring produced by the i -th individual, and X_i are i.i.d. random variables.



Mean of Branching Process

Consider a Galton-Watson process where Z_n represents the population size at generation n .

The population at generation n is the sum of offspring from generation $n - 1$:

$$Z_n = \sum_{i=1}^{Z_{n-1}} X_i$$

where X_i is the number of offspring of the i -th individual, and X_i are i.i.d. random variables with mean $\mathbb{E}[X_i] = \mu$.

Using the linearity of expectation:

$$\mathbb{E}[Z_n] = \mathbb{E}\left[\sum_{i=1}^{Z_{n-1}} X_i\right] = \mathbb{E}[Z_{n-1}] \cdot \mathbb{E}[X_1] = \mu \cdot \mathbb{E}[Z_{n-1}]$$

Thus, iterating this recurrence:

$$\mathbb{E}[Z_n] = \mu^n \cdot Z_0$$

where Z_0 is the initial population size.



Variance of Branching Process

To calculate the variance $\text{Var}(Z_n)$, we apply the law of total variance:

$$\text{Var}(Z_n) = \mathbb{E}[\text{Var}(Z_n \mid Z_{n-1})] + \text{Var}(\mathbb{E}[Z_n \mid Z_{n-1}])$$

First, the conditional variance $\text{Var}(Z_n \mid Z_{n-1})$ is:

$$\text{Var}(Z_n \mid Z_{n-1}) = Z_{n-1} \cdot \text{Var}(X_1)$$

where $\text{Var}(X_1) = \sigma^2$, the variance in the number of offspring per individual.

The conditional expectation $\mathbb{E}[Z_n \mid Z_{n-1}]$ is:

$$\mathbb{E}[Z_n \mid Z_{n-1}] = Z_{n-1} \cdot \mu$$

Therefore, the total variance is:

$$\text{Var}(Z_n) = \sigma^2 \cdot \mathbb{E}[Z_{n-1}] + \mu^2 \cdot \text{Var}(Z_{n-1})$$

Iterating this recurrence:

$$\text{Var}(Z_n) = Z_0 \cdot \sigma^2 \cdot \frac{\mu^n - 1}{\mu - 1}$$



Types of Branching Processes

- **Supercritical Branching Process** ($\mu > 1$): The expected number of offspring per individual is greater than 1, and the population has a chance of growing indefinitely.
- **Critical Branching Process** ($\mu = 1$): The expected number of offspring per individual is equal to 1, and the population is expected to stay roughly constant.
- **Subcritical Branching Process** ($\mu < 1$): The expected number of offspring per individual is less than 1, and the population is expected to die out.



Applications of Branching Processes

- **Epidemiology:** Used to model the spread of infectious diseases in populations.
- **Population Biology:** Modeling the growth or extinction of species over time.
- **Physics:** Describing reactions and particle interactions in nuclear and particle physics.
- **Computer Science:** Studying recursive algorithms and data structures like search trees.

