


# An Introduction to PAC-Bayesian Analysis

Benjamin Guedj   John Shawe-Taylor

Supervised Learning  
December 9–13, 2019


Learning is to be able to generalise

Learning is to be able to generalise



[Figure from Wikipedia]


# Learning is to be able to generalise



From **examples**, what can a system  
**learn** about the **underlying**  
**phenomenon**?

[Figure from Wikipedia]

# Learning is to be able to generalise




From **examples**, what can a system **learn** about the **underlying phenomenon**?

Memorising the already seen data is usually bad → **overfitting**

[Figure from Wikipedia]

# Learning is to be able to generalise



From **examples**, what can a system **learn** about the **underlying phenomenon**?

Memorising the already seen data is usually bad → **overfitting**

**Generalisation** is the ability to 'perform' well on **unseen data**.

[Figure from Wikipedia]

Statistical Learning Theory is about high confidence

# Statistical Learning Theory is about high confidence

For a fixed algorithm, function class and sample size, generating random samples  $\longrightarrow$  distribution of test errors



# Statistical Learning Theory is about high confidence

For a fixed algorithm, function class and sample size, generating random samples  $\longrightarrow$  distribution of test errors

- Focusing on the mean of the error distribution?
  - ▷ can be misleading: learner only has **one** sample

# Statistical Learning Theory is about high confidence

For a fixed algorithm, function class and sample size, generating random samples  $\longrightarrow$  distribution of test errors

- Focusing on the mean of the error distribution?
  - ▷ can be misleading: learner only has **one** sample
- **Statistical Learning Theory**: **tail of the distribution**
  - ▷ finding bounds which hold with high probability over random samples of size  $m$

# Statistical Learning Theory is about high confidence

For a fixed algorithm, function class and sample size, generating random samples  $\longrightarrow$  distribution of test errors

- Focusing on the mean of the error distribution?
  - ▷ can be misleading: learner only has **one** sample
- **Statistical Learning Theory**: **tail of the distribution**
  - ▷ finding bounds which hold with high probability over random samples of size  $m$
- Compare to a statistical test – at **99%** confidence level
  - ▷ chances of the conclusion not being true are less than **1%**

# Statistical Learning Theory is about high confidence

For a fixed algorithm, function class and sample size, generating random samples  $\longrightarrow$  distribution of test errors


- Focusing on the mean of the error distribution?
  - ▷ can be misleading: learner only has **one** sample
- **Statistical Learning Theory**: **tail of the distribution**
  - ▷ finding bounds which hold with high probability over random samples of size  $m$
- Compare to a statistical test – at **99%** confidence level
  - ▷ chances of the conclusion not being true are less than **1%**
- **PAC**: probably approximately correct [59]
  - Use a ‘confidence parameter’  $\delta$ :  $\mathbb{P}^m[\text{large error}] \leq \delta$   
 $\delta$  is the probability of being misled by the training set

# Statistical Learning Theory is about high confidence

For a fixed algorithm, function class and sample size, generating random samples  $\longrightarrow$  distribution of test errors

- Focusing on the mean of the error distribution?
  - ▷ can be misleading: learner only has **one** sample
- **Statistical Learning Theory**: **tail of the distribution**
  - ▷ finding bounds which hold with high probability over random samples of size  $m$
- Compare to a statistical test – at **99%** confidence level
  - ▷ chances of the conclusion not being true are less than **1%**
- **PAC**: probably approximately correct [59]
  - Use a ‘confidence parameter’  $\delta$ :  $\mathbb{P}^m[\text{large error}] \leq \delta$   
 $\delta$  is the probability of being misled by the training set
- Hence **high confidence**:  $\mathbb{P}^m[\text{approximately correct}] \geq 1 - \delta$

# Error distribution picture



# Mathematical formalization

# Mathematical formalization

Learning algorithm  $A : \mathcal{Z}^m \rightarrow \mathcal{H}$

- $\mathcal{Z} = \mathcal{X} \times \mathcal{Y}$   
 $\mathcal{X}$  = set of inputs  
 $\mathcal{Y}$  = set of outputs (e.g. labels)
- $\mathcal{H}$  = hypothesis class  
= set of **predictors**  
(e.g. classifiers)



# Mathematical formalization

Learning algorithm  $A : \mathcal{Z}^m \rightarrow \mathcal{H}$

- $\mathcal{Z} = \mathcal{X} \times \mathcal{Y}$   
 $\mathcal{X}$  = set of inputs  
 $\mathcal{Y}$  = set of outputs (e.g. labels)
- $\mathcal{H}$  = hypothesis class  
= set of **predictors**  
(e.g. classifiers)

**Training set** (aka **sample**):  $S_m = ((X_1, Y_1), \dots, (X_m, Y_m))$   
a finite sequence of **input-output examples**.

# Mathematical formalization

**Learning algorithm**  $A : \mathcal{Z}^m \rightarrow \mathcal{H}$

- $\mathcal{Z} = \mathcal{X} \times \mathcal{Y}$   
 $\mathcal{X}$  = set of inputs  
 $\mathcal{Y}$  = set of outputs (e.g. labels)
- $\mathcal{H}$  = hypothesis class  
= set of **predictors**  
(e.g. classifiers)

**Training set** (aka **sample**):  $S_m = ((X_1, Y_1), \dots, (X_m, Y_m))$   
a finite sequence of **input-output examples**.

**Classical assumptions:**

- A **data-generating distribution**  $\mathbb{P}$  over  $\mathcal{Z}$ .
- Learner doesn't know  $\mathbb{P}$ , only sees the training set.
- The training set **examples are *i.i.d.*** from  $\mathbb{P}$ :  $S_m \sim \mathbb{P}^m$

# Mathematical formalization

**Learning algorithm**  $A : \mathcal{Z}^m \rightarrow \mathcal{H}$

- $\mathcal{Z} = \mathcal{X} \times \mathcal{Y}$   
 $\mathcal{X}$  = set of inputs  
 $\mathcal{Y}$  = set of outputs (e.g. labels)
- $\mathcal{H}$  = hypothesis class  
= set of **predictors**  
(e.g. classifiers)

**Training set** (aka **sample**):  $S_m = ((X_1, Y_1), \dots, (X_m, Y_m))$   
a finite sequence of **input-output examples**.

**Classical assumptions:**

- A **data-generating distribution**  $\mathbb{P}$  over  $\mathcal{Z}$ .
  - Learner doesn't know  $\mathbb{P}$ , only sees the training set.
  - The training set **examples are *i.i.d.*** from  $\mathbb{P}$ :  $S_m \sim \mathbb{P}^m$
- ▷ these can be relaxed (mostly beyond the scope of this tutorial)

What to achieve from the sample?

# What to achieve from the sample?

Use the available sample to:

- 1 learn a predictor
- 2 certify the predictor's performance

# What to achieve from the sample?

Use the available sample to:

- 1 learn a predictor
- 2 certify the predictor's performance

## Learning a predictor:

- algorithm driven by some learning principle
- informed by prior knowledge resulting in inductive bias

# What to achieve from the sample?

Use the available sample to:

- 1 learn a predictor
- 2 certify the predictor's performance

## Learning a predictor:

- algorithm driven by some learning principle
- informed by prior knowledge resulting in inductive bias

## Certifying performance:

- what happens beyond the training set
- generalization bounds

# What to achieve from the sample?

Use the available sample to:

- 1 learn a predictor
- 2 certify the predictor's performance

## Learning a predictor:

- algorithm driven by some learning principle
- informed by prior knowledge resulting in inductive bias

## Certifying performance:

- what happens beyond the training set
- generalization bounds

Actually these two goals interact with each other!



## Risk (aka error) measures

# Risk (aka error) measures

A **loss function**  $\ell(h(X), Y)$  is used to measure the discrepancy between a predicted output  $h(X)$  and the true output  $Y$ .

# Risk (aka error) measures

A **loss function**  $\ell(h(X), Y)$  is used to measure the discrepancy between a predicted output  $h(X)$  and the true output  $Y$ .

**Empirical risk:**  $R_{\text{in}}(h) = \frac{1}{m} \sum_{i=1}^m \ell(h(X_i), Y_i)$   
(in-sample)

# Risk (aka error) measures

A **loss function**  $\ell(h(X), Y)$  is used to measure the discrepancy between a predicted output  $h(X)$  and the true output  $Y$ .

**Empirical risk:**  $R_{\text{in}}(h) = \frac{1}{m} \sum_{i=1}^m \ell(h(X_i), Y_i)$   
(in-sample)

**Theoretical risk:**  $R_{\text{out}}(h) = \mathbb{E}[\ell(h(X), Y)]$   
(out-of-sample)

# Risk (aka error) measures

A **loss function**  $\ell(h(X), Y)$  is used to measure the discrepancy between a predicted output  $h(X)$  and the true output  $Y$ .

**Empirical risk:**  $R_{\text{in}}(h) = \frac{1}{m} \sum_{i=1}^m \ell(h(X_i), Y_i)$   
(in-sample)

**Theoretical risk:**  $R_{\text{out}}(h) = \mathbb{E}[\ell(h(X), Y)]$   
(out-of-sample)

## Examples:

- $\ell(h(X), Y) = \mathbf{1}[h(X) \neq Y]$  : **0-1 loss** (classification)
- $\ell(h(X), Y) = (Y - h(X))^2$  : **square loss** (regression)
- $\ell(h(X), Y) = (1 - Yh(X))_+$  : **hinge loss**
- $\ell(h(X), Y) = -\log(h(X))$  : **log loss** (density estimation) TODO

# Generalization

If predictor  $h$  does well on the in-sample  $(X, Y)$  pairs...

...will it still do well on out-of-sample pairs?

# Generalization

If predictor  $h$  does well on the in-sample  $(X, Y)$  pairs...

...will it still do well on out-of-sample pairs?

Generalization gap:  $\Delta(h) = R_{\text{out}}(h) - R_{\text{in}}(h)$

# Generalization

If predictor  $h$  does well on the in-sample  $(X, Y)$  pairs...

...will it still do well on out-of-sample pairs?

Generalization gap:  $\Delta(h) = R_{\text{out}}(h) - R_{\text{in}}(h)$

Upper bounds: w.h.p.  $\Delta(h) \leq \epsilon(m, \delta)$



# Generalization

If predictor  $h$  does well on the in-sample  $(X, Y)$  pairs...

...will it still do well on out-of-sample pairs?

**Generalization gap:**  $\Delta(h) = R_{\text{out}}(h) - R_{\text{in}}(h)$

**Upper bounds:** w.h.p.  $\Delta(h) \leq \epsilon(m, \delta)$

$$\blacktriangleright R_{\text{out}}(h) \leq R_{\text{in}}(h) + \epsilon(m, \delta)$$

# Generalization

If predictor  $h$  does well on the in-sample  $(X, Y)$  pairs...

...will it still do well on out-of-sample pairs?

**Generalization gap:**  $\Delta(h) = R_{\text{out}}(h) - R_{\text{in}}(h)$

**Upper bounds:** w.h.p.  $\Delta(h) \leq \epsilon(m, \delta)$

$$\blacktriangleright R_{\text{out}}(h) \leq R_{\text{in}}(h) + \epsilon(m, \delta)$$

**Lower bounds:** w.h.p.  $\Delta(h) \geq \tilde{\epsilon}(m, \delta)$

# Generalization

If predictor  $h$  does well on the in-sample  $(X, Y)$  pairs...

...will it still do well on out-of-sample pairs?

**Generalization gap:**  $\Delta(h) = R_{\text{out}}(h) - R_{\text{in}}(h)$

**Upper bounds:** w.h.p.  $\Delta(h) \leq \epsilon(m, \delta)$

$$\blacktriangleright R_{\text{out}}(h) \leq R_{\text{in}}(h) + \epsilon(m, \delta)$$

**Lower bounds:** w.h.p.  $\Delta(h) \geq \tilde{\epsilon}(m, \delta)$

**Flavours:**

- |                     |                          |
|---------------------|--------------------------|
| ■ distribution-free | ■ distribution-dependent |
| ■ algorithm-free    | ■ algorithm-dependent    |

# Why you should care about generalisation bounds

# Why you should care about generalisation bounds

Generalisation bounds are a safety check: give a theoretical guarantee on the performance of a learning algorithm on any unseen data.

# Why you should care about generalisation bounds

**Generalisation bounds** are a **safety check**: give a **theoretical guarantee** on the **performance** of a learning algorithm on **any unseen data**.

$$R_{\text{out}}(h) \leq R_{\text{in}}(h) + \epsilon(m, \delta)$$

Generalisation bounds:

# Why you should care about generalisation bounds

**Generalisation bounds** are a **safety check**: give a **theoretical guarantee** on the **performance** of a learning algorithm on **any unseen data**.

$$R_{\text{out}}(h) \leq R_{\text{in}}(h) + \epsilon(m, \delta)$$

Generalisation bounds:

- may be computed with the **training sample only**, do not depend on any test sample

# Why you should care about generalisation bounds

**Generalisation bounds** are a **safety check**: give a **theoretical guarantee** on the **performance** of a learning algorithm on **any unseen data**.

$$R_{\text{out}}(h) \leq R_{\text{in}}(h) + \epsilon(m, \delta)$$

Generalisation bounds:

- may be computed with the **training sample only**, do not depend on any test sample
- provide a **computable** control on the error on **any unseen data** with prespecified confidence



# Why you should care about generalisation bounds

**Generalisation bounds** are a **safety check**: give a **theoretical guarantee** on the **performance** of a learning algorithm on **any unseen data**.

$$R_{\text{out}}(h) \leq R_{\text{in}}(h) + \epsilon(m, \delta)$$

Generalisation bounds:

- may be computed with the **training sample only**, do not depend on any test sample
- provide a **computable** control on the error on **any unseen data** with prespecified confidence
- explain **why** specific learning algorithms **actually work**

# Why you should care about generalisation bounds

**Generalisation bounds** are a **safety check**: give a **theoretical guarantee** on the **performance** of a learning algorithm on **any unseen data**.

$$R_{\text{out}}(h) \leq R_{\text{in}}(h) + \epsilon(m, \delta)$$

Generalisation bounds:

- may be computed with the **training sample only**, do not depend on any test sample
- provide a **computable** control on the error on **any unseen data** with prespecified confidence
- explain **why** specific learning algorithms **actually work**
- and even lead to **designing new algorithm** which scale to more complex settings

# Before PAC-Bayes

# Before PAC-Bayes

- Single hypothesis  $h$  (building block):

with probability  $\geq 1 - \delta$ ,  $R_{\text{out}}(h) \leq R_{\text{in}}(h) + \sqrt{\frac{1}{2m} \log\left(\frac{1}{\delta}\right)}.$

# Before PAC-Bayes

- Single hypothesis  $h$  (building block):

with probability  $\geq 1 - \delta$ ,  $R_{\text{out}}(h) \leq R_{\text{in}}(h) + \sqrt{\frac{1}{2m} \log\left(\frac{1}{\delta}\right)}.$

- Finite function class  $\mathcal{H}$  (worst-case approach):

w.p.  $\geq 1 - \delta$ ,  $\forall h \in \mathcal{H}, R_{\text{out}}(h) \leq R_{\text{in}}(h) + \sqrt{\frac{1}{2m} \log\left(\frac{|\mathcal{H}|}{\delta}\right)}$

# Before PAC-Bayes

- Single hypothesis  $h$  (building block):

with probability  $\geq 1 - \delta$ ,  $R_{\text{out}}(h) \leq R_{\text{in}}(h) + \sqrt{\frac{1}{2m} \log\left(\frac{1}{\delta}\right)}.$

- Finite function class  $\mathcal{H}$  (worst-case approach):

w.p.  $\geq 1 - \delta$ ,  $\forall h \in \mathcal{H}, R_{\text{out}}(h) \leq R_{\text{in}}(h) + \sqrt{\frac{1}{2m} \log\left(\frac{|\mathcal{H}|}{\delta}\right)}$

- Structural risk minimisation: data-dependent hypotheses  $h_i$  associated with prior weight  $p_i$

w.p.  $\geq 1 - \delta$ ,  $\forall h_i \in \mathcal{H}, R_{\text{out}}(h_i) \leq R_{\text{in}}(h_i) + \sqrt{\frac{1}{2m} \log\left(\frac{1}{p_i \delta}\right)}$

# Before PAC-Bayes

- Single hypothesis  $h$  (building block):

with probability  $\geq 1 - \delta$ ,  $R_{\text{out}}(h) \leq R_{\text{in}}(h) + \sqrt{\frac{1}{2m} \log\left(\frac{1}{\delta}\right)}.$

- Finite function class  $\mathcal{H}$  (worst-case approach):

w.p.  $\geq 1 - \delta$ ,  $\forall h \in \mathcal{H}, R_{\text{out}}(h) \leq R_{\text{in}}(h) + \sqrt{\frac{1}{2m} \log\left(\frac{|\mathcal{H}|}{\delta}\right)}$

- Structural risk minimisation: data-dependent hypotheses  $h_i$  associated with prior weight  $p_i$

w.p.  $\geq 1 - \delta$ ,  $\forall h_i \in \mathcal{H}, R_{\text{out}}(h_i) \leq R_{\text{in}}(h_i) + \sqrt{\frac{1}{2m} \log\left(\frac{1}{p_i \delta}\right)}$

- Uncountably infinite function class: VC dimension, Rademacher complexity...

# Before PAC-Bayes

- Single hypothesis  $h$  (building block):

with probability  $\geq 1 - \delta$ ,  $R_{\text{out}}(h) \leq R_{\text{in}}(h) + \sqrt{\frac{1}{2m} \log\left(\frac{1}{\delta}\right)}.$

- Finite function class  $\mathcal{H}$  (worst-case approach):

w.p.  $\geq 1 - \delta$ ,  $\forall h \in \mathcal{H}$ ,  $R_{\text{out}}(h) \leq R_{\text{in}}(h) + \sqrt{\frac{1}{2m} \log\left(\frac{|\mathcal{H}|}{\delta}\right)}$

- Structural risk minimisation: data-dependent hypotheses  $h_i$  associated with prior weight  $p_i$

w.p.  $\geq 1 - \delta$ ,  $\forall h_i \in \mathcal{H}$ ,  $R_{\text{out}}(h_i) \leq R_{\text{in}}(h_i) + \sqrt{\frac{1}{2m} \log\left(\frac{1}{p_i \delta}\right)}$

- Uncountably infinite function class: VC dimension, Rademacher complexity...

These approaches are suited to analyse the performance of individual functions, and take some account of correlations.



# Before PAC-Bayes

- Single hypothesis  $h$  (building block):

with probability  $\geq 1 - \delta$ ,  $R_{\text{out}}(h) \leq R_{\text{in}}(h) + \sqrt{\frac{1}{2m} \log\left(\frac{1}{\delta}\right)}.$

- Finite function class  $\mathcal{H}$  (worst-case approach):

w.p.  $\geq 1 - \delta$ ,  $\forall h \in \mathcal{H}$ ,  $R_{\text{out}}(h) \leq R_{\text{in}}(h) + \sqrt{\frac{1}{2m} \log\left(\frac{|\mathcal{H}|}{\delta}\right)}$

- Structural risk minimisation: data-dependent hypotheses  $h_i$  associated with prior weight  $p_i$

w.p.  $\geq 1 - \delta$ ,  $\forall h_i \in \mathcal{H}$ ,  $R_{\text{out}}(h_i) \leq R_{\text{in}}(h_i) + \sqrt{\frac{1}{2m} \log\left(\frac{1}{p_i \delta}\right)}$

- Uncountably infinite function class: VC dimension, Rademacher complexity...

These approaches are suited to analyse the performance of individual functions, and take some account of correlations.

→ Extension: PAC-Bayes allows to consider *distributions* over hypotheses.

# The PAC-Bayes framework

# The PAC-Bayes framework

- Before data, fix a distribution  $P \in M_1(\mathcal{H}) \triangleright$  ‘prior’

# The PAC-Bayes framework

- Before data, fix a distribution  $P \in M_1(\mathcal{H}) \triangleright$  ‘prior’
- Based on data, learn a distribution  $Q \in M_1(\mathcal{H}) \triangleright$  ‘posterior’

# The PAC-Bayes framework

- Before data, fix a distribution  $P \in M_1(\mathcal{H}) \triangleright$  ‘prior’
- Based on data, learn a distribution  $Q \in M_1(\mathcal{H}) \triangleright$  ‘posterior’
- Predictions:
  - draw  $h \sim Q$  and predict with the chosen  $h$ .
  - each prediction with a fresh random draw.

# The PAC-Bayes framework

- Before data, fix a distribution  $P \in M_1(\mathcal{H}) \triangleright$  ‘prior’
- Based on data, learn a distribution  $Q \in M_1(\mathcal{H}) \triangleright$  ‘posterior’
- Predictions:
  - draw  $h \sim Q$  and predict with the chosen  $h$ .
  - each prediction with a fresh random draw.


The risk measures  $R_{\text{in}}(h)$  and  $R_{\text{out}}(h)$  are extended by averaging:

$$R_{\text{in}}(Q) \equiv \int_{\mathcal{H}} R_{\text{in}}(h) dQ(h) \quad R_{\text{out}}(Q) \equiv \int_{\mathcal{H}} R_{\text{out}}(h) dQ(h)$$

$\text{KL}(Q||P) = \mathbf{E}_{h \sim Q} \ln \frac{Q(h)}{P(h)}$  is the Kullback-Leibler divergence.

# PAC-Bayes aka Generalised Bayes

# PAC-Bayes aka Generalised Bayes



"Prior": exploration mechanism of  $\mathcal{H}$

"Posterior" is the twisted prior after confronting with data



# PAC-Bayes bounds vs. Bayesian learning

# PAC-Bayes bounds vs. Bayesian learning

- Prior

# PAC-Bayes bounds vs. Bayesian learning

## ■ Prior

- **PAC-Bayes**: bounds hold for any distribution
- **Bayes**: prior choice impacts inference

# PAC-Bayes bounds vs. Bayesian learning

## ■ Prior

- **PAC-Bayes**: bounds hold for any distribution
- **Bayes**: prior choice impacts inference

## ■ Posterior

# PAC-Bayes bounds vs. Bayesian learning

## ■ Prior

- **PAC-Bayes**: bounds hold for any distribution
- **Bayes**: prior choice impacts inference

## ■ Posterior

- **PAC-Bayes**: bounds hold for any distribution
- **Bayes**: posterior uniquely defined by prior and statistical model

# PAC-Bayes bounds vs. Bayesian learning

## ■ Prior

- **PAC-Bayes**: bounds hold for any distribution
- **Bayes**: prior choice impacts inference

## ■ Posterior

- **PAC-Bayes**: bounds hold for any distribution
- **Bayes**: posterior uniquely defined by prior and statistical model

## ■ Data distribution

# PAC-Bayes bounds vs. Bayesian learning

## ■ Prior

- **PAC-Bayes**: bounds hold for any distribution
- **Bayes**: prior choice impacts inference

## ■ Posterior

- **PAC-Bayes**: bounds hold for any distribution
- **Bayes**: posterior uniquely defined by prior and statistical model

## ■ Data distribution

- **PAC-Bayes**: bounds hold for any distribution
- **Bayes**: randomness lies in the noise model generating the output

# A General PAC-Bayesian Theorem

$\Delta$ -function: “distance” between  $R_{\text{in}}(Q)$  and  $R_{\text{out}}(Q)$

Convex function  $\Delta : [0, 1] \times [0, 1] \rightarrow \mathbb{R}$ .

## General theorem

(Bégin et al. [7, 8], Germain [21])

*For any distribution  $D$  on  $\mathcal{X} \times \mathcal{Y}$ , for any set  $\mathcal{H}$  of voters, for any distribution  $P$  on  $\mathcal{H}$ , for any  $\delta \in (0, 1]$ , and for any  $\Delta$ -function, we have, with probability at least  $1 - \delta$  over the choice of  $S \sim D^m$ ,*

$$\forall Q \text{ on } \mathcal{H} : \quad \Delta\left(R_{\text{in}}(Q), R_{\text{out}}(Q)\right) \leq \frac{1}{m} \left[ \text{KL}(Q \| P) + \ln \frac{\mathcal{I}_{\Delta}(m)}{\delta} \right],$$



# A General PAC-Bayesian Theorem

$\Delta$ -function: “distance” between  $R_{\text{in}}(Q)$  and  $R_{\text{out}}(Q)$

Convex function  $\Delta : [0, 1] \times [0, 1] \rightarrow \mathbb{R}$ .

## General theorem

(Bégin et al. [7, 8], Germain [21])

*For any distribution  $D$  on  $\mathcal{X} \times \mathcal{Y}$ , for any set  $\mathcal{H}$  of voters, for any distribution  $P$  on  $\mathcal{H}$ , for any  $\delta \in (0, 1]$ , and for any  $\Delta$ -function, we have, with probability at least  $1 - \delta$  over the choice of  $S \sim D^m$ ,*

$$\forall Q \text{ on } \mathcal{H} : \quad \Delta\left(R_{\text{in}}(Q), R_{\text{out}}(Q)\right) \leq \frac{1}{m} \left[ \text{KL}(Q \| P) + \ln \frac{\mathcal{I}_{\Delta}(m)}{\delta} \right],$$

where

$$\mathcal{I}_{\Delta}(m) = \sup_{r \in [0, 1]} \left[ \sum_{k=0}^m \underbrace{\binom{m}{k} r^k (1-r)^{m-k}}_{\text{Bin}(k; m, r)} e^{m \Delta(\frac{k}{m}, r)} \right].$$

# Proof of the general theorem

## General theorem

$$\Pr_{S \sim D^m} \left( \forall Q \text{ on } \mathcal{H} : \Delta \left( R_{\text{in}}(Q), R_{\text{out}}(Q) \right) \leq \frac{1}{m} \left[ \text{KL}(Q \| P) + \ln \frac{\mathcal{I}_{\Delta}(m)}{\delta} \right] \right) \geq 1 - \delta.$$

## Proof ideas.

### Change of Measure Inequality

For any  $P$  and  $Q$  on  $\mathcal{H}$ , and for any measurable function  $\phi : \mathcal{H} \rightarrow \mathbb{R}$ , we have

$$\begin{aligned} -\ln \left( \mathbf{E}_{h \sim P} e^{\phi(h)} \right) &= -\ln \mathbf{E}_{h \sim Q} \left( \frac{P(h)}{Q(h)} e^{\phi(h)} \right) \\ &\leq \mathbf{E}_{h \sim Q} \ln \left( \frac{Q(h)}{P(h)} \right) - \mathbf{E}_{h \sim Q} \phi(h) \\ &= \text{KL}(Q \| P) - \mathbf{E}_{h \sim Q} \phi(h). \end{aligned}$$

# Proof of the general theorem

## General theorem

$$\Pr_{S \sim D^m} \left( \forall Q \text{ on } \mathcal{H} : \Delta \left( R_{\text{in}}(Q), R_{\text{out}}(Q) \right) \leq \frac{1}{m} \left[ \text{KL}(Q \| P) + \ln \frac{\mathcal{I}_{\Delta}(m)}{\delta} \right] \right) \geq 1 - \delta.$$

## Proof ideas.

### Change of Measure Inequality

For any  $P$  and  $Q$  on  $\mathcal{H}$ , and for any measurable function  $\phi : \mathcal{H} \rightarrow \mathbb{R}$ , we have

$$\begin{aligned} -\ln \left( \mathbf{E}_{h \sim P} e^{\phi(h)} \right) &= -\ln \mathbf{E}_{h \sim Q} \left( \frac{P(h)}{Q(h)} e^{\phi(h)} \right) \\ &\leq \mathbf{E}_{h \sim Q} \ln \left( \frac{Q(h)}{P(h)} \right) - \mathbf{E}_{h \sim Q} \phi(h) \\ &= \text{KL}(Q \| P) - \mathbf{E}_{h \sim Q} \phi(h). \end{aligned}$$

### Markov's inequality

for a random variable  $X$  satisfying  $X \geq 0$

$$\Pr(X \geq a) \leq \frac{\mathbf{E}X}{a} \iff \Pr(X \leq \frac{\mathbf{E}X}{\delta}) \geq 1 - \delta.$$

# Proof of the general theorem

## Probability of observing $k$ misclassifications among $m$ examples

Given a voter  $h$ , consider a **binomial variable** of  $m$  trials with **success**  $R_{\text{out}}(h)$ :

$$\Pr_{S \sim D^m} \left( R_{\text{in}}(h) = \frac{k}{m} \right) = \binom{m}{k} \left( R_{\text{out}}(h) \right)^k \left( 1 - R_{\text{out}}(h) \right)^{m-k} = \text{Bin} \left( k; m, R_{\text{out}}(h) \right)$$

$$\Pr_{S \sim D^m} \left( \forall Q \text{ on } \mathcal{H} : \Delta \left( R_{\text{in}}(Q), R_{\text{out}}(Q) \right) \leq \frac{1}{m} \left[ \text{KL}(Q \| P) + \ln \frac{\mathcal{I}_{\Delta}(m)}{\delta} \right] \right) \geq 1 - \delta.$$

**Proof.**

$$m \cdot \Delta \left( \mathbf{E}_{h \sim Q} R_{\text{in}}(h), \mathbf{E}_{h \sim Q} R_{\text{out}}(h) \right)$$

$$\Pr_{S \sim D^m} \left( \forall Q \text{ on } \mathcal{H} : \Delta \left( R_{\text{in}}(Q), R_{\text{out}}(Q) \right) \leq \frac{1}{m} \left[ \text{KL}(Q \| P) + \ln \frac{J_{\Delta}(m)}{\delta} \right] \right) \geq 1 - \delta.$$

**Proof.**

$$m \cdot \Delta \left( \mathbf{E}_{h \sim Q} R_{\text{in}}(h), \mathbf{E}_{h \sim Q} R_{\text{out}}(h) \right)$$

Jensen's Inequality

$$\leq \mathbf{E}_{h \sim Q} m \cdot \Delta \left( R_{\text{in}}(h), R_{\text{out}}(h) \right)$$

$$\Pr_{S \sim D^m} \left( \forall Q \text{ on } \mathcal{H} : \Delta \left( R_{\text{in}}(Q), R_{\text{out}}(Q) \right) \leq \frac{1}{m} \left[ \text{KL}(Q \| P) + \ln \frac{J_{\Delta}(m)}{\delta} \right] \right) \geq 1 - \delta.$$

**Proof.**

$$m \cdot \Delta \left( \mathbf{E}_{h \sim Q} R_{\text{in}}(h), \mathbf{E}_{h \sim Q} R_{\text{out}}(h) \right)$$

Jensen's Inequality

$$\leq \mathbf{E}_{h \sim Q} m \cdot \Delta \left( R_{\text{in}}(h), R_{\text{out}}(h) \right)$$

Change of measure

$$\leq \text{KL}(Q \| P) + \ln \mathbf{E}_{h \sim P} e^{m \Delta \left( R_{\text{in}}(h), R_{\text{out}}(h) \right)}$$

$$\Pr_{S \sim D^m} \left( \forall Q \text{ on } \mathcal{H} : \Delta(R_{\text{in}}(Q), R_{\text{out}}(Q)) \leq \frac{1}{m} \left[ \text{KL}(Q \| P) + \ln \frac{J_{\Delta}(m)}{\delta} \right] \right) \geq 1 - \delta.$$

**Proof.**

$$m \cdot \Delta \left( \mathbf{E}_{h \sim Q} R_{\text{in}}(h), \mathbf{E}_{h \sim Q} R_{\text{out}}(h) \right)$$

Jensen's Inequality

$$\leq \mathbf{E}_{h \sim Q} m \cdot \Delta(R_{\text{in}}(h), R_{\text{out}}(h))$$

Change of measure

$$\leq \text{KL}(Q \| P) + \ln \mathbf{E}_{h \sim P} e^{m \Delta(R_{\text{in}}(h), R_{\text{out}}(h))}$$

Markov's Inequality

$$\leq_{1-\delta} \text{KL}(Q \| P) + \ln \frac{1}{\delta} \mathbf{E}_{S' \sim D^m} \mathbf{E}_{h \sim P} e^{m \cdot \Delta(R_{\text{in}}(h), R_{\text{out}}(h))}$$



$$\Pr_{S \sim D^m} \left( \forall Q \text{ on } \mathcal{H} : \Delta \left( R_{\text{in}}(Q), R_{\text{out}}(Q) \right) \leq \frac{1}{m} \left[ \text{KL}(Q \| P) + \ln \frac{J_{\Delta}(m)}{\delta} \right] \right) \geq 1 - \delta.$$

**Proof.**

$$m \cdot \Delta \left( \mathbf{E}_{h \sim Q} R_{\text{in}}(h), \mathbf{E}_{h \sim Q} R_{\text{out}}(h) \right)$$

Jensen's Inequality

$$\leq \mathbf{E}_{h \sim Q} m \cdot \Delta \left( R_{\text{in}}(h), R_{\text{out}}(h) \right)$$

Change of measure

$$\leq \text{KL}(Q \| P) + \ln \mathbf{E}_{h \sim P} e^{m \Delta \left( R_{\text{in}}(h), R_{\text{out}}(h) \right)}$$

Markov's Inequality

$$\leq_{1-\delta} \text{KL}(Q \| P) + \ln \frac{1}{\delta} \mathbf{E}_{S' \sim D^m} \mathbf{E}_{h \sim P} e^{m \cdot \Delta \left( R_{\text{in}}(h), R_{\text{out}}(h) \right)}$$

Expectation swap

$$= \text{KL}(Q \| P) + \ln \frac{1}{\delta} \mathbf{E}_{h \sim P} \mathbf{E}_{S' \sim D^m} e^{m \cdot \Delta \left( R_{\text{in}}(h), R_{\text{out}}(h) \right)}$$

$$\Pr_{S \sim D^m} \left( \forall Q \text{ on } \mathcal{H} : \Delta(R_{\text{in}}(Q), R_{\text{out}}(Q)) \leq \frac{1}{m} \left[ \text{KL}(Q \| P) + \ln \frac{\mathcal{I}_\Delta(m)}{\delta} \right] \right) \geq 1 - \delta.$$

**Proof.**

$$m \cdot \Delta \left( \mathbf{E}_{h \sim Q} R_{\text{in}}(h), \mathbf{E}_{h \sim Q} R_{\text{out}}(h) \right)$$

Jensen's Inequality

$$\leq \mathbf{E}_{h \sim Q} m \cdot \Delta(R_{\text{in}}(h), R_{\text{out}}(h))$$

Change of measure

$$\leq \text{KL}(Q \| P) + \ln \mathbf{E}_{h \sim P} e^{m \Delta(R_{\text{in}}(h), R_{\text{out}}(h))}$$

Markov's Inequality

$$\leq_{1-\delta} \text{KL}(Q \| P) + \ln \frac{1}{\delta} \mathbf{E}_{S' \sim D^m} \mathbf{E}_{h \sim P} e^{m \cdot \Delta(R_{\text{in}}(h), R_{\text{out}}(h))}$$

Expectation swap

$$= \text{KL}(Q \| P) + \ln \frac{1}{\delta} \mathbf{E}_{h \sim P} \mathbf{E}_{S' \sim D^m} e^{m \cdot \Delta(R_{\text{in}}(h), R_{\text{out}}(h))}$$

Binomial law

$$= \text{KL}(Q \| P) + \ln \frac{1}{\delta} \mathbf{E}_{h \sim P} \sum_{k=0}^m \text{Bin}(k; m, R_{\text{out}}(h)) e^{m \cdot \Delta(\frac{k}{m}, R_{\text{out}}(h))}$$

$$\Pr_{S \sim D^m} \left( \forall Q \text{ on } \mathcal{H} : \Delta \left( R_{\text{in}}(Q), R_{\text{out}}(Q) \right) \leq \frac{1}{m} \left[ \text{KL}(Q \| P) + \ln \frac{\mathcal{I}_{\Delta}(m)}{\delta} \right] \right) \geq 1 - \delta.$$

**Proof.**

$$m \cdot \Delta \left( \mathbf{E}_{h \sim Q} R_{\text{in}}(h), \mathbf{E}_{h \sim Q} R_{\text{out}}(h) \right)$$

Jensen's Inequality

$$\leq \mathbf{E}_{h \sim Q} m \cdot \Delta \left( R_{\text{in}}(h), R_{\text{out}}(h) \right)$$

Change of measure

$$\leq \text{KL}(Q \| P) + \ln \mathbf{E}_{h \sim P} e^{m \Delta \left( R_{\text{in}}(h), R_{\text{out}}(h) \right)}$$

Markov's Inequality

$$\leq_{1-\delta} \text{KL}(Q \| P) + \ln \frac{1}{\delta} \mathbf{E}_{S' \sim D^m} \mathbf{E}_{h \sim P} e^{m \cdot \Delta \left( R_{\text{in}}(h), R_{\text{out}}(h) \right)}$$

Expectation swap

$$= \text{KL}(Q \| P) + \ln \frac{1}{\delta} \mathbf{E}_{h \sim P} \mathbf{E}_{S' \sim D^m} e^{m \cdot \Delta \left( R_{\text{in}}(h), R_{\text{out}}(h) \right)}$$

Binomial law

$$= \text{KL}(Q \| P) + \ln \frac{1}{\delta} \mathbf{E}_{h \sim P} \sum_{k=0}^m \text{Bin}(k; m, R_{\text{out}}(h)) e^{m \cdot \Delta \left( \frac{k}{m}, R_{\text{out}}(h) \right)}$$

Supremum over risk

$$\leq \text{KL}(Q \| P) + \ln \frac{1}{\delta} \sup_{r \in [0,1]} \left[ \sum_{k=0}^m \text{Bin}(k; m, r) e^{m \Delta \left( \frac{k}{m}, r \right)} \right]$$

$$\Pr_{S \sim D^m} \left( \forall Q \text{ on } \mathcal{H} : \Delta \left( R_{\text{in}}(Q), R_{\text{out}}(Q) \right) \leq \frac{1}{m} \left[ \text{KL}(Q \| P) + \ln \frac{\mathcal{J}_{\Delta}(m)}{\delta} \right] \right) \geq 1 - \delta.$$

**Proof.**

$$m \cdot \Delta \left( \mathbf{E}_{h \sim Q} R_{\text{in}}(h), \mathbf{E}_{h \sim Q} R_{\text{out}}(h) \right)$$

Jensen's Inequality

$$\leq \mathbf{E}_{h \sim Q} m \cdot \Delta \left( R_{\text{in}}(h), R_{\text{out}}(h) \right)$$

Change of measure

$$\leq \text{KL}(Q \| P) + \ln \mathbf{E}_{h \sim P} e^{m \Delta \left( R_{\text{in}}(h), R_{\text{out}}(h) \right)}$$

Markov's Inequality

$$\leq_{1-\delta} \text{KL}(Q \| P) + \ln \frac{1}{\delta} \mathbf{E}_{S' \sim D^m} \mathbf{E}_{h \sim P} e^{m \cdot \Delta \left( R_{\text{in}}(h), R_{\text{out}}(h) \right)}$$

Expectation swap

$$= \text{KL}(Q \| P) + \ln \frac{1}{\delta} \mathbf{E}_{h \sim P} \mathbf{E}_{S' \sim D^m} e^{m \cdot \Delta \left( R_{\text{in}}(h), R_{\text{out}}(h) \right)}$$

Binomial law

$$= \text{KL}(Q \| P) + \ln \frac{1}{\delta} \mathbf{E}_{h \sim P} \sum_{k=0}^m \text{Bin}(k; m, R_{\text{out}}(h)) e^{m \cdot \Delta \left( \frac{k}{m}, R_{\text{out}}(h) \right)}$$

Supremum over risk

$$\leq \text{KL}(Q \| P) + \ln \frac{1}{\delta} \sup_{r \in [0,1]} \left[ \sum_{k=0}^m \text{Bin}(k; m, r) e^{m \Delta \left( \frac{k}{m}, r \right)} \right]$$

$$= \text{KL}(Q \| P) + \ln \frac{1}{\delta} \mathcal{J}_{\Delta}(m).$$

□

## General theorem

$$\Pr_{S \sim D^m} \left( \forall Q \text{ on } \mathcal{H} : \Delta \left( R_{\text{in}}(Q), R_{\text{out}}(Q) \right) \leq \frac{1}{m} \left[ \text{KL}(Q \| P) + \ln \frac{\mathcal{I}_{\Delta}(m)}{\delta} \right] \right) \geq 1 - \delta.$$

## Corollary

[...] with probability at least  $1 - \delta$  over the choice of  $S \sim D^m$ , for all  $Q$  on  $\mathcal{H}$  :

$$\text{(a)} \quad \text{kl} \left( R_{\text{in}}(Q), R_{\text{out}}(Q) \right) \leq \frac{1}{m} \left[ \text{KL}(Q \| P) + \ln \frac{2\sqrt{m}}{\delta} \right], \quad \text{Langford and Seeger [31]}$$

$$\text{kl}(q, p) \stackrel{\text{def}}{=} q \ln \frac{q}{p} + (1 - q) \ln \frac{1 - q}{1 - p}$$

## General theorem

$$\Pr_{S \sim D^m} \left( \forall Q \text{ on } \mathcal{H} : \Delta(R_{\text{in}}(Q), R_{\text{out}}(Q)) \leq \frac{1}{m} \left[ \text{KL}(Q \| P) + \ln \frac{J_{\Delta}(m)}{\delta} \right] \right) \geq 1 - \delta.$$

## Corollary

[...] with probability at least  $1 - \delta$  over the choice of  $S \sim D^m$ , for all  $Q$  on  $\mathcal{H}$  :

$$\text{(a)} \quad \text{kl}(R_{\text{in}}(Q), R_{\text{out}}(Q)) \leq \frac{1}{m} \left[ \text{KL}(Q \| P) + \ln \frac{2\sqrt{m}}{\delta} \right], \quad \text{Langford and Seeger [31]}$$

$$\text{(b)} \quad R_{\text{out}}(Q) \leq R_{\text{in}}(Q) + \sqrt{\frac{1}{2m} \left[ \text{KL}(Q \| P) + \ln \frac{2\sqrt{m}}{\delta} \right]}, \quad \text{McAllester [40, 43]}$$

$$\text{kl}(q, p) \stackrel{\text{def}}{=} q \ln \frac{q}{p} + (1 - q) \ln \frac{1 - q}{1 - p} \geq 2(q - p)^2,$$

## General theorem

$$\Pr_{S \sim D^m} \left( \forall Q \text{ on } \mathcal{H} : \Delta(R_{\text{in}}(Q), R_{\text{out}}(Q)) \leq \frac{1}{m} \left[ \text{KL}(Q \| P) + \ln \frac{J_{\Delta}(m)}{\delta} \right] \right) \geq 1 - \delta.$$

## Corollary

[...] with probability at least  $1 - \delta$  over the choice of  $S \sim D^m$ , for all  $Q$  on  $\mathcal{H}$  :

(a)  $\text{kl}(R_{\text{in}}(Q), R_{\text{out}}(Q)) \leq \frac{1}{m} \left[ \text{KL}(Q \| P) + \ln \frac{2\sqrt{m}}{\delta} \right], \quad \text{Langford and Seeger [31]}$

(b)  $R_{\text{out}}(Q) \leq R_{\text{in}}(Q) + \sqrt{\frac{1}{2m} \left[ \text{KL}(Q \| P) + \ln \frac{2\sqrt{m}}{\delta} \right]}, \quad \text{McAllester [40, 43]}$

(c)  $R_{\text{out}}(Q) \leq \frac{1}{1 - e^{-c}} (c \cdot R_{\text{in}}(Q) + \frac{1}{m} [\text{KL}(Q \| P) + \ln \frac{1}{\delta}]), \quad \text{Catoni [11]}$

$$\text{kl}(q, p) \stackrel{\text{def}}{=} q \ln \frac{q}{p} + (1 - q) \ln \frac{1 - q}{1 - p} \geq 2(q - p)^2,$$

$$\Delta_c(q, p) \stackrel{\text{def}}{=} -\ln[1 - (1 - e^{-c}) \cdot p] - c \cdot q,$$

## General theorem

$$\Pr_{S \sim D^m} \left( \forall Q \text{ on } \mathcal{H} : \Delta \left( R_{\text{in}}(Q), R_{\text{out}}(Q) \right) \leq \frac{1}{m} \left[ \text{KL}(Q \| P) + \ln \frac{J_{\Delta}(m)}{\delta} \right] \right) \geq 1 - \delta.$$

## Corollary

[...] with probability at least  $1 - \delta$  over the choice of  $S \sim D^m$ , for all  $Q$  on  $\mathcal{H}$  :

$$(a) \quad \text{kl} \left( R_{\text{in}}(Q), R_{\text{out}}(Q) \right) \leq \frac{1}{m} \left[ \text{KL}(Q \| P) + \ln \frac{2\sqrt{m}}{\delta} \right], \quad \text{Langford and Seeger [31]}$$

$$(b) \quad R_{\text{out}}(Q) \leq R_{\text{in}}(Q) + \sqrt{\frac{1}{2m} \left[ \text{KL}(Q \| P) + \ln \frac{2\sqrt{m}}{\delta} \right]}, \quad \text{McAllester [40, 43]}$$

$$(c) \quad R_{\text{out}}(Q) \leq \frac{1}{1 - e^{-c}} \left( c \cdot R_{\text{in}}(Q) + \frac{1}{m} \left[ \text{KL}(Q \| P) + \ln \frac{1}{\delta} \right] \right), \quad \text{Catoni [11]}$$

$$(d) \quad R_{\text{out}}(Q) \leq R_{\text{in}}(Q) + \frac{1}{\lambda} \left[ \text{KL}(Q \| P) + \ln \frac{1}{\delta} + f(\lambda, m) \right]. \quad \text{Alquier et al. [4]}$$

$$\text{kl}(q, p) \stackrel{\text{def}}{=} q \ln \frac{q}{p} + (1 - q) \ln \frac{1 - q}{1 - p} \geq 2(q - p)^2,$$

$$\Delta_c(q, p) \stackrel{\text{def}}{=} -\ln[1 - (1 - e^{-c}) \cdot p] - c \cdot q,$$

$$\Delta_\lambda(q, p) \stackrel{\text{def}}{=} \frac{\lambda}{m} (p - q).$$



# Proof of the Langford/Seeger bound

Follows immediately from General Theorem by choosing  $\Delta(q, p) = \text{kl}(q, p)$ .

# Proof of the Langford/Seeger bound

Follows immediately from General Theorem by choosing  $\Delta(q, p) = \text{kl}(q, p)$ .

■ Indeed, in that case we have

$$\begin{aligned} \mathbf{E}_{S \sim D^m} \mathbf{E}_{h \sim P} e^{m \Delta(R_S(h), R(h))} &= \mathbf{E}_{h \sim P} \mathbf{E}_{S \sim D^m} \left( \frac{R_S(h)}{R(h)} \right)^{m R_S(h)} \left( \frac{1 - R_S(h)}{1 - R(h)} \right)^{m(1 - R_S(h))} \\ &= \mathbf{E}_{h \sim P} \sum_{k=0}^m \Pr_{S \sim D^m} \left( R_S(h) = \frac{k}{m} \right) \left( \frac{\frac{k}{m}}{R(h)} \right)^k \left( \frac{1 - \frac{k}{m}}{1 - R(h)} \right)^{m-k} \\ &= \sum_{k=0}^m \binom{m}{k} (k/m)^k (1 - k/m)^{m-k}, \\ &\leq 2\sqrt{m}. \end{aligned} \tag{1}$$

□

# Proof of the Langford/Seeger bound

Follows immediately from General Theorem by choosing  $\Delta(q, p) = \text{kl}(q, p)$ .

■ Indeed, in that case we have

$$\begin{aligned}
 \mathbf{E}_{S \sim D^m} \mathbf{E}_{h \sim P} e^{m \Delta(R_S(h), R(h))} &= \mathbf{E}_{h \sim P} \mathbf{E}_{S \sim D^m} \left( \frac{R_S(h)}{R(h)} \right)^{m R_S(h)} \left( \frac{1 - R_S(h)}{1 - R(h)} \right)^{m(1 - R_S(h))} \\
 &= \mathbf{E}_{h \sim P} \sum_{k=0}^m \Pr_{S \sim D^m} (R_S(h) = \frac{k}{m}) \left( \frac{\frac{k}{m}}{R(h)} \right)^k \left( \frac{1 - \frac{k}{m}}{1 - R(h)} \right)^{m-k} \\
 &= \sum_{k=0}^m \binom{m}{k} (k/m)^k (1 - k/m)^{m-k}, \\
 &\leq 2\sqrt{m}.
 \end{aligned} \tag{1}$$

□

■ Note that, in Line (1) of the proof,  $\Pr_{S \sim D^m} (R_S(h) = \frac{k}{m})$  is replaced by the probability mass function of the binomial.

# Proof of the Langford/Seeger bound

Follows immediately from General Theorem by choosing  $\Delta(q, p) = \text{kl}(q, p)$ .

■ Indeed, in that case we have

$$\begin{aligned} \mathbf{E}_{S \sim D^m} \mathbf{E}_{h \sim P} e^{m \Delta(R_S(h), R(h))} &= \mathbf{E}_{h \sim P} \mathbf{E}_{S \sim D^m} \left( \frac{R_S(h)}{R(h)} \right)^{m R_S(h)} \left( \frac{1 - R_S(h)}{1 - R(h)} \right)^{m(1 - R_S(h))} \\ &= \mathbf{E}_{h \sim P} \sum_{k=0}^m \Pr_{S \sim D^m} (R_S(h) = \frac{k}{m}) \left( \frac{\frac{k}{m}}{R(h)} \right)^k \left( \frac{1 - \frac{k}{m}}{1 - R(h)} \right)^{m-k} \\ &= \sum_{k=0}^m \binom{m}{k} (k/m)^k (1 - k/m)^{m-k}, \\ &\leq 2\sqrt{m}. \end{aligned} \tag{1}$$

□

■ Note that, in Line (1) of the proof,  $\Pr_{S \sim D^m} (R_S(h) = \frac{k}{m})$  is replaced by the probability mass function of the binomial.

■ This is **only true if** the examples of  $S$  are drawn iid. (i.e.,  $S \sim D^m$ )

# Proof of the Langford/Seeger bound

Follows immediately from General Theorem by choosing  $\Delta(q, p) = \text{kl}(q, p)$ .

■ Indeed, in that case we have

$$\begin{aligned}
 \mathbb{E}_{S \sim D^m} \mathbb{E}_{h \sim P} e^{m \Delta(R_S(h), R(h))} &= \mathbb{E}_{h \sim P} \mathbb{E}_{S \sim D^m} \left( \frac{R_S(h)}{R(h)} \right)^{m R_S(h)} \left( \frac{1 - R_S(h)}{1 - R(h)} \right)^{m(1 - R_S(h))} \\
 &= \mathbb{E}_{h \sim P} \sum_{k=0}^m \Pr_{S \sim D^m} (R_S(h) = \frac{k}{m}) \left( \frac{\frac{k}{m}}{R(h)} \right)^k \left( \frac{1 - \frac{k}{m}}{1 - R(h)} \right)^{m-k} \\
 &= \sum_{k=0}^m \binom{m}{k} (k/m)^k (1 - k/m)^{m-k}, \\
 &\leq 2\sqrt{m}.
 \end{aligned} \tag{1}$$

□

- Note that, in Line (1) of the proof,  $\Pr_{S \sim D^m} (R_S(h) = \frac{k}{m})$  is replaced by the probability mass function of the binomial.
- This is **only true** if the examples of  $S$  are drawn iid. (i.e.,  $S \sim D^m$ )
- So this result is no longer valid in the non iid case, even if General Theorem is.

# Linear classifiers

- We will choose the prior and posterior distributions to be Gaussians with unit variance.

# Linear classifiers


- We will choose the prior and posterior distributions to be Gaussians with unit variance.
- The prior  $P$  will be centered at the origin with unit variance

# Linear classifiers

- We will choose the prior and posterior distributions to be Gaussians with unit variance.
- The prior  $P$  will be centered at the origin with unit variance
- The specification of the centre for the posterior  $Q(\mathbf{w}, \mu)$  will be by a unit vector  $\mathbf{m}$  and a scale factor  $\mu$ .




## PAC-Bayes Bound for SVM (1/2)



■ **Prior**  $P$  is Gaussian  $\mathcal{N}(0, 1)$




# PAC-Bayes Bound for SVM (1/2)




- **Prior**  $P$  is Gaussian  $\mathcal{N}(0, 1)$
- Posterior is in the **direction**  $w$
- 
-

# PAC-Bayes Bound for SVM (1/2)



- **Prior**  $P$  is Gaussian  $\mathcal{N}(0, 1)$
- Posterior is in the **direction**  $w$
- at **distance**  $\mu$  from the origin
-

# PAC-Bayes Bound for SVM (1/2)



- **Prior**  $P$  is Gaussian  $\mathcal{N}(0, 1)$
- Posterior is in the **direction**  $w$
- at **distance**  $\mu$  from the origin
- **Posterior**  $Q$  is Gaussian

## PAC-Bayes Bound for SVM (2/2)

**Linear classifiers** performance may be bounded by

$$\text{KL}(\hat{Q}_S(\mathbf{w}, \mu) \| \boxed{Q_D(\mathbf{w}, \mu)}) \leq \frac{\text{KL}(P \| Q(\mathbf{w}, \mu)) + \ln \frac{m+1}{\delta}}{m}$$

# PAC-Bayes Bound for SVM (2/2)

**Linear classifiers** performance may be bounded by

$$\text{KL}(\hat{Q}_S(\mathbf{w}, \mu) \| \boxed{Q_{\mathcal{D}}(\mathbf{w}, \mu)}) \leq \frac{\text{KL}(P \| Q(\mathbf{w}, \mu)) + \ln \frac{m+1}{\delta}}{m}$$

- $Q_{\mathcal{D}}(\mathbf{w}, \mu)$  true performance of the stochastic classifier

# PAC-Bayes Bound for SVM (2/2)

**Linear classifiers** performance may be bounded by

$$\text{KL}(\hat{Q}_S(\mathbf{w}, \mu) \| \boxed{Q_{\mathcal{D}}(\mathbf{w}, \mu)}) \leq \frac{\text{KL}(P \| Q(\mathbf{w}, \mu)) + \ln \frac{m+1}{\delta}}{m}$$

- $Q_{\mathcal{D}}(\mathbf{w}, \mu)$  true performance of the stochastic classifier
- SVM is deterministic classifier that exactly corresponds to  $\text{sgn}(\mathbb{E}_{\mathbf{c} \sim Q(\mathbf{m}\mathbf{w}, \mu)}[\mathbf{c}(\mathbf{x})])$  as centre of the Gaussian gives the same classification as halfspace with more weight.

# PAC-Bayes Bound for SVM (2/2)

**Linear classifiers** performance may be bounded by

$$\text{KL}(\hat{Q}_S(\mathbf{w}, \mu) \| \boxed{Q_{\mathcal{D}}(\mathbf{w}, \mu)}) \leq \frac{\text{KL}(P \| Q(\mathbf{w}, \mu)) + \ln \frac{m+1}{\delta}}{m}$$

- $Q_{\mathcal{D}}(\mathbf{w}, \mu)$  true performance of the stochastic classifier
- SVM is deterministic classifier that exactly corresponds to  $\text{sgn}(\mathbb{E}_{\mathbf{c} \sim Q(\mathbf{m}\mathbf{w}, \mu)}[\mathbf{c}(\mathbf{x})])$  as centre of the Gaussian gives the same classification as halfspace with more weight.
- Hence its error bounded by  $2Q_{\mathcal{D}}(\mathbf{m}\mathbf{w}, \mu)$ , since as observed above if  $\mathbf{x}$  misclassified at least half of  $\mathbf{c} \sim Q$  err.



## PAC-Bayes Bound for SVM (2/2)

**Linear classifiers** performance may be bounded by

$$\text{KL}(\hat{Q}_S(\mathbf{w}, \mu) \| Q_{\mathcal{D}}(\mathbf{w}, \mu)) \leq \frac{\text{KL}(P \| Q(\mathbf{w}, \mu)) + \ln \frac{m+1}{\delta}}{m}$$

## PAC-Bayes Bound for SVM (2/2)

**Linear classifiers** performance may be bounded by

$$\text{KL}(\hat{Q}_S(\mathbf{w}, \mu) \| Q_{\mathcal{D}}(\mathbf{w}, \mu)) \leq \frac{\text{KL}(P \| Q(\mathbf{w}, \mu)) + \ln \frac{m+1}{\delta}}{m}$$

- $\hat{Q}_S(\mathbf{w}, \mu)$  stochastic measure of the training error

## PAC-Bayes Bound for SVM (2/2)

**Linear classifiers** performance may be bounded by

$$\text{KL}(\hat{Q}_S(\mathbf{w}, \mu) \| Q_{\mathcal{D}}(\mathbf{w}, \mu)) \leq \frac{\text{KL}(P \| Q(\mathbf{w}, \mu)) + \ln \frac{m+1}{\delta}}{m}$$

- $\hat{Q}_S(\mathbf{w}, \mu)$  stochastic measure of the training error
- $\hat{Q}_S(\mathbf{w}, \mu) = \mathbb{E}_m[\tilde{F}(\mu\gamma(\mathbf{x}, y))]$

# PAC-Bayes Bound for SVM (2/2)

**Linear classifiers** performance may be bounded by

$$\text{KL}(\hat{Q}_S(\mathbf{w}, \mu) \| Q_{\mathcal{D}}(\mathbf{w}, \mu)) \leq \frac{\text{KL}(P \| Q(\mathbf{w}, \mu)) + \ln \frac{m+1}{\delta}}{m}$$

- $\hat{Q}_S(\mathbf{w}, \mu)$  stochastic measure of the training error
- $\hat{Q}_S(\mathbf{w}, \mu) = \mathbb{E}_m[\tilde{F}(\mu\gamma(\mathbf{x}, y))]$
- $\gamma(\mathbf{x}, y) = (y\mathbf{w}^T\phi(\mathbf{x})) / (\|\phi(\mathbf{x})\| \|\mathbf{w}\|)$

# PAC-Bayes Bound for SVM (2/2)

**Linear classifiers** performance may be bounded by

$$\text{KL}(\hat{Q}_S(\mathbf{w}, \mu) \| Q_{\mathcal{D}}(\mathbf{w}, \mu)) \leq \frac{\text{KL}(P \| Q(\mathbf{w}, \mu)) + \ln \frac{m+1}{\delta}}{m}$$

- $\hat{Q}_S(\mathbf{w}, \mu)$  stochastic measure of the training error
- $\hat{Q}_S(\mathbf{w}, \mu) = \mathbb{E}_m[\tilde{F}(\mu\gamma(\mathbf{x}, y))]$
- $\gamma(\mathbf{x}, y) = (y\mathbf{w}^T \phi(\mathbf{x})) / (\|\phi(\mathbf{x})\| \|\mathbf{w}\|)$
- $\tilde{F}(t) = 1 - \frac{1}{\sqrt{2\pi}} \int_{-\infty}^t e^{-x^2/2} dx$

## PAC-Bayes Bound for SVM (2/2)

**Linear classifiers** performance may be bounded by

$$\text{KL}(\hat{Q}_S(\mathbf{w}, \mu) \| Q_{\mathcal{D}}(\mathbf{w}, \mu)) \leq \frac{\boxed{\text{KL}(P \| Q(\mathbf{w}, \mu))} + \ln \frac{m+1}{\delta}}{m}$$

# PAC-Bayes Bound for SVM (2/2)

**Linear classifiers** performance may be bounded by

$$\text{KL}(\hat{Q}_S(\mathbf{w}, \mu) \| Q_{\mathcal{D}}(\mathbf{w}, \mu)) \leq \frac{\boxed{\text{KL}(P \| Q(\mathbf{w}, \mu))} + \ln \frac{m+1}{\delta}}{m}$$

- Prior  $P \equiv$  Gaussian centered on the origin

## PAC-Bayes Bound for SVM (2/2)

**Linear classifiers** performance may be bounded by

$$\text{KL}(\hat{Q}_S(\mathbf{w}, \mu) \| Q_{\mathcal{D}}(\mathbf{w}, \mu)) \leq \frac{\text{KL}(P \| Q(\mathbf{w}, \mu)) + \ln \frac{m+1}{\delta}}{m}$$

- Prior  $P \equiv$  Gaussian centered on the origin
- Posterior  $Q \equiv$  Gaussian along  $\mathbf{w}$  at a distance  $\mu$  from the origin



## PAC-Bayes Bound for SVM (2/2)

**Linear classifiers** performance may be bounded by

$$\text{KL}(\hat{Q}_S(\mathbf{w}, \mu) \| Q_{\mathcal{D}}(\mathbf{w}, \mu)) \leq \frac{\boxed{\text{KL}(P \| Q(\mathbf{w}, \mu))} + \ln \frac{m+1}{\delta}}{m}$$

- Prior  $P \equiv$  Gaussian centered on the origin
- Posterior  $Q \equiv$  Gaussian along  $\mathbf{w}$  at a distance  $\mu$  from the origin
- $\text{KL}(P \| Q) = \mu^2/2$

## PAC-Bayes Bound for SVM (2/2)

**Linear classifiers** performance may be bounded by

$$\text{KL}(\hat{Q}_S(\mathbf{w}, \mu) \| Q_D(\mathbf{w}, \mu)) \leq \frac{\text{KL}(P \| Q(\mathbf{w}, \mu)) + \ln \frac{m+1}{\delta}}{m}$$

## PAC-Bayes Bound for SVM (2/2)

**Linear classifiers** performance may be bounded by

$$\text{KL}(\hat{Q}_S(\mathbf{w}, \mu) \| Q_D(\mathbf{w}, \mu)) \leq \frac{\text{KL}(P \| Q(\mathbf{w}, \mu)) + \ln \frac{m+1}{\delta}}{m}$$

■  $\delta$  is the confidence

# PAC-Bayes Bound for SVM (2/2)

**Linear classifiers** performance may be bounded by

$$\text{KL}(\hat{Q}_S(\mathbf{w}, \mu) \| Q_D(\mathbf{w}, \mu)) \leq \frac{\text{KL}(P \| Q(\mathbf{w}, \mu)) + \ln \frac{m+1}{\delta}}{m}$$

- $\delta$  is the confidence
- The bound holds with probability  $1 - \delta$  over the random i.i.d. selection of the training data.

# Form of the SVM bound

- Note that bound holds for all posterior distributions so that we can choose  $\mu$  to optimise the bound

# Form of the SVM bound

- Note that bound holds for all posterior distributions so that we can choose  $\mu$  to optimise the bound
- If we define the inverse of the KL by

$$\text{KL}^{-1}(q, A) = \max\{p : \text{KL}(q\|p) \leq A\}$$

then have with probability at least  $1 - \delta$

$$\Pr(\langle \mathbf{w}, \phi(\mathbf{x}) \rangle \neq y) \leq 2 \min_{\mu} \text{KL}^{-1} \left( \mathbb{E}_m[\tilde{F}(\mu\gamma(\mathbf{x}, y))], \frac{\mu^2/2 + \ln \frac{m+1}{\delta}}{m} \right)$$

# Gives SVM Optimisation

## ■ Primal form:


$$\begin{aligned} \min_{\mathbf{w}, \xi_i} \quad & \left[ \frac{1}{2} \|\mathbf{w}\|^2 + C \sum_{i=1}^m \xi_i \right] \\ \text{s.t.} \quad & y_i \mathbf{w}^T \phi(\mathbf{x}_i) \geq 1 - \xi_i \quad i = 1, \dots, m \\ & \xi_i \geq 0 \quad i = 1, \dots, m \end{aligned}$$

## ■ Dual form:

$$\begin{aligned} \max_{\alpha} \quad & \left[ \sum_{i=1}^m \alpha_i - \frac{1}{2} \sum_{i,j=1}^m \alpha_i \alpha_j y_i y_j \kappa(\mathbf{x}_i, \mathbf{x}_j) \right] \\ \text{s.t.} \quad & 0 \leq \alpha_i \leq C \quad i = 1, \dots, m \end{aligned}$$

where  $\kappa(\mathbf{x}_i, \mathbf{x}_j) = \langle \phi(\mathbf{x}_i), \phi(\mathbf{x}_j) \rangle$  and  $\langle \mathbf{w}, \phi(\mathbf{x}) \rangle = \sum_{i=1}^m \alpha_i y_i \kappa(\mathbf{x}_i, \mathbf{x})$ .

## Slack variable conversion





# Data- or distribution-dependent priors

- An important component in the PAC-Bayes analysis is the choice of the **prior distribution**

# Data- or distribution-dependent priors

- An important component in the PAC-Bayes analysis is the choice of the **prior distribution**
- The results hold whatever the choice of prior, provided that it is chosen *before* seeing the data sample

# Data- or distribution-dependent priors

- An important component in the PAC-Bayes analysis is the choice of the **prior distribution**
- The results hold whatever the choice of prior, provided that it is chosen *before* seeing the data sample
- Are there ways we can choose a 'better' prior?

# Data- or distribution-dependent priors

- An important component in the PAC-Bayes analysis is the choice of the **prior distribution**
- The results hold whatever the choice of prior, provided that it is chosen *before* seeing the data sample
- Are there ways we can choose a 'better' prior?
- Will explore:

# Data- or distribution-dependent priors

- An important component in the PAC-Bayes analysis is the choice of the **prior distribution**
- The results hold whatever the choice of prior, provided that it is chosen *before* seeing the data sample
- Are there ways we can choose a 'better' prior?
- Will explore:
  - using part of the data to *learn the prior* for SVMs, but also more interestingly and more generally

# Data- or distribution-dependent priors

- An important component in the PAC-Bayes analysis is the choice of the **prior distribution**
- The results hold whatever the choice of prior, provided that it is chosen *before* seeing the data sample
- Are there ways we can choose a 'better' prior?
- Will explore:
  - using part of the data to *learn the prior* for SVMs, but also more interestingly and more generally
  - defining the prior in terms of the *data generating distribution* (aka *localised PAC-Bayes*).

## Learning the prior (1/3)

- Bound depends on the **distance between prior and posterior**

## Learning the prior (1/3)

- Bound depends on the **distance between prior and posterior**
- Better prior (closer to posterior) would lead to **tighter bound**



## Learning the prior (1/3)

- Bound depends on the **distance between prior and posterior**
- Better prior (closer to posterior) would lead to **tighter bound**
- **Learn** the prior  $P$  with part of the data


# Learning the prior (1/3)

- Bound depends on the **distance between prior and posterior**
- Better prior (closer to posterior) would lead to **tighter bound**
- **Learn** the prior  $P$  with part of the data
- Introduce the learnt prior **in the bound**

# Learning the prior (1/3)

- Bound depends on the **distance between prior and posterior**
- Better prior (closer to posterior) would lead to **tighter bound**
- **Learn** the prior  $P$  with part of the data
- Introduce the learnt prior **in the bound**
- Compute stochastic error with **remaining data**


## New prior for the SVM (3/3)



- Solve SVM with **subset of patterns**




## New prior for the SVM (3/3)




- Solve SVM with **subset of patterns**
- Prior in the **direction  $w_r$**
- 
-

## New prior for the SVM (3/3)



- Solve SVM with **subset of patterns**
- Prior in the **direction  $w_r$**
- **Posterior** like PAC-Bayes Bound
-

## New prior for the SVM (3/3)



- Solve SVM with **subset of patterns**
- Prior in the **direction  $w_r$**
- **Posterior** like PAC-Bayes Bound
- **New bound** depends on  $KL(P||Q)$

## New Bound for the SVM (2/3)

SVM performance may be **tightly** bounded by

$$\text{KL}(\hat{Q}_S(\mathbf{w}, \mu) \| \boxed{Q_{\mathcal{D}}(\mathbf{w}, \mu)}) \leq \frac{0.5 \|\mu \mathbf{w} - \eta \mathbf{w}_r\|^2 + \ln \frac{(m-r+1)J}{\delta}}{m-r}$$



## New Bound for the SVM (2/3)

SVM performance may be **tightly** bounded by

$$\text{KL}(\hat{Q}_S(\mathbf{w}, \mu) \| \boxed{Q_{\mathcal{D}}(\mathbf{w}, \mu)}) \leq \frac{0.5 \|\mu \mathbf{w} - \eta \mathbf{w}_r\|^2 + \ln \frac{(m-r+1)J}{\delta}}{m-r}$$

- $Q_{\mathcal{D}}(\mathbf{w}, \mu)$  true performance of the classifier

## New Bound for the SVM (2/3)

SVM performance may be **tightly** bounded by

$$\text{KL}(\hat{Q}_S(\mathbf{w}, \mu) \| Q_{\mathcal{D}}(\mathbf{w}, \mu)) \leq \frac{0.5 \|\mu \mathbf{w} - \eta \mathbf{w}_r\|^2 + \ln \frac{(m-r+1)J}{\delta}}{m-r}$$

## New Bound for the SVM (2/3)

SVM performance may be **tightly** bounded by

$$\text{KL}(\hat{Q}_S(\mathbf{w}, \mu) \| Q_{\mathcal{D}}(\mathbf{w}, \mu)) \leq \frac{0.5 \|\mu \mathbf{w} - \eta \mathbf{w}_r\|^2 + \ln \frac{(m-r+1)J}{\delta}}{m-r}$$

- $\hat{Q}_S(\mathbf{w}, \mu)$  stochastic measure of the training error on remaining data

$$\hat{Q}(\mathbf{w}, \mu)_S = \mathbb{E}_{m-r}[\tilde{F}(\mu \gamma(\mathbf{x}, y))]$$

## New Bound for the SVM (2/3)

SVM performance may be **tightly** bounded by

$$\text{KL}(\hat{Q}_S(\mathbf{w}, \mu) \| Q_{\mathcal{D}}(\mathbf{w}, \mu)) \leq \frac{0.5 \|\mu \mathbf{w} - \eta \mathbf{w}_r\|^2 + \ln \frac{(m-r+1)J}{\delta}}{m-r}$$

## New Bound for the SVM (2/3)

SVM performance may be **tightly** bounded by

$$\text{KL}(\hat{Q}_S(\mathbf{w}, \mu) \| Q_{\mathcal{D}}(\mathbf{w}, \mu)) \leq \frac{0.5 \|\mu \mathbf{w} - \eta \mathbf{w}_r\|^2 + \ln \frac{(m-r+1)J}{\delta}}{m-r}$$

- $0.5 \|\mu \mathbf{w} - \eta \mathbf{w}_r\|^2$  distance between prior and posterior

## New Bound for the SVM (2/3)

SVM performance may be **tightly** bounded by

$$\text{KL}(\hat{Q}_S(\mathbf{w}, \mu) \| Q_{\mathcal{D}}(\mathbf{w}, \mu)) \leq \frac{0.5 \|\mu \mathbf{w} - \eta \mathbf{w}_r\|^2 + \ln \frac{(m-r+1)J}{\delta}}{m-r}$$

## New Bound for the SVM (2/3)

SVM performance may be **tightly** bounded by

$$\text{KL}(\hat{Q}_S(\mathbf{w}, \mu) \| Q_{\mathcal{D}}(\mathbf{w}, \mu)) \leq \frac{0.5 \|\mu \mathbf{w} - \eta \mathbf{w}_r\|^2 + \ln \frac{(m-r+1)J}{\delta}}{m-r}$$

- Penalty term only dependent on the remaining data  $m-r$

# Prior-SVM

- New bound proportional to  $\|\mu\mathbf{w} - \eta\mathbf{w}_r\|^2$



# Prior-SVM

- New bound proportional to  $\|\mu\mathbf{w} - \eta\mathbf{w}_r\|^2$
- Classifier that **optimises the bound**

# Prior-SVM

- New bound proportional to  $\|\mu\mathbf{w} - \eta\mathbf{w}_r\|^2$
- Classifier that **optimises the bound**
- Optimisation problem to determine the **p-SVM**

$$\begin{aligned} \min_{\mathbf{w}, \xi_i} & \left[ \frac{1}{2} \|\mathbf{w} - \mathbf{w}_r\|^2 + C \sum_{i=1}^{m-r} \xi_i \right] \\ \text{s.t.} \quad & y_i \mathbf{w}^T \phi(\mathbf{x}_i) \geq 1 - \xi_i & i = 1, \dots, m-r \\ & \xi_i \geq 0 & i = 1, \dots, m-r \end{aligned}$$

# Prior-SVM

- New bound proportional to  $\|\mu\mathbf{w} - \eta\mathbf{w}_r\|^2$
- Classifier that **optimises the bound**
- Optimisation problem to determine the **p-SVM**

$$\begin{aligned} \min_{\mathbf{w}, \xi_i} & \left[ \frac{1}{2} \|\mathbf{w} - \mathbf{w}_r\|^2 + C \sum_{i=1}^{m-r} \xi_i \right] \\ \text{s.t.} \quad & y_i \mathbf{w}^T \phi(\mathbf{x}_i) \geq 1 - \xi_i & i = 1, \dots, m-r \\ & \xi_i \geq 0 & i = 1, \dots, m-r \end{aligned}$$

- The p-SVM is only solved with the **remaining points**

# Bound for p-SVM

- 1 Determine the **prior** with a subset of the training examples to obtain  $\mathbf{w}_r$

# Bound for p-SVM

- 1 Determine the **prior** with a subset of the training examples to obtain  $\mathbf{w}_r$
- 2 Solve **p-SVM** and obtain  $\mathbf{w}$

# Bound for p-SVM

- 1 Determine the **prior** with a subset of the training examples to obtain  $\mathbf{w}_r$
- 2 Solve **p-SVM** and obtain  $\mathbf{w}$
- 3 **Margin** for the stochastic classifier  $\hat{Q}_s$

$$\gamma(\mathbf{x}_j, y_j) = \frac{y_j \mathbf{w}^T \phi(\mathbf{x}_j)}{\|\phi(\mathbf{x}_j)\| \|\mathbf{w}\|} \quad j = 1, \dots, m - r$$

# Bound for p-SVM

- 1 Determine the **prior** with a subset of the training examples to obtain  $\mathbf{w}_r$
- 2 Solve **p-SVM** and obtain  $\mathbf{w}$
- 3 **Margin** for the stochastic classifier  $\hat{Q}_s$

$$\gamma(\mathbf{x}_j, y_j) = \frac{y_j \mathbf{w}^T \phi(\mathbf{x}_j)}{\|\phi(\mathbf{x}_j)\| \|\mathbf{w}\|} \quad j = 1, \dots, m - r$$

- 4 **Linear search** to obtain the optimal value of  $\mu$ . This introduces an insignificant extra penalty term

# Bound for $\eta$ -prior-SVM

- Prior is elongated along the line of  $\mathbf{w}_r$  but spherical with variance 1 in other directions



# Bound for $\eta$ -prior-SVM

- Prior is elongated along the line of  $\mathbf{w}_r$  but spherical with variance 1 in other directions
- Posterior again on the line of  $\mathbf{w}$  at a distance  $\mu$  chosen to optimise the bound.

# Bound for $\eta$ -prior-SVM

- Prior is elongated along the line of  $\mathbf{w}_r$  but spherical with variance 1 in other directions
- Posterior again on the line of  $\mathbf{w}$  at a distance  $\mu$  chosen to optimise the bound.
- Resulting bound depends on a benign parameter  $\tau$  determining the variance in the direction  $\mathbf{w}_r$

$$\text{KL}(\hat{Q}_{S \setminus R}(\mathbf{w}, \mu) \| Q_{\mathcal{D}}(\mathbf{w}, \mu)) \leqslant \frac{0.5(\ln(\tau^2) + \tau^{-2} - 1 + P_{\mathbf{w}_r}^{\parallel}(\mu \mathbf{w} - \mathbf{w}_r)^2 / \tau^2 + P_{\mathbf{w}_r}^{\perp}(\mu \mathbf{w})^2) + \ln(\frac{m-r+1}{\delta})}{m-r}$$

# $\eta$ -Prior-SVM

- Consider using a prior distribution  $P$  that is elongated in the direction of  $\mathbf{w}_r$

# $\eta$ -Prior-SVM

- Consider using a prior distribution  $P$  that is elongated in the direction of  $\mathbf{w}_r$
- This will mean that there is low penalty for large projections onto this direction

## $\eta$ -Prior-SVM

- Consider using a prior distribution  $P$  that is elongated in the direction of  $\mathbf{w}_r$
- This will mean that there is low penalty for large projections onto this direction
- Translates into an optimisation:

$$\min_{\mathbf{v}, \eta, \xi_i} \left[ \frac{1}{2} \|\mathbf{v}\|^2 + C \sum_{i=1}^{m-r} \xi_i \right]$$

# $\eta$ -Prior-SVM

- Consider using a prior distribution  $P$  that is elongated in the direction of  $\mathbf{w}_r$
- This will mean that there is low penalty for large projections onto this direction
- Translates into an optimisation:

$$\min_{\mathbf{v}, \eta, \xi_i} \left[ \frac{1}{2} \|\mathbf{v}\|^2 + C \sum_{i=1}^{m-r} \xi_i \right]$$

- subject to

$$\begin{aligned} y_i(\mathbf{v} + \eta \mathbf{w}_r)^T \phi(\mathbf{x}_i) &\geq 1 - \xi_i & i = 1, \dots, m-r \\ \xi_i &\geq 0 & i = 1, \dots, m-r \end{aligned}$$

# Model Selection with the new bound: setup

- Comparison of 10-fold Xvalidation, PAC-Bayes Bound and the Prior PAC-Bayes Bound

# Model Selection with the new bound: setup

- Comparison of 10-fold Xvalidation, PAC-Bayes Bound and the Prior PAC-Bayes Bound
- UCI datasets



# Model Selection with the new bound: setup

- Comparison of 10-fold Xvalidation, PAC-Bayes Bound and the Prior PAC-Bayes Bound
- UCI datasets
- Select  $C$  and  $\sigma$  that lead to minimum Classification Error (CE)

# Model Selection with the new bound: setup

- Comparison of 10-fold Xvalidation, PAC-Bayes Bound and the Prior PAC-Bayes Bound
- UCI datasets
- Select  $C$  and  $\sigma$  that lead to minimum Classification Error (CE)
  - For 10-F XV select the pair that minimize the validation error

# Model Selection with the new bound: setup

- Comparison of 10-fold Xvalidation, PAC-Bayes Bound and the Prior PAC-Bayes Bound
- UCI datasets
- Select  $C$  and  $\sigma$  that lead to minimum Classification Error (CE)
  - For 10-F XV select the pair that minimize the validation error
  - For PAC-Bayes Bound and Prior PAC-Bayes Bound select the pair that minimize the bound

# Results

		Classifier					
Problem		SVM				$\eta$ Prior SVM	
		2FCV	10FCV	PAC	PrPAC	PrPAC	$\tau$ -PrPAC
digits	Bound	–	–	0.175	0.107	0.050	<b>0.047</b>
	TE	<b>0.007</b>	<b>0.007</b>	<b>0.007</b>	0.014	0.010	0.009
waveform	Bound	–	–	0.203	0.185	0.178	<b>0.176</b>
	TE	0.090	0.086	<b>0.084</b>	0.088	0.087	0.086
pima	Bound	–	–	0.424	0.420	0.428	<b>0.416</b>
	TE	0.244	0.245	<b>0.229</b>	<b>0.229</b>	0.233	0.233
ringnorm	Bound	–	–	0.203	0.110	0.053	<b>0.050</b>
	TE	<b>0.016</b>	<b>0.016</b>	0.018	0.018	<b>0.016</b>	<b>0.016</b>
spam	Bound	–	–	0.254	0.198	0.186	<b>0.178</b>
	TE	0.066	<b>0.063</b>	0.067	0.077	0.070	0.072
Average	TE	0.0846	0.0834	0.081	0.0852	0.0832	0.0832

# Take home messages

- Bounds are remarkably tight: for final column average factor between bound and TE is under 3.

# Take home messages

- Bounds are remarkably tight: for final column average factor between bound and TE is under 3.
- Model selection from the bounds is as good as 10FCV: in fact all but one of the PAC-Bayes model selections give better averages for TE.

# Take home messages

- Bounds are remarkably tight: for final column average factor between bound and TE is under 3.
- Model selection from the bounds is as good as 10FCV: in fact all but one of the PAC-Bayes model selections give better averages for TE.
- The better bounds do not appear to give better model selection - best model selection is from the simplest bound.
  - A. Ambroladze, E. Parrado-Hernández, and J. Shawe-Taylor. Tighter PAC-Bayes bounds. In *Advances in Neural Information Processing Systems* 18, (2006) Pages 9-16.
  - P. Germain, A. Lacasse, F. Laviolette and M. Marchand. PAC-Bayesian learning of linear classifiers, in *Proceedings of the 26th International Conference on Machine Learning (ICML'09, Montréal, Canada.)*. ACM Press (2009), 382, Pages 453-460.

# Distribution-defined priors



# Distribution-defined priors

- Consider  $P$  and  $Q$  are Gibbs-Boltzmann distributions

$$P(h) := \frac{1}{Z'} e^{-\gamma \text{risk}(h)} \quad Q(h) := \frac{1}{Z} e^{-\gamma \text{risk}_S(h)}$$

# Distribution-defined priors

- Consider  $P$  and  $Q$  are Gibbs-Boltzmann distributions

$$P(h) := \frac{1}{Z'} e^{-\gamma \text{risk}(h)} \quad Q(h) := \frac{1}{Z} e^{-\gamma \hat{\text{risk}}_S(h)}$$

- These distributions are hard to work with since we cannot apply the bound to a single weight vector, but the bounds can be very tight:

$$KL_+(\hat{Q}_S(\gamma) \| Q_D(\gamma)) \leq \frac{1}{m} \left( \frac{\gamma}{\sqrt{m}} \sqrt{\ln \frac{8\sqrt{m}}{\delta}} + \frac{\gamma^2}{4m} + \ln \frac{4\sqrt{m}}{\delta} \right)$$

with the only uncertainty the dependence on  $\gamma$ .

# Distribution-defined priors

- Consider  $P$  and  $Q$  are Gibbs-Boltzmann distributions

$$P(h) := \frac{1}{Z'} e^{-\gamma \text{risk}(h)} \quad Q(h) := \frac{1}{Z} e^{-\gamma \hat{\text{risk}}_S(h)}$$

- These distributions are hard to work with since we cannot apply the bound to a single weight vector, but the bounds can be very tight:

$$KL_+(\hat{Q}_S(\gamma) \| Q_D(\gamma)) \leq \frac{1}{m} \left( \frac{\gamma}{\sqrt{m}} \sqrt{\ln \frac{8\sqrt{m}}{\delta}} + \frac{\gamma^2}{4m} + \ln \frac{4\sqrt{m}}{\delta} \right)$$

with the only uncertainty the dependence on  $\gamma$ .

- O. Catoni. A PAC-Bayesian approach to adaptive classification. Preprint n.840, Laboratoire de Probabilités et Modèles Aléatoires, Universités Paris 6 and Paris 7, 2003.
- G. Lever, F. Laviolette, J. Shawe-Taylor. Distribution-Dependent PAC-Bayes Priors. Proceedings of the 21st International Conference on Algorithmic Learning Theory (ALT 2010), 119-133.

# Observations

- We cannot compute the prior distribution  $P$  or even sample from it:

# Observations

- We cannot compute the prior distribution  $P$  or even sample from it:
  - Note that this would not be possible to consider in normal Bayesian inference;

# Observations

- We cannot compute the prior distribution  $P$  or even sample from it:
  - Note that this would not be possible to consider in normal Bayesian inference;
  - Trick here is that the error measures only depend on the posterior  $Q$ , while the bound depends on KL between posterior and prior: an estimate of this KL is made without knowing the prior explicitly

# Observations

- We cannot compute the prior distribution  $P$  or even sample from it:
  - Note that this would not be possible to consider in normal Bayesian inference;
  - Trick here is that the error measures only depend on the posterior  $Q$ , while the bound depends on KL between posterior and prior: an estimate of this KL is made without knowing the prior explicitly
- the Gibbs distributions are hard to sample from so not easy to work with this bound.

## Other distribution defined priors

- An alternative distribution defined prior for an SVM is to place symmetrical Gaussian at the weight vector:  
 $\mathbf{w}_p = \mathbb{E}_{(\mathbf{x}, y) \sim D}(y \boldsymbol{\phi}(\mathbf{x}))$  to give distributions that are easier to work with, but results not impressive...



## Other distribution defined priors

- An alternative distribution defined prior for an SVM is to place symmetrical Gaussian at the weight vector:  
 $\mathbf{w}_p = \mathbb{E}_{(\mathbf{x}, y) \sim D}(y \boldsymbol{\phi}(\mathbf{x}))$  to give distributions that are easier to work with, but results not impressive...
- What if we were to take the expected weight vector returned from a random training set of size  $m$ : then the KL between posterior and prior is related to the concentration of weight vectors from different training sets

## Other distribution defined priors

- An alternative distribution defined prior for an SVM is to place symmetrical Gaussian at the weight vector:  
 $\mathbf{w}_p = \mathbb{E}_{(\mathbf{x}, y) \sim D}(y \boldsymbol{\phi}(\mathbf{x}))$  to give distributions that are easier to work with, but results not impressive...
- What if we were to take the expected weight vector returned from a random training set of size  $m$ : then the KL between posterior and prior is related to the concentration of weight vectors from different training sets
- This is connected to stability...

# Outline

- stability

# Stability

Uniform **hypothesis sensitivity**  $\beta$  at sample size  $m$ :

$$\|A(z_{1:m}) - A(z'_{1:m})\| \leq \beta \sum_{i=1}^m \mathbf{1}[z_i \neq z'_i]$$

$(z_1, \dots, z_m)$

■  $A(z_{1:m}) \in \mathcal{H}$  normed space

■  $w_m = A(z_{1:m})$  ‘weight vector’

$(z'_1, \dots, z'_m)$

■ Lipschitz

■ smoothness

Uniform **loss sensitivity**  $\beta$  at sample size  $m$ :

$$|\ell(A(z_{1:m}), z) - \ell(A(z'_{1:m}), z)| \leq \beta \sum_{i=1}^m \mathbf{1}[z_i \neq z'_i]$$

■ worst-case

■ distribution-insensitive

■ data-insensitive

■ Open: data-dependent?

# Generalization from Stability

If  $A$  has sensitivity  $\beta$  at sample size  $m$ , then for any  $\delta \in (0, 1)$ ,

$$\text{w.p.} \geq 1 - \delta, \quad R_{\text{out}}(h) \leq R_{\text{in}}(h) + \epsilon(\beta, m, \delta)$$

# Generalization from Stability

If  $A$  has sensitivity  $\beta$  at sample size  $m$ , then for any  $\delta \in (0, 1)$ ,

$$\text{w.p.} \geq 1 - \delta, \quad R_{\text{out}}(h) \leq R_{\text{in}}(h) + \epsilon(\beta, m, \delta)$$

(e.g. Bousquet & Elisseeff)

# Generalization from Stability

If  $A$  has sensitivity  $\beta$  at sample size  $m$ , then for any  $\delta \in (0, 1)$ ,

$$\text{w.p.} \geq 1 - \delta, \quad R_{\text{out}}(h) \leq R_{\text{in}}(h) + \epsilon(\beta, m, \delta)$$

(e.g. Bousquet & Elisseeff)

- the intuition is that if individual examples do not affect the loss of an algorithm then it will be concentrated

# Generalization from Stability

If  $A$  has sensitivity  $\beta$  at sample size  $m$ , then for any  $\delta \in (0, 1)$ ,

$$\text{w.p.} \geq 1 - \delta, \quad R_{\text{out}}(h) \leq R_{\text{in}}(h) + \epsilon(\beta, m, \delta)$$

(e.g. Bousquet & Elisseeff)

- the intuition is that if individual examples do not affect the loss of an algorithm then it will be concentrated
- can be applied to kernel methods where  $\beta$  is related to the regularisation constant, but bounds are quite weak



# Generalization from Stability

If  $A$  has sensitivity  $\beta$  at sample size  $m$ , then for any  $\delta \in (0, 1)$ ,

$$\text{w.p.} \geq 1 - \delta, \quad R_{\text{out}}(h) \leq R_{\text{in}}(h) + \epsilon(\beta, m, \delta)$$

(e.g. Bousquet & Elisseeff)

- the intuition is that if individual examples do not affect the loss of an algorithm then it will be concentrated
- can be applied to kernel methods where  $\beta$  is related to the regularisation constant, but bounds are quite weak
- question: algorithm output is highly concentrated  
 $\implies$  stronger results?

# Stability + PAC-Bayes I

If  $A$  has uniform hypothesis stability  $\beta$  at sample size  $n$ , then for any  $\delta \in (0, 1)$ , **w.p.**  $\geq 1 - 2\delta$ ,

$$\text{KL}(R_{\text{in}}(Q) \| R_{\text{out}}(Q)) \leq \frac{\frac{n\beta^2}{2\sigma^2} \left(1 + \sqrt{\frac{1}{2} \log\left(\frac{1}{\delta}\right)}\right)^2 + \log\left(\frac{n+1}{\delta}\right)}{n}$$

Gaussian randomization

- $P = \mathcal{N}(\mathbb{E}[W_n], \sigma^2 I)$
- $Q = \mathcal{N}(W_n, \sigma^2 I)$
- $\text{KL}(Q \| P) = \frac{1}{2\sigma^2} \|W_n - \mathbb{E}[W_n]\|^2$

Main proof components:

- **w.p.**  $\geq 1 - \delta$ ,  $\text{KL}(R_{\text{in}}(Q) \| R_{\text{out}}(Q)) \leq \frac{\text{KL}(Q \| Q_0) + \log\left(\frac{n+1}{\delta}\right)}{n}$
- **w.p.**  $\geq 1 - \delta$ ,  $\|W_n - \mathbb{E}[W_n]\| \leq \sqrt{n} \beta \left(1 + \sqrt{\frac{1}{2} \log\left(\frac{1}{\delta}\right)}\right)$

# A flexible framework

# A flexible framework

Since 1997, PAC-Bayes has been successfully used in **many** machine learning settings (this list is by no means exhaustive).

**Statistical learning theory** *Audibert and Bousquet [6], Catoni [9, 10], Guedj [25], Guedj and Pujol [27], Maurer [39], McAllester [41, 42, 44, 45], Mhammedi et al. [46], Seeger [51, 52], Shawe-Taylor and Williamson [56], Thiemann et al. [58]*

**SVMs & linear classifiers** *Germain et al. [19], Langford and Shawe-Taylor [32], McAllester [44]*

**Supervised learning algorithms** reinterpreted as bound minimizers  
*Ambroladze et al. [5], Germain et al. [22], Shawe-Taylor and Hadoon [57]*

**High-dimensional regression** *Alquier and Biau [1], Alquier and Lounici [2], Guedj and Robbiano [24], Guedj and Alquier [26], Li et al. [35]*

**Classification** *Catoni [9, 10], Lacasse et al. [30], Langford and Shawe-Taylor [32], Parrado-Hernández et al. [49]*

# A flexible framework

**Transductive learning, domain adaptation** *Bégin et al. [7], Derbeko et al. [12], Germain et al. [20], Nozawa et al. [48]*

**Non-iid or heavy-tailed data** *Alquier and Guedj [3], Holland [29], Lever et al. [34], Seldin et al. [54, 55]*

**Density estimation** *Higgs and Shawe-Taylor [28], Seldin and Tishby [53]*

**Reinforcement learning** *Fard and Pineau [16], Fard et al. [17], Ghavamzadeh et al. [23], Seldin et al. [54, 55]*

**Sequential learning** *Gerchinovitz [18], Li et al. [36]*

**Algorithmic stability, differential privacy** *Dziugaite and Roy [13, 14], London [37], London et al. [38], Rivasplata et al. [50]*

**Deep neural networks** *Dziugaite and Roy [15], Letarte et al. [33], Neyshabur et al. [47], Zhou et al. [60]*

...

# References I

- [1] P. Alquier and G. Biau. Sparse single-index model. *Journal of Machine Learning Research*, 14:243–280, 2013.
- [2] P. Alquier and K. Lounici. PAC-Bayesian theorems for sparse regression estimation with exponential weights. *Electronic Journal of Statistics*, 5:127–145, 2011.
- [3] Pierre Alquier and Benjamin Guedj. Simpler PAC-Bayesian bounds for hostile data. *Machine Learning*, 107(5):887–902, 2018.
- [4] Pierre Alquier, James Ridgway, and Nicolas Chopin. On the properties of variational approximations of Gibbs posteriors. *ArXiv e-prints*, 2015. URL <http://arxiv.org/abs/1506.04091>.
- [5] A. Ambroladze, E. Parrado-Hernández, and J. Shawe-taylor. Tighter PAC-Bayes bounds. In *Advances in Neural Information Processing Systems, NIPS*, pages 9–16, 2007.
- [6] Jean-Yves Audibert and Olivier Bousquet. Combining PAC-Bayesian and generic chaining bounds. *Journal of Machine Learning Research*, 2007.
- [7] Luc Bégin, Pascal Germain, François Laviolette, and Jean-François Roy. PAC-Bayesian theory for transductive learning. In *AISTATS*, 2014.
- [8] Luc Bégin, Pascal Germain, François Laviolette, and Jean-François Roy. PAC-Bayesian bounds based on the Rényi divergence. In *AISTATS*, 2016.
- [9] O. Catoni. *Statistical Learning Theory and Stochastic Optimization*. École d'Été de Probabilités de Saint-Flour 2001. Springer, 2004.
- [10] O. Catoni. *PAC-Bayesian Supervised Classification: The Thermodynamics of Statistical Learning*, volume 56 of *Lecture notes – Monograph Series*. Institute of Mathematical Statistics, 2007.
- [11] Olivier Catoni. *PAC-Bayesian supervised classification: the thermodynamics of statistical learning*, volume 56. Inst. of Mathematical Statistics, 2007.
- [12] Philip Derbeko, Ran El-Yaniv, and Ron Meir. Explicit learning curves for transduction and application to clustering and compression algorithms. *J. Artif. Intell. Res. (JAIR)*, 22, 2004.
- [13] G. K. Dziugaite and D. M. Roy. Data-dependent PAC-Bayes priors via differential privacy. In *NeurIPS*, 2018.
- [14] G. K. Dziugaite and D. M. Roy. Entropy-SGD optimizes the prior of a PAC-Bayes bound: Generalization properties of Entropy-SGD and data-dependent priors. In *International Conference on Machine Learning*, pages 1376–1385, 2018.

# References II

- [15] Gintare K. Dziugaite and Daniel M. Roy. Computing nonvacuous generalization bounds for deep (stochastic) neural networks with many more parameters than training data. In *Proceedings of Uncertainty in Artificial Intelligence (UAI)*, 2017.
- [16] Mahdi Milani Fard and Joelle Pineau. PAC-Bayesian model selection for reinforcement learning. In *Advances in Neural Information Processing Systems (NIPS)*, 2010.
- [17] Mahdi Milani Fard, Joelle Pineau, and Csaba Szepesvári. PAC-Bayesian Policy Evaluation for Reinforcement Learning. In *UAI, Proceedings of the Twenty-Seventh Conference on Uncertainty in Artificial Intelligence*, pages 195–202, 2011.
- [18] S. Gerchinovitz. *Prédiction de suites individuelles et cadre statistique classique : étude de quelques liens autour de la régression parcimonieuse et des techniques d'agrégation*. PhD thesis, Université Paris-Sud, 2011.
- [19] P. Germain, A. Lacasse, F. Laviolette, and M. Marchand. PAC-Bayesian learning of linear classifiers. In *Proceedings of the 26th Annual International Conference on Machine Learning, ICML*, 2009.
- [20] P. Germain, A. Habrard, F. Laviolette, and E. Morvant. A new PAC-Bayesian perspective on domain adaptation. In *Proceedings of International Conference on Machine Learning*, volume 48, 2016.
- [21] Pascal Germain. *Généralisations de la théorie PAC-bayésienne pour l'apprentissage inductif, l'apprentissage transductif et l'adaptation de domaine*. PhD thesis, Université Laval, 2015.
- [22] Pascal Germain, Alexandre Lacasse, Mario Marchand, Sara Shanian, and François Laviolette. From PAC-Bayes bounds to KL regularization. In *Advances in Neural Information Processing Systems*, pages 603–610, 2009.
- [23] M. Ghavamzadeh, S. Mannor, J. Pineau, and A. Tamar. Bayesian reinforcement learning: A survey. *Foundations and Trends in Machine Learning*, 8(5-6):359–483, 2015.
- [24] B. Guedj and S. Robbiano. PAC-Bayesian high dimensional bipartite ranking. *Journal of Statistical Planning and Inference*, 196:70 – 86, 2018. ISSN 0378-3758.
- [25] Benjamin Guedj. A primer on PAC-Bayesian learning. *arXiv:1901.05353*, 2019. To appear in the Proceedings of the French Mathematical Society.
- [26] Benjamin Guedj and Pierre Alquier. PAC-Bayesian estimation and prediction in sparse additive models. *Electron. J. Statist.*, 7: 264–291, 2013.
- [27] Benjamin Guedj and Louis Pujol. Still no free lunches: the price to pay for tighter PAC-Bayes bounds. *arXiv preprint arXiv:1910.04460*, 2019.

# References III

- [28] Matthew Higgs and John Shawe-Taylor. A PAC-Bayes bound for tailored density estimation. In *Proceedings of the International Conference on Algorithmic Learning Theory (ALT)*, 2010.
- [29] Matthew J Holland. PAC-Bayes under potentially heavy tails. *arXiv:1905.07900*, 2019. To appear in NeurIPS.
- [30] A. Lacasse, F. Laviolette, M. Marchand, P. Germain, and N. Usunier. PAC-Bayes bounds for the risk of the majority vote and the variance of the Gibbs classifier. In *Advances in Neural information processing systems*, pages 769–776, 2007.
- [31] John Langford and Matthias Seeger. Bounds for averaging classifiers. Technical report, Carnegie Mellon, Departement of Computer Science, 2001.
- [32] John Langford and John Shawe-Taylor. PAC-Bayes & margins. In *Advances in Neural Information Processing Systems (NIPS)*, 2002.
- [33] Gaël Letarte, Pascal Germain, Benjamin Guedj, and François Laviolette. Dichotomize and Generalize: PAC-Bayesian Binary Activated Deep Neural Networks. *arXiv:1905.10259*, 2019. To appear at NeurIPS.
- [34] G. Lever, F. Laviolette, and J. Shawe-Taylor. Distribution-dependent PAC-Bayes priors. In *International Conference on Algorithmic Learning Theory*, pages 119–133. Springer, 2010.
- [35] C. Li, W. Jiang, and M. Tanner. General oracle inequalities for Gibbs posterior with application to ranking. In *Conference on Learning Theory*, pages 512–521, 2013.
- [36] Le Li, Benjamin Guedj, and Sébastien Loustau. A quasi-Bayesian perspective to online clustering. *Electron. J. Statist.*, 12(2): 3071–3113, 2018.
- [37] B. London. A PAC-Bayesian analysis of randomized learning with application to stochastic gradient descent. In *Advances in Neural Information Processing Systems*, pages 2931–2940, 2017.
- [38] B. London, B. Huang, B. Taskar, and L. Getoor. PAC-Bayesian collective stability. In *Artificial Intelligence and Statistics*, pages 585–594, 2014.
- [39] A. Maurer. A note on the PAC-Bayesian Theorem. *arXiv preprint cs/0411099*, 2004.
- [40] D. A. McAllester. PAC-Bayesian stochastic model selection. *Machine Learning*, 51(1):5–21, 2003.
- [41] David McAllester. Some PAC-Bayesian theorems. In *Proceedings of the International Conference on Computational Learning Theory (COLT)*, 1998.



# References IV

- [42] David McAllester. Some PAC-Bayesian theorems. *Machine Learning*, 37, 1999.
- [43] David McAllester. Some PAC-Bayesian theorems. *Machine Learning*, 37(3), 1999.
- [44] David McAllester. PAC-Bayesian stochastic model selection. *Machine Learning*, 51(1), 2003.
- [45] David McAllester. Simplified PAC-Bayesian margin bounds. In *COLT*, 2003.
- [46] Zakaria Mhammedi, Peter D. Grunwald, and Benjamin Guedj. PAC-Bayes Un-Expected Bernstein Inequality. *arXiv preprint arXiv:1905.13367*, 2019. Accepted at NeurIPS 2019.
- [47] B. Neyshabur, S. Bhojanapalli, D. A. McAllester, and N. Srebro. Exploring generalization in deep learning. In *Advances in Neural Information Processing Systems*, pages 5947–5956, 2017.
- [48] Kento Nozawa, Pascal Germain, and Benjamin Guedj. PAC-Bayesian contrastive unsupervised representation learning. *arXiv preprint arXiv:1910.04464*, 2019.
- [49] E. Parrado-Hernández, A. Ambroladze, J. Shawe-Taylor, and S. Sun. PAC-Bayes bounds with data dependent priors. *Journal of Machine Learning Research*, 13:3507–3531, 2012.
- [50] O. Rivasplata, E. Parrado-Hernandez, J. Shawe-Taylor, S. Sun, and C. Szepesvari. PAC-Bayes bounds for stable algorithms with instance-dependent priors. In *Advances in Neural Information Processing Systems*, pages 9214–9224, 2018.
- [51] M. Seeger. PAC-Bayesian generalization bounds for gaussian processes. *Journal of Machine Learning Research*, 3:233–269, 2002.
- [52] M. Seeger. *Bayesian Gaussian Process Models: PAC-Bayesian Generalisation Error Bounds and Sparse Approximations*. PhD thesis, University of Edinburgh, 2003.
- [53] Y. Seldin and N. Tishby. PAC-Bayesian analysis of co-clustering and beyond. *Journal of Machine Learning Research*, 11: 3595–3646, 2010.
- [54] Y. Seldin, F. Laviolette, N. Cesa-Bianchi, J. Shawe-Taylor, and P. Auer. PAC-Bayesian inequalities for martingales. *IEEE Transactions on Information Theory*, 58(12):7086–7093, 2012.
- [55] Yevgeny Seldin, Peter Auer, François Laviolette, John Shawe-Taylor, and Ronald Ortner. PAC-Bayesian analysis of contextual bandits. In *Advances in Neural Information Processing Systems (NIPS)*, 2011.

# References V

- [56] J. Shawe-Taylor and R. C. Williamson. A PAC analysis of a Bayes estimator. In *Proceedings of the 10th annual conference on Computational Learning Theory*, pages 2–9. ACM, 1997. doi: 10.1145/267460.267466.
- [57] John Shawe-Taylor and David Hardoon. Pac-bayes analysis of maximum entropy classification. In *Proceedings on the International Conference on Artificial Intelligence and Statistics (AISTATS)*, 2009.
- [58] Niklas Thiemann, Christian Igel, Olivier Wintenberger, and Yevgeny Seldin. A Strongly Quasiconvex PAC-Bayesian Bound. In *International Conference on Algorithmic Learning Theory, ALT*, pages 466–492, 2017.
- [59] L. G. Valiant. A theory of the learnable. *Communications of the ACM*, 27(11):1134–1142, 1984.
- [60] Wenda Zhou, Victor Veitch, Morgane Austern, Ryan P. Adams, and Peter Orbanz. Non-vacuous generalization bounds at the imagenet scale: a PAC-bayesian compression approach. In *ICLR*, 2019.