## Fitting Models to Data in Ecology and Evolution

#### Samraat Pawar

Department of Life Sciences (Silwood Park)

Imperial College London

October 25, 2019

What does "modelling data" mean to you?


 Mechanistic models aim to explain the PROCESSES or MECHANISMS underlying PATTERNS or PHENOMENA in empirical data

- Mechanistic models aim to explain the PROCESSES or MECHANISMS underlying PATTERNS or PHENOMENA in empirical data
  - These models have a THEORETICAL BASIS


- Mechanistic models aim to explain the PROCESSES or MECHANISMS underlying PATTERNS or PHENOMENA in empirical data
  - These models have a THEORETICAL BASIS
- Empirical/Phenomenological models establish the existence of STATISTICALLY SIGNIFICANT, NON-RANDOM PATTERNS or PHENOMENA in empirical data

- Mechanistic models aim to explain the PROCESSES or MECHANISMS underlying PATTERNS or PHENOMENA in empirical data
  - These models have a THEORETICAL BASIS
- Empirical/Phenomenological models establish the existence of STATISTICALLY SIGNIFICANT, NON-RANDOM PATTERNS or PHENOMENA in empirical data
  - They make no assumptions about the processes or mechanisms that generate the patterns

- Mechanistic models aim to explain the PROCESSES or MECHANISMS underlying PATTERNS or PHENOMENA in empirical data
  - These models have a THEORETICAL BASIS
- Empirical/Phenomenological models establish the existence of STATISTICALLY SIGNIFICANT, NON-RANDOM PATTERNS or PHENOMENA in empirical data
  - They make no assumptions about the processes or mechanisms that generate the patterns
  - That is, these models lack a THEORETICAL BASIS




source: https://www.cds.caltech.edu/~murray/amwiki/images/8/8f/LHgraph.gif



**SOURCE**: https://www.cds.caltech.edu/~murray/amwiki/images/8/8f/LHgraph.gif

• Mechanistic model: The Lynx-Hare Cycle is driven by density-dependent population growth in hares



source: https://www.cds.caltech.edu/~murray/amwiki/images/8/8f/LHgraph.gif

- Mechanistic model: The Lynx-Hare Cycle is driven by density-dependent population growth in hares
- Phenomenological model: The Lynx and Hare Cycles have a significant asynchrony (period shift) of x years

• It's not really one vs. the other, Both types of models play a role in science (and Biology)

- It's not really one vs. the other, Both types of models play a role in science (and Biology)
- Phenomenological model-fitting reveals patterns in data that generate HYPOTHESES

- It's not really one vs. the other, Both types of models play a role in science (and Biology)
- Phenomenological model-fitting reveals patterns in data that generate HYPOTHESES
  - These can be tested using further model fitting

- It's not really one vs. the other, Both types of models play a role in science (and Biology)
- Phenomenological model-fitting reveals patterns in data that generate HYPOTHESES
  - These can be tested using further model fitting
  - Example: Whether climatic temperature affects the Lynx-Hare cycle (using Generalized Linear Model-fitting)

- It's not really one vs. the other; Both types of models play a role in science (and Biology)
- Phenomenological model-fitting reveals patterns in data that generate HYPOTHESES
  - These can be tested using further model fitting
  - Example: Whether climatic temperature affects the Lynx-Hare cycle (using Generalized Linear Model-fitting)
- Mechanistic model-fitting tries to validate a mechanistic model that can explain the observed phenomenological pattern and generate MORE ACCURATE, MECHANISTIC HYPOTHESES

- It's not really one vs. the other; Both types of models play a role in science (and Biology)
- Phenomenological model-fitting reveals patterns in data that generate HYPOTHESES
  - These can be tested using further model fitting
  - Example: Whether climatic temperature affects the Lynx-Hare cycle (using Generalized Linear Model-fitting)
- Mechanistic model-fitting tries to validate a mechanistic model that can explain the observed phenomenological pattern and generate MORE ACCURATE, MECHANISTIC HYPOTHESES
  - Example: How climatic temperature drives the Lynx-Hare cycle

- It's not really one vs. the other, Both types of models play a role in science (and Biology)
- Phenomenological model-fitting reveals patterns in data that generate HYPOTHESES
  - These can be tested using further model fitting
  - Example: Whether climatic temperature affects the Lynx-Hare cycle (using Generalized Linear Model-fitting)
- Mechanistic model-fitting tries to validate a mechanistic model that can explain the observed phenomenological pattern and generate MORE ACCURATE, MECHANISTIC HYPOTHESES
  - Example: How climatic temperature drives the Lynx-Hare cycle
- Ultimately, successful, EMPIRICALLY-GROUNDED mechanistic models are the best path towards a THEORY in any scientific discipline (including ecology and evolution)

#### Individuals

Mechanisms ⇒
Metabolic rate,
Temperature response,
Growth rate



#### Interactions

Mechanisms ⇒
Consumer-resource
interactions, Competition,



#### Communities

Mechanisms ⇒

Trophic cascades, Bottom-up & Top-down regulation



6/19

 Do most ecological studies perform phenomenological or mechanistic modelling (or model-fitting)?

- Do most ecological studies perform phenomenological or mechanistic modelling (or model-fitting)?
- The answer is mostly Phenomenological

- Do most ecological studies perform phenomenological or mechanistic modelling (or model-fitting)?
- The answer is mostly Phenomenological

- Do most ecological studies perform phenomenological or mechanistic modelling (or model-fitting)?
- The answer is mostly Phenomenological Why?

- Do most ecological studies perform phenomenological or mechanistic modelling (or model-fitting)?
- The answer is mostly Phenomenological Why?
  - Partly because we are still establishing the existence of GENERAL PATTERNS/PHENOMENA,

- Do most ecological studies perform phenomenological or mechanistic modelling (or model-fitting)?
- The answer is mostly Phenomenological Why?
  - Partly because we are still establishing the existence of GENERAL PATTERNS/PHENOMENA,
  - ... and partly because we are (or are forced to be) interested in FORECASTING rather than EXPLAINING.

- Do most ecological studies perform phenomenological or mechanistic modelling (or model-fitting)?
- The answer is mostly Phenomenological Why?
  - Partly because we are still establishing the existence of GENERAL PATTERNS/PHENOMENA,
  - ... and partly because we are (or are forced to be) interested in FORECASTING rather than EXPLAINING.
- So the big question is, can we FORECAST WITHOUT FXPI AINING?

- Do most ecological studies perform phenomenological or mechanistic modelling (or model-fitting)?
- The answer is mostly Phenomenological Why?
  - Partly because we are still establishing the existence of GENERAL PATTERNS/PHENOMENA,
  - ... and partly because we are (or are forced to be) interested in FORECASTING rather than EXPLAINING.
- So the big question is, can we FORECAST WITHOUT EXPLAINING?
  - For example, disease outbreaks: Do we really need to care about the underlying mechanisms if we can predict a future event using phenomenological modelling (e.g., Machine-learning of time series patterns)?

Somewhat subjective!

- Somewhat subjective!
- For example, the Ricker model can be thought of as mechanistic:

$$N_{t+1} = N_t e^{r\left(1 - \frac{N_t}{k}\right)}$$

- Somewhat subjective!
- For example, the Ricker model can be thought of as mechanistic:

$$N_{t+1} = N_t e^{r\left(1 - \frac{N_t}{k}\right)}$$

• What is the mechanism?

- Somewhat subjective!
- For example, the Ricker model can be thought of as mechanistic:

$$N_{t+1} = N_t e^{r\left(1 - \frac{N_t}{k}\right)}$$

• What is the mechanism?

- Somewhat subjective!
- For example, the Ricker model can be thought of as mechanistic:

$$N_{t+1} = N_t e^{r\left(1 - \frac{N_t}{k}\right)}$$


- What is the mechanism? Density dependence through scramble competition (Brannstrom & Sumpter 2005)
- If the Ricker model and another model with contest competition were compared with data — some would call it mechanistic modelling because one is trying to get at the underlying mechanism, scramble or contest competition

- Somewhat subjective!
- For example, the Ricker model can be thought of as mechanistic:

$$N_{t+1} = N_t e^{r\left(1 - \frac{N_t}{k}\right)}$$


- What is the mechanism? Density dependence through scramble competition (Brannstrom & Sumpter 2005)
- If the Ricker model and another model with contest competition were compared with data — some would call it mechanistic modelling because one is trying to get at the underlying mechanism, scramble or contest competition
- But is this REALLY mechanistic? What are r and k really?

 Proponents of Ecological Metabolic Theory (AKA "Metabolic Theory of Ecology") argue that we have not progressed far enough towards mechanistic modelling because metabolism has been ignored




 The mechanistic basis of thermal performance curves (https://youtu.be/6n8fCuDwn74)

 The mechanistic basis of thermal performance curves (https://youtu.be/6n8fCuDwn74)



 The mechanistic basis of thermal performance curves (https://youtu.be/6n8fCuDwn74)



$$B = B_0 e^{-\frac{E}{kT}} f(T, T_{pk}, E_D)$$

T = temperature (K)

 $k = \text{Boltzmann constant (eV K}^{-1})$ 


E = Activation energy (eV)

 $T_{pk}$  = Temperature of peak performance  $E_D$  = Deactivation energy (eV)

(J H van't Hoff 1884, S Arrhenius 1889)

### EXAMPLE OF A FUNDAMENTAL MECHANISM: **METABOLIC RATE**

 The mechanistic basis of thermal performance curves (https://youtu.be/6n8fCuDwn74)



$$B = B_0 e^{-\frac{E}{kT}} f(T, T_{pk}, E_D)$$

T = temperature (K)

 $k = \text{Boltzmann constant (eV K}^{-1})$ 

E = Activation energy (eV)

 $T_{pk}$  = Temperature of peak performance  $\dot{E}_D$  = Deactivation energy (eV)

(J H van't Hoff 1884, S Arrhenius 1889)

- Surely there is more to thermal responses?
- What about alternative models?

• If possible, use biological knowledge to construct models

- If possible, use biological knowledge to construct models
- See if the models "agree well" with data

- If possible, use biological knowledge to construct models
- See if the models "agree well" with data
- Whichever model "agrees best" is most likely to have the right mechanisms

- If possible, use biological knowledge to construct models
- See if the models "agree well" with data
- Whichever model "agrees best" is most likely to have the right mechanisms
- That's the one that's best for predictions (e.g. population cycles), estimating rates (e.g. population or individual growth rates), etc

- If possible, use biological knowledge to construct models
- See if the models "agree well" with data
- Whichever model "agrees best" is most likely to have the right mechanisms
- That's the one that's best for predictions (e.g. population cycles), estimating rates (e.g. population or individual growth rates), etc
- Don't use models you already know have the wrong mechanisms just because they are popular!

- If possible, use biological knowledge to construct models
- See if the models "agree well" with data
- Whichever model "agrees best" is most likely to have the right mechanisms
- That's the one that's best for predictions (e.g. population cycles), estimating rates (e.g. population or individual growth rates), etc
- Don't use models you already know have the wrong mechanisms just because they are popular!
- Phenomenological/statistical models often perform better than mechanistic ones.

- If possible, use biological knowledge to construct models
- See if the models "agree well" with data
- Whichever model "agrees best" is most likely to have the right mechanisms
- That's the one that's best for predictions (e.g. population cycles), estimating rates (e.g. population or individual growth rates), etc
- Don't use models you already know have the wrong mechanisms just because they are popular!
- Phenomenological/statistical models often perform better than mechanistic ones.

- If possible, use biological knowledge to construct models
- See if the models "agree well" with data
- Whichever model "agrees best" is most likely to have the right mechanisms
- That's the one that's best for predictions (e.g. population cycles), estimating rates (e.g. population or individual growth rates), etc
- Don't use models you already know have the wrong mechanisms just because they are popular!
- Phenomenological/statistical models often perform better than mechanistic ones. Why?

- If possible, use biological knowledge to construct models
- See if the models "agree well" with data
- Whichever model "agrees best" is most likely to have the right mechanisms
- That's the one that's best for predictions (e.g. population cycles), estimating rates (e.g. population or individual growth rates), etc
- Don't use models you already know have the wrong mechanisms just because they are popular!
- Phenomenological/statistical models often perform better than mechanistic ones. Why? — because they have lest restrictive assumptions

#### MODELS: HOW TO BUILD THEM?

- It's an art, takes practice (Levins' paper on the strategy of model building in biology)
- Build models one mechanism at a time in biology, it means start at the right level of organization!
- Always consider an alternative that is more parsimonious, even if it is phenomenological!
- For example, the Boltzmann-Arrhenius model is a good first try describe and uncover mechanisms underlying individual level "traits" that are rates (e.g., fecundity or development rate)
- The next step would be to include species interactions with temperature dependence of individuals (or go in an evolutionary direction)

#### FITTING MODELS TO DATA

#### Multiple ways to do it:

- Least Squares methods
  - Linear
  - Non-linear
- Likelihood-based methods
  - Maximum Likelihood Estimation (MLE)
  - Bayesian
- Artificial intelligence and Machine learning
  - Focus in on maximizing ability to discover pattern and predict at the cost of mechanistic insights

#### **METHODS YOU CAN USE**

- Least squares: along with Linear Model fitting, Non-linear Least Squares (NLLS) fitting is a particularly versatile and powerful approach because many mechanisms in biology and inherently non-linear
- MLE/Bayesian methods: more robust if you are able to calculate the likelihood function analytically or numerically.
- Al/machine Learning: most versatile for large amounts of noisy data

 Ideally, several competing (meaningful, not just null) hypotheses (mathematical models) should be fitted to data and compared using statistical theory

15/19

- Ideally, several competing (meaningful, not just null) hypotheses (mathematical models) should be fitted to data and compared using statistical theory
- This is an advance over the traditional "null hypothesis" approach in Biology

- Ideally, several competing (meaningful, not just null) hypotheses (mathematical models) should be fitted to data and compared using statistical theory
- This is an advance over the traditional "null hypothesis" approach in Biology
- Necessary for developing the advancement of Biology from from an observational and axiomatic discipline to one with general theories.

- Ideally, several competing (meaningful, not just null) hypotheses (mathematical models) should be fitted to data and compared using statistical theory
- This is an advance over the traditional "null hypothesis" approach in Biology
- Necessary for developing the advancement of Biology from from an observational and axiomatic discipline to one with general theories.
- Necessary for understanding the mechanisms underlying biological patterns/phenomena

 It's all about the "Likelihood" of a model: the set of parameter values of the model (θ) given outcomes (x), equals the probability of those observed outcomes given those parameter values, that is,

$$\mathcal{L}(\theta|\mathbf{x}) = P(\mathbf{x}|\theta)$$

- The easiest thing to do for you is to use information theory (including AIC and BIC) to compare models.
- Both AIC and BIC use the *estimated likelihoods of a model*: AIC:  $-2 \ln[\mathcal{L}(\theta|x)] + 2p$ Small sample AIC (AICc):  $-2 \ln[\mathcal{L}(\theta|x)] + 2p$ BIC (Schwartz criterion):  $-2 \ln[\mathcal{L}(\theta|x)] + p \ln(n)$ (where n = sample size, p number of free parameters)
- The lower the AIC or BIC, the better.

This is how you calculate AIC and BIC (using python syntax):

This is how you calculate AIC and BIC (using python syntax):

- residuals = Observations Predictions
- rss = sum(residuals \*\* 2)
- Then, AIC is n \* log((2 \* pi) / n) + n + 2 + n \* log(rss) + 2 \* p (note n and p!)
- And BIC is n + n \* log(2 \* pi) + n \* log(rss / n) + (log(n)) \* (p + 1)
- That is,  $\mathcal{L}(\theta|x) = -\frac{n}{2/\ln(RSS/n)}$
- For both AIC and BIC, If model A has AIC lower by 2-3 or more than model B, it's better — Differences of less than 2-3 don't really matter

This is how you calculate AIC and BIC (using python syntax):

- residuals = Observations Predictions
- rss = sum(residuals \*\* 2)
- Then, AIC is n \* log((2 \* pi) / n) + n + 2 + n \* log(rss) + 2 \* p (note n and p!)
- And BIC is n + n \* log(2 \* pi) + n \* log(rss / n) + (log(n)) \* (p + 1)
- That is,  $\mathcal{L}(\theta|x) = -\frac{n}{2/\ln(RSS/n)}$
- For both AIC and BIC, If model A has AIC lower by 2-3 or more than model B, it's better — Differences of less than 2-3 don't really matter

#### Also note that:

 R<sup>2</sup> = 1 - (rss/tss), where tss is total sum of squares: tss = sum((Observations - mean(Predictions)) \*\* 2) (a useful measure of goodness of fit)

### **COMPARING AND SELECTING MODELS: MORE STUFF**

- You can also calculate Akaike Weights, which is very useful/important when comparing > 2 models. These weights can then be used to perform model averaging.
- Model selection using the Likelihood-Ratio test (LRT) is another option when you are comparing 2 models.
- Adjusted R<sup>2</sup> can be used to get a rigorous "idea" about how alternative models are performing.
- Very often, you will end up doing model simplification, especially in for linear least squares model fitting — starting with a complex model and then dropping terms till you have found a the most parsimonious version of the original model. There are functions in R to do this (of course!).

#### READINGS

- Levins, R. (1966) The strategy of model building in population biology. Am. Sci. 54, 421–431.
- Johnson, J. B. & Omland, K. S. (2004) Model selection in ecology and evolution. Trends Ecol. Evol. 19, 101–108.
- Bolker, B. M. et al. (2013) Strategies for fitting nonlinear ecological models in R, AD Model Builder, and BUGS. Methods Ecol. Evol. 4, 501–512.
- Additional readings on the TheMulQuaBio git repository