

Some Prolog Practice Questions

Define the following predicates in Prolog using any auxiliary predicates you wish.

1. `subList(L1, L2)` to mean every element in list L1 is also in list L2.

You can assume both arguments are grounded in the call.

E.g.

`subList([1,2,3], [1,1,3,2,3,4])` and `subList([1,1,4,3], [5,1,3,2,3,4])` should both succeed.

2. `difference(L1, L2, L)` to mean L consists of all the elements in L1 that are not in L2.

You can assume both arguments L1 and L2 are grounded in the call.

E.g. `difference([1,1,2,3, 5, 5], [1,3,2, 3,4], L)` should give `L=[5, 5]`. The repetition in the list L is fine and you may get the same answer more than once. That too is fine.

`difference([a, c], [], L)` should give `L=[a,c]`.

3. `sift(L, N, Result)` to mean Result is list L but with all occurrences of elements greater than N removed. You can assume both arguments L and N are grounded in the call.

E.g. `sift([1,4,3,6,8], 3, X)` should give `X=[1,3]`.

4) `common(L1, L2, I)`

to mean I is the list of the common elements of lists L1 and L2.

You can assume both arguments L1 and L2 are grounded in the call. The resulting list I should have no repeated elements. The order of the elements in list I is not important. If L1 and L2 have no common elements then the output I should be the empty list [].

E.g.

`common([1,1,4,2,5], [1,1,7,2,3,4,4,8], I)` should give the answer `I=[1,2,4]`, but the order of the elements in I does not matter.

`common([1,2], [4,8], I)` should give the answer `I=[]`.

5. `delete(L, Result)`

Result is list L with every other element deleted.

Example:

?- delete([1,2,3,4], R).

R=[1,3]

6. process(L1, L2, Consistent, Inconsistent)

where **L1** is a given list of items of the form (Name, Number), and **L2** is a given list of items of the form (Name, Number, MoreInfo). Then the output **Consistent** should be those items (Name, Number, MoreInfo) in **L2** that agree on (Name, Number) with list **L1**, and **Inconsistent** should be whatever is left over from list **L2**.

E.g. Suppose **L1** has (Name , Age) items and **L2** has (Name, Age, Marital_status) items.

Then **Consistent** should be those items (Name, Age, Marital_status) where for the same Name **L1** provides the same Age.

E.g.

process([(mary, 20), (john, 30), (pete, 40)], [(mary, 20, single), (pete, 40, single), (joe, 35, widowed), (john, 35, married)], C, I)

should give the answer

C= [(mary, 20, single), (pete, 40, single)]

I= [(john, 35, married), (joe, 35, widowed)].

The order of the elements in **C** and **I** is not important.

7. Duplicate the elements of a list.

Example:

?- dupli([a,b,c,c,d],X).

X = [a,a,b,b,c,c,c,c,d,d]

8. Duplicate the elements of a list a given number of times.

Example:

?- dupli([a,b,c],3,X).

X = [a,a,a,b,b,b,c,c,c]

9. Drop every N'th element from a list.

Example:

?- drop([a,b,c,d,e,f,g,h,i,k],3,X).

X = [a,b,d,e,g,h,k]

10. Split a list into two parts; the length of the first part is given.

Example:

?- split([a,b,c,d,e,f,g,h,i,k],3,L1,L2).

$L1 = [a,b,c]$
 $L2 = [d,e,f,g,h,i,k]$

The Bratko book has many exercises.