Commonly in dogs, cats and goats

Predisposing causes:
- Pendulous ear, excessive hair in the ear canal
- Inadequate air circulation & excess moisture

Commonly organisms: Pseudomonas spp.
Staphylococcus spp.
Proteus spp.
Malassezia pachydermatis (syn. Pityrosporum canis) Ear mites
- Otodectes cynotis
- Psoroptes cuniculi- rabbit, horse, goat, deer
- Raillietia auris- horse
- Otobius megnini – ruminant, pigs, horses dogs

Otitis externa

Gross findings

– Hyperemia, accumulation of serum, cerumen, leukocytes & epithelial debris
- Pseudomonas: Yellow- red (hemorrhage) pus,
ulcers & foal smell
- Malassezia: Greasy dull dark brown, sour smell
- Mites: A lot of dry dark crumbly accumulation

Histopathology

- Commonly chronic lesion
- Epidermal hyperplasia with acanthosis
- Lymphocytes, plasma cells, neutrophils, mast cells
in dermis
- Dermal fibrosis
- Sebaceous & ceruminous gland hyperplasia

Otitis externa

- Commonly in feeder pigs
- Cause: bacteria; Hemolytic streptococci & Pasteurella hemolytica
- Dog & cat; chronic otitis externa

- Continuing spread from the middle ears
- Commonly focal suppurative meningitis/encephalitis at pons

Otitis media

Otitis interna