

MAC 0460 / 5832

Introduction to Machine Learning

18 — Support Vector Machines (SVM) – Part II

- hyperplane • margin • margin violation •
- QP problems • dual problems • kernel trick •

IME/USP (21/06/2021)

- Binary linear classification: The linearly separable case
- hard-margin SVM: Maximum margin formulation
- Binary linear classification: The non-linearly separable case
- soft-margin SVM: allows margin violation
- hard-margin/soft-margin SVM is a QP problem
- Dual of hard-margin/soft-margin SVM is also QP
- How to solve QP problems
- Non-linear classification: the kernel trick

Dual QP optimization

Recall primal and dual formulation

When we discussed regularization, we started with the following optimization problem

$$\begin{aligned} & \underset{\mathbf{w}}{\text{minimize}} && E_{in}(\mathbf{w}) \\ & \text{subject to:} && \mathbf{w}^T \mathbf{w} \leq C \end{aligned}$$

and we ended up solving the following problem

$$\underset{\mathbf{w}}{\text{minimize}} \quad E_{aug}(\mathbf{w}) = E_{in}(\mathbf{w}) + \lambda \mathbf{w}^T \mathbf{w}$$

That is, we started with a problem with constraints on \mathbf{w} and ended up with a problem without such constraints (only one constraint: $\lambda \geq 0$)

Introduction to the dual formulation, first with one constraint

QP with one constraint:

$$\begin{array}{ll}\underset{\mathbf{u}}{\text{minimize}} & \frac{1}{2}\mathbf{u}^T Q \mathbf{u} + \mathbf{p}^T \mathbf{u} \\ \text{subject to:} & \mathbf{a}^T \mathbf{u} \geq c \quad (\text{one constraint})\end{array}$$

Fact: If there is an optimal solution \mathbf{u}^* for the above problem, then \mathbf{u}^* is also an optimal solution of the following problem:

$$\underset{\mathbf{u}}{\text{minimize}} \quad \frac{1}{2}\mathbf{u}^T Q \mathbf{u} + \mathbf{p}^T \mathbf{u} + \max_{\alpha \geq 0} \alpha(c - \mathbf{a}^T \mathbf{u})$$

Why? Since $\mathbf{a}^T \mathbf{u} \geq c \iff c - \mathbf{a}^T \mathbf{u} \leq 0$, then $\max_{\alpha \geq 0} \alpha(c - \mathbf{a}^T \mathbf{u}) = 0$

Introduction to the dual formulation, first with one constraint

Dual formulation

$$\underset{\mathbf{u}}{\text{minimize}} \quad \frac{1}{2} \mathbf{u}^T Q \mathbf{u} + \mathbf{p}^T \mathbf{u} + \max_{\alpha \geq 0} \alpha (c - \mathbf{a}^T \mathbf{u})$$

The term $\alpha(c - \mathbf{a}^T \mathbf{u})$ forces $c - \mathbf{a}^T \mathbf{u}$ to stay negative – i.e., to satisfy the constraint $\mathbf{a}^T \mathbf{u} \geq c$ (because this helps to minimize the cost function). On the other hand, α is chosen so as to maximize $\alpha(c - \mathbf{a}^T \mathbf{u})$ (to avoid $c - \mathbf{a}^T \mathbf{u}$ going to $-\infty$)

There is no constraints

We have a min-max optimization problem

Lagrangian

$$\mathcal{L}(\mathbf{u}, \alpha) = \frac{1}{2} \mathbf{u}^T Q \mathbf{u} + \mathbf{p}^T \mathbf{u} + \alpha(c - \mathbf{a}^T \mathbf{u})$$

Min-max optimization

$$\min_{\mathbf{u}} \max_{\alpha \geq 0} \mathcal{L}(\mathbf{u}, \alpha)$$

Strong duality

$$\min_{\mathbf{u}} \max_{\alpha \geq 0} \mathcal{L}(\mathbf{u}, \alpha) = \max_{\alpha \geq 0} \min_{\mathbf{u}} \mathcal{L}(\mathbf{u}, \alpha)$$

Theorem 8.7 (KKT). For a feasible convex QP-problem in *primal* form,

$$\begin{array}{ll} \underset{\mathbf{u} \in \mathbb{R}^L}{\text{minimize:}} & \frac{1}{2} \mathbf{u}^T \mathbf{Q} \mathbf{u} + \mathbf{p}^T \mathbf{u} \\ \text{subject to:} & \mathbf{a}_m^T \mathbf{u} \geq c_m \quad (m = 1, \dots, M), \end{array}$$

define the Lagrange function

$$\mathcal{L}(\mathbf{u}, \boldsymbol{\alpha}) = \frac{1}{2} \mathbf{u}^T \mathbf{Q} \mathbf{u} + \mathbf{p}^T \mathbf{u} + \sum_{m=1}^M \alpha_m (c_m - \mathbf{a}_m^T \mathbf{u}).$$

The solution \mathbf{u}^* is optimal for the primal if and only if $(\mathbf{u}^*, \boldsymbol{\alpha}^*)$ is a solution to the dual optimization problem

$$\max_{\boldsymbol{\alpha} \geq \mathbf{0}} \min_{\mathbf{u}} \mathcal{L}(\mathbf{u}, \boldsymbol{\alpha}).$$

KKT conditions

The optimal $(\mathbf{u}^*, \boldsymbol{\alpha}^*)$ satisfies the Karush-Kuhn-Tucker (KKT) conditions:

(i) *Primal and dual constraints:*

$$\mathbf{a}_m^T \mathbf{u}^* \geq c_m \quad \text{and} \quad \alpha_m \geq 0 \quad (m = 1, \dots, M).$$

(ii) *Complementary slackness:*

$$\alpha_m^* (\mathbf{a}_m^T \mathbf{u}^* - c_m) = 0.$$

(iii) *Stationarity with respect to \mathbf{u} :*

$$\nabla_{\mathbf{u}} \mathcal{L}(\mathbf{u}, \boldsymbol{\alpha})|_{\mathbf{u}=\mathbf{u}^*, \boldsymbol{\alpha}=\boldsymbol{\alpha}^*} = \mathbf{0}.$$

Solving the dual QP optimization problems

Dual: characterized by the Lagrangean \mathcal{L}

It is a min-max problem

$$\min_{\mathbf{u}} \max_{\alpha \geq 0} \mathcal{L}(\mathbf{u}, \alpha) = \max_{\alpha \geq 0} \min_{\mathbf{u}} \mathcal{L}(\mathbf{u}, \alpha)$$

Two-step optimization:

1. We fix α and optimize $\min_{\mathbf{u}} \mathcal{L}(\mathbf{u}, \alpha)$
2. Then, we fix \mathbf{u} and optimize $\max_{\alpha} \mathcal{L}(\mathbf{u}, \alpha)$

Dual formulation for the hard-margin SVM

Primal formulation:

$$\begin{aligned} & \underset{\mathbf{w}, b}{\text{minimize}} && \frac{1}{2} \mathbf{w}^T \mathbf{w} \\ & \text{subject to} && y_n (\mathbf{w}^T \mathbf{x}_n + b) \geq 1, n = 1, \dots, N \end{aligned}$$

Lagrangian function

$$\begin{aligned} \mathcal{L}(b, \mathbf{w}, \boldsymbol{\alpha}) &= \frac{1}{2} \mathbf{w}^T \mathbf{w} + \sum_{n=1}^N \alpha_n (1 - y_n (\mathbf{w}^T \mathbf{x}_n + b)) \\ &= \frac{1}{2} \mathbf{w}^T \mathbf{w} - \sum_{n=1}^N \alpha_n y_n \mathbf{w}^T \mathbf{x}_n - b \sum_{n=1}^N \alpha_n y_n + \sum_{n=1}^N \alpha_n. \end{aligned}$$

Solution of the dual hard-margin SVM

$$\begin{aligned}\mathcal{L}(b, \mathbf{w}, \boldsymbol{\alpha}) &= \frac{1}{2} \mathbf{w}^T \mathbf{w} + \sum_{n=1}^N \alpha_n (1 - y_n (\mathbf{w}^T \mathbf{x}_n + b)) \\ &= \frac{1}{2} \mathbf{w}^T \mathbf{w} - \sum_{n=1}^N \alpha_n y_n \mathbf{w}^T \mathbf{x}_n - b \sum_{n=1}^N \alpha_n y_n + \sum_{n=1}^N \alpha_n.\end{aligned}$$

1. Minimize $\mathcal{L}(b, \mathbf{w}, \boldsymbol{\alpha})$ with respect to (b, \mathbf{w}) :

$$\frac{\partial \mathcal{L}}{\partial b} = -\sum_{n=1}^N \alpha_n y_n \quad \text{and} \quad \frac{\partial \mathcal{L}}{\partial \mathbf{w}} = \mathbf{w} - \sum_{n=1}^N \alpha_n y_n \mathbf{x}_n.$$

Computing the zero:

$$\sum_{n=1}^N \alpha_n y_n = 0; \quad \mathbf{w} = \sum_{n=1}^N \alpha_n y_n \mathbf{x}_n.$$

Solution of the dual hard-margin SVM

$$\begin{aligned}\mathcal{L}(b, \mathbf{w}, \boldsymbol{\alpha}) &= \frac{1}{2} \mathbf{w}^T \mathbf{w} + \sum_{n=1}^N \alpha_n (1 - y_n (\mathbf{w}^T \mathbf{x}_n + b)) \\ &= \frac{1}{2} \mathbf{w}^T \mathbf{w} - \sum_{n=1}^N \alpha_n y_n \mathbf{w}^T \mathbf{x}_n - b \sum_{n=1}^N \alpha_n y_n + \sum_{n=1}^N \alpha_n.\end{aligned}$$

$$\sum_{n=1}^N \alpha_n y_n = 0; \quad \mathbf{w} = \sum_{n=1}^N \alpha_n y_n \mathbf{x}_n.$$

The Lagrangean is reduced to a function on $\boldsymbol{\alpha}$ only:

$$\mathcal{L}(\boldsymbol{\alpha}) = -\frac{1}{2} \sum_{n=1}^N \sum_{m=1}^N y_n y_m \alpha_n \alpha_m \mathbf{x}_n^T \mathbf{x}_m + \sum_{n=1}^N \alpha_n.$$

Solution of the dual hard-margin SVM

The Lagrangean is reduced to a function on α only:

$$\mathcal{L}(\alpha) = -\frac{1}{2} \sum_{n=1}^N \sum_{m=1}^N y_n y_m \alpha_n \alpha_m \mathbf{x}_n^T \mathbf{x}_m + \sum_{n=1}^N \alpha_n.$$

2. Now we would like to maximize \mathcal{L} : Minimize $-\mathcal{L}$ with respect to α

$$\begin{aligned} \underset{\alpha \in \mathbb{R}^N}{\text{minimize:}} \quad & \frac{1}{2} \sum_{m=1}^N \sum_{n=1}^N y_n y_m \alpha_n \alpha_m \mathbf{x}_n^T \mathbf{x}_m - \sum_{n=1}^N \alpha_n \\ \text{subject to:} \quad & \sum_{n=1}^N y_n \alpha_n = 0 \\ & \alpha_n \geq 0 \quad (n = 1, \dots, N). \end{aligned}$$

This is a standard QP and we can solve it using Solvers!

Recall: Standard form of QP problems

M inequality constraints and Q positive semi-definite

$$\begin{aligned} & \underset{\mathbf{u}}{\text{minimize}} && \frac{1}{2} \mathbf{u}^T Q \mathbf{u} + \mathbf{p}^T \mathbf{u} \\ & \text{subject to:} && \mathbf{a}_m^T \mathbf{u} \geq c_m \quad (m = 1, \dots, M) \end{aligned}$$

In matrix form

$$\begin{aligned} & \underset{\mathbf{u}}{\text{minimize}} && \frac{1}{2} \mathbf{u}^T Q \mathbf{u} + \mathbf{p}^T \mathbf{u} \\ & \text{subject to:} && A \mathbf{u} \geq \mathbf{c} \end{aligned}$$

QP solvers can be used to compute the optimal solution \mathbf{u}^* :

$$\mathbf{u}^* \leftarrow \text{QP}(Q, \mathbf{p}, A, \mathbf{c})$$

Solution of the dual hard-margin SVM

Minimization of $-\mathcal{L}$ with respect to α is a standard QP:

$$\begin{aligned} \underset{\alpha \in \mathbb{R}^N}{\text{minimize:}} \quad & \frac{1}{2} \alpha^T Q_D \alpha - \mathbf{1}_N^T \alpha \\ \text{subject to:} \quad & A_D \alpha \geq \mathbf{0}_{N+2}, \end{aligned} \quad (8.)$$

where Q_D and A_D (D for dual) are given by:

$$Q_D = \begin{bmatrix} y_1 y_1 \mathbf{x}_1^T \mathbf{x}_1 & \dots & y_1 y_N \mathbf{x}_1^T \mathbf{x}_N \\ y_2 y_1 \mathbf{x}_2^T \mathbf{x}_1 & \dots & y_2 y_N \mathbf{x}_2^T \mathbf{x}_N \\ \vdots & \vdots & \vdots \\ y_N y_1 \mathbf{x}_N^T \mathbf{x}_1 & \dots & y_N y_N \mathbf{x}_N^T \mathbf{x}_N \end{bmatrix} \quad \text{and} \quad A_D = \begin{bmatrix} \mathbf{y}^T \\ -\mathbf{y}^T \\ \mathbf{I}_{N \times N} \end{bmatrix}$$

$$\alpha^* \leftarrow QP(Q_D, -\mathbf{1}, A_D, \mathbf{0})$$

Computing (b^*, \mathbf{w}^*) from α^*

From the stationarity condition, we derived $\mathbf{w} = \sum_{n=1}^N \alpha_n y_n \mathbf{x}_n$.

$$\Rightarrow \text{Thus, } \mathbf{w}^* = \sum_{n=1}^N y_n \alpha_n^* \mathbf{x}_n$$

At least one α_s^* is strictly positive (suposing we have examples from both classes). From the complementary slackness, we know that

$$\alpha_m^* (\mathbf{a}_m^T \mathbf{u}^* - c_m) = 0.$$

\Rightarrow Thus equality holds for sample s :

$$y_s (\mathbf{w}^{*T} \mathbf{x}_s + b^*) = 1 \Rightarrow b^* = y_s - \sum_{n=1}^N y_n \alpha_n^* \mathbf{x}_n^T \mathbf{x}_s$$

Hard-Margin SVM with Dual QP

1: Construct Q_D and A_D as in Exercise 8.11

$$Q_D = \begin{bmatrix} y_1 y_1 \mathbf{x}_1^T \mathbf{x}_1 & \dots & y_1 y_N \mathbf{x}_1^T \mathbf{x}_N \\ y_2 y_1 \mathbf{x}_2^T \mathbf{x}_1 & \dots & y_2 y_N \mathbf{x}_2^T \mathbf{x}_N \\ \vdots & \vdots & \vdots \\ y_N y_1 \mathbf{x}_N^T \mathbf{x}_1 & \dots & y_N y_N \mathbf{x}_N^T \mathbf{x}_N \end{bmatrix} \quad \text{and} \quad A_D = \begin{bmatrix} \mathbf{y}^T \\ -\mathbf{y}^T \\ \mathbf{I}_{N \times N} \end{bmatrix}.$$

2: Use a QP-solver to optimize the dual problem:

$$\boldsymbol{\alpha}^* \leftarrow \text{QP}(Q_D, -\mathbf{1}_N, A_D, \mathbf{0}_{N+2}).$$

3: Let s be a support vector for which $\alpha_s^* > 0$. Compute b^* ,

$$b^* = y_s - \sum_{\alpha_n^* > 0} y_n \alpha_n^* \mathbf{x}_n^T \mathbf{x}_s.$$

4: Return the final hypothesis

$$g(\mathbf{x}) = \text{sign} \left(\sum_{\alpha_n^* > 0} y_n \alpha_n^* \mathbf{x}_n^T \mathbf{x} + b^* \right).$$

There are software packages for solving optimization problems

Python Notebook `svm_cvxpy.ipynb`

- uses CVXPY (<https://www.cvxpy.org/>)
- presents solutions for the toy example
 - using a solver for standard optimization
 - using a solver for the standard QP formulation
(Linear Hard-Margin SVM with QP)
 - using a solver for the dual QP formulation
(Hard Margin SVM with Dual QP)


Interpretation of the solution

Support vectors: $\alpha_s > 0 \implies y_s(\mathbf{w}^{*T} \mathbf{x}_s + b^*) = 1$


Weights: $\mathbf{w}^* = \sum_{n=1}^N y_n \alpha_n^* \mathbf{x}_n$

Bias: $b^* = y_s - \mathbf{w}^{*T} \mathbf{x}_s = y_s - \sum_{n=1}^N y_n \alpha_n^* \mathbf{x}_n^T \mathbf{x}_s$

Hypothesis: $g(\mathbf{x}) = \text{sign}\left(\sum_{\alpha_n^* > 0} y_n \alpha_n^* \mathbf{x}_n^T \mathbf{x} + b^*\right)$



(a) All data



(b) Only support vectors

Solution of the soft-margin SVM

The soft-margin SVM is also a QP

Only with more constraints

Thus the same discussion on QP and dual QP holds

The new optimization

$$\text{Minimize} \quad \frac{1}{2} \mathbf{w}^\top \mathbf{w} + C \sum_{n=1}^N \xi_n$$

$$\text{subject to} \quad y_n (\mathbf{w}^\top \mathbf{x}_n + b) \geq 1 - \xi_n \quad \text{for } n = 1, \dots, N$$

$$\text{and} \quad \xi_n \geq 0 \quad \text{for } n = 1, \dots, N$$

$$\mathbf{w} \in \mathbb{R}^d, \quad b \in \mathbb{R}, \quad \boldsymbol{\xi} \in \mathbb{R}^N$$

Lagrange formulation

$$\mathcal{L}(\mathbf{w}, b, \xi, \alpha, \beta) = \frac{1}{2} \mathbf{w}^\top \mathbf{w} + C \sum_{n=1}^N \xi_n - \sum_{n=1}^N \alpha_n (y_n (\mathbf{w}^\top \mathbf{x}_n + b) - 1 + \xi_n) - \sum_{n=1}^N \beta_n \xi_n$$

Minimize w.r.t. \mathbf{w} , b , and ξ and maximize w.r.t. each $\alpha_n \geq 0$ and $\beta_n \geq 0$

$$\nabla_{\mathbf{w}} \mathcal{L} = \mathbf{w} - \sum_{n=1}^N \alpha_n y_n \mathbf{x}_n = \mathbf{0}$$


$$\frac{\partial \mathcal{L}}{\partial b} = - \sum_{n=1}^N \alpha_n y_n = 0$$

$$\frac{\partial \mathcal{L}}{\partial \xi_n} = C - \alpha_n - \beta_n = 0$$


Hard × soft margin

Optimization of \mathcal{L} with respect to α :

Hard-margin


$$\begin{aligned} & \underset{\alpha \in \mathbb{R}^N}{\text{minimize:}} && \frac{1}{2} \sum_{m=1}^N \sum_{n=1}^N y_n y_m \alpha_n \alpha_m \mathbf{x}_n^T \mathbf{x}_m - \sum_{n=1}^N \alpha_n \\ & \text{subject to:} && \sum_{n=1}^N y_n \alpha_n = 0 \\ & && \alpha_n \geq 0 \quad (n = 1, \dots, N). \end{aligned}$$

Soft-margin (it is also a QP problem)



$$\begin{aligned} & \min_{\alpha} && \frac{1}{2} \alpha^T Q_D \alpha - \mathbf{1}^T \alpha \\ & \text{subject to} && \mathbf{y}^T \alpha = 0; . \\ & && \mathbf{0} \leq \alpha \leq C \cdot \mathbf{1}. \end{aligned}$$

Interpretation of C

$$\text{minimize } \frac{1}{2} \mathbf{w}^T \mathbf{w} + C \sum_{i=1}^N \xi_i$$

$$0 \leq \alpha_n \leq C$$

Interpretation of C



$0 < \alpha_n^* < C \implies \mathbf{x}_n$ is a support vector

$\alpha_n^* = 0 \implies \mathbf{x}_n$ is beyond the margin on the right side

$\alpha_n^* = C \implies \mathbf{x}_n$ is inside the margin or in the wrong side

When the data is linearly separable, there exists C such that the soft-margin SVM solution is exactly the same solution of the hard-margin SVM


Types of support vectors

margin support vectors ($0 < \alpha_n < C$)

$$y_n(\mathbf{w}^\top \mathbf{x}_n + b) = 1 \quad (\xi_n = 0)$$

non-margin support vectors ($\alpha_n = C$)

$$y_n(\mathbf{w}^\top \mathbf{x}_n + b) < 1 \quad (\xi_n > 0)$$



The Kernel trick


Motivation

Soft-margin SVM could be used to solve non-linear cases


Would we get good solutions for both examples below ?

Two types of non-separable

slightly:




seriously:




\mathbf{z} instead of \mathbf{x}

$$\mathcal{L}(\alpha) = \sum_{n=1}^N \alpha_n - \frac{1}{2} \sum_{n=1}^N \sum_{m=1}^N y_n y_m \alpha_n \alpha_m \mathbf{z}_n^\top \mathbf{z}_m$$



$\mathcal{X} \rightarrow \mathcal{Z}$



What is the problem?

When we map data $\mathbf{x} \in \mathbb{R}^d$ to $\mathbf{z} \in \mathbb{R}^{\tilde{d}}$, $\tilde{d} \gg d$

huge \tilde{d}

What do we need from the \mathcal{Z} space?

$$\mathcal{L}(\boldsymbol{\alpha}) = \sum_{n=1}^N \alpha_n - \frac{1}{2} \sum_{n=1}^N \sum_{m=1}^N y_n y_m \alpha_n \alpha_m \mathbf{z}_n^\top \mathbf{z}_m$$

Constraints: $\alpha_n \geq 0$ for $n = 1, \dots, N$ and $\sum_{n=1}^N \alpha_n y_n = 0$

$$g(\mathbf{x}) = \text{sign}(\mathbf{w}^\top \mathbf{z} + b)$$

need $\mathbf{z}_n^\top \mathbf{z}$

where $\mathbf{w} = \sum_{\mathbf{z}_n \text{ is SV}} \alpha_n y_n \mathbf{z}_n$

and $b: y_m (\mathbf{w}^\top \mathbf{z}_m + b) = 1$ need $\mathbf{z}_n^\top \mathbf{z}_m$

Is there any kernel function $K()$ satisfying

$$K_{\Phi}(\mathbf{x}, \mathbf{x}') = \Phi(\mathbf{x})^T \Phi(\mathbf{x}')$$

and such that computation is more efficient than computing $\mathbf{z}^T \mathbf{z}' = \Phi(\mathbf{x})^T \Phi(\mathbf{x}')$?

If there is such $K()$, then

$$\mathbf{Q}_D = \begin{bmatrix} y_1 y_1 \mathbf{x}_1^T \mathbf{x}_1 & \dots & y_1 y_N \mathbf{x}_1^T \mathbf{x}_N \\ y_2 y_1 \mathbf{x}_2^T \mathbf{x}_1 & \dots & y_2 y_N \mathbf{x}_2^T \mathbf{x}_N \\ \vdots & \vdots & \vdots \\ y_N y_1 \mathbf{x}_N^T \mathbf{x}_1 & \dots & y_N y_N \mathbf{x}_N^T \mathbf{x}_N \end{bmatrix} \quad \mathbf{Q}_D = \begin{bmatrix} y_1 y_1 K_{11} & \dots & y_1 y_N K_{1N} \\ y_2 y_1 K_{21} & \dots & y_2 y_N K_{2N} \\ \vdots & \vdots & \vdots \\ y_N y_1 K_{N1} & \dots & y_N y_N K_{NN} \end{bmatrix}$$

Kernel K would be equivalent to mapping \mathbf{x} to \mathbf{z} and applying dual SVM on \mathbf{z} , but without explicitly computing \mathbf{z} !

Hard-Margin SVM with Kernel

- 1: Construct Q_D from the kernel K , and A_D :

$$Q_D = \begin{bmatrix} y_1 y_1 K_{11} & \dots & y_1 y_N K_{1N} \\ y_2 y_1 K_{21} & \dots & y_2 y_N K_{2N} \\ \vdots & \vdots & \vdots \\ y_N y_1 K_{N1} & \dots & y_N y_N K_{NN} \end{bmatrix} \quad \text{and} \quad A_D = \begin{bmatrix} \mathbf{y}^T \\ -\mathbf{y}^T \\ \mathbf{I}_{N \times N} \end{bmatrix},$$

where $K_{mn} = K(\mathbf{x}_m, \mathbf{x}_n)$. (K is called the *Gram* matrix.)

- 2: Use a QP-solver to optimize the dual problem:

$$\boldsymbol{\alpha}^* \leftarrow \text{QP}(Q_D, -\mathbf{1}_N, A_D, \mathbf{0}_{N+2}).$$

- 3: Let s be any support vector for which $\alpha_s^* > 0$. Compute

$$b^* = y_s - \sum_{\alpha_n^* > 0} y_n \alpha_n^* K(\mathbf{x}_n, \mathbf{x}_s).$$

- 4: Return the final hypothesis

$$g(\mathbf{x}) = \text{sign} \left(\sum_{\alpha_n^* > 0} y_n \alpha_n^* K(\mathbf{x}_n, \mathbf{x}) + b^* \right).$$

The final hypothesis

Express $g(\mathbf{x}) = \text{sign}(\mathbf{w}^T \mathbf{z} + b)$ in terms of $K(-, -)$

$$\mathbf{w} = \sum_{\mathbf{z}_n \text{ is SV}} \alpha_n y_n \mathbf{z}_n \implies g(\mathbf{x}) = \text{sign} \left(\sum_{\alpha_n > 0} \alpha_n y_n K(\mathbf{x}_n, \mathbf{x}) + b \right)$$

$$\text{where } b = y_m - \sum_{\alpha_n > 0} \alpha_n y_n K(\mathbf{x}_n, \mathbf{x}_m)$$

for any support vector ($\alpha_m > 0$)

Examples of kernel

- Linear: $K(\mathbf{x}, \mathbf{x}') = \mathbf{x}^T \mathbf{x}'$
- Polynomial of order Q : $K(\mathbf{x}, \mathbf{x}') = (\zeta + \gamma \mathbf{x}^T \mathbf{x}')^Q$, $\zeta, \gamma > 0$
- Gaussian RBF: $K(\mathbf{x}, \mathbf{x}') = \exp(-\gamma \|\mathbf{x} - \mathbf{x}'\|^2)$, $\gamma > 0$

Polynomial kernel

$$\mathbf{x} = (x_1, \dots, x_d)$$


$$\mathbf{z} = \Phi(\mathbf{x}) = (1, x_1, \dots, x_d, x_1 x_1, x_1 x_2, \dots, x_2 x_1, \dots, \dots, x_d x_d)$$

Dimension of \mathbf{z} : $\tilde{d} = 1 + d + d^2$


$$\begin{aligned}\Phi(\mathbf{x})^T \Phi(\mathbf{x}') &= 1 + \sum_{i=1}^d x_i x'_i + \sum_{i=1}^d \sum_{j=1}^d x_i x_j x'_i x'_j \\ &= 1 + \mathbf{x}^T \mathbf{x}' + \left(\sum_{i=1}^d x_i x'_i \right) \left(\sum_{j=1}^d x_j x'_j \right) \\ &= 1 + \mathbf{x}^T \mathbf{x}' + (\mathbf{x}^T \mathbf{x}')^2 = (1 + \mathbf{x}^T \mathbf{x})^2\end{aligned}$$

Computational complexity: from $\mathcal{O}(\tilde{d})$ to $\mathcal{O}(d)$


Example: polynomial kernel (degree 2)



$$(1 + 0.001\mathbf{x}^T\mathbf{x}')^2$$




$$1 + (\mathbf{x}^T\mathbf{x}') + (\mathbf{x}^T\mathbf{x}')^2$$



$$(1 + 1000\mathbf{x}^T\mathbf{x}')^2$$

$$K(\mathbf{x}, \mathbf{x}') = (\zeta + \gamma\mathbf{x}^T\mathbf{x}')^Q$$

Example: polynomial kernel (degree 10)




$$K(\mathbf{x}, \mathbf{x}') = \exp(-\gamma \|\mathbf{x} - \mathbf{x}'\|^2) \quad (\gamma > 0)$$

Expanding it for the case when $d = 1$


$$\begin{aligned} K(x, x') &= \exp(-\|x - x'\|^2) \\ &= \exp(-(x)^2) \cdot \exp(2xx') \cdot \exp(-(x')^2) \\ &= \exp(-(x)^2) \cdot \left(\sum_{k=0}^{\infty} \frac{2^k (x)^k (x')^k}{k!} \right) \cdot \exp(-(x')^2), \\ \Phi(x) &= \exp(-x^2) \cdot \left(1, \sqrt{\frac{2^1}{1!}}x, \sqrt{\frac{2^2}{2!}}x^2, \sqrt{\frac{2^3}{3!}}x^3, \dots \right) \end{aligned}$$

That means $\tilde{d} = \infty$!


Example: Gaussian-RBF kernel



$$\exp(-1\|\mathbf{x} - \mathbf{x}'\|^2)$$




$$\exp(-10\|\mathbf{x} - \mathbf{x}'\|^2)$$




$$\exp(-100\|\mathbf{x} - \mathbf{x}'\|^2)$$

Example: linear \times Gaussian-RBF kernels



(a) linear classifier



(b) Gaussian-RBF kernel

How do we know that \mathcal{Z} exists ...

... for a given $K(\mathbf{x}, \mathbf{x}')$? valid kernel

Three approaches:

1. By construction
2. Math properties (*Mercer's condition*)
3. Who cares? 😊

Design your own kernel

$K(\mathbf{x}, \mathbf{x}')$ is a valid kernel iff

1. It is symmetric and 2. The matrix:
- $$\begin{bmatrix} K(\mathbf{x}_1, \mathbf{x}_1) & K(\mathbf{x}_1, \mathbf{x}_2) & \dots & K(\mathbf{x}_1, \mathbf{x}_N) \\ K(\mathbf{x}_2, \mathbf{x}_1) & K(\mathbf{x}_2, \mathbf{x}_2) & \dots & K(\mathbf{x}_2, \mathbf{x}_N) \\ \dots & \dots & \dots & \dots \\ K(\mathbf{x}_N, \mathbf{x}_1) & K(\mathbf{x}_N, \mathbf{x}_2) & \dots & K(\mathbf{x}_N, \mathbf{x}_N) \end{bmatrix}$$

is **positive semi-definite**

for any $\mathbf{x}_1, \dots, \mathbf{x}_N$ (Mercer's condition)

Summary

- SVM solves the maximum margin problem
- The optimization problem, in its primal form, can be written as a QP problem
- Its dual formulation (Lagrangian) also reduces to a QP problem
- Primal: d variables and N constraints
- Dual: N variables (α_n) and $N + 1$ constraints
- Dual is more interesting when we use the kernel trick (very large \tilde{d})
- Dual finds the support vectors

The long journey ends here ...

If you got to this point, you are a survivor :-)