



source: best_walk_params.json (study ground_walk_v4b_kine_pf_gain 2.18) on fitted refit - edges drawn from the COMPILED network object (build_network.py), representative knee column x46/side | network size (live): 46 MN + 46x(3) sensory per side, +12 rhythm/pattern cells + 5 load-sharing INs per side; both sides + 8 shared = 410 neurons, 376 inputs



Deng et al./Nourse 2023 Table A6 (rat hindlimb SNS) - reference values:
HC->IN 2.75 uS exc | IN->HC 2.75 uS inh (RG+PF mutual inhibition IS LAMINATED through dedicated INs; no direct HC->HC synapses, no mutual excitation)
RGHC->PFHC 0.10 uS exc | PF->MN 1.5-4.9 uS exc (per joint)
PF->Ia 0.50 uS exc (phase gate) | IaIN->MN 2.0 uS inh | IaIN<->IaIN 0.5 inh
MN->RC 0.50 uS exc | RC->MN 0.50 uS inh | RC->RC 0.5 inh | RC->IAIN 0.5 inh
IbIN->MN 0.59 uS EXCITATORY (positive force feedback) | PF->Ib 2.0 uS shunt
MN->muscle: act = $1/(1+\exp(s(x0-V))) + y0$, $s=0.153$, $x0=-70$ mV (Fig 6B)
feedback: MuJoCo muscle TENSION formatted as Ia and Ib input (no II)