




CONTACT SENSORS (heel/toe) + spiking-conversion OUTCOMES

SPIKING CONVERSION OUTCOMES (headless AnimatSimulator, 2026-10-02 goal 3 - topology preserved verbatim, classes converted):
W2L v1 (thr -55/-59, decay 10 ms): RHYTHM DEAD, 2 spikes/5 s | W2L v2 (calibrated thr): dead | W2L v3 (decay 30 ms): dead
BipRG v1: asymmetric latch, R half-centers 442-448 Hz | Biped v1: SYNCHRONOUS co-bursting 11.8 Hz both HCs (same-leg xcorr r=1.000)
Biped v2 (calibrated): 0 spikes | ABLATION (spiking thresholds ONLY, graded synapses kept): RHYTHM PRESERVED -> the SYNAPSE conversion breaks the rhythm;
a spiking version needs per-spike amplitudes/decays re-tuned as a rate code (Li 2023 model: natively spiking, tuned that way from birth, 0.77 Hz bursts).

every neuron drawn with a spike glyph: the spiking copies convert ALL neurons (v1 threshold 0.5, no v1 MN template 2 per-discrete lateralized W2L from the gradient peaks) to a non-spiking chemical synapses to the spiking chemical (validate.py: connexion sets identical). Reports: reports_spiking_20261002/goal3_animatlab_spiking.md.