W2L108 R · central feedback connection key

g: maximum conductance (uS); Erev, threshold, saturation: mV. Full IDs in CSV/SVG.

Key	Source → target	Named type	g	Erev	lo	hi
R01	R Hip flx Ia → R Hip PF flx	Ia_HipFlx_to_HipPFflx_Excite	0.01	-40	-60	-40
R02	R Hip ext Ib → R Hip PF ext	Ib_HipExt_to_HipPFext_Excite	0.01	-40	-60	-40
R03	R Hip flx II → R Hip PF flx	II_HipFlx_to_HipPFflx_Excite	0.01	-40	-60	-40
R04	R heel contact → R RG ext	Heel_Ipsi_RGext_Excite	0.01	-40	-60	-40
R05	R heel contact → R Hip PF ext	Heel_Ipsi_HipPFext_Excite	0.01	-40	-60	-40
R06	R heel contact → R Knee PF ext	Heel_Ipsi_KneePFext_Excite	0.01	-40	-60	-40
R07	R Hip flx Ia → R RG flx	Ia_HipFlx_to_RGflx_Excite	0.01	-40	-60	-40
R08	R Knee flx Ia → R RG flx	Ia_KneeFlx_to_RGflx_Excite	0.01	-40	-60	-40
R09	R Hip ext Ib → R RG ext	Ib_HipExt_to_RGext_Excite	0.01	-40	-60	-40
R10	R Knee ext Ib → R RG ext	Ib_KneeExt_to_RGext_Excite	0.01	-40	-60	-40
R11	R Hip flx II → R RG flx	II_HipFlx_to_RGflx_Excite	0.01	-40	-60	-40
R12	R Knee flx II → R RG flx	II_KneeFlx_to_RGflx_Excite	0.01	-40	-60	-40
R13	R Knee flx Ia → R Knee PF flx	Ia_KneeFlx_to_KneePFflx_Excite	0.01	-40	-60	-40
R14	R Knee ext Ib → R Knee PF ext	Ib_KneeExt_to_KneePFext_Excite	0.01	-40	-60	-40
R15	R Knee flx II → R Knee PF flx	II_KneeFlx_to_KneePFflx_Excite	0.01	-40	-60	-40
R16	L heel contact → R RG flx	Heel_Contra_RGflx_Excite	0.01	-40	-60	-40
R17	L V3 → R RG ext IN	V3 contralateral RG ext IN, weak positive	0.02749	-40	-60	-25

Source: W2L108 verified standalone; receptor conversion and MN wiring are separate layers.