
W2L108 · left central sensory feedback

Half-centres and interneurons shown explicitly; one source neuron per sensory box



Triangle base meets its receiving object; triangle point lies on the incoming line.

Filled circle: inhibition at target rest. Crossings without dots are not junctions.

Ia / II / Ib sensory boxes each represent one source afferent neuron, not a muscle bundle.

All direct afferent targets here are HCs; opposite-side V3 targets the RG extensor IN.

Other homonymous II and toe routes belong to motor/reflex circuits, not this central panel.

W2L108 L · central feedback connection key


g: maximum conductance (uS); Erev, threshold, saturation: mV. Full IDs in CSV/SVG.

Key	Source → target	Named type	g	Erev	lo	hi
L01	L Hip flx Ia → L Hip PF flx	Ia_HipFlx_to_HipPFflx_Excite	0.01	-40	-60	-40
L02	L Hip ext Ib → L Hip PF ext	Ib_HipExt_to_HipPFext_Excite	0.01	-40	-60	-40
L03	L Hip flx II → L Hip PF flx	II_HipFlx_to_HipPFflx_Excite	0.01	-40	-60	-40
L04	L heel contact → L RG ext	Heel_Ipsi_RGext_Excite	0.01	-40	-60	-40
L05	L heel contact → L Hip PF ext	Heel_Ipsi_HipPFext_Excite	0.01	-40	-60	-40
L06	L heel contact → L Knee PF ext	Heel_Ipsi_KneePFext_Excite	0.01	-40	-60	-40
L07	L Hip flx Ia → L RG flx	Ia_HipFlx_to_RGflx_Excite	0.01	-40	-60	-40
L08	L Knee flx Ia → L RG flx	Ia_KneeFlx_to_RGflx_Excite	0.01	-40	-60	-40
L09	L Hip ext Ib → L RG ext	Ib_HipExt_to_RGext_Excite	0.01	-40	-60	-40
L10	L Knee ext Ib → L RG ext	Ib_KneeExt_to_RGext_Excite	0.01	-40	-60	-40
L11	L Hip flx II → L RG flx	II_HipFlx_to_RGflx_Excite	0.01	-40	-60	-40
L12	L Knee flx II → L RG flx	II_KneeFlx_to_RGflx_Excite	0.01	-40	-60	-40
L13	L Knee flx Ia → L Knee PF flx	Ia_KneeFlx_to_KneePFflx_Excite	0.01	-40	-60	-40
L14	L Knee ext Ib → L Knee PF ext	Ib_KneeExt_to_KneePFext_Excite	0.01	-40	-60	-40
L15	L Knee flx II → L Knee PF flx	II_KneeFlx_to_KneePFflx_Excite	0.01	-40	-60	-40
L16	R heel contact → L RG flx	Heel_Contra_RGflx_Excite	0.01	-40	-60	-40
L17	R V3 → L RG ext IN	V3 contralateral RG ext IN, weak positive	0.02749	-40	-60	-25

Source: W2L108 verified standalone; receptor conversion and MN wiring are separate layers.

W2L108 · right central sensory feedback

Half-centres and interneurons shown explicitly; one source neuron per sensory box



Triangle base meets its receiving object; triangle point lies on the incoming line.

Filled circle: inhibition at target rest. Crossings without dots are not junctions.

Ia / II / Ib sensory boxes each represent one source afferent neuron, not a muscle bundle.

All direct afferent targets here are HCs; opposite-side V3 targets the RG extensor IN.

Other homonymous II and toe routes belong to motor/reflex circuits, not this central panel.

W2L108 R · central feedback connection key


g: maximum conductance (uS); Erev, threshold, saturation: mV. Full IDs in CSV/SVG.

Key	Source → target	Named type	g	Erev	lo	hi
R01	R Hip flx Ia → R Hip PF flx	Ia_HipFlx_to_HipPFflx_Excite	0.01	-40	-60	-40
R02	R Hip ext Ib → R Hip PF ext	Ib_HipExt_to_HipPFext_Excite	0.01	-40	-60	-40
R03	R Hip flx II → R Hip PF flx	II_HipFlx_to_HipPFflx_Excite	0.01	-40	-60	-40
R04	R heel contact → R RG ext	Heel_Ipsi_RGext_Excite	0.01	-40	-60	-40
R05	R heel contact → R Hip PF ext	Heel_Ipsi_HipPFext_Excite	0.01	-40	-60	-40
R06	R heel contact → R Knee PF ext	Heel_Ipsi_KneePFext_Excite	0.01	-40	-60	-40
R07	R Hip flx Ia → R RG flx	Ia_HipFlx_to_RGflx_Excite	0.01	-40	-60	-40
R08	R Knee flx Ia → R RG flx	Ia_KneeFlx_to_RGflx_Excite	0.01	-40	-60	-40
R09	R Hip ext Ib → R RG ext	Ib_HipExt_to_RGext_Excite	0.01	-40	-60	-40
R10	R Knee ext Ib → R RG ext	Ib_KneeExt_to_RGext_Excite	0.01	-40	-60	-40
R11	R Hip flx II → R RG flx	II_HipFlx_to_RGflx_Excite	0.01	-40	-60	-40
R12	R Knee flx II → R RG flx	II_KneeFlx_to_RGflx_Excite	0.01	-40	-60	-40
R13	R Knee flx Ia → R Knee PF flx	Ia_KneeFlx_to_KneePFflx_Excite	0.01	-40	-60	-40
R14	R Knee ext Ib → R Knee PF ext	Ib_KneeExt_to_KneePFext_Excite	0.01	-40	-60	-40
R15	R Knee flx II → R Knee PF flx	II_KneeFlx_to_KneePFflx_Excite	0.01	-40	-60	-40
R16	L heel contact → R RG flx	Heel_Contra_RGflx_Excite	0.01	-40	-60	-40
R17	L V3 → R RG ext IN	V3 contralateral RG ext IN, weak positive	0.02749	-40	-60	-25

Source: W2L108 verified standalone; receptor conversion and MN wiring are separate layers.

w2lvar · left sensory feedback into RG

Muscle bundles are labelled by channel count; actual receiving interneurons stay explicit.




A thick ×N line stands for N distinct source-muscle connections; the member list is in the companion key.

Triangles have their base on the receiving object. Filled circles are inhibitory terminals.

HC = half-centre · IN = interneuron · Dashed destination: reference to the same HC, not an added cell.

w2lvar · left sensory feedback into PF

Muscle bundles are labelled by channel count; actual receiving interneurons stay explicit.



A thick ×N line stands for N distinct source-muscle connections; the member list is in the companion key.

Triangles have their base on the receiving object. Filled circles are inhibitory terminals.

HC = half-centre · IN = interneuron · Dashed destination: reference to the same HC, not an added cell.

w2lvar · exact members of the muscle bundles

Ia: 19 source channels

Ia_add_brev_l
Ia_add_long_l
Ia_add_mag1_l
Ia_add_mag2_l
Ia_bifemlh_l
Ia_bifemsh_l
Ia_ext_dig_l
Ia_ext_hal_l
Ia_extobl_l
Ia_grac_l
Ia_iliacus_l
Ia_intobl_l
Ia_pect_l
Ia_psoas_l
Ia_sar_l
Ia_semimem_l
Ia_semiten_l
Ia_tfl_l
Ia_tib_ant_l

II: 19 source channels

II_add_brev_l
II_add_long_l
II_add_mag1_l
II_add_mag2_l
II_bifemlh_l
II_bifemsh_l
II_ext_dig_l
II_ext_hal_l
II_extobl_l
II_grac_l
II_iliacus_l
II_intobl_l
II_pect_l
II_psoas_l
II_sar_l
II_semimem_l
II_semiten_l
II_tfl_l
II_tib_ant_l


Ib: 27 source channels

Ib_add_mag3_l
Ib_ercspn_l
Ib_flex_dig_l
Ib_flex_hal_l
Ib_gem_l
Ib_glut_max1_l
Ib_glut_max2_l
Ib_glut_max3_l
Ib_glut_med1_l
Ib_glut_med2_l
Ib_glut_med3_l
Ib_glut_min1_l
Ib_glut_min2_l
Ib_glut_min3_l
Ib_lat_gas_l
Ib_med_gas_l
Ib_per_brev_l
Ib_per_long_l
Ib_per_tert_l
Ib_peri_l
Ib_quad_fem_l
Ib_rect_fem_l
Ib_soleus_l
Ib_tib_post_l
Ib_vas_int_l
Ib_vas_lat_l
Ib_vas_med_l

Each bundle expands into these separate muscle-channel connections, not one summed synapse.
Endpoints, individual conductances, and edge IDs remain in the companion connection keys.

syn6 · left sensory feedback into RG

Muscle bundles are labelled by channel count; actual receiving interneurons stay explicit.




Saved heel/toe physical input gains are zero: the drawn chemical paths exist but receive no contact drive.

Triangles have their base on the receiving object. Filled circles are inhibitory terminals.

HC = half-centre · IN = interneuron · Dashed destination: reference to the same HC, not an added cell.

syn6 · left sensory feedback into PF

Muscle bundles are labelled by channel count; actual receiving interneurons stay explicit.



Saved heel/toe physical input gains are zero: the drawn chemical paths exist but receive no contact drive.

Triangles have their base on the receiving object. Filled circles are inhibitory terminals.

HC = half-centre · IN = interneuron · Dashed destination: reference to the same HC, not an added cell.

s3k · left sensory feedback into RG / PF

Muscle bundles are labelled by channel count; actual receiving interneurons stay explicit.

Ia
No direct RG/PF route
in this saved condition

II
No direct RG/PF route
in this saved condition

Ib
No direct RG/PF route
in this saved condition

heel
No direct RG/PF route
in this saved condition

toe
No direct RG/PF route
in this saved condition

No direct Ia, II, Ib, heel, or toe
chemical projection into RG/PF
in this saved s3k standing condition.

A thick ×N line stands for N distinct source-muscle connections; the member list is in the companion key.

Triangles have their base on the receiving object. Filled circles are inhibitory terminals.

HC = half-centre · IN = interneuron · Dashed destination: reference to the same HC, not an added cell.