
side l: cycle mean NMF synergy coefficients (n=1 complete stride)
stance
swing
normalized PF voltage
1.0
0.8
0.6
0.4
0.2
0.0
S1
S2
S3
S4
S5
S6
S7
S8
0
20
40
60
80
100
stance-rescaled gait phase (%)
NMF spatial gains (extracted synergies; functional labels provisional)
S1 hip ext. / biartic. knee flx.
S2 knee ext. / ankle DF
S3 posterior chain / ankle PF
S4 hip flex. + frontal stabilization
S5 trunk-pelvis / ankle DF / hip abd.
S6 hip ext. / knee flx. residual
S7
S8
glut_med1_l
glut_med2_l
glut_med3_l
glut_min1_l
glut_min2_l
glut_min3_l
semimem_l
semiten_l
bifemlh_l
bifemsh_l
sar_l
add_long_l
add_brev_l
add_mag1_l
add_mag2_l
add_mag3_l
tfl_l
pect_l
grac_l
glut_max1_l
glut_max2_l
glut_max3_l
iliacus_l
psoas_l
rect_fem_l
vas_med_l
vas_int_l
vas_lat_l
med_gas_l
lat_gas_l
soleus_l
tib_post_l
flex_dig_l
flex_hal_l
tib_ant_l
per_brev_l
per_long_l
per_tert_l
ext_dig_l
ext_hal_l
ercspn_l
intobl_l
extobl_l
activation gain
1.0
0.8
0.6
0.4
0.2
0.0