
Shared PF source · two muscle-specific reflex and physical loops



AD = map / adapter. Brown arrows: physical signals. Open triangles: excitation. Filled circles: inhibition.

Spindle and tendon symbols show encoded signals; muscle→spindle→AD→Ia/II and muscle→force sensor→AD→Ib are explicit.


MN→AD→muscle carries activation. Each muscle retains its own MN, Ia, II, Ib, interneurons, and Renshaw cell.

Gray dashed PF→MN routes have effective g = 0. Central afferent→RG/PF projections are detailed in the preceding RG/PF sheets.

VM and VL share PF drive, but retain separate loops. The four antagonist destinations are grouped; every line to that group represents ×4 synapses.

SM = semimembranosus; ST = semitendinosus; BFLH/BFSH = biceps femoris long/short head. Their own local loops are outside this crop.

Biarticular muscles · PF inputs, local reflexes, and physical maps



AD = map / adapter. Brown arrows: physical signals. Open triangles: excitation. Filled circles: inhibition.

Spindle and tendon symbols show encoded signals; muscle→spindle→AD→Ia/II and muscle→force sensor→AD→Ib are explicit.

MN→AD→muscle carries activation. Each muscle retains its own MN, Ia, II, Ib, interneurons, and Renshaw cell.

Gray dashed PF→MN routes have effective g = 0. Central afferent→RG/PF projections are detailed in the preceding RG/PF sheets.

RF: hip flexion + knee extension. SM: hip extension + knee flexion. Both PF inputs terminate on the same muscle MN.