



Excitation: triangle base on destination · Inhibition: filled circle · Crossings: semicircular bridges

Internal RG/PF wiring omitted. Opposite-side sources are placed beside their RG destinations.


w2lvar · RG, Hip PF, and K&A PF



×N = separate muscle-channel synapses; their individual destination parameters are retained in the key.


K&A is the shared knee–ankle PF pair. This page uses one set of destination objects, without duplicate HCs.

w2lvar · Ia and II pathways




Same RG / Hip PF / K&A PF arrangement as the overview. Gray lines retain the RG→PF context.

w2lvar · lb, heel, and toe pathways



Same RG / Hip PF / K&A PF arrangement as the overview. Gray lines retain the RG→PF context.


syn6 · RG and six NMF pattern-formation channels



Red dashed: the saved heel→flexor-IN branch disputed in Comment 30; the source circuit has not been changed.


S1–S6 are PF channels, not hip/knee/ankle pairs. Saved contact-input gains are zero in this standing condition.

s3k · RG and four PF phase channels




Saved standing condition: actual descending, contralateral, and RG→PF connections. Internal motifs omitted.

Rat134 · curated receptor-geometry condition




Separate knee and ankle PF modules are retained here. These are source connections, not assumed human-model pathways.

RF / HAM · four-cell PF subnetwork (saved left-side projection)




Joint PF → RF/HAM pair → motor neurons. Expanded four-cell connections shown; other afferents are outside this saved projection.



Population readouts drive the chemical pathways; these are not soma-to-soma spike-event arrows. Internal motifs omitted.

Gait2392 TimingZeroSFA · central spiking populations and PF channels



RG and S1–S6 are spiking populations; INs remain graded. Chemical pathways use spike-derived proxy voltage. Red branch remains disputed.

Source / destination / synapse: what the labels mean

II_tib_ant names one type-II afferent neuron, not one globally fixed synapse.

A connection is defined by its source, destination, conductance, reversal, and presynaptic window.

The same afferent can therefore have different parameters at different destinations.

The checked left/right homologues below match; diagram labels omit the side suffix.

Source → destination	g (uS)	Erev (mV)	lo / hi (mV)	L = R
II_tib_ant → RG_F	0.0278947	8.0	0 / 5	yes
II_tib_ant → InF	0.0278947	8.0	0 / 5	yes
II_tib_ant → PF_HIP-F	0.0278947	8.0	0 / 5	yes
II_tib_ant → PF_KNEE-F	0.0278947	8.0	0 / 5	yes
II_tib_ant → IIX_tib_ant	1	8.0	0 / 5	yes
II_tib_ant → IIIN_tib_ant	1	8.0	0 / 5	yes
IIIN_tib_ant → MN_med_gas	0.133333	-5.0	0 / 5	yes
IIIN_tib_ant → MN_lat_gas	0.133333	-5.0	0 / 5	yes
IIIN_tib_ant → MN_soleus	0.133333	-5.0	0 / 5	yes
IIIN_tib_ant → MN_tib_post	0.133333	-5.0	0 / 5	yes
IIIN_tib_ant → MN_flex_dig	0.133333	-5.0	0 / 5	yes
IIIN_tib_ant → MN_flex_hal	0.133333	-5.0	0 / 5	yes
IIX_tib_ant → MN_tib_ant	0.4	8.0	0 / 5	yes
IIIN_tib_ant → MN_per_brev	0.133333	-5.0	0 / 5	yes
IIIN_tib_ant → MN_per_long	0.133333	-5.0	0 / 5	yes
IIIN_tib_ant → MN_per_tert	0.133333	-5.0	0 / 5	yes

A ×N line bundles N separately parameterized connections; it is not a single summed synapse.

Ankle channels are included in the muscle bundles where present. Complete members and edge IDs accompany the drawings.