ADHD Week 9 - Completion Quiz

Warning: You have already made the maximum number of submissions. Additional submissions will not count for credit. You are welcome to try it as a learning exercise.

■ In accordance with the Coursera Honor Code, I (Matthew Kramer) certify that the answers here are my own work.

Question 1

The Vanderbilt ADHD rating scale should be completed by:

- A. Parents
- B. Teachers
- C. Patients
- O. A and B
- E. A and C
- F. All of the above

Question 2

Which of the following about stimulant medications is true?

- A. Stimulants are addictive when used as prescribed
- B. Stimulants increase aggression
- C. Stimulants alone can cause Tourette's Syndrome
- D. Most doctors overdose ADHD patients with stimulants
- E. For kids with ADHD, taking stimulants early in life increases their risk of later substance abuse

- F. Stimulant medications don't typically help ADHD patients perform better academically
- G. None of the above

Question 3

The Vanderbilt rating scale requires parents and teachers to comment on patient's

- A. Symptoms
- B. Academic performance
- C. Classroom behavior
- D. All of the above

Question 4

Of all the possible side effects listed, which are the most commonly experienced by patients taking stimulant medications?

- A. Tics and heart rate changes
- B. Headache and abdominal pain
- C. Insomnia and appetite changes
- D. Nausea and constipation
- E. Nervousness and crying

Question 5

On average, how much are weight and height (wt, ht) INITIALLY slowed/delayed by stimulant medications?

A. 1-4 lb, 1 cm

- B. 1-4 lb, 2 cm
- O. 3-6 lb, 1 cm
- D. 3-6 lb, 2 cm
- E. Weight is gained and height may actually increase on stimulants

Question 6

What mechanism of action do methylphenidate and amphetamine derivatives share?

- A. Releasing transmitter from vesicle
- B. Blocking re-uptakeinto loading vesicle
- C. Increasing release of neurotransmitter into the synaptic cleft
- D. Blocking re-uptake of neurotransmitter into pre-synaptic terminal

Question 7

Please select all the following medications that are some form of methylphenidate.

- A. Daytrana
- B. Adderall
- C. Concerta
- D. Ritalin
- E. Vyvanse
- F. Focalin

Question 8

TRUE OR FALSE: After a large review of the literature, the use of stimulant medications for ADHD was linked to an increased risk of cardiovascular illness.

True			
False			
In accordance with the Coursera Honor Code, I (Matthew Kramer) certify that the			
answers here are my own	work.		

Submit Answers Save Answers

You cannot submit your work until you agree to the Honor Code. Thanks!