

This document sketches out the calculation of autocorrelation functions.

First, suppose we have calculate the left eigenvectors and eigenvalues:

$$T^T \phi_i = \lambda \phi_i$$

Suppose that  $\pi$  is the equilibrium population. Then, we can normalize the eigenvectors such that:

$$\phi_i^T \pi^{-1} \phi_j = \delta_{ij}$$

Above, we denote  $\pi^{-1}$  to be a diagonal matrix with elements  $\pi_i^{-1}$ . The autocorrelation function of the observable  $f_i$  can be denoted:

$$E(f(z_t)f(z_0)) = \sum_{i,j} f_i P(z_0 = i) f_j P(z_t = j | z_0 = i) = \sum_{i,j} f_i f_j \pi_i T_{ij} =$$

We know that

$$T_{ab} = \sum_k \lambda_k (\psi_k)_a (\phi_k)_b = \sum_k \lambda_k (\pi_a)^{-1} (\phi_k)_a (\phi_k)_b$$

Thus,

$$E(f(z_t)f(z_0)) = \sum_{i,j,k} f_i f_j \lambda_k (\phi_k)_i (\phi_k)_j = \sum_k \lambda_k s_k^2$$

Where

$$s_k = \sum_i f_i (\phi_k)_i$$