

A Book of High School and Engineering Common Core Mathematical Formulae

Agnij Mallick

December, 2020

Contents

I	Algebra	7
1	Logarithm	8
1.1	Basic Formulae	8
1.2	Series	8
2	Complex Numbers	9
2.1	Basic Formulae	9
2.2	Arithmetic Operation of Complex Number	9
2.3	Euler's Formula	9
2.4	Trigonometric Ratios in Complex Form	10
2.5	De Moivre's Formula	10
2.6	Application of Euler's and De Moivre's Formula	10
2.7	Roots of Unity	10
2.8	Important Relations of Complex Numbers	10
3.1.2	Important Relation	11
3.2	Geometric Progression (G.P.)	11
3.2.1	The Value of 'r'	11
3.2.2	Sum of a G.P. Series	11
3.5.1	Riemann Zeta Function	12
3.5.2	Riemann's Infinite Series as an Integration	12
4	Test of Convergence of Infinite Series	13
4.1	Definition	13
4.2	Tests of Convergence	13
4.2.1	Comparison Test	13
4.2.2	Limit Form	13
4.2.3	Integral Test or Maclaurin-Cauchy Test	13
4.2.4	Ratio Test	14
4.2.5	D'Alembert's Ratio Test	14
4.2.6	Rabbe's Test	14
4.2.7	Cauchy's Root Test	14
4.2.8	Logarithmic Test	15
5.1.1	Minor and Cofactor	16
5.2	Properties of Determinants	16

5.3	Cramer's Rule	17
5.3.1	Consistency Test	17
6.2	Multiplication of Two Matrices	18
6.2.1	Multiplicative Properties	18
6.3	Adjoint of a Matrix	18
6.4	Martin's Rule	19
7.4	Pascal's Rule	20
8	Boolean Algebra	21
9	Remainder Theorems	22
9.1	Remainder Theorem	22
9.2	Euler's Remainder Theorem	22
9.2.1	Euler's Totient Function	22
9.3	Wilson Theorem	23
II	Co-ordinate Geometry	25
10	2-D Co-ordinate Geometry	26
10.1	Distance between Two Points	26
10.2	Section Formula	26
10.2.1	Corollary: Mid - Point Formula	26
11	Triangles	27
11.1	Centroid of a Traiangle	27
11.2	Area of Triangle	27
11.2.1	Determinant Method	27
11.2.2	Heron's Formula	27
11.3	Incircle of a Triangle	28
11.4	Circumcircle of a Triangle	28
12	Straight Line	29
12.1	Equation of Straight Line Passing Through (x_0, y_0) and Slope m	29
12.2	Distance Between Two Points on a Line	29
12.3	Angle Between Two Lines	29
12.4	Distance of a Point from a Line	30
12.5	Angle Bisector of a Line	30
12.6	Equation of a Straight Line Passing through the Intersection of Two Lines	30
12.7	Relative Position of Points w.r.t. a Line	30
12.8	Ratio of Division of Line Segment	30
13	General Theory of Second Degree Equation	31

14 Conics	32
14.1 Parametric Form of Conics	32
14.1.1 Hyperbola	32
14.1.2 Ellipse	32
14.1.3 Parabola	32
14.2 Equation form of Conics	32
14.2.1 Parabola	32
14.2.2 Ellipse and Hyperbola	33
15 Circles	34
15.1 Locus Form	34
15.2 Diameter Form	34
15.3 General Form	34
15.4 Important Relations	34
15.5 Common for Two Circles	35
16 Vectors	36
16.1 Modulus of a Vector	36
16.2 Sum of Vectors	36
16.3 Product of Vectors	36
16.3.1 Dot Product	36
16.3.2 Cross Product	36
16.4 Test of Co-planarity	37
17 3D - Space	38
17.1 Line segments in 3D - Space	38
17.1.1 Distance between Two Points	38
17.1.2 Section Formula of a Line Segment Divided in the ratio $m : n$	38
17.2 Line in 3D - Space	38
17.2.1 Angle between Two Lines	38
17.2.2 Skew and Co-planar Lines	39
17.2.3 Distance between Lines	39
17.3 Triangular Plane	39
17.3.1 Centroid of a Triangle	39
18 3D - Plane	40
18.1 Angle Between Two Planes	40
18.2 Distance of a Point from a Plane	40
18.2.1 Catesian Form	40
18.2.2 Vector Form	41
III Statistics	42
19 Descriptive Statistics	43
19.1 Measure of Location	43
19.1.1 Mean	43
19.1.2 Median	43
19.1.3 Mode	43

19.1.4	Quartile	43
19.2	Measure of Spread	43
19.2.1	Variance	43
19.2.2	Sample Variance	44
19.2.3	Standard Deviation and Sample Standard	44
19.2.4	Co-efficient of Variance	44
19.3	Skewness	44
19.3.1	Types of Skewness	44
19.3.2	Measure of Skewness	44
19.4	Kurtosis	45
19.4.1	Type of Kurtosis	45
20	Hypothesis Testing	46
20.1	T-Test	46
20.2	χ^2 Test	46
21	Research and Survey Design	47
21.1	Population Covariance	47
21.2	Sample Covariance	47
21.3	Bravais-Pearson Correlation Co-efficient	47
21.4	Spearman's Rank Correlation Co-efficient	47
22	Estimation of Regression Function	48
22.1	Sum of Squares Error	49
22.1.1	R^2 : Coefficient of Determination	49
22.1.2	\bar{R}^2 : Coefficient of Determination	50
22.2	T-Test	50
22.3	F-Test	50
22.4	Test for Heteroskedasticity	50
22.4.1	Definition	50
22.4.2	Durbin-Watson Test	50
23	Dummy Variables	51
23.1	Dummy Variable	51
23.2	Slope Dummy Variable	51
23.3	Slope & Dummy Variable	52
23.4	Multi-Categories Dummy Variable	53
24	Logistic Regression	54
IV	Trigonometry	55
25	Circular Trigonometric Functions	56
25.1	Trigonometric Ratios of Standard Angles	56
25.2	Negative Angle Formula	56
25.3	Sum of Angles Formula	57
25.4	Difference of Angles Formula	57
25.5	Multiples and Sub-multiples of π and $\frac{\pi}{2}$	57
25.6	$(\frac{\pi}{2} \pm \theta)$ Formula	58

25.7 $(\frac{\pi}{4} \pm \theta)$ Formula	58
25.8 Trigonometric Identities	58
25.9 Double Angle Formula	59
25.10 Triple Angle Formula	59
25.11 Sum and Product of Two Ratios	59
25.12 General Solutions	60
25.13 Taylor Series Expansion of Trigonometric Ratios	60
26 Inverse Circular Trigonometric Function	61
26.1 Definition of Inverse Circular Trigonometric Function	61
26.1.1 For $\sin x$	61
26.1.2 For $\cos x$	61
26.1.3 For $\tan x$	61
26.1.4 For $\cot x$	62
26.1.5 For $\csc x$	62
26.2 Negative Arguments	62
26.3 Reciprocal Relations	63
26.4 I.T.F. Identities	63
26.5 Sum of Two Angles	63
26.6 Difference of Two Angles	63
26.7 Interconversion of Ratios	63
26.8 Miscellaneous Relations	64
27 Hyperbolic Trigonometric Functions	65
27.1 Definition	65
27.2 Identities	65
27.3 Inverse Hyperbolic Function	66
27.4 Relation to Circular Trigonometric Functions	66
V Calculus	67
28 Limits	68
28.1 L'Hospital Rule	68
29 Differentiation	70
29.1 Differentiation by First Principle	70
29.2 Standard Differentiation Formulae	70
29.2.1 Circular Trigonometric Functions	71
29.2.2 Inverse Circular Trigonometric Functions	71
29.2.3 Hyperbolic Trigonometric Function	71
29.2.4 Inverse Hyperbolic Trigonometric Function	72
29.3 Rules of Differentiation	72
29.4 Chain Rule	72
30 Successive Differentiation	73
30.1 Leibnitz's Theorem	73

31 Partial Derivative	74
31.1 Chain Rule	74
31.2 Euler's Theorem	74
32 Application of Differential	75
32.1 Rolle's Theorem	75
32.2 Mean Value Theorem or LaGrange's Theorem	75
32.3 Cauchy's Mean Value Theorem	75
32.4 Maxima and Minima	76
32.4.1 Maxima	76
32.4.2 Minima	76
32.5 Taylor's Theorem	76
32.5.1 Remainder Term	76
32.5.2 Conditions for Validity of Expansion	77
32.5.3 Taylor's Theorem for Two Variables	77
32.6 Maclaurin's Series	77
32.6.1 Maclaurin's Series with Two Variables	77
32.7 Curvature	77
32.7.1 Radius of Curvature	78
32.7.2 Newton's Formula	78
32.7.3 Tangent at Origin	78
32.8 Asymptotes	78
32.8.1 Asymptote of Algebraic Curves	79
33 Integration	80
33.1 General Formulae	80
33.2 Circular Trigonometric Functions	81
33.3 Inverse Circular Trigonometric Function	81
33.4 Standard Integrals	82
33.5 Special Forms	82
33.5.1 Integration by Part	82
34 Definite Integration	83
34.1 Properties of Definite Integration	83
34.2 Approximation	83
34.3 Sum of Infinite Series as a Definite Integral	83
35 Reduction Formula	84
35.1 Circular Trigonometric Functions	84
35.2 Special Definite Integrations	84
35.3 Recursive Forms in Circular Trigonometric Functions	85
36 β and Γ Functions	86
36.1 Important Relations between $\beta(m, n)$ and $\Gamma(n)$ Functions	86
37 Multiple Integrals	87
37.1 Two Variables	87
37.2 Three Variables	87

Part I

Algebra

Chapter 1

Logarithm

1.1 Basic Formulae

For $a^x = b$:

$$\log_a x, \forall x \leq 0 \text{ is undefined} \quad (1.1)$$

$$\log_a b = x, \text{ if } a^x = b, a \neq 1 \quad (1.2)$$

$$\log_b a^m = m \log_b a, \text{ for } a^m = b \quad (1.3)$$

$$a^{\log_a x} = x \quad (1.4)$$

$$a^{\log_b c} = c^{\log_b a} \quad (1.5)$$

$$\frac{1}{\log_a b} = \log_b a \quad (1.6)$$

$$\log_c(ab) = \log_c a + \log_c b \quad (1.7)$$

$$\log_c \left(\frac{a}{b} \right) = \log_c a - \log_c b \quad (1.8)$$

$$|\log_a x| = \begin{cases} -\log_a x, & \text{if } 0 < x < 1 \\ \log_a x, & \text{if } 1 \leq x < \infty \end{cases} \quad (1.9)$$

1.2 Series

$$e^x = 1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots \infty \quad (1.10)$$

$$= \lim_{n \rightarrow \infty} \sum_{i=0}^n \frac{x^i}{i!} \quad (1.11)$$

$$\log(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \cdots \infty \quad (1.12)$$

$$= \lim_{n \rightarrow \infty} \sum_{i=1}^n (-1)^{(i-1)} \frac{x^i}{i} \quad (1.13)$$

Chapter 2

Complex Numbers

2.1 Basic Formulae

For $z = x + iy$,

$$|z| = \sqrt{x^2 + y^2} \quad (2.1)$$

$$\tan \theta = \frac{y}{x} \quad (2.2)$$

$$\bar{z} = x - iy \quad (2.3)$$

2.2 Arithmetic Operation of Complex Number

For two complex numbers $z_1 = x_1 + iy_1$ and $z_2 = x_2 + iy_2$:

$$z_1 + z_2 = (x_1 + x_2) + i(y_1 + y_2) \quad (2.4)$$

$$z_1 z_2 = (x_1 x_2 - y_1 y_2) + i(x_1 y_2 + x_2 y_1) \quad (2.5)$$

$$|z_1 z_2| = |z_1| \cdot |z_2| \quad (2.6)$$

$$\frac{z_1}{z_2} = \frac{(x_1 x_2 + y_1 y_2) + i(x_2 y_1 - x_1 y_2)}{a_2^2 + b_2^2} \quad (2.7)$$

$$\left| \frac{z_1}{z_2} \right| = \frac{|z_1|}{|z_2|} \quad (2.8)$$

2.3 Euler's Formula

$$z = r e^{i\theta}, \text{ where} \quad (2.9)$$

$$r = |z| \quad (2.10)$$

$$e^{i\theta} = \cos \theta + i \sin \theta \quad (2.11)$$

$$\theta = \arctan \left(\frac{y}{x} \right) \quad (2.12)$$

2.4 Trigonometric Ratios in Complex Form

$$e^{i\theta} + e^{-i\theta} = 2 \cos \theta \quad (2.13)$$

$$\Rightarrow \cos \theta = \frac{e^{i\theta} + e^{-i\theta}}{2} \quad (2.14)$$

$$e^{i\theta} - e^{-i\theta} = 2 \sin \theta \quad (2.15)$$

$$\Rightarrow \sin \theta = \frac{e^{i\theta} - e^{-i\theta}}{2} \quad (2.16)$$

2.5 De Moivre's Formula

According to DeMoivre's Formula:

$$(\cos \theta + i \sin \theta)^n = \cos(n\theta) + i \sin(n\theta) \quad (2.17)$$

Proof

$$\begin{aligned} \cos \theta + i \sin \theta &= e^{i\theta} \\ \Rightarrow (\cos \theta + i \sin \theta)^n &= e^{n(i\theta)} \\ &= \cos(n\theta) + i \sin(n\theta) \\ &\text{Q.E.D.} \end{aligned}$$

2.6 Application of Euler's and De Moivre's Formula

For $z_1 = |r_1| e^{i\theta_1}$ and $z_2 = |r_2| e^{i\theta_2}$

$$z_1 z_2 = (r_1 r_2) e^{i(\theta_1 + \theta_2)} \quad (2.18)$$

$$\frac{z_1}{z_2} = \left(\frac{r_1}{r_2} \right) e^{i(\theta_1 - \theta_2)} \quad (2.19)$$

2.7 Roots of Unity

$$\sqrt[n]{1} = e^{i \frac{2k\pi}{n}}, \text{ where } k \in [0, n-1] \quad (2.20)$$

2.8 Important Relations of Complex Numbers

$$|z_1 + z_2| \leq |z_1| + |z_2| \quad (2.21)$$

$$|z_1 - z_2| \leq |z_1| + |z_2| \quad (2.22)$$

$$|z_1 - z_2| \geq ||z_1| - |z_2|| \quad (2.23)$$

$$|z_1 + z_2| \geq ||z_1| - |z_2|| \quad (2.24)$$

$$|z_1 + z_2|^2 = 2 \left(|z_1|^2 + |z_2|^2 \right) \quad (2.25)$$

3.1.2 Important Relation

If the three terms a, b, c are in A.P., then

$$2b = a + c \quad (3.2)$$

3.2 Geometric Progression (G.P.)

An geometric sequence is $a, ar, ar^2, \dots \infty$ or $t_n = ar^{n-1}$, where a is the first term, r is the common ratio, and n is the n^{th} -term. An geometric series is $a + ar + ar^2 + \dots \infty$.

3.2.1 The Value of 'r'

$$r = \frac{t_2}{t_1} = \frac{t_3}{t_2} = \dots = \frac{t_n}{t_{n-1}} \quad (3.3)$$

3.2.2 Sum of a G.P. Series

For a definite G.P. series, where there are n terms in the series, the sum of the series is:

$$S_n = \frac{a|r^n - 1|}{|r - 1|} \quad (3.4)$$

For an infite G.P. series the sum of the series is defined for $r < 1$. Sum of such a series is:

$$S_\infty = \frac{a}{1 - r} \quad (3.5)$$

3.5.1 Riemann Zeta Function

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} \quad (3.13)$$

3.5.2 Riemann's Infinite Series as an Integration

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=r_1}^{r_2} f\left(\frac{i}{n}\right) = \int_{\frac{r_1}{n}}^{\frac{r_2}{n}} f(x) dx \quad (3.14)$$

Chapter 4

Test of Convergence of Infinite Series

If $a_1, a_2, a_3, \dots, a_n$ is a sequence by a_n and their sum of series is S_n , then the following apply.

4.1 Definition

If

$$\lim_{n \rightarrow \infty} S_n = l$$

where l is a finite value, the series S_n is said to converge. A non-convergent series is called a divergent series.

4.2 Tests of Convergence

4.2.1 Comparison Test

If u_n and v_n are two positive series, then:

1. (a) v_n converges
(b) $u_n \leq v_n \forall n$ Then u_n converges.
2. (a) v_n diverges
(b) $u_n \geq v_n \forall n$ Then u_n diverges.

4.2.2 Limit Form

If

$$\lim_{x \rightarrow \infty} \frac{u_n}{v_n} = l$$

where l is a finite quantity $\neq 0$, then u_n and v_n converge and diverge together.

4.2.3 Integral Test or Maclaurin-Cauchy Test

For a series

$$\sum_{i=N}^{\infty} f(x), \text{ where } N \in \mathbb{Z} \quad (4.1)$$

will only converge if the improper integral

$$\int_N^{\infty} f(x)dx \quad (4.2)$$

is finite.

If the improper integral is finite, the upper and lower limit of the infinite series is given by:

$$\int_N^{\infty} f(x)dx \leq \sum_{i=N}^{\infty} f(x) \leq f(N) + \int_N^{\infty} f(x)dx \quad (4.3)$$

4.2.4 Ratio Test

If, for two series $\sum u_n$ and $\sum v_n$:

1. (a) $\sum v_n$ converges
(b) from or after a particular term $\frac{u_n}{u_{n+1}} > \frac{v_n}{v_{n+1}}$, then u_n converges.
2. (a) $\sum v_n$ diverges
(b) from or after a particular term $\frac{u_n}{u_{n+1}} < \frac{v_n}{v_{n+1}}$, then u_n diverges.

4.2.5 D'Alembert's Ratio Test

$$\lim_{n \rightarrow \infty} \frac{u_n}{u_{n+1}} = \lambda \quad (4.4)$$

- series converges if $\lambda < 1$
- series diverges if $\lambda > 1$
- fails if $\lambda = 1$

4.2.6 Rabbe's Test

$$\lim_{n \rightarrow \infty} n \left[\frac{u_n}{u_{n+1}} - 1 \right] = \kappa \quad (4.5)$$

- series converges if $\kappa < 1$
- series diverges if $\kappa > 1$
- fails if $\kappa = 1$

4.2.7 Cauchy's Root Test

$$\lim_{n \rightarrow \infty} |u_n| = \lambda \quad (4.6)$$

- series converges for $\lambda < 1$
- series diverges for $\lambda > 1$
- test fails for $\lambda = 1$

4.2.8 Logarithmic Test

$$\lim_{n \rightarrow \infty} n \log \left(\frac{u_n}{u_{n+1}} \right) = \kappa \quad (4.7)$$

- series converges for $\kappa < 1$
- series diverges for $\kappa > 1$
- test fails for $\kappa = 1$

5.1.1 Minor and Cofactor

For a third order determinant

$$\begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}$$

Minor

$$M(a_{11}) = M_{11} = \begin{vmatrix} b_2 & c_2 \\ b_3 & c_3 \end{vmatrix} \quad (5.2)$$

i.e., all the terms of determinant except those in the same row and columns as the one of which the minor is being calculated.

Cofactor

$$C_{ij} = (-1)^{i+j} M_{ij} \quad (5.3)$$

5.2 Properties of Determinants

1. Transposing a determinant does not alter its value.

$$\Delta = \Delta^T \quad (5.4)$$

2. If rows and columns are interchanges m times, the value of the new determinant is

$$\Delta' = (-1)^m \Delta \quad (5.5)$$

3. If two parallel lines are equal, then $\Delta = 0$

4. For $\Delta = \begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}$ and $\Delta_1 = \begin{vmatrix} ka_1 & kb_1 & kc_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}$, then $\Delta_1 = k\Delta$

5. For $\Delta = \begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}$ and $\Delta_1 = \begin{vmatrix} ka_1 & b_1 & c_1 \\ ka_2 & b_2 & c_2 \\ ka_3 & b_3 & c_3 \end{vmatrix}$, then $\Delta_1 = k\Delta$

6. For $C_n \rightarrow k_1 C_l + k_2 C_m + k_3 C_n$ or $R_n \rightarrow k_1 R_l + k_2 R_m + k_3 R_n$, $\Delta' = \Delta$

5.3 Cramer's Rule

For a system of equations:

$$a_1x + b_1y + c_1z = d_1$$

$$a_2x + b_2y + c_2z = d_2$$

$$a_3x + b_3y + c_3z = d_3$$

the following determinants are defined:

$$D = \begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}$$

$$D_x = \begin{vmatrix} d_1 & b_1 & c_1 \\ d_2 & b_2 & c_2 \\ d_3 & b_3 & c_3 \end{vmatrix}$$

$$D_y = \begin{vmatrix} a_1 & d_1 & c_1 \\ a_2 & d_2 & c_2 \\ a_3 & d_3 & c_3 \end{vmatrix}$$

$$D_z = \begin{vmatrix} a_1 & b_1 & d_1 \\ a_2 & b_2 & d_2 \\ a_3 & b_3 & d_3 \end{vmatrix}$$

The solution for the system of equations is:

$$x = \frac{D_x}{D} \tag{5.6}$$

$$y = \frac{D_y}{D} \tag{5.7}$$

$$z = \frac{D_z}{D} \tag{5.8}$$

5.3.1 Consistency Test

1. If $D \neq 0$, the system is consistent and has unique solutions.
2. If $D = D_x = D_y = D_z = 0$, the system may or may not be consistent and it will have infinite solutions and the system will be dependent.
3. If $D = 0$ and at least one of D_x, D_y, D_z is non zero, the system is inconsistent

6.2 Multiplication of Two Matrices

If

$$C_{m \times p} = A_{m \times n} \cdot B_{n \times p} \quad (6.2)$$

then,

$$c_{ik} = \sum_{j=1}^n a_{ij} \cdot b_{jk} \quad (6.3)$$

6.2.1 Multiplicative Properties

1. Multiplication of matrices is associative, hence $(AB)C = A(BC)$.
2. $AI = A$
3. $A \cdot A^{-1} = I$
4. $A \cdot (\text{adj } A) = (\text{adj } A) \cdot A = |A|I$
5. $A^{-1} = \frac{1}{|A|}(\text{adj } A)^t$
6. $(AB)^t = B^t A^t$

6.3 Adjoint of a Matrix

$$\text{adj } A = \begin{bmatrix} M_{11} & M_{12} & \cdots & M_{1n} \\ M_{21} & M_{22} & \cdots & M_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ M_{m1} & M_{m2} & \cdots & M_{mn} \end{bmatrix}^t, \text{ where } M_{ij} \text{ is the minor of } a_{ij} \quad (6.4)$$

6.4 Martin's Rule

For a system of equation,

$$\begin{aligned}a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n &= b_1 \\a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n &= b_2 \\&\vdots \\a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n &= b_n\end{aligned}$$

The system can be written as:

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix} \quad (6.5)$$

$$\Rightarrow AX = B \quad (6.6)$$

$$\Rightarrow X = A^{-1}B \quad (6.7)$$

7.4 Pascal's Rule

For $1 \leq k \leq n$ and $k, n \in \mathbb{N}$:

$$\binom{n}{k} + \binom{n}{k-1} = \binom{n+1}{k} \quad (7.7)$$

Chapter 8

Boolean Algebra

Let B be a set of a, b, c with operations sum $(+)$ and product (\cdot) .
Then B is said to belong to the Boolean Structure if the following conditions are satisfied:

Table 8.1: Properties of Boolean Algebraic Structure

Property	Name of Property
$a + b \in B$ $a \cdot b \in B$	Closure Property
$a + b = b + a$ $a \cdot b = b \cdot a$	Associative Law
$a(b + c) = ab + ac$ $a + bc = (a + b)(a + c)$	Commutative Law
$\{0, 1\} \in B$ $a + 0 = a$ $a + 1 = 1$ $a \cdot 0 = 0$ $a \cdot 1 = a$	Laws of 1 and 0
$a + ab = a$ $a(a + b) = a$	Absorption Law
$(a + b)' = (a'b')$	De'Morgan's Law

Chapter 9

Remainder Theorems

9.1 Remainder Theorem

If a function $f(x)$ is divided by a binomial $x - a$, then the remainder is provided by $f(a)$.

$$\frac{f(x)}{x - a} \equiv f(a) \pmod{(x - a)} \quad (9.1)$$

Worked Example

Find the remainder when $f(x) = x^3 - 4x^2 - 7x + 10$ is divided by $(x - 2)$.
The remainder:

$$R = (x^3 - 4x^2 - 7x + 10) \pmod{(x - 2)}$$

is given by:

$$\begin{aligned} R = f(2) &= (2)^3 - 4(2)^2 - 7(2) + 10 \\ &= 8 - 16 - 14 + 10 = -12 \end{aligned}$$

9.2 Euler's Remainder Theorem

According to Euler's Remainder Theorem, if x and n are two co-prime numbers:

$$x^{\varphi(n)} \equiv 1 \pmod{n}, x, n \in \mathbb{Z}^+ \quad (9.2)$$

where, $\varphi(n)$ is Euler's totient function.

9.2.1 Euler's Totient Function

For a number defined as:

$$n = \prod_{i=1}^r a_i^{b_i} \quad (9.3)$$

then Euler's totient function is defined as:

$$\begin{aligned}\varphi(n) &= n \cdot \left[\left(1 - \frac{1}{a_1}\right) \cdot \left(1 - \frac{1}{a_2}\right) \cdot \left(1 - \frac{1}{a_3}\right) \cdots \right] \\ &= n \prod_{i=1}^r \left(1 - \frac{1}{a_i}\right)\end{aligned}\tag{9.4}$$

Worked Example

Find the remainder if 3^{76} is divided by 35.

Since:

$$35 = 5^1 \times 7^1$$

Hence the totient quotient of 35 is:

$$\begin{aligned}\varphi(35) &= 35 \cdot \left(1 - \frac{1}{5}\right) \cdot \left(1 - \frac{1}{7}\right) \\ &= 35 \times \frac{4}{5} \times \frac{6}{7} \\ &= 24\end{aligned}$$

Hence Euler's Theorem yields:

$$\begin{aligned}3^{24} &\equiv 1 \pmod{35} \\ 3^{76} &\equiv 3^{24 \times 3 + 4} \\ &\equiv (3^{24})^3 \times 3^4 \pmod{35} \\ &\equiv (1)^3 \times 3^4 \pmod{35} \\ &\equiv 81 \pmod{35} \\ &\equiv 11 \pmod{35}\end{aligned}$$

The remainder when 3^{76} is divided by 35 is 11.

9.3 Wilson Theorem

According to Wilson Theorem:

$$(n-1)! \equiv -1 \pmod{n}\tag{9.5}$$

Worked Example

Find the remainder when $28!$ is divided by 31.

By Wilson's Theorem:

$$\begin{array}{ll} 30! & \equiv -1 \pmod{31} \\ \Rightarrow 30 \cdot 29 \cdot 28! & \equiv -1 \pmod{31} \\ \text{Let } 28! \pmod{31} & = x \\ \Rightarrow (-1) \cdot (-2) \cdot x & \equiv 30 \pmod{31} \\ \Rightarrow 2x & = 30 \\ \Rightarrow x & = 15 \end{array}$$

The remainder when $28!$ is divided by 31 is 15.

Part II

Co-ordinate Geometry

Chapter 10

2-D Co-ordinate Geometry

For the ordered pairs, $A(x_1, y_1)$ and $B(x_2, y_2)$:

10.1 Distance between Two Points

$$AB = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2} \quad (10.1)$$

(10.2)

10.2 Section Formula

If point C divides AB in the ratio $m : n$:

$$C = \left(\frac{nx_1 + mx_2}{m + n}, \frac{ny_1 + my_2}{m + n} \right) \quad (10.3)$$

10.2.1 Corollary: Mid - Point Formula

If C is the mid-point of AB , and $m : n = 1 : 1$:

$$C = \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right) \quad (10.4)$$

(10.5)

Chapter 11

Triangles

For a triangle defined with three vertices $A(x_1, y_1)$, $B(x_2, y_2)$, $C(x_3, y_3)$ and corresponding sides of length a, b, c , then:

11.1 Centroid of a Traiangle

$$\text{Centroid of } \triangle ABC = \left(\frac{x_1 + x_2 + x_3}{3}, \frac{y_1 + y_2 + y_3}{3} \right) \quad (11.1)$$

$$(11.2)$$

11.2 Area of Triangle

11.2.1 Determinant Method

$$\text{Area of } \triangle ABC = \frac{1}{2} \begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix} \quad (11.3)$$

11.2.2 Heron's Formula

For a triangle, the semi-perimeter, s , is defined as:

$$s = \frac{a + b + c}{2}$$

The area of the triangle can be defined as:

$$\text{Area of } \triangle ABC = \sqrt{s \cdot (s - a) \cdot (s - b) \cdot (s - c)} \quad (11.4)$$

11.3 Incircle of a Triangle

The radius, r , and centre of incircle, o , is:

$$o = \left(\frac{ax_1 + bx_2 + cx_3}{a + b + c}, \frac{ay_1 + by_2 + cy_3}{a + b + c} \right) \quad (11.5)$$

$$r = \sqrt{\frac{(s - a) \cdot (s - b) \cdot (s - c)}{s}} \quad (11.6)$$

$$(11.7)$$

11.4 Circumcircle of a Triangle

The radius, R , and centre, O , of circumcircle is defined as:

$$O = \left(\frac{x_1 \sin 2A + x_2 \sin 2B + x_3 \sin 2C}{\sin 2A + \sin 2B + \sin 2C}, \frac{y_1 \sin 2A + y_2 \sin 2B + y_3 \sin 2C}{\sin 2A + \sin 2B + \sin 2C} \right) \quad (11.8)$$

$$2R = \frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C} \quad (11.9)$$

Chapter 12

Straight Line

A straight line can be defined as:

$$y = mx + c \quad (12.1)$$

$$\frac{x}{a} + \frac{y}{b} = 1, \text{ where } a \text{ and } b \text{ are the intercepts at } x \text{ and } y \text{ axes respectively} \quad (12.2)$$

$$x \cos \alpha + y \sin \alpha = p \text{ (Normal Form)} \quad (12.3)$$

$$Ax + By + C = 0 \text{ (General Form)} \quad (12.4)$$

12.1 Equation of Straight Line Passing Through (x_0, y_0) and Slope m

$$(y - y_0) = m(x - x_0) \quad (12.5)$$

12.2 Distance Between Two Points on a Line

$$\frac{y_1 - y_2}{\sin \theta} = \frac{x_1 - x_2}{\cos \theta} = \gamma \quad (12.6)$$

$$\theta = \tan^{-1} m \quad (12.7)$$

12.3 Angle Between Two Lines

For two lines with slopes m_1, m_2 , the angle between them, θ :

$$\theta = \arctan \left(\frac{m_1 - m_2}{1 + m_1 m_2} \right) \quad (12.8)$$

12.4 Distance of a Point from a Line

Line: $ax + by + c = 0$ Point: (g, h)

$$S = \frac{ag + bh + c}{\sqrt{a^2 + b^2}} \quad (12.9)$$

12.5 Angle Bisector of a Line

For the two lines: $a_1x + b_1y + c_1 = 0$ and $a_2x + b_2y + c_2 = 0$, the angle bisector is:

$$\frac{a_1x + b_1y + c_1}{\sqrt{a_1^2 + b_1^2}} = \frac{a_2x + b_2y + c_2}{\sqrt{a_2^2 + b_2^2}} \quad (12.10)$$

If the sign of c_1 and c_2 is the same, then the equation obtained is the internal bisector.

12.6 Equation of a Straight Line Passing through the Intersection of Two Lines

$$(a_1x + b_1y + c_1) + k(a_2x + b_2y + c_2) = 0 \quad \forall k \in \mathbb{R} \quad (12.11)$$

12.7 Relative Position of Points w.r.t. a Line

For the points (x_1, y_1) and (x_2, y_2) :

$$\begin{aligned} k_1 &= ax_1 + by_1 + c \\ k_2 &= ax_2 + by_2 + c \end{aligned}$$

If k_1 and k_2 have the same sign, they are on the same side of a line, otherwise on opposite sides.

12.8 Ratio of Division of Line Segment

For any line, $f(x, y) = 0$, the ratio in which it divides (x_1, y_1) and (x_2, y_2) is given by:

$$r = -\frac{f(x_1, y_1)}{f(x_2, y_2)} \quad (12.12)$$

If $\begin{cases} r > 0, \text{ then division is internal} \\ r < 0, \text{ then division is external} \end{cases}$.

Chapter 13

General Theory of Second Degree Equation

For any general equation of the form:

$$ax^2 + by^2 + 2gx + 2fy + 2hxy + c = 0 \quad (13.1)$$

Δ is defined as:

$$\Delta = \begin{vmatrix} a & h & g \\ h & b & f \\ g & f & c \end{vmatrix} \quad (13.2)$$

If $\Delta = 0$ then the equation is a pair of straight lines. If $a + b = 0$ then the lines are \perp .

If the $\Delta \neq 0$:

1. $a = b, h = 0 \rightarrow$ circle
2. $h^2 = ab \rightarrow$ parabola
3. $h^2 < ab \rightarrow$ ellipse
4. $h^2 > ab \rightarrow$ hyperbola

Chapter 14

Conics

The four conic sections are: circle, parabola, ellipse, and hyperbola. Circle has been done separately in the next chapter.

14.1 Parametric Form of Conics

14.1.1 Hyperbola

$$x = a \sec \theta \quad (14.1)$$

$$y = b \tan \theta \quad (14.2)$$

14.1.2 Ellipse

$$x = a \cos \phi \quad (14.3)$$

$$y = b \sin \phi \quad (14.4)$$

14.1.3 Parabola

$$x = at^2 \quad (14.5)$$

$$y = 2at \quad (14.6)$$

14.2 Equation form of Conics

14.2.1 Parabola

Table 14.1: Properties of a Parabola

Property	$y^2 = 4ax$	$x^2 = 4ay$
Axis	$y = 0$	$x = 0$
Eccentricity	1	1
Directrix	$x + a = 0$	$y + a = 0$
Focus	$(a, 0)$	$(0, a)$
Vertex	$(0, 0)$	$(0, 0)$
Length of latus rectum	$ 4a $	$ 4a $
Equation of latus rectum	$x - a = 0$	$y - a = 0$

14.2.2 Ellipse and Hyperbola

For $a > b$:

Table 14.2: Properties of Ellipse and Hyperbola

Property	$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ Ellipse	$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ Hyperbola
Length of Major Axis	$2a$	$2a$
Length of Minor Axis	$2b$	$2b$
Equation of Major Axis	$x = 0$	$x = 0$
Equation of Minor Axis	$y = 0$	$y = 0$
Eccentricity e	$\sqrt{1 - \frac{b^2}{a^2}}$	$\sqrt{1 + \frac{b^2}{a^2}}$
Vertices	$(\pm a, 0)$	$(\pm a, 0)$
Foci	$(\pm ae, 0)$	$(\pm ae, 0)$
Equation of Directrix	$x \pm \frac{a}{e} = 0$	$x = \pm \frac{a}{e}$
Length of latus rectum	$\frac{2b^2}{a}$	$\frac{2b^2}{a}$
Equation of latus rectum	$x \pm ae = 0$	
Centre	$(0, 0)$	$(0, 0)$

Chapter 15

Circles

15.1 Locus Form

$$(x - g)^2 + (y - h)^2 = r^2 \quad (15.1)$$

where the centre is (g, h) and the radius is r .

15.2 Diameter Form

$$(x - a)(x - c) + (y - b)(y - d) = 0 \quad (15.2)$$

where (a, b) and (c, d) are the two ends of the diameter.

15.3 General Form

If the equation of a circle is in the form:

$$x^2 + y^2 + 2gx + 2fy + c = 0 \quad (15.3)$$

Then the following is true about the circle:

1. centre of the circle is $(-g, -f)$
2. radius of circle is $\sqrt{g^2 + f^2 - c}$

15.4 Important Relations

1. If the circle passes through the origin, $g = 0, f = 0$.
2. If the circle touches the x-axis $c = g^2$.
3. If the circle touches the y-axis $c = f^2$.

15.5 Common for Two Circles

1. The common chord passing between two circles S_1 and S_2 are:

$$S_1 - S_2 = 0 \tag{15.4}$$

2. Circles passing through the intersection of two circles is:

$$S_2 + k(S_1 - S_2) = 0 \quad \forall k \in \mathbb{R} \tag{15.5}$$

Chapter 16

Vectors

Let two vectors be $\vec{a} = a\hat{i} + b\hat{j} + c\hat{k}$ and $\vec{b} = x\hat{i} + y\hat{j} + z\hat{k}$:

16.1 Modulus of a Vector

For a vector \vec{a} , the modulus of the vector is:

$$|\vec{a}| = \sqrt{a^2 + b^2 + c^2} \quad (16.1)$$

16.2 Sum of Vectors

The sum of two vectors is:

$$|\vec{a} + \vec{b}| = \sqrt{|\vec{a}|^2 + |\vec{b}|^2 + 2|\vec{a}||\vec{b}|\cos\theta} \quad (16.2)$$

$$\vec{a} + \vec{b} = (a + x)\hat{i} + (b + y)\hat{j} + (c + z)\hat{k} \quad (16.3)$$

The direction of the resultant vector is:

$$\tan\alpha = \frac{b\sin\theta}{a + b\cos\theta} \quad (16.4)$$

where, θ is the angle between the two vectors.

16.3 Product of Vectors

16.3.1 Dot Product

$$\vec{a} \cdot \vec{b} = |a||b|\cos\theta \quad (16.5)$$

$$\vec{a} \cdot \vec{b} = ax + by + cz \quad (16.6)$$

16.3.2 Cross Product

$$\vec{a} \times \vec{b} = |a||b|\sin\theta\hat{n} \quad (16.7)$$

$$(16.8)$$

where \hat{n} is a vector $\perp \vec{a}, \vec{b}$.

$$\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a & b & c \\ x & y & z \end{vmatrix} \quad (16.9)$$

16.4 Test of Co-planarity

Three vectors are called co-planar if:

$$\lambda \vec{a} + \mu \vec{b} = \vec{c} \quad (16.10)$$

$$(\vec{a} \times \vec{b}) \cdot \vec{c} = 0 \quad (16.11)$$

Chapter 17

3D - Space

17.1 Line segments in 3D - Space

For points defined in a 3D space as $A(x_1, y_1, z_1)$ and $B(x_2, y_2, z_2)$:

17.1.1 Distance between Two Points

$$AB = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2 + (z_1 - z_2)^2} \quad (17.1)$$

17.1.2 Section Formula of a Line Segment Divided in the ratio $m : n$

$$P = \left(\frac{nx_1 + mx_2}{m + n}, \frac{ny_1 + my_2}{m + n}, \frac{nz_1 + mz_2}{m + n} \right) \quad (17.2)$$

17.2 Line in 3D - Space

For a line which is defined as $\vec{a} = a\hat{i} + b\hat{j} + c\hat{k}$:

1. Line numbers of the line is

$$< a, b, c > \quad (17.3)$$

2. The line cosines are:

$$< \frac{a}{\sqrt{a^2 + b^2 + c^2}}, \frac{b}{\sqrt{a^2 + b^2 + c^2}}, \frac{c}{\sqrt{a^2 + b^2 + c^2}} > \quad (17.4)$$

$$= < l, m, n > \quad (17.5)$$

17.2.1 Angle between Two Lines

$$\cos \theta = \frac{a_1 a_2 + b_1 b_2 + c_1 c_2}{\sqrt{a_1^2 + b_1^2 + c_1^2} \sqrt{a_2^2 + b_2^2 + c_2^2}} \quad (17.6)$$

$$\Rightarrow \cos \theta = l_1 l_2 + m_1 m_2 + n_1 n_2 \quad (17.7)$$

When two lines are \perp , $l_1 l_2 + m_1 m_2 + n_1 n_2 = 0$.

When two lines are \parallel $\frac{l_1}{l_2} = \frac{m_1}{m_2} = \frac{n_1}{n_2} = 1$.

17.2.2 Skew and Co-planar Lines

Let there be two lines r_1 and r_2 ,

$$r_1 = a_1 + \mu b_1 r_2 = a_2 + \lambda b_2 \quad (17.8)$$

17.2.3 Distance between Lines

The shortest distance between r_1 and r_2

$$S = \left| \frac{(\vec{a}_1 - \vec{a}_2) \cdot (\vec{b}_1 \times \vec{b}_2)}{|\vec{b}_1 \times \vec{b}_2|} \right| \quad (17.9)$$

If $S = 0$, the lines intersect.

Cartesian Form

For two lines defined as $\frac{x - x_1}{a_1} = \frac{y - y_1}{b_1} = \frac{z - z_1}{c_1}$ and $\frac{x - x_2}{a_2} = \frac{y - y_2}{b_2} = \frac{z - z_2}{c_2}$:

$$S = \begin{vmatrix} x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} \quad (17.10)$$

Distance Between Parallel Lines

$$S = \left| \frac{\vec{b} \cdot (\vec{a}_2 - \vec{a}_1)}{|\vec{b}|} \right| \quad (17.11)$$

Distance of a Point to a Line

For a point, (x_1, y_1, z_1) the distance to a line $\frac{x - \alpha}{l} = \frac{y - \beta}{m} = \frac{z - \gamma}{n}$:

$$S = \left(\left| \frac{x_1 - \alpha}{l} - \frac{y_1 - \beta}{m} \right| + \left| \frac{y_1 - \beta}{m} - \frac{z_1 - \gamma}{n} \right| + \left| \frac{z_1 - \gamma}{n} - \frac{x_1 - \alpha}{l} \right| \right)^{\frac{1}{2}} \quad (17.12)$$

17.3 Triangular Plane

17.3.1 Centroid of a Triangle

$$G = \left(\frac{x_1 + x_2 + x_3}{3}, \frac{y_1 + y_2 + y_3}{3}, \frac{z_1 + z_2 + z_3}{3} \right) \quad (17.13)$$

Chapter 18

3D - Plane

A plane in 3-D space can be defined as:

1. Cartesian Form:

$$ax + by + cz + d = 0 \quad (18.1)$$

2. Vectorial Form:

$$\vec{r} \cdot \vec{n} = p \quad (18.2)$$

, where \vec{r} is a line on the plane, \vec{n} is a normal to the plane, and p is perpendicular distance to the plane from the origin.

18.1 Angle Between Two Planes

For two planes, $\vec{r}_1 \cdot \vec{n}_1 = p_1$ and $\vec{r}_2 \cdot \vec{n}_2 = p_2$, the angle between the planes, θ is:

$$\cos \theta = \frac{\vec{n}_1 \cdot \vec{n}_2}{|\vec{n}_1||\vec{n}_2|} \quad (18.3)$$

In the Cartesian Form:

$$\cos \theta = \frac{a_1 a_2 + b_1 b_2 + c_1 c_2}{\sqrt{a_1^2 + b_1^2 + c_1^2} \sqrt{a_2^2 + b_2^2 + c_2^2}} \quad (18.4)$$

18.2 Distance of a Point from a Plane

18.2.1 Cartesian Form

For the point (p, q, r) and the plane, $ax + by + cz + d = 0$:

$$S = \frac{ap + bq + cr + d}{\sqrt{a^2 + b^2 + c^2}} \quad (18.5)$$

18.2.2 Vector Form

For the point $\vec{g} = p\hat{i} + q\hat{j} + r\hat{k}$ and the plane $\vec{r} \cdot (a\hat{i} + b\hat{j} + c\hat{k}) + d = 0$:

$$S = \frac{(a\hat{i} + b\hat{j} + c\hat{k}) \cdot (p\hat{i} + q\hat{j} + r\hat{k})}{\sqrt{a^2 + b^2 + c^2}} \quad (18.6)$$

$$\Rightarrow S = \frac{(a\hat{i} + b\hat{j} + c\hat{k}) \cdot \vec{g}}{|a\hat{i} + b\hat{j} + c\hat{k}|} \quad (18.7)$$

Part III

Statistics

Chapter 19

Descriptive Statistics

19.1 Measure of Location

19.1.1 Mean

$$\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i \quad (19.1)$$

19.1.2 Median

For odd number of elements in a dataset:

$$\tilde{x} = x_{\frac{n+1}{2}} \quad (19.2)$$

For even number of elements in a dataset:

$$\tilde{x} = \frac{x_{\frac{n}{2}} + x_{(\frac{n}{2}+1)}}{2} \quad (19.3)$$

19.1.3 Mode

$$Mo(x) = \max(f(x_i)) \quad (19.4)$$

19.1.4 Quartile

Measure of percentage of elements less than or equal to a term

19.2 Measure of Spread

19.2.1 Variance

Variance measured on the whole population

$$\sigma^2 = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2 \quad (19.5)$$

19.2.2 Sample Variance

Variance measured on a sample population

$$s^2 = \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2 \quad (19.6)$$

19.2.3 Standard Deviation and Sample Standard

$$\sigma = \sqrt{\sigma^2} \quad (19.7)$$

$$s = \sqrt{s^2} \quad (19.8)$$

19.2.4 Co-efficient of Variance

$$v = \frac{s}{\bar{x}} \quad (19.9)$$

19.3 Skewness

19.3.1 Types of Skewness

Name	Other Name	Characteristic
Right Skew	Positive Skew	Data concentrated on the lower side
Symmetric Distribution	Normal Distribution	Data distributed evenly
Left Skew	Negative Skew	Data concentrated on the higher side

19.3.2 Measure of Skewness

Skewness is measured by the Moment Co-efficient of Skewness.

$$g_m = \frac{m_3}{s^3}, \text{ where} \quad (19.10)$$

$$m_3 = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^3 \quad (19.11)$$

Type of Skewness

The type of skewness from the value is g_m is:

Value of g_m	Type
$g_m = 0$	Symmetric
$g_m > 0$	Positive Skew
$g_m < 0$	Negative Skew

Value of g_m	Degree
$ g_m > 1$	High Skewness
$0.5 < g_m \leq 1$	Moderate Skewness
$ g_m \leq 0.5$	Low Skewness

Degree of Skewness

The degree of skewness from the value is g_m is:

19.4 Kurtosis

Kurtosis is the measure of peakedness of data. Fisher's kurtosis measure is defined as:

$$\gamma = \frac{m_4}{s^4}, \text{ where} \quad (19.12)$$

$$m_4 = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^4 \quad (19.13)$$

19.4.1 Type of Kurtosis

The types of kurtosis from the value of γ are:

Value of γ	Type
$\gamma = 0$	Normal Distribution or Mesokurtic
$\gamma < 0$	Flattened or Platykurtic
$\gamma > 0$	Peaked or Lepokurtic

Chapter 20

Hypothesis Testing

20.1 T-Test

$$T = \frac{\bar{X} - \mu}{\frac{s}{\sqrt{n}}} \quad (20.1)$$

where:

\bar{X} = Sample Mean

μ = Assumed Mean

s = Number of Samples

n = Number of observations

If $T < t_c$ the H_0 is not rejected. t_c is a functions of level of significance (α) and degrees of freedom ($v = n - 1$).

20.2 χ^2 Test

$$\chi^2 = \sum_i \sum_j \frac{(h_{ij}^o - h_{ij}^e)^2}{h_{ij}^e} \quad (20.2)$$

where:

h_e = Expected Value

h_o = Actual Value

If $\chi^2 < \chi_c^2$ then H_0 is not rejected. χ_c is a functions of level of significance (α) and degrees of freedom ($v = (i - 1)(j - 1)$).

Chapter 21

Research and Survey Design

21.1 Population Covariance

$$\text{Cov}(x, y) = \frac{1}{n} \sum_{i=1}^n (x_i - \mu_x)(y_i - \mu_y) \quad (21.1)$$

21.2 Sample Covariance

$$\text{Cov}(x, y) = \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y}) \quad (21.2)$$

21.3 Bravais-Pearson Correlation Co-efficient

$$r = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sqrt{\sum_{i=1}^n (x_i - \bar{x})^2} \cdot \sqrt{\sum_{i=1}^n (y_i - \bar{y})^2}} \quad (21.3)$$

$$= \frac{\text{Cov}(x, y)}{\sqrt{\text{Var}(x) \cdot \text{Var}(y)}} \quad (21.4)$$

$$= \frac{\text{Cov}(x, y)}{\sigma_x \cdot \sigma_y} \quad (21.5)$$

21.4 Spearman's Rank Correlation Co-efficient

$$r_s = 1 - \frac{6 \sum d_i^2}{n(n^2 - 1)} \quad (21.6)$$

$$d_i = R(X_i) - R(Y_i) \quad (21.7)$$

Chapter 22

Estimation of Regression Function

For the regression functions:

$$Y_i = \beta_0 + \beta_1 X_1 \quad (22.1)$$

$$\hat{Y}_i = \hat{\beta}_0 + \hat{\beta}_1 X_1 \quad (22.2)$$

$$(22.3)$$

where Y_i is the observed dependent variable (DV), \hat{Y}_i is the estimated DV, and X_i is the independent variable (IV).

$$u_i = Y_i - \hat{Y}_i \quad (22.4)$$

$$\Rightarrow Y_i = \hat{Y}_i + u_i \quad (22.5)$$

$$\Rightarrow Y_i = \hat{\beta}_0 + \hat{\beta}_1 X_1 + u_i \quad (22.6)$$

$$Y_i = \beta_0 + \beta_1 X_i + \epsilon_i \quad (22.7)$$

The objective function is:

$$\min_{u_i} \sum u_i = \min \sum_i \left[Y_i - \hat{\beta}_0 - \hat{\beta}_1 X_i \right]^2$$

Since the regression function passes through: (\bar{X}, \bar{Y})

$$\beta_0 = \bar{Y} - \hat{\beta}_1 \bar{X}$$

$$\begin{aligned} \min_{u_i} \sum u_i &= \min \sum_i \left[Y_i - \bar{Y} + \hat{\beta}_1 \bar{X} - \hat{\beta}_1 X_i \right]^2 \\ &= \min \sum_i \left[(Y_i - \bar{Y}) - \hat{\beta}_1 (X_i - \bar{X}) \right]^2 \\ &= \min \sum_i \left[(Y_i - \bar{Y})^2 - 2 \cdot (Y_i - \bar{Y}) \cdot \hat{\beta}_1 (X_i - \bar{X}) + \hat{\beta}_1^2 (X_i - \bar{X})^2 \right] \\ &= \min \left[\sum_i (Y_i - \bar{Y})^2 - 2 \cdot \hat{\beta}_1 \sum_i (Y_i - \bar{Y}) \cdot (X_i - \bar{X}) + \hat{\beta}_1^2 \sum_i (X_i - \bar{X})^2 \right] \\ \Rightarrow u_i^{\beta_1} &= -2 \sum_i (Y_i - \bar{Y}) + 2 \hat{\beta}_1 \sum_i (X_i - \bar{X})^2 = 0 \quad (\text{For optima Conditions}) \end{aligned}$$

$$\Rightarrow \hat{\beta}_1 = \frac{\sum_i (Y_i - \bar{Y})(X_i - \bar{X})}{\sum_i (X_i - \bar{X})^2}$$

$$\Rightarrow \hat{\beta}_0 = \bar{Y} - \hat{\beta}_1 \bar{X}$$

22.1 Sum of Squares Error

$$TSS = \sum_i (Y_i - \bar{Y})^2 \quad (22.8)$$

$$= \underbrace{\sum_i (\hat{Y}_i - \bar{Y})^2}_{\text{Explained Sum of Square Error (ESS)}} + \underbrace{\sum_i u_i^2}_{\text{Residual Sum of Squares Error (RSS)}} \quad (22.9)$$

22.1.1 R^2 : Coefficient of Determination

$$R^2 = \frac{ESS}{TSS} \quad (22.10)$$

$$= 1 - \frac{RSS}{TSS} \quad (22.11)$$

$$= 1 - \frac{\sum_i u_i^2}{\sum_i (Y_i - \bar{Y})^2} \quad (22.12)$$

For a regression analysis with single IV:

$$\sqrt{R^2} = v$$

22.1.2 \bar{R}^2 : Coefficient of Determination

$$\bar{R}^2 = 1 - \frac{\frac{\sum_i u_i^2}{(N - K - 1)}}{\frac{\sum_i (Y_i - \bar{Y})^2}{(N - 1)}} \quad (22.13)$$

where, N is the number of observations and K is the number of independent variables.

22.2 T-Test

Test for statistical significance of a single IV.

$$T = \frac{\hat{\beta}_1 - 0}{S_e(\hat{\beta}_1)} \quad (22.14)$$

22.3 F-Test

Test for statistical significance of all IVs together.

$$F = \frac{\frac{\text{ESS}}{(K - 1)}}{\frac{\text{RSS}}{(N - K)}} \quad (F \geq F_c, H_0 \text{ is rejected})$$

22.4 Test for Heteroskedasticity**22.4.1 Definition**

$$\sigma_{\epsilon_i} \forall \epsilon_i \in [X_a, X_b] = \sigma_{\epsilon_i} \forall \epsilon_i \in [X_{b+1}, X_c]$$

22.4.2 Durbin-Watson Test

$$d_e = \frac{\sum_{t=2}^n (\hat{u}_t - \hat{u}_{t-1})^2}{\sum_{t=2}^n \hat{u}_t^2} \quad (22.15)$$

For the H_0 : No autocorrelation:

d	H_0
$0 \leq d_e \leq d_L$ & $(4 - d_L) \leq d_e \leq 4$	Rejected
$d_L < d_e \leq d_U$ & $(4 - d_U) < d_e \leq (4 - d_L)$	Decision Free Zone
$d_L < d_e < D_U$	Not rejected


Chapter 23

Dummy Variables

23.1 Dummy Variable

$$P_i = \beta_0 + \beta_1 S_i + \beta_2 D_i + \epsilon_i \quad (23.1)$$


$$E(P_i) = \begin{cases} (\hat{\beta}_0 + \hat{\beta}_2) + \hat{\beta}_1 S_i, & D_i = 1 \\ \hat{\beta}_0 + \hat{\beta}_1 S_i, & D_i = 0 \end{cases} \quad (23.2)$$



23.2 Slope Dummy Variable

$$P_i = \beta_0 + \beta_1 S_i + \beta_2 (S_i \cdot D_i) + \epsilon_i \quad (23.3)$$


$$E(P_i) = \begin{cases} \hat{\beta}_0 + (\hat{\beta}_1 + \hat{\beta}_2) S_i, & D_i = 1 \\ \hat{\beta}_0 + \hat{\beta}_1 S_i, & D_i = 0 \end{cases} \quad (23.4)$$



23.3 Slope & Dummy Variable

$$P_i = \beta_0 + \beta_1 S_i + \beta_2 D_i + \beta_3 S_i D_i + \epsilon_i \tag{23.5}$$

$$E(P_i) = \begin{cases} (\hat{\beta}_0 + \hat{\beta}_2) + (\hat{\beta}_1 + \hat{\beta}_3) S_i, & D_i = 1 \\ \hat{\beta}_0 + \hat{\beta}_1 S_i, & D_i = 0 \end{cases} \tag{23.6}$$



23.4 Multi-Categories Dummy Variable

$$P_0 = b_0 \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} + b_1 \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} + b_2 \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} + b_3 \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \quad (23.7)$$

Leads to Perfect Multicollinearity

Alternatives

- B_n captures the mean of each category, but F-Test is impossible

$$y = \beta_1 D_{1i} + \beta_2 D_{2i} + \beta_3 D_{3i} \quad (23.8)$$

- Computer drops automatically drops a variable

$$y = \beta_0 + \beta_1 D_{1i} + \beta_2 D_{2i} + \beta_3 D_{3i} \quad (23.9)$$

- Manually dropping a variable

$$y = \beta_0 + \beta_1 D_{1i} + \beta_2 D_{2i} \quad (23.10)$$

Chapter 24

Logistic Regression

For $Y_i \in \{0, 1\}$:

$$z_k = \beta_0 + \sum_{j=1}^n \beta_{jk} x_j + \epsilon_k, \beta_j \rightarrow \text{Logit Coefficient} \quad (24.1)$$

$$p = \frac{\exp^k}{1 + \exp^k} = \frac{1}{1 + \exp^{-k}} \quad (24.2)$$

where p is the probability of $y = 1$.

Part IV

Trigonometry

Chapter 25

Circular Trigonometric Functions

25.1 Trigonometric Ratios of Standard Angles

Table 25.1: Trigonometric Ratios of Standard Angles

θ	$\sin \theta$	$\cos \theta$	$\tan \theta$
0°	0	1	0
15°	$\frac{1}{4}$	$\frac{1}{4(2-\sqrt{3})}$	$2 - \sqrt{3}$
18°	$\frac{\sqrt{5}-1}{4}$	$\frac{\sqrt{10+2\sqrt{5}}}{4}$	$\frac{\sqrt{5}-1}{\sqrt{10+2\sqrt{5}}}$
30°	$\frac{1}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{1}{\sqrt{3}}$
36°	$\frac{\sqrt{5}+1}{4}$	$\frac{\sqrt{10-2\sqrt{5}}}{4}$	$\frac{\sqrt{5}+1}{\sqrt{10-2\sqrt{5}}}$
45°	$\frac{1}{\sqrt{2}}$	$\frac{1}{\sqrt{2}}$	1
60°	$\frac{1}{\sqrt{3}}$	$\frac{1}{2}$	$\sqrt{3}$
72°	$\frac{\sqrt{10+2\sqrt{5}}}{4}$	$\frac{\sqrt{5}-1}{4}$	$\frac{\sqrt{10+2\sqrt{5}}}{\sqrt{5}-1}$
90°	1	0	∞

For any given triangle:

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C} = 2R \quad (25.1)$$

, where R is the radius of circumcircle. Refer to 11.4.

25.2 Negative Angle Formula

$$\sin(-\theta) = -\sin \theta \quad (25.2)$$

$$\cos(-\theta) = \cos \theta \quad (25.3)$$

$$\tan(-\theta) = -\tan \theta \quad (25.4)$$

$$\csc(-\theta) = -\csc \theta \quad (25.5)$$

$$\sec(-\theta) = \sec \theta \quad (25.6)$$

$$\cot(-\theta) = -\cot \theta \quad (25.7)$$

25.3 Sum of Angles Formula

$$\sin(A + B) = \sin A \cos B + \cos A \sin B \quad (25.8)$$

$$\cos(A + B) = \cos A \cos B - \sin A \sin B \quad (25.9)$$

$$\tan(A + B) = \frac{\tan A + \tan B}{1 - \tan A \tan B} \quad (25.10)$$

25.4 Difference of Angles Formula

$$\sin(A - B) = \sin A \cos B - \cos A \sin B \quad (25.11)$$

$$\cos(A - B) = \cos A \cos B + \sin A \sin B \quad (25.12)$$

$$\tan(A - B) = \frac{\tan A - \tan B}{1 + \tan A \tan B} \quad (25.13)$$

25.5 Multiples and Sub-multiples of π and $\frac{\pi}{2}$

$$\forall k \in \mathbb{Z}$$

$$\sin \left[(4k + 1) \frac{\pi}{2} \right] = 1 \quad (25.14)$$

$$\sin \left[(4k - 1) \frac{\pi}{2} \right] = -1 \quad (25.15)$$

$$\sin k\pi = 0 \quad (25.16)$$

$$\sin \left[(2k + 1) \frac{\pi}{2} \right] = 0 \quad (25.17)$$

$$\sin \left[(2k - 1) \frac{\pi}{2} \right] = 0 \quad (25.18)$$

$$\sin k\pi = (-1)^k \quad (25.19)$$

25.6 $\left(\frac{\pi}{2} \pm \theta\right)$ Formula

$$\sin\left(\frac{\pi}{2} - \theta\right) = \cos \theta \quad (25.20)$$

$$\sin\left(\frac{\pi}{2} + \theta\right) = \cos \theta \quad (25.21)$$

$$\cos\left(\frac{\pi}{2} - \theta\right) = \sin \theta \quad (25.22)$$

$$\cos\left(\frac{\pi}{2} + \theta\right) = -\sin \theta \quad (25.23)$$

$$\tan\left(\frac{\pi}{2} - \theta\right) = \cot \theta \quad (25.24)$$

$$\tan\left(\frac{\pi}{2} + \theta\right) = -\cot \theta \quad (25.25)$$

$$\cot\left(\frac{\pi}{2} - \theta\right) = \tan \theta \quad (25.26)$$

$$\cot\left(\frac{\pi}{2} + \theta\right) = -\tan \theta \quad (25.27)$$

$$\csc\left(\frac{\pi}{2} - \theta\right) = \sec \theta \quad (25.28)$$

$$\csc\left(\frac{\pi}{2} + \theta\right) = \sec \theta \quad (25.29)$$

$$\sec\left(\frac{\pi}{2} - \theta\right) = \csc \theta \quad (25.30)$$

$$\sec\left(\frac{\pi}{2} + \theta\right) = -\csc \theta \quad (25.31)$$

25.7 $\left(\frac{\pi}{4} \pm \theta\right)$ Formula

$$\tan\left(\frac{\pi}{4} + \theta\right) = \frac{1 + \tan \theta}{1 - \tan \theta} \quad (25.32)$$

$$\tan\left(\frac{\pi}{4} - \theta\right) = \frac{1 - \tan \theta}{1 + \tan \theta} \quad (25.33)$$

25.8 Trigonometric Identities

$$\sin^2 \theta + \cos^2 \theta = 1 \quad (25.34)$$

$$\tan^2 \theta + 1 = \sec^2 \theta \quad (25.35)$$

$$\cot^2 \theta + 1 = \csc^2 \theta \quad (25.36)$$

25.9 Double Angle Formula

$$\sin 2\theta = 2 \sin \theta \cos \theta \quad (25.37)$$

$$= \frac{2 \tan \theta}{1 + \tan^2 \theta} \quad (25.38)$$

$$\cos 2\theta = \cos^2 \theta - \sin^2 \theta \quad (25.39)$$

$$= 2 \cos^2 \theta - 1 \quad (25.40)$$

$$= 1 - 2 \sin^2 \theta \quad (25.41)$$

$$= \frac{1 - \tan^2 \theta}{1 + \tan^2 \theta} \quad (25.42)$$

$$\tan 2\theta = \frac{2 \tan \theta}{1 - \tan^2 \theta} \quad (25.43)$$

25.10 Triple Angle Formula

$$\sin 3\theta = 3 \sin \theta - 4 \sin^3 \theta \quad (25.44)$$

$$\cos 3\theta = 4 \cos^3 \theta - 3 \cos \theta \quad (25.45)$$

$$\tan 3\theta = \frac{3 \tan \theta - \tan^3 \theta}{1 - 3 \tan^2 \theta} \quad (25.46)$$

25.11 Sum and Product of Two Ratios

For $A > B$:

$$\sin A + \sin B = 2 \sin \left(\frac{A+B}{2} \right) \cos \left(\frac{A-B}{2} \right) \quad (25.47)$$

$$\sin A - \sin B = 2 \cos \left(\frac{A+B}{2} \right) \sin \left(\frac{A-B}{2} \right) \quad (25.48)$$

$$2 \sin A \cos B = \sin(A+B) + \sin(A-B) \quad (25.49)$$

$$2 \cos A \sin B = \sin(A+B) - \sin(A-B) \quad (25.50)$$

$$\cos A + \cos B = 2 \cos \left(\frac{A+B}{2} \right) \cos \left(\frac{A-B}{2} \right) \quad (25.51)$$

$$\cos A - \cos B = -2 \sin \left(\frac{A+B}{2} \right) \sin \left(\frac{A-B}{2} \right) \quad (25.52)$$

$$2 \cos A \cos B = \cos(A+B) + \cos(A-B) \quad (25.53)$$

$$2 \cos A \sin B = \cos(A+B) - \cos(A-B) \quad (25.54)$$

$$\sin(A-B) \sin(A+B) = \sin^2 A - \sin^2 B \quad (25.55)$$

$$\cos(A-B) \cos(A+B) = \cos^2 A - \sin^2 B \quad (25.56)$$

$$\tan(A-B) \tan(A+B) = \frac{\tan^2 A - \tan^2 B}{1 - \tan^2 A \tan^2 B} \quad (25.57)$$

25.12 General Solutions

If $\sin \theta = \sin \alpha$:

$$\theta = n\pi + (-1)^n \alpha \quad (25.58)$$

$n \in \mathbb{Z}$

If $\cos \theta = \cos \alpha$:

$$\theta = 2n\pi \pm \alpha \quad (25.59)$$

$n \in \mathbb{Z}$

If $\tan \theta = \tan \alpha$:

$$\theta = n\pi \pm \alpha \quad (25.60)$$

$n \in \mathbb{Z}$

25.13 Taylor Series Expansion of Trigonometric Ratios

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \cdots \infty = \sum_{i=1}^{\infty} (-1)^{i+1} \frac{x^{2i-1}}{(2i-1)!} \quad (25.61)$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \cdots \infty = \sum_{i=0}^{\infty} (-1)^i \frac{x^{2i}}{(2i)!} \quad (25.62)$$

Chapter 26

Inverse Circular Trigonometric Function

26.1 Definition of Inverse Circular Trigonometric Function

26.1.1 For $\sin x$

$y = \arcsin x$ iff $x = \sin y$, then:

1. $y \in [-\frac{\pi}{2}, \frac{\pi}{2}]$
2. domain of $x \in [-1, 1]$
3. $\sin(\arcsin x) = x, \forall x \in [-1, 1]$
4. $\arcsin(\sin y) = y, \forall y \in [-\frac{\pi}{2}, \frac{\pi}{2}]$
5. $\sin x$ is a strictly increasing in the domain $[-\frac{\pi}{2}, \frac{\pi}{2}]$ and one-one.

26.1.2 For $\cos x$

$y = \arccos x$ iff $x = \cos y$, then:

1. $y \in [0, \pi]$
2. domain of $x \in [-1, 1]$
3. $\cos(\arccos x) = x, \forall x \in [-1, 1]$
4. $\arccos(\cos y) = y, \forall y \in [0, \pi]$
5. $\cos x$ is a strictly decreasing in the domain $[0, \pi]$ and one-one.

26.1.3 For $\tan x$

$y = \arctan x$ iff $x = \tan y$, then:

1. $y \in (-\frac{\pi}{2}, \frac{\pi}{2})$
2. domain of $x \in \mathbb{R}$
3. $\tan(\arctan x) = x, \forall x \in \mathbb{R}$

4. $\arctan(\tan y) = y, \forall y \in [-\frac{\pi}{2}, \frac{\pi}{2}]$
5. $\tan x$ is a strictly increasing in the domain $[-\frac{\pi}{2}, \frac{\pi}{2}]$ and one-one.

26.1.4 For $\cot x$

$y = \cot^{-1} x$ iff $x = \cot y$, then:

1. $y \in (0, \pi)$
2. domain of $x \in \mathbb{R}$
3. $\cot(\cot^{-1} x) = x, \forall x \in \mathbb{R}$
4. $\cot^{-1}(\cot y) = y, \forall y \in (0, \pi)$
5. $\cot x$ is a strictly decreasing in the domain $(0, \pi)$ and one-one.

For $\sec x$

$y = \sec^{-1} x$ iff $x = \sec y$

1. $y \in \{[0, \frac{\pi}{2}) \cup (\frac{\pi}{2}, \pi]\}$
2. $x \in \{(-\infty, -1] \cup [1, \infty)\}$
3. $\sec(\sec^{-1} x) = x, \forall |x| \geq 1$
4. $\sec^{-1}(\sec y) = y, \forall y \in \{[0, \frac{\pi}{2}) \cup (\frac{\pi}{2}, \pi]\}$

26.1.5 For $\csc x$

$y = \csc^{-1} x$ iff $x = \csc y$

1. $y \in \{[-\frac{\pi}{2}, 0) \cup (0, \frac{\pi}{2}]\}$
2. $x \in \{(-\infty, -1] \cup [1, \infty)\}$
3. $\csc(\csc^{-1} x) = x, \forall |x| \geq 1$
4. $\csc^{-1}(\csc y) = y, \forall y \in \{[-\frac{\pi}{2}, 0) \cup (0, \frac{\pi}{2}]\}$

26.2 Negative Arguments

$$\arcsin(-x) = -\arcsin x \quad (26.1)$$

$$\arctan(-x) = -\arctan x \quad (26.2)$$

$$\csc^{-1}(-x) = -\csc^{-1} x \quad (26.3)$$

$$\arccos(-x) = \pi - \arccos x \quad (26.4)$$

$$\cot^{-1}(-x) = \pi - \cot^{-1} x \quad (26.5)$$

$$\sec^{-1}(-x) = \pi - \sec^{-1} x \quad (26.6)$$

26.3 Reciprocal Relations

$$\csc^{-1} x = \arcsin \frac{1}{x} \quad (26.7)$$

$$\sec^{-1} x = \arccos \frac{1}{x} \quad (26.8)$$

$$\sec^{-1} x = \begin{cases} \arctan \frac{1}{x}, & x > 0 \\ \pi + \arctan \frac{1}{x}, & x < 0 \end{cases} \quad (26.9)$$

26.4 I.T.F. Identities

$$\arcsin x + \arccos x = \frac{\pi}{2}, \quad |x| \leq 1 \quad (26.10)$$

$$\arctan x + \cot^{-1} x = \frac{\pi}{2}, \quad x \in \mathbb{R} \quad (26.11)$$

$$\sec^{-1} x + \csc^{-1} x = \frac{\pi}{2}, \quad |x| \geq 1 \quad (26.12)$$

26.5 Sum of Two Angles

$$\arctan x + \arctan y = \arctan \left(\frac{x+y}{1-xy} \right) \quad (26.13)$$

$$\arcsin x + \arcsin y = \arcsin \left(y\sqrt{1-x^2} + x\sqrt{1-y^2} \right) \quad (26.14)$$

$$\arccos x + \arccos y = \arccos \left(xy - \sqrt{1-x^2}\sqrt{1-y^2} \right) \quad (26.15)$$

26.6 Difference of Two Angles

$$\arctan x - \arctan y = \arctan \left(\frac{x-y}{1+xy} \right) \quad (26.16)$$

$$\arcsin x - \arcsin y = \arcsin \left(x\sqrt{1-y^2} - y\sqrt{1-x^2} \right) \quad (26.17)$$

$$\arccos x - \arccos y = \arccos \left(xy + \sqrt{1-x^2}\sqrt{1-y^2} \right) \quad (26.18)$$

26.7 Interconversion of Ratios

$$\arcsin x = \arccos \sqrt{1-x^2} \quad (26.19)$$

$$= \arctan \left(\frac{x}{\sqrt{1-x^2}} \right) \quad (26.20)$$

$$\arccos x = \arcsin \sqrt{1 - x^2} \quad (26.21)$$

$$= \arctan \left(\frac{\sqrt{1 - x^2}}{x} \right) \quad (26.22)$$

$$2 \arctan x = \arcsin \left(\frac{2x}{1 + x^2} \right) \quad (26.23)$$

$$= \arccos \left(\frac{1 - x^2}{1 + x^2} \right) \quad (26.24)$$

$$= \arctan \left(\frac{2x}{1 - x^2} \right) \quad (26.25)$$

26.8 Miscellaneous Relations

$$\cos(\arcsin x) = \sin(\arccos x) = \sqrt{1 - x^2} \quad (26.26)$$

$$\arctan x = \frac{\pi}{2} - \arctan \left(\frac{1}{x} \right), x > 1 \quad (26.27)$$

Chapter 27

Hyperbolic Trigonometric Functions

27.1 Definition

Hyperbolic trigonometric functions are defined such that $(\cosh t, \sinh t)$ form the right half of an equilateral hyperbola. The functions are defined as follows:

$$\sinh x = \frac{\exp(x) - \exp(-x)}{2} \quad (27.1)$$

$$\cosh x = \frac{\exp(x) + \exp(-x)}{2} \quad (27.2)$$

$$\tanh x = \frac{\sinh x}{\cosh x} = \frac{\exp(x) - \exp(-x)}{\exp(x) + \exp(-x)} \quad (27.3)$$

$$\coth x = \frac{1}{\tanh x} = \frac{\exp(x) + \exp(-x)}{\exp(x) - \exp(-x)} \quad (27.4)$$

$$\operatorname{csch} x = \frac{1}{\sinh x} = \frac{2}{\exp(x) - \exp(-x)} \quad (27.5)$$

$$\operatorname{sech} x = \frac{1}{\cosh x} = \frac{2}{\exp(x) + \exp(-x)} \quad (27.6)$$

27.2 Identities

$$\coth^2 x - \sinh^2 x = 1 \quad (27.7)$$

$$\tanh^2 x + \operatorname{sech}^2 x = 1 \quad (27.8)$$

$$\coth^2 x - \operatorname{csch}^2 x = 1 \quad (27.9)$$

27.3 Inverse Hyperbolic Function

$$\sinh^{-1} z = \ln(z + \sqrt{z^2 + 1}) \quad (27.10)$$

$$\cosh^{-1} z = \ln(z \pm \sqrt{z^2 - 1}) \quad (27.11)$$

$$\tanh^{-1} z = \frac{1}{2} \ln \left(\frac{1+z}{1-z} \right) \quad (27.12)$$

$$\coth^{-1} z = \frac{1}{2} \ln \left(\frac{z+1}{z-1} \right) \quad (27.13)$$

$$\operatorname{csch}^{-1} z = \ln \left(\frac{1 \pm \sqrt{z^2 + 1}}{z} \right) \quad (27.14)$$

$$\operatorname{sech}^{-1} z = \ln \left(\frac{1 \pm \sqrt{1 - z^2}}{2} \right) \quad (27.15)$$

27.4 Relation to Circular Trigonometric Functions

$$\sinh(z) = -i \sin(iz) \quad (27.16)$$

$$\cosh(z) = \cos(iz) \quad (27.17)$$

$$\tanh(z) = -i \tan(iz) \quad (27.18)$$

$$\operatorname{csch}(z) = i \csc(iz) \quad (27.19)$$

$$\operatorname{sech}(z) = \sec(iz) \quad (27.20)$$

$$\coth(z) = i \cot(iz) \quad (27.21)$$

Part V

Calculus

Chapter 28

Limits

$$\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1 \quad (28.1)$$

$$\lim_{x \rightarrow 0} \frac{\tan x}{x} = 1 \quad (28.2)$$

$$\lim_{x \rightarrow 0} \cos x = 1 \quad (28.3)$$

$$\lim_{x \rightarrow a} \frac{x^n - a^n}{x - a} = na^{n-1} \quad (28.4)$$

$$\lim_{x \rightarrow a} f(x)g(x) = \lim_{x \rightarrow a} f(x) \cdot \lim_{x \rightarrow a} g(x) \quad (28.5)$$

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow a} f(x)}{\lim_{x \rightarrow a} g(x)}, \lim_{x \rightarrow a} g(x) \neq 0 \quad (28.6)$$

$$\lim_{x \rightarrow 0} \exp(x) = 1 \quad (28.7)$$

$$\lim_{x \rightarrow a} \exp(x) = \exp(c) \quad (28.8)$$

$$\lim_{x \rightarrow 0} \frac{\exp(x) - 1}{x} = 1 \quad (28.9)$$

$$\lim_{x \rightarrow a} c^x = c^a \quad (28.10)$$

$$\lim_{x \rightarrow a} \ln x = \ln a \quad (28.11)$$

$$\lim_{x \rightarrow 0} (1 + x)^{\frac{1}{x}} = e \quad (28.12)$$

$$\lim_{x \rightarrow 0} \frac{\ln(1 + x)}{x} = 1 \quad (28.13)$$

$$\lim_{x \rightarrow 0} \frac{a^x - 1}{x} = \ln a \quad (28.14)$$

$$\lim_{n \rightarrow \infty} \frac{x^n}{n!} = 0, \forall x \in \mathbb{R} \quad (28.15)$$

28.1 L'Hospital Rule

If:

$$L = \lim_{x \rightarrow a} \frac{f(x)}{g(x)}$$

is such that $f(a) = 0$ and $g(a) = 0$, or $f(a) = \infty$ and $g(a) = \infty$, then:

$$L = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$$

Chapter 29

Differentiation

29.1 Differentiation by First Principle

$$\frac{df(x)}{dx} = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \quad (29.1)$$

29.2 Standard Differentiation Formulae

$$\frac{dk}{dx} = 0 \quad (29.2)$$

$$\frac{dx^n}{dx} = nx^{n-1} \quad (29.3)$$

$$\frac{da^x}{dx} = \ln a \cdot a^x \quad (29.4)$$

$$\frac{d \exp(x)}{dx} = \exp(x) \quad (29.5)$$

$$\frac{d \ln x}{dx} = \frac{1}{x} \quad (29.6)$$

$$\frac{d\sqrt{x}}{dx} = \frac{1}{2\sqrt{2}} \quad (29.7)$$

$$(29.8)$$

29.2.1 Circular Trigonometric Functions

$$\frac{d \sin x}{dx} = \cos x \quad (29.9)$$

$$\frac{d \cos x}{dx} = -\sin x \quad (29.10)$$

$$\frac{d \tan x}{dx} = \sec^2 x \quad (29.11)$$

$$\frac{d \sec x}{dx} = \sec x \cdot \tan x \quad (29.12)$$

$$\frac{d \csc x}{dx} = -\csc x \cdot \cot x \quad (29.13)$$

$$\frac{d \cot x}{dx} = -\csc^2 x \quad (29.14)$$

29.2.2 Inverse Circular Trigonometric Functions

$$\frac{d \arcsin x}{dx} = \frac{1}{\sqrt{1-x^2}}, |x| \leq 1 \quad (29.15)$$

$$\frac{d \arccos x}{dx} = -\frac{1}{\sqrt{1-x^2}}, |x| \leq 1 \quad (29.16)$$

$$\frac{d \arctan x}{dx} = \frac{1}{1+x^2} \quad (29.17)$$

$$\frac{d \cot^{-1} x}{dx} = -\frac{1}{1+x^2} \quad (29.18)$$

$$\frac{d \sec^{-1} x}{dx} = \frac{1}{x \cdot \sqrt{x^2-1}}, |x| \geq 1 \quad (29.19)$$

$$\frac{d \csc^{-1} x}{dx} = -\frac{1}{x \cdot \sqrt{x^2-1}}, |x| \geq 1 \quad (29.20)$$

29.2.3 Hyperbolic Trigonometric Function

$$\frac{d \sinh x}{dx} = \cosh x \quad (29.21)$$

$$\frac{d \cosh x}{dx} = \sinh x \quad (29.22)$$

$$\frac{d \tanh x}{dx} = 1 - \tanh^2 x = \operatorname{sech}^2(x) \quad (29.23)$$

$$\frac{d \coth x}{dx} = 1 - \coth^2 x = -\operatorname{csch}^2(x) \quad (29.24)$$

$$\frac{d[\operatorname{sech}(x)]}{dx} = -\tanh x \operatorname{sech} x \quad (29.25)$$

$$\frac{d[\operatorname{csch}(x)]}{dx} = -\coth x \operatorname{csch} x \quad (29.26)$$

29.2.4 Inverse Hyperbolic Trigonometric Function

$$\frac{d \sinh x}{dx} = \frac{1}{\sqrt{x^2 + 1}} \quad (29.27)$$

$$\frac{d \cosh x}{dx} = \frac{1}{\sqrt{x^2 - 1}} \quad (29.28)$$

$$\frac{d \tanh x}{dx} = \frac{1}{1 - x^2} \quad (29.29)$$

$$\frac{d \coth x}{dx} = \frac{1}{1 - x^2} \quad (29.30)$$

$$\frac{d[\operatorname{sech}(x)]}{dx} = \frac{1}{x\sqrt{1 - x^2}} \quad (29.31)$$

$$\frac{d[\operatorname{csch}(x)]}{dx} = \frac{1}{|x|\sqrt{1 + x^2}} \quad (29.32)$$

29.3 Rules of Differentiation

$$\frac{d[c \cdot f(x)]}{dx} = c \cdot \frac{df(x)}{dx} \quad (29.33)$$

$$\frac{d[f_1(x) + f_2(x)]}{dx} = \frac{d[f_1(x)]}{dx} + \frac{d[f_2(x)]}{dx} \quad (29.34)$$

$$\frac{d[f_1 \cdot f_2]}{dx} = f_1 \cdot f_2' + f_2 \cdot f_1' \quad (29.35)$$

$$\frac{d\left(\frac{f_1}{f_2}\right)}{dx} = \frac{f_2 \cdot f_1' - f_1 \cdot f_2'}{f_2^2} \quad (29.36)$$

29.4 Chain Rule

If two functions are defined as $z = f(y)$ and $y = g(x)$:

$$\frac{dz}{dx} = \frac{dz}{dy} \cdot \frac{dy}{dx} \quad (29.37)$$

If two functions are defined as $x = f(\theta)$ and $y = g(\theta)$:

$$\frac{d^2 y}{dx^2} = \left[\frac{d}{d\theta} \left(\frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}} \right) \right] \frac{d\theta}{dx} \quad (29.38)$$

Chapter 30

Successive Differentiation

$$D^n(ax+b)^m = m(m-1) \cdots (m-n+1)a^n(ax+b)^{m-n} \quad (30.1)$$

$$D^n \left(\frac{1}{ax+b} \right) = \frac{(-1)^n \cdot n! \cdot a^n}{(ax+b)^{n+1}} \quad (30.2)$$

$$D^n \ln(ax+b) = \frac{(-1)^{n-1} \cdot (n-1)! \cdot a^n}{(ax+b)^n}, n \geq 2 \quad (30.3)$$

$$D^n(a^{mx}) = m^n (\ln a)^n \cdot a^{mx} \quad (30.4)$$

$$D^n(e^{mx}) = m^n e^{mx} \quad (30.5)$$

$$D^n \sin(ax+b) = a^n \sin(ax+b+n\frac{\pi}{2}) \quad (30.6)$$

$$D^n \cos(ax+b) = a^n \cos(ax+b+n\frac{\pi}{2}) \quad (30.7)$$

$$D^n[e^{ax} \sin(bx+c)] = (a^2+b^2)^{\frac{n}{2}} e^{ax} \sin(bx+c+n \arctan \frac{b}{a}) \quad (30.8)$$

$$D^n[e^{ax} \cos(bx+c)] = (a^2+b^2)^{\frac{n}{2}} e^{ax} \cos(bx+c+n \arctan \frac{b}{a}) \quad (30.9)$$

30.1 Leibnitz's Theorem

For two functions u and v of x , the successive differentiation of their product is defined as:

$$(uv)_n = {}^nC_0 u_n v + {}^nC_1 u_{n-1} v_1 + \cdots + {}^nC_n u v_n \quad (30.10)$$

$$= \sum_{i=0}^n {}^nC_i u_{n-i} v_i \quad (30.11)$$

Chapter 31

Partial Derivative

If $f(x, y)$ is a function of (x, y) , then $\frac{\partial f(x, y)}{\partial x}$ is the differentiation of $f(x, y)$ w.r.t. x , keeping all other parameters constant.

31.1 Chain Rule

If f is a function of u and v , which are functions of x and y , then:

$$\frac{\partial f}{\partial x} = \frac{\partial f}{\partial u} \frac{\partial u}{\partial x} + \frac{\partial f}{\partial v} \frac{\partial v}{\partial x} \quad (31.1)$$

$$\frac{\partial f}{\partial y} = \frac{\partial f}{\partial u} \frac{\partial u}{\partial y} + \frac{\partial f}{\partial v} \frac{\partial v}{\partial y} \quad (31.2)$$

$$(31.3)$$

If f is a function of x and y , which are functions of t , then:

$$\frac{df}{dt} = \frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt} \quad (31.4)$$

31.2 Euler's Theorem

For a homogeneous function ¹, $f(x_i)$ of degree n :

$$\sum x_i \frac{\partial f}{\partial x_i} = n f(x_i) \quad (31.5)$$

¹Homogeneous functions are defined as $f(ax, ay) = a^\kappa f(x, y)$, where κ is the degree of homogeneity. E.g. $f(x, y) = x^2 + y^2$, then $f(tx, ty) = t^2(x^2 + y^2)$, and the degree of homogeneity is 2.

Chapter 32

Application of Differential

32.1 Rolle's Theorem

For a function $f(x)$:

1. is continuous in $[a, b]$
2. is differentiable in (a, b)
3. $f(a) = f(b)$,

then there exists a point $x = c$ such that $f'(c) = 0$, $c \in (a, b)$

32.2 Mean Value Theorem or LaGrange's Theorem

For a function $f(x)$:

1. is continuous in $[a, b]$
2. is differentiable in (a, b) ,

then there exists a point $x = c$ such that $f'(c) = \frac{f(b) - f(a)}{b - a}$, $c \in (a, b)$, i.e., the tangent is parallel to the line joining the the points $(a, f(a))$ and $(b, f(b))$.

32.3 Cauchy's Mean Value Theorem

For a function $f(x)$ and $g(x)$:

1. are continuous in $[a, b]$
2. are differentiable in (a, b)
3. $g'(x) \neq 0$ in (a, b) ,

then there exists a point $c \in (a, b)$, such that $\frac{f(x)}{g(x)} = \frac{f(b) - f(a)}{g(b) - g(a)}$.

32.4 Maxima and Minima

32.4.1 Maxima

For the local maxima of a function $f(x)$:

1. $f'(c) = 0$ and

$$\lim_{\epsilon \rightarrow c^-} f'(\epsilon) > 0$$

$$\lim_{\epsilon \rightarrow c^+} f'(\epsilon) < 0$$

OR

2. $f'(c) = 0$ and $f''(x) < 0$,

then $f(c)$ is the local maxima point of the function $f(x)$.

32.4.2 Minima

For the local minima of a function $f(x)$:

1. $f'(c) = 0$ and

$$\lim_{\epsilon \rightarrow c^-} f'(\epsilon) < 0$$

$$\lim_{\epsilon \rightarrow c^+} f'(\epsilon) > 0$$

OR

2. $f'(c) = 0$ and $f''(x) > 0$,

then $f(c)$ is the local minima point of the function $f(x)$.

32.5 Taylor's Theorem

For a function which is differentiable n times:

$$f(a+h) = f(a) + h \cdot f'(a) + \frac{h^2}{2!} \cdot f''(a) + \cdots + \frac{h^{n-1}}{(n-1)!} \cdot f^{n-1}(a) + \frac{h^n}{x!} \cdot R_n \quad (32.1)$$

where R_n is the remainder term.

32.5.1 Remainder Term

LeGrange's Form

$$R_n = f^n(a + \theta h), \theta \in (0, 1) \quad (32.2)$$

Cauchy's Form

$$R_n = n(1-\theta)^{n-1} f^n(a + \theta h), \theta \in (0, 1) \quad (32.3)$$

32.5.2 Conditions for Validity of Expansion

For validity of Taylor Expansion, the condition

$$\lim_{n \rightarrow \infty} R_n = 0 \quad (32.4)$$

needs to be satisfied either where R_n is the remainder term in either LeGrange's Form or Cauchy's Form. If the condition is satisfied in a certain domain, then the expansion is valid within that domain only.

32.5.3 Taylor's Theorem for Two Variables

$$\begin{aligned} f(a+x, b+y) = & f(x, y) + \left(a \frac{\partial}{\partial x} + b \frac{\partial}{\partial y} \right) f(x, y) + \frac{1}{2!} \left(a^2 \frac{\partial^2}{\partial x^2} + b^2 \frac{\partial^2}{\partial y^2} \right) f(x, y) + \cdots + \\ & \frac{1}{n!} \left(a^n \frac{\partial^n}{\partial x^n} + b^n \frac{\partial^n}{\partial y^n} \right) f(x + \theta a, y + \theta b), \theta \in (0, 1) \end{aligned} \quad (32.5)$$

32.6 Maclaurin's Series

$$f(x) = f(0) + x f'(0) + \frac{1}{2!} x^2 f''(0) + \frac{1}{3!} x^3 f'''(0) + \cdots \infty \quad (32.6)$$

$$= \sum_{i=0}^{\infty} \frac{1}{i!} x^i f^{(i)}(0) \quad (32.7)$$

32.6.1 Maclaurin's Series with Two Variables

$$f(a, b) = f(0, 0) + \left(a \frac{\delta}{\delta x} + b \frac{\delta}{\delta y} \right) f(0, 0) + \frac{1}{2!} \left(a^2 \frac{\delta^2}{\delta x^2} + b^2 \frac{\delta^2}{\delta y^2} \right) f(0, 0) + \cdots \infty \quad (32.8)$$

$$= \sum_{i=0}^{\infty} \frac{1}{n!} \left(a^i \frac{\delta^i}{\delta x^i} + b^i \frac{\delta^i}{\delta y^i} \right) f(0, 0) \quad (32.9)$$

32.7 Curvature

Curvature is the rate of change of direction w.r.t. arc. Mathematically:

$$\text{Curvature} = \frac{d(\text{direction})}{d(\text{arc})} \quad (32.10)$$

$$\lim_{\Delta s \rightarrow 0} \frac{\Delta \psi}{\Delta s} = \frac{d\psi}{ds} \quad (32.11)$$

32.7.1 Radius of Curvature

Cartesian Form

For a curve $y = f(x)$:

$$\rho = \frac{(1 + y'^2)^{\frac{3}{2}}}{y''} \quad (32.12)$$

However, this formula fails for $y' \rightarrow \infty$.

Parametric Form

For a curve defined as $x = \phi(t)$ and $y = \psi(t)$:

$$\rho = \frac{(\ddot{x}^2 + \ddot{y}^2)^{\frac{3}{2}}}{x\ddot{y} - y\ddot{x}} \quad (32.13)$$

32.7.2 Newton's Formula

1. If the curve passes through origin, and the tangent at origin is the x-axis:

$$\rho = \lim_{\substack{x \rightarrow 0 \\ y \rightarrow 0}} \frac{x^2}{2y} \quad (32.14)$$

2. If the curve passes through origin, and the tangent at origin is the y-axis:

$$\rho = \lim_{\substack{x \rightarrow 0 \\ y \rightarrow 0}} \frac{y^2}{2x} \quad (32.15)$$

3. If the curve passes through origin and $ax + by + c = 0$ is the tangent at origin:

$$\rho(0, 0) = \frac{1}{2} \sqrt{a^2 + b^2} \lim_{\substack{x \rightarrow 0 \\ y \rightarrow 0}} \frac{a^2 + y^2}{ax + by} \quad (32.16)$$

32.7.3 Tangent at Origin

For a curve

$$\sum c_i x^j y^k = 0, i \in \mathbb{N} \text{ and } j, k \in \mathbb{Z} - \{0\} \quad (32.17)$$

The curve passes through origin $c = 0$. Then the lowest degree term equated to x gives the tangent at origin.

32.8 Asymptotes

If the distance between a line P and a curve $f(x)$, s is such that $s \rightarrow 0$, as $x \rightarrow \infty$, then P is the asymptote of $f(x)$. For asymptotes not parallel to x-axis:

Let $y = mx + c$ be the asymptote of the function $y = f(x)$, then:

$$m = \lim_{x \rightarrow \infty} \frac{y}{x} \quad (32.18)$$

$$c = \lim_{x \rightarrow \infty} (y - mx) \quad (32.19)$$

32.8.1 Asymptote of Algebraic Curves

For an algebraic curve, passing through origin, defined as:

$$\begin{aligned}
 & (a_0x^n + a_1x^{n-1}y^1 + \cdots + a_{n-1}xy^{n-1} + a_ny^n) \\
 & + (b_0x^{n-1} + b_1x^{n-2}y^1 + \cdots + b_{n-1}xy^{n-2} + b_ny^{n-1}) + \cdots = 0 \\
 & \Rightarrow x^n\phi_n\left(\frac{y}{x}\right) + x^{n-1}\phi_{n-1}\left(\frac{y}{x}\right) + \cdots + x\phi_1\left(\frac{y}{x}\right) = 0
 \end{aligned} \tag{32.20}$$

The asymptote(s) defined as $y = mx + c$,

1. m is the solution for the equation

$$\phi_n(m) = 0 \tag{32.21}$$

- 2.

$$c = -\frac{\phi_{n-1}(m)}{\phi_n(m)} \tag{32.22}$$

where c is a finite value.

Chapter 33

Integration

33.1 General Formulae

$$\int nx^{n-1}dx = x^n + A \quad (33.1)$$

$$\int x^n dx = \frac{1}{n+1}x^{n+1} + A \quad (33.2)$$

$$\int e^x dx = e^x + A \quad (33.3)$$

$$\int \frac{1}{x} dx = \ln x + A \quad (33.4)$$

$$\int \ln x dx = x(\ln x - 1) + A \quad (33.5)$$

33.2 Circular Trigonometric Functions

$$\int \sin x dx = -\cos x + A \quad (33.6)$$

$$\int \cos x dx = \sin x + A \quad (33.7)$$

$$\int \sec^2 x dx = \tan x + A \quad (33.8)$$

$$\int \csc^2 x dx = -\cot x + A \quad (33.9)$$

$$\int \sec x \tan x dx = \sec x + A \quad (33.10)$$

$$\int \csc x \cot x dx = -\csc x + A \quad (33.11)$$

$$\int \sec x dx = \ln(\sec x + \tan x) + A \quad (33.12)$$

$$\int \csc x dx = -\ln(\csc x + \cot x) + A \quad (33.13)$$

$$\int \tan x dx = -\ln(\cos x) + A \quad (33.14)$$

$$= \ln(\sec x) + A \quad (33.15)$$

$$\int \cot x dx = \ln(\sin x) + A \quad (33.16)$$

33.3 Inverse Circular Trigonometric Function

$$\int \frac{1}{\sqrt{1-x^2}} dx = \sin^{-1} x + A \quad (33.17)$$

$$\int \frac{-1}{\sqrt{1-x^2}} dx = \cos^{-1} x + A \quad (33.18)$$

$$\int \frac{1}{1+x^2} dx = \tan^{-1} x + A \quad (33.19)$$

$$\int \frac{-1}{1+x^2} dx = \cot^{-1} x + A = -\tan^{-1} x + A \quad (33.20)$$

$$\int \frac{1}{x\sqrt{x^2-1}} dx = \sec^{-1} x + A = -\csc^{-1} x + A \quad (33.21)$$

$$\int \frac{-1}{x\sqrt{x^2-1}} dx = \csc^{-1} x + A = -\sec^{-1} x + A \quad (33.22)$$

33.4 Standard Integrals

$$\int \frac{dx}{\sqrt{a^2 - x^2}} = \sin^{-1} \frac{x}{a} + A \quad (33.23)$$

$$\int \frac{dx}{\sqrt{x^2 + a^2}} = \ln(x + \sqrt{x^2 + a^2}) + A \quad (33.24)$$

$$\int \frac{dx}{a^2 + x^2} = \frac{1}{a} \tan^{-1} \frac{x}{a} + A \quad (33.25)$$

$$\int \frac{dx}{\sqrt{x^2 - a^2}} = \ln(x + \sqrt{x^2 - a^2}) + A \quad (33.26)$$

$$\int \frac{dx}{x\sqrt{x^2 - a^2}} = \frac{1}{a} \sec^{-1} \frac{x}{a} + A \quad (33.27)$$

$$\int \sqrt{a^2 - x^2} dx = \frac{x\sqrt{a^2 - x^2}}{2} + \frac{a^2}{2} \sin^{-1} \frac{x}{a} + A \quad (33.28)$$

$$\int \sqrt{a^2 + x^2} dx = \frac{x\sqrt{a^2 + x^2}}{2} + \frac{a^2}{2} \ln(x + \sqrt{x^2 + a^2}) + A \quad (33.29)$$

$$\int \sqrt{x^2 - a^2} dx = \frac{x\sqrt{x^2 - a^2}}{2} - \frac{a^2}{2} \ln(x + \sqrt{a^2 - x^2}) + A \quad (33.30)$$

$$\int \frac{dx}{x^2 - a^2} = \frac{1}{2a} \ln \left| \frac{x - a}{x + a} \right| + A \quad (33.31)$$

$$\int \frac{dx}{a^2 - x^2} = \frac{1}{2a} \ln \left| \frac{a + x}{a - x} \right| + A \quad (33.32)$$

1

33.5 Special Forms

For a function $f(x)$:

$$\int [f(x)]^n f'(x) dx = \begin{cases} \frac{[f(x)]^{n+1}}{n+1} + A, n \neq -1 \\ \ln|f(x)| + A, n = -1 \end{cases} \quad (33.33)$$

33.5.1 Integration by Part

For two functions $u(x)$ and $v(x)$:

$$\int u(x)v(x)dx = u(x) \left[\int v(x)dx \right] - \int \left[\frac{du(x)}{dx} \left(\int v(x)dx \right) dx \right] \quad (33.34)$$

¹ a is a constant $\in \mathbb{R}$

Chapter 34

Definite Integration

For a function $f(x)$ for which $\int f(x)dx = F(x) + A$,

$$\int_a^b f(x)dx = F(b) - F(a) \quad (34.1)$$

34.1 Properties of Definite Integration

$$\int_a^b f(x)dx = \int_a^b f(t)dt \quad (34.2)$$

$$\int_a^b f(x)dx = - \int_b^a f(x)dx \quad (34.3)$$

$$\int_a^b f(x)dx = \int_a^c f(x)dx + \int_c^b f(x)dx \quad (34.4)$$

$$\int_a^b f(x)dx = \int_a^b f(a+b-x)dx \quad (34.5)$$

$$\int_0^a f(x)dx = \int_0^a f(a-x)dx \quad (34.6)$$

$$\int_0^{2a} f(x)dx = \begin{cases} 2 \int_0^a f(x)dx, & f(2a-x) = f(x) \\ 0, & f(2a-x) = f-(x) \end{cases} \quad (34.7)$$

$$\int_{-a}^a f(x)dx = \begin{cases} 2 \int_0^a f(x)dx, & f(x) \text{ is even} \\ 0, & f(x) \text{ is odd} \end{cases} \quad (34.8)$$

34.2 Approximation

$$f(a)(b-a) \leq \int_a^b f(x)dx \leq f(b)(b-a) \quad (34.9)$$

34.3 Sum of Infinite Series as a Definite Integral

Refer to 3.5.2.

Chapter 35

Reduction Formula

35.1 Circular Trigonometric Functions

$$\int \sin^n x dx = -\frac{1}{n} \sin^{n-1} x \cos x + \frac{n-1}{n} \int \sin^{n-2} x dx \quad (35.1)$$

$$\int \cos^n x dx = \frac{1}{n} \cos^{n-1} x \sin x + \frac{n-1}{n} \int \cos^{n-2} x dx \quad (35.2)$$

$$\int_0^{\frac{\pi}{2}} \sin^n x dx = \int_0^{\frac{\pi}{2}} \cos^n x dx = \begin{cases} \frac{(n-1) \cdot (n-3) \cdots 3 \cdot 1}{n \cdot (n-2) \cdots 4 \cdot 2} \left(\frac{\pi}{2}\right), n \rightarrow \text{even} \\ \frac{(n-1) \cdot (n-3) \cdots 4 \cdot 2}{n \cdot (n-2) \cdots 3 \cdot 1}, n \rightarrow \text{odd} \end{cases} \quad (35.3)$$

$$\int \sin^m x \cos^n x dx = \frac{-\sin^{m-1} x \cos^{n+1} x}{m+n} + \frac{m-1}{m+n} \int \sin^{m-2} x \cos^n x dx \quad (35.4)$$

35.2 Special Definite Integrations

For $I(m, n) = \int_0^{\frac{\pi}{2}} \sin^m x \cos^n x dx$:

$$I(m, n) = \frac{[(m-1) \cdot (m-3) \cdots 3 \cdot 1][(n-1) \cdot (n-3) \cdots 3 \cdot 1]}{(m+n) \cdot (m+n-1) \cdots (4) \cdot (2)} \cdot \frac{\pi}{2} \rightarrow m, n \text{ are even} \quad (35.5)$$

$$I(m, n) = \frac{[(m-1) \cdot (m-3) \cdots (2 \text{ or } 1)][(n-1) \cdot (n-3) \cdots (2 \text{ or } 1)]}{(m+n) \cdot (m+n-1) \cdots (2 \text{ or } 1)} \rightarrow \text{all other cases} \quad (35.6)$$

$$(35.7)$$

35.3 Recursive Forms in Circular Trigonometric Functions

$$\begin{aligned} I_n &= \int \tan^n x dx \\ \Rightarrow I_n &= \frac{\tan^{n-2} x}{n-1} - I_{n-2} \end{aligned} \tag{35.8}$$

$$\begin{aligned} I_n &= \int \cot^n x dx \\ \Rightarrow I_n &= -\frac{\cot^{n-2} x}{n-1} - I_{n-2} \end{aligned} \tag{35.9}$$

$$\begin{aligned} I_n &= \int \sec^n x dx \\ \Rightarrow I_n &= \frac{\sec^{n-2} x \tan x}{n-1} + \frac{n-2}{n-1} I_{n-2} \end{aligned} \tag{35.10}$$

$$\begin{aligned} I_n &= \int \csc^n x dx \\ \Rightarrow I_n &= -\frac{\csc^{n-2} x \cot x}{n-1} + \frac{n-2}{n-1} I_{n-2} \end{aligned} \tag{35.11}$$

$$I_n = \int x^n e^{ax} dx \tag{35.12}$$

$$\int x^n e^{ax} dx = \frac{x^n e^{ax}}{a} - \frac{n}{a} I_{n-2} \tag{35.13}$$

$$I(m, n) = \int x^m (\ln x)^n dx \tag{35.14}$$

$$\int x^m (\ln x)^n dx = \frac{x^{m+1}}{m+1} (\ln x)^n - \frac{n}{m+1} I_{m, n-1} \tag{35.15}$$

Chapter 36

β and Γ Functions

For $m, n > 0$:

$$\beta(m, n) = \int_0^1 x^{m-1} (1-x)^{n-1} dx \quad (36.1)$$

$$= 2 \int_0^{\frac{\pi}{2}} \sin^{2m-1} x \cos^{2n-1} x dx \Gamma(n) = \int_0^\infty e^{-x} x^{n-1} dx \quad (36.2)$$

36.1 Important Relations between $\beta(m, n)$ and $\Gamma(n)$ Functions

$$\Gamma(n) = \frac{\Gamma(n+1)}{n} \quad (36.3)$$

$$\Gamma(1) = 1 \quad (36.4)$$

$$\Gamma\left(\frac{1}{2}\right) = \sqrt{\pi} \approx 1.772 \quad (36.5)$$

$$\Gamma(n+1) = n!, n \in \mathbb{N} \quad (36.6)$$

$$\Gamma(m)\Gamma\left(m + \frac{1}{2}\right) = \sqrt{\pi}\Gamma(2m) \quad (36.7)$$

$$\Gamma(m)\Gamma(m-1) = \pi \csc(m\pi) \quad (36.8)$$

$$\beta(m, n) = \beta(n, m) \quad (36.9)$$

$$\beta(m, n) = \frac{\Gamma(m)\Gamma(n)}{\Gamma(m+n)} \quad (36.10)$$

$$\beta\left(\frac{1}{2}, \frac{1}{2}\right) = \pi \quad (36.11)$$

$$\int_0^{\frac{\pi}{2}} \sin^p x \cos^q x = \frac{1}{2} \frac{\Gamma\left(\frac{p+1}{2}\right) \Gamma\left(\frac{q+1}{2}\right)}{\Gamma\left(\frac{p+2}{2}\right)} \quad (36.12)$$

$$(36.13)$$

Chapter 37

Multiple Integrals

37.1 Two Variables

For

$$I = \iint_R f(x, y) dx dy \quad (37.1)$$

The following substitution are made:

$$x = g(r, \theta) \quad (37.2)$$

$$y = h(r, \theta) \quad (37.3)$$

$$\Rightarrow dx dy = |J| dr d\theta \quad (37.4)$$

Where J is the Jacobian, defined as:

$$J = \begin{vmatrix} \frac{\partial x}{\partial r} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial r} & \frac{\partial y}{\partial \theta} \end{vmatrix} \quad (37.5)$$

The equivalent integral is:

$$I = \iint_{R_1} f(g(r, \theta), h(r, \theta)) |J| dr d\theta \quad (37.6)$$

37.2 Three Variables

For

$$I = \iiint_R f(x, y, z) dx dy dz \quad (37.7)$$

The following substitution are made:

$$x = g(r, \theta, \phi) \quad (37.8)$$

$$y = h(r, \theta, \phi) \quad (37.9)$$

$$z = k(r, \theta, \phi) dx dy dz = |J| dr d\theta d\phi \quad (37.10)$$

Where J is the Jacobian, defined as:

$$J = \begin{vmatrix} \frac{\partial x}{\partial r} & \frac{\partial y}{\partial r} & \frac{\partial z}{\partial r} \\ \frac{\partial x}{\partial \theta} & \frac{\partial y}{\partial \theta} & \frac{\partial z}{\partial \theta} \\ \frac{\partial x}{\partial \phi} & \frac{\partial y}{\partial \phi} & \frac{\partial z}{\partial \phi} \end{vmatrix} \quad (37.11)$$

The equivalent integral is:

$$I = \iiint_{R_1} f(g(r, \theta, \phi), h(r, \theta, \phi), k(r, \theta, \phi)) |J| dr d\theta d\phi \quad (37.12)$$