

Mathematical Formula Sheet

A Book of High School and Engineering Common Course Mathematical
Formulae

Agnij Mallick

December, 2020

Contents

I	Algebra	1
1	Logarithm	2
1.1	Basic Formulae	2
1.2	Series	3
2	Complex Number	4
2.1	Basic Formulae	4
2.2	Arithmetic Operation of Complex Number	4
2.3	Euler's Formula	4
2.4	Trigonometric Ratios in Complex Form	5
2.5	De Moivre's Formula	5
2.6	Application of Euler's and De Moivre's Formula	5
2.7	Roots of Unity	5
2.8	Important Relations of Complex Numbers	5
3	Progression	6
3.1	Arithmetic Progression (A.P.)	6
3.1.1	Sum of A.P. Series	6
3.1.2	Important Relation	6
3.2	Geometric Progression (G.P.)	6
3.2.1	The Value of 'r'	7
3.2.2	Sum of a G.P. Series	7
3.2.3	Important relations	7
3.3	Harmonic Progression (H.P.)	7
3.4	Arithmetico-Geometric Progression (A.G.P.)	7
3.4.1	Sum of A.G.P.:	8
3.5	Special Series	8
3.5.1	Riemann Zeta Function	8
3.5.2	Riemann's Infinite Series as an Integration	8
4	Test of Convergence of Infinite Series	9
4.1	Definition	9
4.2	Tests of Convergence	9
4.2.1	Comparison Test	9
4.2.2	Limit Form	10
4.2.3	Integral Test or Maclaurin-Cauchy Test	10

4.2.4	Ratio Test	10
4.2.5	D'Alembert's Ratio Test	11
4.2.6	Rabbe's Test	11
4.2.7	Cauchy's Root Test	11
4.2.8	Logarithmic Test	11
5	Determinants	12
5.1	Definition	12
5.1.1	Minor and Cofactor	12
5.2	Important Properties	12
5.3	Cramer's Rule	13
5.3.1	Consistency Test	14
6	Matrices	15
6.1	Sum of Two Matrices	15
6.2	Multiplication of Two Matrices	15
6.2.1	Multiplicative Properties	16
6.3	Adjoint of a Matrix	16
6.4	Martin's Rule	16
7	Binomial Theorem	17
7.1	Expansion of a binomial expression	17
7.2	Trinomial Expansion	17
7.3	Properties of Coefficients	18
7.4	Pascal's Rule	18
8	Boolean Algebra	19
9	Remainder Theorems	20
9.1	Remainder Theorem	20
9.2	Euler's Remainder Theorem	20
9.2.1	Euler's Totient Function	21
9.3	Wilson Theorem	22
II	Co-Ordinate Geometry	23
10	2-D Co-ordinate Geometry	24
11	Triangles	25
12	Straight Line	26
13	General Theory of Second Degree Equation	28

14 Conics	29
14.1 Parabola	29
14.2 Ellipse and Hyperbola	29
14.3 Parametric Form of Conics	30
14.3.1 Hyperbola	30
14.3.2 Ellipse	30
14.3.3 Parabola	30
15 Circles	31
15.1 Locus Form	31
15.2 Diameter Form	31
15.3 General Form	31
15.4 Important Relations	31
16 Vectors	33
16.1 Modulus of a Vector	33
16.2 Sum of Vectors	33
16.3 Product of Vectors	33
16.3.1 Dot Product	33
16.3.2 Cross Product	34
16.4 Test of Co-planarity	34
17 3-D Geometry	35
17.1 Distance between two points $A(x_1, y_1, z_1)$ and $B(x_2, y_2, z_2)$:	35
17.2 Section Formula of a Line Segment Divided in the ratio $m : n$	35
17.3 Centroid of a Triangle	35
18 Line in 3-D Space	36
18.1 Angle between Two Lines	36
18.2 Skew and Co-planar Lines	36
18.3 Distances	37
18.3.1 The shortest distance between r_1 and r_2	37
18.3.2 Cartesian Form	37
18.3.3 Distance Between Parallel Lines	37
18.3.4 Distance of a Point to a Line	37
19 3-D Plane	38
19.1 Angle Between Two Planes	38
19.2 Distance of a Point from a Plane	38
19.2.1 Cartesian Form	38

19.2.2 Vector Form	39
III Statistics	40
20 Statistics	41
21 Lines of Regression	42
21.1 Karl Pearson's Co-efficient of Correlation (20.5)	42
21.1.1 Degree of Correlation	43
IV Trigonometry	44
22 Circular Trigonometric Functions	45
22.1 Negative Angle Formula	46
22.2 Sum of Angles Formula	46
22.3 Difference of Angles Formula	46
22.4 Multiples and Sub-multiples of π and $\frac{\pi}{2}$	47
22.5 $(\frac{\pi}{2} \pm \theta)$ Formula	47
22.6 $(\frac{\pi}{4} \pm \theta)$ Formula	48
22.7 Trigonometric Identities	48
22.8 Double Angle Formula	48
22.9 Triple Angle Formula	48
22.10 Sum and Product of Two Ratios	49
22.11 General Solutions	49
22.12 Taylor Series Expansion of Trigonometric Ratios	50
23 Inverse Circular Trigonometric Function	51
23.1 Definition of Inverse Circular Trigonometric Function	51
23.1.1 For $\sin x$	51
23.1.2 For $\cos x$	51
23.1.3 For $\tan x$	52
23.1.4 For $\cot x$	52
23.1.5 For $\csc x$	52
23.2 Negative Arguments	53
23.3 Reciprocal Relations	53
23.4 I.T.F. Identities	53
23.5 Sum of Two Angles	53
23.6 Difference of Two Angles	54
23.7 Interconversion of Ratios	54
23.8 Miscellaneous Relations	54
24 Hyperbolic Trigonometric Function	55
24.1 Definition	55
24.2 Identities	56
24.3 Inverse Hyperbolic Function	56
24.4 Relation to Circular Trigonometric Functions	56

V	Calculus	57
25	Limits	58
25.1	L'Hospital Rule	59
26	Differentiation	60
26.1	Differentiation by First Principle	60
26.2	Standard Differentiation Formulae	60
26.2.1	Circular Trigonometric Functions	61
26.2.2	Inverse Circular Trigonometric Functions	61
26.2.3	Hyperbolic Trigonometric Function	62
26.2.4	Inverse Hyperbolic Trigonometric Function	62
26.3	Rules of Differentiation	63
26.4	Chain Rule	63
27	Successive Differentiation	64
27.1	Leibnitz's Theorem	64
28	Partial Derivative	65
28.1	Chain Rule	65
28.2	Euler's Theorem	65
29	Application of Differentiation	66
29.1	Rolle's Theorem	66
29.2	Mean Value Theorem or LaGrange's Theorem	66
29.3	Cauchy's Mean Value Theorem	66
29.4	Maxima and Minima	67
29.4.1	Maxima	67
29.4.2	Minima	67
29.5	Taylor's Theorem	67
29.5.1	Remainder Term	68
29.5.2	Conditions for Validity of Expansion	68
29.5.3	Taylor's Theorem for Two Variables	68
29.6	Maclaurin's Series	68
29.6.1	Maclaurin's Series with Two Variables	69
29.7	Curvature	69
29.7.1	Radius of Curvature	69
29.7.2	Newton's Formula	69
29.7.3	Tangent at Origin	70
29.8	Asymptotes	70
29.8.1	Asymptote of Algebraic Curves	70
30	Integration	72
30.1	General Formulae	72
30.2	Circular Trigonometric Functions	73
30.3	Inverse Circular Trigonometric Function	74
30.4	Standard Integrals	75
30.5	Special Forms	75

30.5.1	Integration by Part	76
31	Definite Integral	77
31.1	Definition	77
31.2	Properties of Definite Integration	78
31.3	Approximation	78
31.4	Sum of Infinite Series as a Definite Integral	78
32	Reduction Formulae	79
33	Multiple Integrals	82
33.1	Two Variables	82
33.2	Three Variables	82
34	Differential Equation	84
34.1	1 st Order, 1 st Degree Differential Equation	84
34.2	2 nd Order, 1 st Degree Differential Equation	84
34.3	Special Cases of Differential Equation	85
34.3.1	Definition of Inverse Operator	85
34.3.2	Special Cases	85
34.4	Method of Variation of Parameters	87
34.5	Singular and Ordinary Point	87
35	Beta and Gamma Functions	89
35.1	Important Relations between $\beta(m, n)$ and $\Gamma(n)$ Functions	89
36	Laplace Transformations	91
36.1	Basic Transformations	91
36.2	Important Relations	92
36.3	Convolution	92
36.4	Laplace Transforms of Differentials	92

Part I

Algebra

Chapter 1

Logarithm

1.1 Basic Formulae

For $a^x = b$:

$$\log_a x, \text{ for all } x \leq 0 \text{ is undefined} \quad (1.1)$$

$$\log_a b = x, \text{ } bax \neq 1, a \neq 1 \quad (1.2)$$

$$\log_b a^m = m \log_b a, \text{ for } a^m = b \quad (1.3)$$

$$a^{\log_a x} = x \quad (1.4)$$

$$a^{\log_b c} = c^{\log_b a} \quad (1.5)$$

$$\frac{1}{\log_a b} = \log_b a \quad (1.6)$$

$$\log_c(ab) = \log_c a + \log_c b \quad (1.7)$$

$$\log_c\left(\frac{a}{b}\right) = \log_c a - \log_c b \quad (1.8)$$

$$|\log_a x| = \begin{cases} -\log_a x, & \text{if } 0 < x < 1 \\ \log_a x, & \text{if } 1 \leq x < \infty \end{cases} \quad (1.9)$$

1.2 Series

$$e^x = 1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots \infty = \lim_{n \rightarrow \infty} \sum_{i=0}^n \frac{x^i}{i!} \quad (1.10)$$

$$\log(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots \infty = \lim_{n \rightarrow \infty} \sum_{i=1}^n (-1)^{(i-1)} \frac{x^i}{i} \quad (1.11)$$

Chapter 2

Complex Number

2.1 Basic Formulae

For $z = x + iy$,

$$|z| = \sqrt{x^2 + y^2} \quad (2.1)$$

$$\tan \theta = \frac{y}{x} \quad (2.2)$$

$$\bar{z} = x - iy \quad (2.3)$$

2.2 Arithmetic Operation of Complex Number

For two complex numbers $z_1 = x_1 + iy_1$ and $z_2 = x_2 + iy_2$:

$$z_1 + z_2 = (x_1 + x_2) + i(y_1 + y_2) \quad (2.4)$$

$$z_1 z_2 = (x_1 x_2 - y_1 y_2) + i(x_1 y_2 + x_2 y_1) \quad (2.5)$$

$$|z_1 z_2| = |z_1| |z_2| \quad (2.6)$$

$$\frac{z_1}{z_2} = \frac{(x_1 x_2 + y_1 y_2) + i(x_2 y_1 - x_1 y_2)}{a_2^2 + b_2^2} \quad (2.7)$$

$$\left| \frac{z_1}{z_2} \right| = \frac{|z_1|}{|z_2|} \quad (2.8)$$

2.3 Euler's Formula

$$z = r e^{i\theta}, \text{ where } r = |z|, e^{i\theta} = \cos \theta + i \sin \theta, \text{ and } \theta = \tan^{-1} \frac{y}{x} \quad (2.9)$$

2.4 Trigonometric Ratios in Complex Form

$$e^{i\theta} + e^{-i\theta} = 2 \cos \theta \quad (2.10)$$

$$\Rightarrow \cos \theta = \frac{e^{i\theta} + e^{-i\theta}}{2} \quad (2.11)$$

$$e^{i\theta} - e^{-i\theta} = 2 \sin \theta \quad (2.12)$$

$$\Rightarrow \sin \theta = \frac{e^{i\theta} - e^{-i\theta}}{2} \quad (2.13)$$

2.5 De Moivre's Formula

$$(\cos \theta + i \sin \theta)^n = \cos(n\theta) + i \sin(n\theta) \quad (2.14)$$

2.6 Application of Euler's and De Moivre's Formula

For $z_1 = |r_1|e^{i\theta_1}$ and $z_2 = |r_2|e^{i\theta_2}$

$$z_1 z_2 = r_1 r_2 e^{i(\theta_1 + \theta_2)} \frac{z_1}{z_2} = \frac{r_1}{r_2} e^{i(\theta_1 - \theta_2)} \quad (2.15)$$

2.7 Roots of Unity

$$\sqrt[n]{1} = e^{i \frac{2k\pi}{n}}, \text{ where } k \in [0, n-1] \quad (2.16)$$

2.8 Important Relations of Complex Numbers

$$|z_1 + z_2| \leq |z_1| + |z_2| \quad (2.17)$$

$$|z_1 - z_2| \leq |z_1| + |z_2| \quad (2.18)$$

$$|z_1 - z_2| \geq ||z_1| - |z_2|| \quad (2.19)$$

$$|z_1 + z_2| \geq ||z_1| - |z_2|| \quad (2.20)$$

$$|z_1 + z_2|^2 = 2(|z_1|^2 + |z_2|^2) \quad (2.21)$$

Chapter 3

Progression

3.1 Arithmetic Progression (A.P.)

An arithmetic sequence is $a, a + d, a + 2d, \dots \infty$ or $t_n = a + (n - 1)d$, where a is the first term, d is the common difference, and n is the n^{th} -term.

An arithmetic series is $a + (a + d) + (a + 2d) + \dots \infty$.

3.1.1 Sum of A.P. Series

$$\begin{aligned} S_n &= a + (a + d) + \dots + (a + \overline{n - 2}d) + (a + \overline{n - 1}d) \\ S_n &= (a + \overline{n - 1}d) + (a + \overline{n - 2}d + \dots + (a + d) + a \\ &\Rightarrow 2S_n = n(2a + \overline{n - 1}d) \\ &\Rightarrow S_n = \frac{n}{2}(2a + \overline{n - 1}d) \end{aligned} \quad (3.1)$$

3.1.2 Important Relation

If the three terms a, b, c are in A.P., then

$$2b = a + c \quad (3.2)$$

3.2 Geometric Progression (G.P.)

An geometric sequence is $a, ar, ar^2, \dots \infty$ or $t_n = ar^{n-1}$, where a is the first term, r is the common ratio, and n is the n^{th} -term.

An geometric series is $a + ar + ar^2 + \dots \infty$.

3.2.1 The Value of 'r'

$$r = \frac{t_2}{t_1} = \frac{t_3}{t_2} = \dots = \frac{t_n}{t_{n-1}} \quad (3.3)$$

3.2.2 Sum of a G.P. Series

For a definite G.P. series, where there are n terms in the series, the sum of the series is:

$$S_n = \frac{a|r^n - 1|}{|r - 1|} \quad (3.4)$$

For an infite G.P. series the sum of the series is defined for $r < 1$. Sum of such a series is:

$$S_\infty = \frac{a}{1 - r} \quad (3.5)$$

3.2.3 Important relations

If the three terms a, b, c are in G.P., then:

$$b^2 = ac \quad (3.6)$$

3.3 Harmonic Progression (H.P.)

If a, b, c are terms of an H.P. then $\frac{1}{a}, \frac{1}{b}, \frac{1}{c}$ are in A.P.

$$\therefore \frac{2}{b} = \frac{1}{a} + \frac{1}{c} \quad (3.7)$$

$$\Rightarrow b = \frac{2ac}{a + c} \quad (3.8)$$

3.4 Arithmetico-Geometric Progression (A.G.P.)

Sequence $a, (a + d)r, (a + 2d)r^2, \dots, (a + \overline{n - 1}d)r^{n-1}$, where $a \rightarrow$ first term of A.G.P., $d \rightarrow$ common difference, and $r \rightarrow$ common ratio.

3.4.1 Sum of A.G.P.:

For an infinite A.G.P. series, the sum is defined for $r < 1$:

$$S_\infty = \frac{a}{1 - r} + \frac{dr}{(1 - r)^2} \quad (3.9)$$

3.5 Special Series

For $n \in \mathbb{N}$

$$1 + 2 + 3 + \dots + (n - 1) + n = \frac{n(n - 1)}{2} \quad (3.10)$$

$$1^2 + 2^2 + 3^2 + \dots + (n - 1)^2 + n^2 = \frac{n(n + 1)(2n + 1)}{6} \quad (3.11)$$

$$1^3 + 2^3 + 3^3 + \dots + (n - 1)^3 + n^3 = \left[\frac{n(n - 1)}{2} \right]^2 \quad (3.12)$$

3.5.1 Riemann Zeta Function

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} \quad (3.13)$$

3.5.2 Riemann's Infinite Series as an Integration

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=r_1}^{r_2} f\left(\frac{i}{n}\right) = \int_{\frac{r_1}{n}}^{\frac{r_2}{n}} f(x) dx \quad (3.14)$$

Chapter 4

Test of Convergence of Infinite Series

If $a_1, a_2, a_3, \dots, a_n$ is a sequence by a_n and their sum of series is S_n , then the following apply.

4.1 Definition

If

$$\lim_{n \rightarrow \infty} S_n = l$$

where l is a finite value, the series S_n is said to converge. A non-convergent series is called a divergent series.

4.2 Tests of Convergence

4.2.1 Comparison Test

If u_n and v_n are two positive series, then:

1. (a) v_n converges
(b) $u_n \leq v_n \forall n$ Then u_n converges.
2. (a) v_n diverges
(b) $u_n \geq v_n \forall n$ Then u_n diverges.

4.2.2 Limit Form

If

$$\lim_{x \rightarrow \infty} \frac{u_n}{v_n} = l$$

where l is a finite quantity $\neq 0$, then u_n and v_n converge and diverge together.

4.2.3 Integral Test or Maclaurin-Cauchy Test

For a series

$$\sum_{i=N}^{\infty} f(x), \text{ where } N \in \mathbb{Z} \quad (4.1)$$

will only converge if the improper integral

$$\int_N^{\infty} f(x)dx \quad (4.2)$$

is finite.

If the improper integral is finite, the upper and lower limit of the infinite series is given by:

$$\int_N^{\infty} f(x)dx \leq \sum_{i=N}^{\infty} f(x) \leq f(N) + \int_N^{\infty} f(x)dx \quad (4.3)$$

4.2.4 Ratio Test

If, for two series $\sum u_n$ and $\sum v_n$:

1. (a) $\sum v_n$ converges
 - (b) from or after a particular term $\frac{u_n}{u_{n+1}} > \frac{v_n}{v_{n+1}}$, then u_n converges.
2. (a) $\sum v_n$ diverges
 - (b) from or after a particular term $\frac{u_n}{u_{n+1}} < \frac{v_n}{v_{n+1}}$, then u_n diverges.

4.2.5 D'Alembert's Ratio Test

$$\lim_{n \rightarrow \infty} \frac{u_n}{u_{n+1}} = \lambda \quad (4.4)$$

- series converges if $\lambda < 1$
- series diverges if $\lambda > 1$
- fails if $\lambda = 1$

4.2.6 Rabbe's Test

$$\lim_{n \rightarrow \infty} n \left[\frac{u_n}{u_{n+1}} - 1 \right] = \kappa \quad (4.5)$$

- series converges if $\kappa < 1$
- series diverges if $\kappa > 1$
- fails if $\kappa = 1$

4.2.7 Cauchy's Root Test

$$\lim_{n \rightarrow \infty} |u_n| = \lambda \quad (4.6)$$

- series converges for $\lambda < 1$
- series diverges for $\lambda > 1$
- test fails for $\lambda = 1$

4.2.8 Logarithmic Test

$$\lim_{n \rightarrow \infty} n \log \left(\frac{u_n}{u_{n+1}} \right) = \kappa \quad (4.7)$$

- series converges for $\kappa < 1$
- series diverges for $\kappa > 1$
- test fails for $\kappa = 1$

Chapter 5

Determinants

5.1 Definition

For a determinant:

$$\Delta = \begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix} = a_1 \begin{vmatrix} b_2 & c_2 \\ b_3 & c_3 \end{vmatrix} - b_1 \begin{vmatrix} a_2 & c_2 \\ a_3 & c_3 \end{vmatrix} + c_1 \begin{vmatrix} a_2 & b_2 \\ a_3 & b_3 \end{vmatrix} \quad (5.1)$$

5.1.1 Minor and Cofactor

For a third order determinant $\begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}$, the minor of a_{11} is $M_{11} = \begin{vmatrix} b_2 & c_2 \\ b_3 & c_3 \end{vmatrix}$,

i.e., all the terms of the determinant except those in the same row and columns as the one of which the minor is being calculated.

$$\text{Cofactor } C_{ij} = (-1)^{i+j} M_{ij}$$

5.2 Important Properties

1. Transposing a determinant does not alter its value.
2. If rows and columns are interchanges m times, the value of the new determinant is

$$\Delta' = (-1)^m \Delta \quad (5.2)$$

3. If two parallel lines are equal, then $\Delta = 0$

4. For $\Delta = \begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}$ and $\Delta_1 = \begin{vmatrix} ka_1 & kb_1 & kc_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}$, then $\Delta_1 = k\Delta$

5. For $\Delta = \begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}$ and $\Delta_1 = \begin{vmatrix} ka_1 & b_1 & c_1 \\ ka_2 & b_2 & c_2 \\ ka_3 & b_3 & c_3 \end{vmatrix}$, then $\Delta_1 = k\Delta$

6. For $C_n \rightarrow k_1C_l + k_2C_m + k_3C_n$ or $R_n \rightarrow k_1R_l + k_2R_m + k_3R_n$, $\Delta' = \Delta$

5.3 Cramer's Rule

For a system of equations:

$$a_1x + b_1y + c_1z = d_1$$

$$a_2x + b_2y + c_2z = d_2$$

$$a_3x + b_3y + c_3z = d_3$$

the following determinants are defined:

$$D = \begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}$$

$$D_x = \begin{vmatrix} d_1 & b_1 & c_1 \\ d_2 & b_2 & c_2 \\ d_3 & b_3 & c_3 \end{vmatrix}$$

$$D_y = \begin{vmatrix} a_1 & d_1 & c_1 \\ a_2 & d_2 & c_2 \\ a_3 & d_3 & c_3 \end{vmatrix}$$

$$D_z = \begin{vmatrix} a_1 & b_1 & d_1 \\ a_2 & b_2 & d_2 \\ a_3 & b_3 & d_3 \end{vmatrix}$$

The solution of the system of equations is:

$$x = \frac{D_x}{D} \tag{5.3}$$

$$y = \frac{D_y}{D} \tag{5.4}$$

$$z = \frac{D_z}{D} \tag{5.5}$$

5.3.1 Consistency Test

1. If $D \neq 0$, the system is consistent and has unique solutions.
2. If $D = D_x = D_y = D_z = 0$, the system may or may not be consistent and it will have infinite solutions and the system will be dependent.
3. If $D = 0$ and at least one of D_x, D_y, D_z is non zero, the system is inconsistent

Chapter 6

Matrices

For a matrix,

$$A_{m \times n} = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

and where I_p is an identity matrix of the p^{th} order, the following relations are applicable.

6.1 Sum of Two Matrices

$$A_{m \times n} + B_{m \times n} = \begin{bmatrix} a_{11} + b_{11} & a_{12} + b_{12} & \cdots & a_{1n} + b_{1n} \\ a_{21} + b_{21} & a_{22} + b_{21} & \cdots & a_{2n} + b_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} + b_{m1} & a_{m2} + b_{m2} & \cdots & a_{mn} + b_{mn} \end{bmatrix} \quad (6.1)$$

6.2 Multiplication of Two Matrices

If

$$C_{m \times p} = A_{m \times n} \cdot B_{n \times p}$$

then,

$$c_{ik} = \sum_{j=1}^n a_{ij} b_{jk} \quad (6.2)$$

6.2.1 Multiplicative Properties

1. Multiplication of matrices is associative, hence $(AB)C = A(BC)$.

2. $AI = A$
3. $A \cdot A^{-1} = I$
4. $A \cdot (\text{adj} A) = (\text{adj} A) \cdot A = |A|I$
5. $A^{-1} = \frac{1}{|A|}(\text{adj} A)^t$
6. $(AB)^t = B^t A^t$

6.3 Adjoint of a Matrix

$$\text{adj} A = \begin{bmatrix} M_{11} & M_{12} & \cdots & M_{1n} \\ M_{21} & M_{22} & \cdots & M_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ M_{m1} & M_{m2} & \cdots & M_{mn} \end{bmatrix}^t, \text{ where } M_{ij} \text{ is the minor of } a_{ij} \quad (6.3)$$

6.4 Martin's Rule

For a system of equation,

$$\begin{aligned} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n &= b_1 \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n &= b_2 \\ &\vdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n &= b_n \end{aligned}$$

The system can be written as:

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix} \quad (6.4)$$

$$\Rightarrow AX = B \quad (6.5)$$

$$\Rightarrow X = A^{-1}B \quad (6.6)$$

Chapter 7

Binomial Theorem

For a binomial expansion $(a + b)^n$, there are $(n + 1)$ terms and $(a + b + c)^n$ has $\frac{(n + 1)(n + 2)}{2}$ terms.

7.1 Expansion of a binomial expression

$$\begin{aligned}(a + b)^n &= {}^nC_0 a^n b^0 + {}^nC_1 a^{n-1} b^1 + {}^nC_2 a^{n-2} b^2 + \\ &\quad \dots + {}^nC_n a^0 b^n \quad \forall n \in \mathbb{N} \\ &= \sum_{i=0}^n {}^nC_i a^{n-i} b^i \quad \forall n \in \mathbb{N}\end{aligned}\tag{7.1}$$

$$\begin{aligned}(a + b)^n &= a^n b^0 + n a^{n-1} b + \frac{n(n-1)}{2!} a^{n-2} b^2 \\ &\quad + \dots + \frac{n(n-1) \dots 3 \cdot 2 \cdot 1}{n!} a^0 b^n + \dots \infty \quad \forall n \in \mathbb{R}\end{aligned}\tag{7.2}$$

7.2 Trinomial Expansion

For $(a + b + c)^n$:

$$\begin{aligned}(a + b + c)^n &= \sum \frac{n!}{i!j!k!} a^i b^j c^k \\ &\quad \forall (i + j + k) = n; i, j, k, n \in \mathbb{N}\end{aligned}\tag{7.3}$$

7.3 Properties of Coefficients

$$\text{Sum of Co-efficients: } C_0 + C_1 + C_2 + \cdots + C_{n-1} + C_n = 2^n \quad (7.4)$$

$$\text{Sum of Odd Co-efficients: } C_0 + C_2 + C_4 + \cdots + C_{2n-3} + C_{2n-1} = 2^{n-1} \quad (7.5)$$

$$C_0 - C_1 + C_2 - \cdots + C_{2n-1} - C_{2n} = 0 \quad (7.6)$$

7.4 Pascal's Rule

For $1 \leq k \leq n$ and $k, n \in \mathbb{N}$:

$$\binom{n}{k} + \binom{n}{k-1} = \binom{n+1}{k} \quad (7.7)$$

Chapter 8

Boolean Algebra

Let B be a set of a, b, c with operations sum $(+)$ and product (\cdot) . Then B is said to belong to the Boolean Structure if the following conditions are satisfied:

Property	Name of Property
$a + b \in B$ $a \cdot b \in B$	Closure Property
$a + b = b + a$ $a \cdot b = b \cdot a$	Associative Law
$a(b + c) = ab + ac$ $a + bc = (a + b)(a + c)$	Commutative Law
$\{0, 1\} \in B$ $a + 0 = a$ $a + 1 = 1$ $a \cdot 0 = 0$ $a \cdot 1 = a$	Laws of 1 and 0
$a + ab = a$ $a(a + b) = a$	Absorption Law
$(a + b)' = (a'b')$	De'Morgan's Law

Table 8.1: Properties of Boolean Algebraic Structure

Chapter 9

Remainder Theorems

9.1 Remainder Theorem

If a function $f(x)$ is divided by a binomial $x - a$, then the remainder is provided by $f(a)$.

$$\frac{f(x)}{x - a} \equiv f(a) \pmod{(x - a)} \quad (9.1)$$

Worked Example

Find the remainder when $f(x) = x^3 - 4x^2 - 7x + 10$ is divided by $(x - 2)$.

The remainder:

$$R = (x^3 - 4x^2 - 7x + 10) \pmod{(x - 2)}$$

is given by:

$$\begin{aligned} R &= f(2) = (2)^3 - 4(2)^2 - 7(2) + 10 \\ &= 8 - 16 - 14 + 10 = -12 \end{aligned}$$

9.2 Euler's Remainder Theorem

According to Euler's Remainder Theorem, if x and n are two co-prime numbers:

$$x^{\varphi(n)} \equiv 1 \pmod{n}, x, n \in \mathbb{Z}^+ \quad (9.2)$$

where, $\varphi(n)$ is Euler's totient function.

9.2.1 Euler's Totient Function

For a number defined as:

$$n = \prod_{i=1}^r a_i^{b_i} \quad (9.3)$$

then Euler's totient function is defined as:

$$\begin{aligned} \varphi(n) &= n \cdot \left[\left(1 - \frac{1}{a_1}\right) \cdot \left(1 - \frac{1}{a_2}\right) \cdot \left(1 - \frac{1}{a_3}\right) \cdots \right] \\ &= n \prod_{i=1}^r \left(1 - \frac{1}{a_i}\right) \end{aligned} \quad (9.4)$$

Worked Example

Find the remainder if 3^{76} is divided by 35.

Since:

$$35 = 5^1 \times 7^1$$

Hence the totient quotient of 35 is:

$$\begin{aligned} \varphi(35) &= 35 \cdot \left(1 - \frac{1}{5}\right) \cdot \left(1 - \frac{1}{7}\right) \\ &= 35 \times \frac{4}{5} \times \frac{6}{7} \\ &= 24 \end{aligned}$$

Hence Euler's Theorem yields:

$$\begin{aligned} 3^{24} &\equiv 1 \pmod{35} \\ 3^{76} &\equiv 3^{24 \times 3 + 4} \\ &\equiv (3^{24})^3 \times 3^4 \pmod{35} \\ &\equiv (1)^3 \times 3^4 \pmod{35} \\ &\equiv 81 \pmod{35} \\ &\equiv 11 \pmod{35} \end{aligned}$$

The remainder when 3^{76} is divided by 35 is 11.

9.3 Wilson Theorem

According to Wilson Theorem:

$$(n-1)! \equiv -1 \pmod{n} \quad (9.5)$$

Worked Example

Find the remainder when $28!$ is divided by 31.

By Wilson's Theorem:

$$\begin{aligned}
 30! &\equiv -1 \pmod{31} \\
 \Rightarrow 30 \cdot 29 \cdot 28! &\equiv -1 \pmod{31} \\
 \text{Let } 28! \pmod{31} &= x \\
 \Rightarrow (-1) \cdot (-2) \cdot x &\equiv 30 \pmod{31} \\
 \Rightarrow 2x &= 30 \\
 \Rightarrow x &= 15
 \end{aligned}$$

The remainder when $28!$ is divided by 31 is 15.

Part II

Co-Ordinate Geometry

Chapter 10

2-D Co-ordinate Geometry

For the ordered pairs, $A(x_1, y_1)$ and $B(x_2, y_2)$:

$$AB = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2} \quad (10.1)$$

$$\text{Mid point of AB} = \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right) \quad (10.2)$$

$$\text{Point C, which divides AB in the ratio } m : n = \left(\frac{nx_1 + mx_2}{m + n}, \frac{ny_1 + my_2}{m + n} \right) \quad (10.3)$$

Chapter 11

Triangles

For a triangle defined with three vertices $A(x_1, y_1), B(x_2, y_2), C(x_3, y_3)$ and corresponding sides of length a, b, c , then:

$$\text{Centroid of } \triangle ABC = \left(\frac{x_1 + x_2 + x_3}{3}, \frac{y_1 + y_2 + y_3}{3} \right) \quad (11.1)$$

$$\text{Area of } \triangle ABC = \frac{1}{2} \begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix} \quad (11.2)$$

For a triangle, the semiperimeter, s , is defined as:

$$s = \frac{a + b + c}{2}$$

Then the radius, r , and centre of incircle, o , is:

$$o = \left(\frac{ax_1 + bx_2 + cx_3}{a + b + c}, \frac{ay_1 + by_2 + cy_3}{a + b + c} \right) \quad (11.3)$$

$$r = \sqrt{\frac{(s - a)(s - b)(s - c)}{s}} \quad (11.4)$$

The radius, R , and centre, O , of circumcircle is defined as:

$$O = \left(\frac{x_1 \sin 2A + x_2 \sin 2B + x_3 \sin 2C}{\sin 2A + \sin 2B + \sin 2C}, \frac{y_1 \sin 2A + y_2 \sin 2B + y_3 \sin 2C}{\sin 2A + \sin 2B + \sin 2C} \right) \quad (11.5)$$

$$2R = \frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C} \quad (11.6)$$

Chapter 12

Straight Line

A straight line can be defined as:

$$y = mx + c \quad (12.1)$$

$$\frac{x}{a} + \frac{y}{b} = 1, \text{ where } a \text{ and } b \text{ are the intercepts at x and y axes respectively} \quad (12.2)$$

$$x \cos \alpha + y \sin \alpha = p \text{ (Normal Form)} \quad (12.3)$$

$$Ax + By + C = 0 \text{ (General Form)} \quad (12.4)$$

Equation of Straight Line Passing Through (x_0, y_0) and Slope m

$$(y - y_0) = m(x - x_0) \quad (12.5)$$

Distance Between Two Points on a Line

$$\frac{y_1 - y_2}{\sin \theta} = \frac{x_1 - x_2}{\cos \theta} = \gamma \quad (12.6)$$

$$\theta = \tan^{-1} m \quad (12.7)$$

Angle Between Two Lines

For two lines with slopes m_1, m_2 , the angle between them, θ :

$$\theta = \arctan\left(\frac{m_1 - m_2}{1 + m_1 m_2}\right) \quad (12.8)$$

Distance of a Point from a Line

Line: $ax + by + c = 0$ Point: (g, h)

$$S = \frac{ag + bh + c}{\sqrt{a^2 + b^2}} \quad (12.9)$$

Angle Bisector of a Line For the two lines: $a_1x + b_1y + c_1 = 0$ and $a_2x + b_2y + c_2 = 0$, the angle bisector is:

$$\frac{a_1x + b_1y + c_1}{\sqrt{a_1^2 + b_1^2}} = \frac{a_2x + b_2y + c_2}{\sqrt{a_2^2 + b_2^2}} \quad (12.10)$$

If the sign of c_1 and c_2 is the same, then the equation obtained is the internal bisector.

Equation of a Straight Line Passing through the Intersection of Two Lines

$$(a_1x + b_1y + c_1) + k(a_2x + b_2y + c_2) = 0 \quad \forall k \in \mathbb{R} \quad (12.11)$$

Relative Position of Points w.r.t. a Line For the points (x_1, y_1) and (x_2, y_2) :

$$\begin{aligned} k_1 &= ax_1 + by_1 + c \\ k_2 &= ax_2 + by_2 + c \end{aligned}$$

If k_1 and k_2 have the same sign, they are on the same side of a line, otherwise on opposite sides.

Ratio of Division of Line Segment For any line, $f(x, y) = 0$, the ratio in which it divides (x_1, y_1) and (x_2, y_2) is given by:

$$r = -\frac{f(x_1, y_1)}{f(x_2, y_2)} \quad (12.12)$$

If $\begin{cases} r > 0, \text{ then division is internal} \\ r < 0, \text{ then division is external} \end{cases}$.

Chapter 13

General Theory of Second Degree Equation

For any general equation of the form:

$$ax^2 + by^2 + 2gx + 2fy + 2hxy + c = 0 \quad (13.1)$$

Δ is defined as:

$$\Delta = \begin{vmatrix} a & h & g \\ h & b & f \\ g & f & c \end{vmatrix} \quad (13.2)$$

If $\Delta = 0$ then the equation is a pair of straight lines. If $a + b = 0$ then the lines are \perp .

If the $\Delta \neq 0$:

1. $a = b, h = 0 \rightarrow$ circle
2. $h^2 = ab \rightarrow$ parabola
3. $h^2 < ab \rightarrow$ ellipse
4. $h^2 > ab \rightarrow$ hyperbola

Chapter 14

Conics

The four conic sections are: circle, parabola, ellipse, and hyperbola. Circle has been done separately in the next chapter.

14.1 Parabola

Property	$y^2 = 4ax$	$x^2 = 4ay$
Axis	$y = 0$	$x = 0$
Eccentricity	1	1
Directrix	$x + a = 0$	$y + a = 0$
Focus	$(a, 0)$	$(0, a)$
Vertex	$(0, 0)$	$(0, 0)$
Length of latus rectum	$ 4a $	$ 4a $
Equation of latus rectum	$x - a = 0$	$y - a = 0$

Table 14.1: Properties of a Parabola

14.2 Ellipse and Hyperbola

For $a > b$:

Property	$\frac{x^2}{a} + \frac{y^2}{b} = 1$ Ellipse	$\frac{x^2}{a} - \frac{y^2}{b} = 1$ Hyperbola
Length of Major Axis	$2a$	$2a$
Length of Minor Axis	$2b$	$2b$
Equation of Major Axis	$x = 0$	$x = 0$
Equation of Minor Axis	$y = 0$	$y = 0$
Eccentricity e	$\sqrt{1 - \frac{b^2}{a^2}}$	$\sqrt{1 + \frac{b^2}{a^2}}$
Vertices	$(\pm a, 0)$	$(\pm a, 0)$
Foci	$(\pm ae, 0)$	$(\pm ae, 0)$
Equation of Directrix	$x \pm \frac{a}{e} = 0$	$x = \pm \frac{a}{e}$
Length of latus rectum	$\frac{2b^2}{a}$	$\frac{2b^2}{a}$
Equation of latus rectum	$x \pm ae = 0$	
Centre	$(0, 0)$	$(0, 0)$

Table 14.2: Properties of Ellipse and Hyperbola

14.3 Parametric Form of Conics

14.3.1 Hyperbola

$$x = a \sec \theta \quad (14.1)$$

$$y = b \tan \theta \quad (14.2)$$

14.3.2 Ellipse

$$x = a \cos \phi \quad (14.3)$$

$$y = b \sin \phi \quad (14.4)$$

14.3.3 Parabola

$$x = at^2 \quad (14.5)$$

$$y = 2at \quad (14.6)$$

Chapter 15

Circles

15.1 Locus Form

$$(x - g)^2 + (y - h)^2 = r^2 \quad (15.1)$$

where the centre is (g, h) and the radius is r .

15.2 Diameter Form

$$(x - a)(x - c) + (y - b)(y - d) = 0 \quad (15.2)$$

where (a, b) and (c, d) are the two ends of the diameter.

15.3 General Form

If the equation of a circle is in the form:

$$x^2 + y^2 + 2gx + 2fy + c = 0 \quad (15.3)$$

Then the following is true about the circle:

1. centre of the circle is $(-g, -f)$
2. radius of circle is $\sqrt{g^2 + f^2 - c}$

15.4 Important Relations

1. If the circle passes through the origin, $g = 0, f = 0$.
2. If the circle touches the x-axis $c = g^2$.
3. If the circle touches the y-axis $c = f^2$.

Common for Two Circles

1. The common chord passing between two circles S_1 and S_2 are:

$$S_1 - S_2 = 0 \quad (15.4)$$

2. Circles passing through the intersection of two circles is:

$$S_2 + k(S_1 - S_2) = 0 \quad \forall k \in \mathbb{R} \quad (15.5)$$

Chapter 16

Vectors

Let two vectors be $\vec{a} = a\hat{i} + b\hat{j} + c\hat{k}$ and $\vec{b} = x\hat{i} + y\hat{j} + z\hat{k}$:

16.1 Modulus of a Vector

For a vector \vec{a} , the modulus of the vector is:

$$|\vec{a}| = \sqrt{a^2 + b^2 + c^2} \quad (16.1)$$

16.2 Sum of Vectors

The sum of two vectors is:

$$|\vec{a} + \vec{b}| = \sqrt{|\vec{a}|^2 + |\vec{b}|^2 + 2|\vec{a}||\vec{b}|\cos\theta} \quad (16.2)$$

$$\vec{a} + \vec{b} = (a + x)\hat{i} + (b + y)\hat{j} + (c + z)\hat{k} \quad (16.3)$$

The direction of the resultant vector is:

$$\tan\alpha = \frac{b\sin\theta}{a + b\cos\theta} \quad (16.4)$$

where, θ is the angle between the two vectors.

16.3 Product of Vectors

16.3.1 Dot Product

$$\vec{a} \cdot \vec{b} = |a||b|\cos\theta \quad (16.5)$$

$$\vec{a} \cdot \vec{b} = ax + by + cz \quad (16.6)$$

16.3.2 Cross Product

$$\vec{a} \times \vec{b} = |a||b| \sin \theta \hat{n} \quad (16.7)$$

where \hat{n} is a vector $\perp \vec{a}, \vec{b}$.

$$\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a & b & c \\ x & y & z \end{vmatrix} \quad (16.8)$$

16.4 Test of Co-planarity

Three vectors are called co-planar if:

$$\lambda \vec{a} + \mu \vec{b} = \vec{c} \quad (16.9)$$

$$(\vec{a} \times \vec{b}) \cdot \vec{c} = 0 \quad (16.10)$$

Chapter 17

3-D Geometry

17.1 Distance between two points $A(x_1, y_1, z_1)$ and $B(x_2, y_2, z_2)$:

$$AB = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2 + (z_1 - z_2)^2} \quad (17.1)$$

17.2 Section Formula of a Line Segment Divided in the ratio $m : n$

$$P = \left(\frac{nx_1 + mx_2}{m + n}, \frac{ny_1 + my_2}{m + n}, \frac{nz_1 + mz_2}{m + n} \right) \quad (17.2)$$

17.3 Centroid of a Triangle

$$G = \left(\frac{x_1 + x_2 + x_3}{3}, \frac{y_1 + y_2 + y_3}{3}, \frac{z_1 + z_2 + z_3}{3} \right) \quad (17.3)$$

Chapter 18

Line in 3-D Space

For a line which is defined as $\vec{a} = a\hat{i} + b\hat{j} + c\hat{k}$:

1. Line numbers of the line is

$$< a, b, c > \quad (18.1)$$

2. The line cosines are:

$$< \frac{a}{\sqrt{a^2 + b^2 + c^2}}, \frac{b}{\sqrt{a^2 + b^2 + c^2}}, \frac{c}{\sqrt{a^2 + b^2 + c^2}} > \quad (18.2)$$

$$=< l, m, n > \quad (18.3)$$

18.1 Angle between Two Lines

$$\cos \theta = \frac{a_1a_2 + b_1b_2 + c_1c_2}{\sqrt{a_1^2 + b_1^2 + c_1^2} \sqrt{a_2^2 + b_2^2 + c_2^2}} \quad (18.4)$$

$$\Rightarrow \cos \theta = l_1l_2 + m_1m_2 + n_1n_2 \quad (18.5)$$

When two lines are \perp , $l_1l_2 + m_1m_2 + n_1n_2 = 0$.

When two lines are \parallel $\frac{l_1}{l_2} = \frac{m_1}{m_2} = \frac{n_1}{n_2} = 1$.

18.2 Skew and Co-planar Lines

Let there be two lines \vec{r}_1 and \vec{r}_2 ,

$$\vec{r}_1 = \vec{a}_1 + \mu \vec{b}_1, \vec{r}_2 = \vec{a}_2 + \lambda \vec{b}_2 \quad (18.6)$$

18.3 Distances

18.3.1 The shortest distance between r_1 and r_2

$$S = \left| \frac{(\vec{a}_1 - \vec{a}_2) \cdot (\vec{b}_1 \times \vec{b}_2)}{|\vec{b}_1 \times \vec{b}_2|} \right| \quad (18.7)$$

If $S = 0$, the lines intersect.

18.3.2 Cartesian Form

For two lines defined as $\frac{x - x_1}{a_1} = \frac{y - y_1}{b_1} = \frac{z - z_1}{c_1}$ and $\frac{x - x_2}{a_2} = \frac{y - y_2}{b_2} = \frac{z - z_2}{c_2}$:

$$S = \begin{vmatrix} x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} \quad (18.8)$$

18.3.3 Distance Between Parallel Lines

$$S = \left| \frac{\vec{b} \cdot (\vec{a}_2 - \vec{a}_1)}{|\vec{b}|} \right| \quad (18.9)$$

18.3.4 Distance of a Point to a Line

For a point, (x_1, y_1, z_1) the distance to a line $\frac{x - \alpha}{l} = \frac{y - \beta}{m} = \frac{z - \gamma}{n}$:

$$S = \left(\left| \begin{vmatrix} x_1 - \alpha & y_1 - \beta \\ l & m \end{vmatrix} \right| + \left| \begin{vmatrix} y_1 - \beta & z_1 - \gamma \\ m & n \end{vmatrix} \right| + \left| \begin{vmatrix} z_1 - \gamma & x_1 - \alpha \\ n & l \end{vmatrix} \right| \right)^{\frac{1}{2}} \quad (18.10)$$

Chapter 19

3-D Plane

A plane in 3-D space can be defined as:

1. Cartesian Form:

$$ax + by + cz + d = 0 \quad (19.1)$$

2. Vectorial Form:

$$\vec{r} \cdot \vec{n} = p \quad (19.2)$$

, where \vec{r} is a line on the plane, \vec{n} is a normal to the plane, and p is perpendicular distance to the plane from the origin.

19.1 Angle Between Two Planes

For two planes, $\vec{r}_1 \cdot \vec{n}_1 = p_1$ and $\vec{r}_2 \cdot \vec{n}_2 = p_2$, the angle between the planes, θ is:

$$\cos \theta = \frac{\vec{n}_1 \cdot \vec{n}_2}{|\vec{n}_1||\vec{n}_2|} \quad (19.3)$$

In the Cartesian Form:

$$\cos \theta = \frac{a_1 a_2 + b_1 b_2 + c_1 c_2}{\sqrt{a_1^2 + b_1^2 + c_1^2} \sqrt{a_2^2 + b_2^2 + c_2^2}} \quad (19.4)$$

19.2 Distance of a Point from a Plane

19.2.1 Cartesian Form

For the point (p, q, r) and the plane, $ax + by + cz + d = 0$:

$$S = \frac{ap + bq + cr + d}{\sqrt{a^2 + b^2 + c^2}} \quad (19.5)$$

19.2.2 Vector Form

For the point $\vec{g} = p\hat{i} + q\hat{j} + r\hat{k}$ and the plane $\vec{r} \cdot (a\hat{i} + b\hat{j} + c\hat{k}) + d = 0$:

$$S = \frac{(a\hat{i} + b\hat{j} + c\hat{k}) \cdot (p\hat{i} + q\hat{j} + r\hat{k})}{\sqrt{a^2 + b^2 + c^2}} \quad (19.6)$$

$$\Rightarrow S = \frac{(a\hat{i} + b\hat{j} + c\hat{k}) \cdot \vec{g}}{|a\hat{i} + b\hat{j} + c\hat{k}|} \quad (19.7)$$

Part III

Statistics

Chapter 20

Statistics

For a set a data $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n), \dots$:

Mean of x :

$$\text{bar } x = \frac{x_1 + x_2 + \dots + x_n}{n} = \frac{1}{n} \sum_{i=1}^n x_i \quad (20.1)$$

Variance of x :

$$\sigma^2 = \frac{(x_1 - \bar{x})^2 + ((x_2 - \bar{x})^2 + \dots + ((x_n - \bar{x})^2}{n} = \sum_{i=1}^n \frac{(x_i - \bar{x})^2}{n} = \frac{\sum x_i^2}{n} - \bar{x}^2 \quad (20.2)$$

Standard Deviation of x :

$$\begin{aligned} \sigma &= \sqrt{\frac{(x_1 - \bar{x})^2 + ((x_2 - \bar{x})^2 + \dots + ((x_n - \bar{x})^2}{n}} \\ &= \sqrt{\sum_{i=1}^n \frac{(x_i - \bar{x})^2}{n}} \\ &= \sqrt{\frac{\sum x_i^2}{n} - \bar{x}^2} \end{aligned} \quad (20.3)$$

Covariance of (x, y) :

$$\text{Cov}(x, y) = \frac{\sum_{i=1}^N (x_i - \bar{x})(y_i - \bar{y})}{N} = \sum xy - \frac{1}{N} \sum x \sum y \quad (20.4)$$

Correlation Co-efficient, $\gamma(x, y)$:

$$\gamma(x, y) = \frac{\text{Cov}(x, y)}{\sigma_x \sigma_y} \quad (20.5)$$

Chapter 21

Lines of Regression

An assumption is made for the line of regression. It is assumed to be:

$$y = ax + b$$

For a given set of data (x_i, y_i) , the solutions of a and b are obtained by solving the following equations simultaneously:

$$\sum y_i = a \sum x_i + nb \quad (21.1)$$

$$\sum x_i y_i = a \sum x_i^2 + b \sum x_i \quad (21.2)$$

If the regressive function is defined as:

$$y = cx^a \quad (21.3)$$

, where c is a constant, then the following conversions are performed:

$$y = cx^a \quad (21.4)$$

$$\Rightarrow \log y = \log c + a \log x \quad (21.5)$$

Making the substitutions $\log y = Y$, $\log x = X$, and $\log c = C$, the required equation becomes:

$$Y = aX + C \quad (21.6)$$

This transformed equation can be solved using the method describes in equations ?? and 21.2.

21.1 Karl Pearson's Co-efficient of Correlation (20.5)

$$r = \rho(x, y) = \frac{Cov(x, y)}{\sigma_x \sigma_y} \quad (21.7)$$

21.1.1 Degree of Correlation

Value	Relation
$0 \leq r < \frac{1}{4}$	Low
$\frac{1}{4} \leq r < \frac{3}{4}$	Moderate
$\frac{3}{4} \leq r \leq 1$	High

Table 21.1: Degree of Correlation

Part IV

Trigonometry

Chapter 22

Circular Trigonometric Functions

θ	$\sin \theta$	$\cos \theta$	$\tan \theta$
0°	0	1	0
15°	$\frac{1}{4}$	$\frac{1}{4(2-\sqrt{3})}$	$2 - \sqrt{3}$
18°	$\frac{\sqrt{5}-1}{4}$	$\frac{\sqrt{10+2\sqrt{5}}}{4}$	$\frac{\sqrt{5}-1}{\sqrt{10+2\sqrt{5}}}$
30°	$\frac{1}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{1}{\sqrt{3}}$
36°	$\frac{\sqrt{5}+1}{4}$	$\frac{\sqrt{10-2\sqrt{5}}}{4}$	$\frac{\sqrt{5}+1}{\sqrt{10-2\sqrt{5}}}$
45°	$\frac{1}{\sqrt{2}}$	$\frac{1}{\sqrt{2}}$	1
60°	$\frac{1}{\sqrt{3}}$	$\frac{1}{2}$	$\sqrt{3}$
72°	$\frac{\sqrt{10+2\sqrt{5}}}{4}$	$\frac{\sqrt{5}-1}{4}$	$\frac{\sqrt{10+2\sqrt{5}}}{\sqrt{5}-1}$
90°	1	0	∞

Table 22.1: Trigonometric Ratios of Standard Angles

For any given triangle:

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C} = 2R \quad (22.1)$$

, where $2R$ is the radius of circumcircle.

22.1 Negative Angle Formula

$$\sin(-\theta) = -\sin \theta \quad (22.2)$$

$$\cos(-\theta) = \cos \theta \quad (22.3)$$

$$\tan(-\theta) = -\tan \theta \quad (22.4)$$

$$\csc(-\theta) = -\csc \theta \quad (22.5)$$

$$\sec(-\theta) = \sec \theta \quad (22.6)$$

$$\cot(-\theta) = -\cot \theta \quad (22.7)$$

22.2 Sum of Angles Formula

$$\sin(A + B) = \sin A \cos B + \cos A \sin B \quad (22.8)$$

$$\cos(A + B) = \cos A \cos B - \sin A \sin B \quad (22.9)$$

$$\tan(A + B) = \frac{\tan A + \tan B}{1 - \tan A \tan B} \quad (22.10)$$

22.3 Difference of Angles Formula

$$\sin(A - B) = \sin A \cos B - \cos A \sin B \quad (22.11)$$

$$\cos(A - B) = \cos A \cos B + \sin A \sin B \quad (22.12)$$

$$\tan(A - B) = \frac{\tan A - \tan B}{1 + \tan A \tan B} \quad (22.13)$$

22.4 Multiples and Sub-multiples of π and $\frac{\pi}{2}$

$$\forall k \in \mathbb{Z}$$

$$\sin\left((4k + 1)\frac{\pi}{2}\right) = 1 \quad (22.14)$$

$$\sin\left((4k - 1)\frac{\pi}{2}\right) = -1 \quad (22.15)$$

$$\sin k\pi = 0 \quad (22.16)$$

$$\sin\left((2k + 1)\frac{\pi}{2}\right) = 0 \quad (22.17)$$

$$\sin\left((2k - 1)\frac{\pi}{2}\right) = 0 \quad (22.18)$$

$$\sin k\pi = (-1)^k \quad (22.19)$$

22.5 $\left(\frac{\pi}{2} \pm \theta\right)$ Formula

$$\sin\left(\frac{\pi}{2} - \theta\right) = \cos \theta \quad (22.20)$$

$$\sin\left(\frac{\pi}{2} + \theta\right) = \cos \theta \quad (22.21)$$

$$\cos\left(\frac{\pi}{2} - \theta\right) = \sin \theta \quad (22.22)$$

$$\cos\left(\frac{\pi}{2} + \theta\right) = -\sin \theta \quad (22.23)$$

$$\tan\left(\frac{\pi}{2} - \theta\right) = \cot \theta \quad (22.24)$$

$$\tan\left(\frac{\pi}{2} + \theta\right) = -\cot \theta \quad (22.25)$$

$$\cot\left(\frac{\pi}{2} - \theta\right) = \tan \theta \quad (22.26)$$

$$\cot\left(\frac{\pi}{2} + \theta\right) = -\tan \theta \quad (22.27)$$

$$\csc\left(\frac{\pi}{2} - \theta\right) = \sec \theta \quad (22.28)$$

$$\csc\left(\frac{\pi}{2} + \theta\right) = \sec \theta \quad (22.29)$$

$$\sec\left(\frac{\pi}{2} - \theta\right) = \csc \theta \quad (22.30)$$

$$\sec\left(\frac{\pi}{2} + \theta\right) = -\csc \theta \quad (22.31)$$

22.6 $\left(\frac{\pi}{4} \pm \theta\right)$ Formula

$$\tan\left(\frac{\pi}{4} + \theta\right) = \frac{1 + \tan \theta}{1 - \tan \theta} \quad (22.32)$$

$$\tan\left(\frac{\pi}{4} - \theta\right) = \frac{1 - \tan \theta}{1 + \tan \theta} \quad (22.33)$$

22.7 Trigonometric Identities

$$\sin^2 \theta + \cos^2 \theta = 1 \quad (22.34)$$

$$\tan^2 \theta + 1 = \sec^2 \theta \quad (22.35)$$

$$\cot^2 \theta + 1 = \csc^2 \theta \quad (22.36)$$

22.8 Double Angle Formula

$$\begin{aligned}\sin 2\theta &= 2 \sin \theta \cos \theta \\ &= \frac{2 \tan \theta}{1 + \tan^2 \theta}\end{aligned}\tag{22.37}$$

$$\begin{aligned}\cos 2\theta &= \cos^2 \theta - \sin^2 \theta \\ &= 2 \cos^2 \theta - 1 \\ &= 1 - 2 \sin^2 \theta \\ &= \frac{1 - \tan^2 \theta}{1 + \tan^2 \theta}\end{aligned}\tag{22.38}$$

$$\tan 2\theta = \frac{2 \tan \theta}{1 - \tan^2 \theta}\tag{22.39}$$

22.9 Triple Angle Formula

$$\sin 3\theta = 3 \sin \theta - 4 \sin^3 \theta\tag{22.40}$$

$$\cos 3\theta = 4 \cos^3 \theta - 3 \cos \theta\tag{22.41}$$

$$\tan 3\theta = \frac{3 \tan \theta - \tan^3 \theta}{1 - 3 \tan^2 \theta}\tag{22.42}$$

22.10 Sum and Product of Two Ratios

For $A > B$:

$$\sin A + \sin B = 2 \sin \left(\frac{A+B}{2} \right) \cos \left(\frac{A-B}{2} \right) \quad (22.43)$$

$$\sin A - \sin B = 2 \cos \left(\frac{A+B}{2} \right) \sin \left(\frac{A-B}{2} \right) \quad (22.44)$$

$$2 \sin A \cos B = \sin(A+B) + \sin(A-B) \quad (22.45)$$

$$2 \cos A \sin B = \sin(A+B) - \sin(A-B) \quad (22.46)$$

$$\cos A + \cos B = 2 \cos \left(\frac{A+B}{2} \right) \cos \left(\frac{A-B}{2} \right) \quad (22.47)$$

$$\cos A - \cos B = -2 \sin \left(\frac{A+B}{2} \right) \sin \left(\frac{A-B}{2} \right) \quad (22.48)$$

$$2 \cos A \cos B = \cos(A+B) + \cos(A-B) \quad (22.49)$$

$$2 \cos A \sin B = \cos(A+B) - \cos(A-B) \quad (22.50)$$

$$\sin(A-B) \sin(A+B) = \sin^2 A - \sin^2 B \quad (22.51)$$

$$\cos(A-B) \cos(A+B) = \cos^2 A - \sin^2 B \quad (22.52)$$

$$\tan(A-B) \tan(A+B) = \frac{\tan^2 A - \tan^2 B}{1 - \tan^2 A \tan^2 B} \quad (22.53)$$

22.11 General Solutions

If $\sin \theta = \sin \alpha$:

$$\theta = n\pi + (-1)^n \alpha \quad (22.54)$$

$n \in \mathbb{Z}$

If $\cos \theta = \cos \alpha$:

$$\theta = 2n\pi \pm \alpha \quad (22.55)$$

$n \in \mathbb{Z}$

If $\tan \theta = \tan \alpha$:

$$\theta = n\pi \pm \alpha \quad (22.56)$$

$n \in \mathbb{Z}$

22.12 Taylor Series Expansion of Trigonometric Ratios

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \cdots \infty = \sum_{i=1}^{\infty} (-1)^{i+1} \frac{x^{2i-1}}{(2i-1)!} \quad (22.57)$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \cdots \infty = \sum_{i=0}^{\infty} (-1)^i \frac{x^{2i}}{(2i)!} \quad (22.58)$$

Chapter 23

Inverse Circular Trigonometric Function

23.1 Definition of Inverse Circular Trigonometric Function

23.1.1 For $\sin x$

$y = \arcsin x$ iff $x = \sin y$, then:

1. $y \in [-\frac{\pi}{2}, \frac{\pi}{2}]$
2. domain of $x \in [-1, 1]$
3. $\sin(\arcsin x) = x, \forall x \in [-1, 1]$
4. $\arcsin(\sin y) = y, \forall y \in [-\frac{\pi}{2}, \frac{\pi}{2}]$
5. $\sin x$ is a strictly increasing in the domain $[-\frac{\pi}{2}, \frac{\pi}{2}]$ and one-one.

23.1.2 For $\cos x$

$y = \arccos x$ iff $x = \cos y$, then:

1. $y \in [0, \pi]$
2. domain of $x \in [-1, 1]$
3. $\cos(\arccos x) = x, \forall x \in [-1, 1]$
4. $\arccos(\cos y) = y, \forall y \in [0, \pi]$
5. $\cos x$ is a strictly decreasing in the domain $[0, \pi]$ and one-one.

23.1.3 For $\tan x$

$y = \arctan x$ iff $x = \tan y$, then:

1. $y \in [-\frac{\pi}{2}, \frac{\pi}{2}]$
2. domain of $x \in \mathbb{R}$
3. $\tan(\arctan x) = x, \forall x \in \mathbb{R}$
4. $\arctan(\tan y) = y, \forall y \in [-\frac{\pi}{2}, \frac{\pi}{2}]$
5. $\tan x$ is a strictly increasing in the domain $[-\frac{\pi}{2}, \frac{\pi}{2}]$ and one-one.

23.1.4 For $\cot x$

$y = \cot^{-1} x$ iff $x = \cot y$, then:

1. $y \in (0, \pi)$
2. domain of $x \in \mathbb{R}$
3. $\cot(\cot^{-1} x) = x, \forall x \in \mathbb{R}$
4. $\cot^{-1}(\cot y) = y, \forall y \in (0, \pi)$
5. $\cot x$ is a strictly decreasing in the domain $(0, \pi)$ and one-one.

For $\sec x$

$y = \sec^{-1} x$ iff $x = \sec y$

1. $y \in \{[0, \frac{\pi}{2}) \cup (\frac{\pi}{2}, \pi]\}$
2. $x \in \{(-\infty, -1] \cup [1, \infty)\}$
3. $\sec(\sec^{-1} x) = x, \forall |x| \geq 1$
4. $\sec^{-1}(\sec y) = y, \forall y \in \{[0, \frac{\pi}{2}) \cup (\frac{\pi}{2}, \pi]\}$

23.1.5 For $\csc x$

$y = \csc^{-1} x$ iff $x = \csc y$

1. $y \in \{[-\frac{\pi}{2}, 0) \cup (0, \frac{\pi}{2}]\}$
2. $x \in \{(-\infty, -1] \cup [1, \infty)\}$
3. $\csc(\csc^{-1} x) = x, \forall |x| \geq 1$
4. $\csc^{-1}(\csc y) = y, \forall y \in \{[-\frac{\pi}{2}, 0) \cup (0, \frac{\pi}{2}]\}$

23.2 Negative Arguments

$$\arcsin(-x) = -\arcsin x \quad (23.1)$$

$$\arctan(-x) = -\arctan x \quad (23.2)$$

$$\csc^{-1}(-x) = -\csc^{-1} x \quad (23.3)$$

$$\arccos(-x) = \pi - \arccos x \quad (23.4)$$

$$\cot^{-1}(-x) = \pi - \cot^{-1} x \quad (23.5)$$

$$\sec^{-1}(-x) = \pi - \sec^{-1} x \quad (23.6)$$

23.3 Reciprocal Relations

$$\csc^{-1} x = \arcsin \frac{1}{x} \quad (23.7)$$

$$\sec^{-1} x = \arccos \frac{1}{x} \quad (23.8)$$

$$\sec^{-1} x = \begin{cases} \arctan \frac{1}{x}, & x > 0 \\ \pi + \arctan \frac{1}{x}, & x < 0 \end{cases} \quad (23.9)$$

23.4 I.T.F. Identities

$$\arcsin x + \arccos x = \frac{\pi}{2}, |x| \leq 1 \quad (23.10)$$

$$\arctan x + \cot^{-1} x = \frac{\pi}{2}, x \in \mathbb{R} \quad (23.11)$$

$$\sec^{-1} x + \csc^{-1} x = \frac{\pi}{2}, |x| \geq 1 \quad (23.12)$$

23.5 Sum of Two Angles

$$\arctan x + \arctan y = \arctan \left(\frac{x+y}{1-xy} \right) \quad (23.13)$$

$$\arcsin x + \arcsin y = \arcsin(y\sqrt{1-x^2} + x\sqrt{1-y^2}) \quad (23.14)$$

$$\arccos x + \arccos y = \arccos(xy - \sqrt{1-x^2}\sqrt{1-y^2}) \quad (23.15)$$

23.6 Difference of Two Angles

$$\arctan x - \arctan y = \arctan \left(\frac{x - y}{1 + xy} \right) \quad (23.16)$$

$$\arcsin x - \arcsin y = \arcsin(x\sqrt{1 - y^2} - y\sqrt{1 - x^2}) \quad (23.17)$$

$$\arccos x - \arccos y = \arccos(xy + \sqrt{1 - x^2}\sqrt{1 - y^2}) \quad (23.18)$$

23.7 Interconversion of Ratios

$$\begin{aligned} \arcsin x &= \arccos \sqrt{1 - x^2} \\ &= \arctan \left(\frac{x}{\sqrt{1 - x^2}} \right) \end{aligned} \quad (23.19)$$

$$\begin{aligned} \arccos x &= \arcsin \sqrt{1 - x^2} \\ &= \arctan \left(\frac{\sqrt{1 - x^2}}{x} \right) \end{aligned} \quad (23.20)$$

$$\begin{aligned} 2 \arctan x &= \arcsin \left(\frac{2x}{1 + x^2} \right) \\ &= \arccos \left(\frac{1 - x^2}{1 + x^2} \right) \\ &= \arctan \left(\frac{2x}{1 - x^2} \right) \end{aligned} \quad (23.21)$$

23.8 Miscellaneous Relations

$$\cos(\arcsin x) = \sin(\arccos x) = \sqrt{1 - x^2} \quad (23.22)$$

$$\arctan x = \frac{\pi}{2} - \arctan \left(\frac{1}{x} \right), x > 1 \quad (23.23)$$

Chapter 24

Hyperbolic Trigonometric Function

24.1 Definition

Hyperbolic trigonometric functions are defined such that $(\cosh t, \sinh t)$ form the right half of an equilateral hyperbola. The functions are defined as follows:

$$\sinh x = \frac{\exp(x) - \exp(-x)}{2} \quad (24.1)$$

$$\cosh x = \frac{\exp(x) + \exp(-x)}{2} \quad (24.2)$$

$$\tanh x = \frac{\sinh x}{\cosh x} = \frac{\exp(x) - \exp(-x)}{\exp(x) + \exp(-x)} \quad (24.3)$$

$$\coth x = \frac{1}{\tanh x} = \frac{\exp(x) + \exp(-x)}{\exp(x) - \exp(-x)} \quad (24.4)$$

$$csch x = \frac{1}{\sinh x} = \frac{2}{\exp(x) - \exp(-x)} \quad (24.5)$$

$$sech x = \frac{1}{\cosh x} = \frac{2}{\exp(x) + \exp(-x)} \quad (24.6)$$

24.2 Identities

$$\coth^2 x - \sinh^2 x = 1 \quad (24.7)$$

$$\tanh^2 x + sech^2 x = 1 \quad (24.8)$$

$$\coth^2 x - csch^2 x = 1 \quad (24.9)$$

24.3 Inverse Hyperbolic Function

$$\sinh^{-1} z = \ln(z + \sqrt{z^2 + 1}) \quad (24.10)$$

$$\cosh^{-1} z = \ln(z \pm \sqrt{z^2 - 1}) \quad (24.11)$$

$$\tanh^{-1} z = \frac{1}{2} \ln \left(\frac{1+z}{1-z} \right) \quad (24.12)$$

$$\coth^{-1} z = \frac{1}{2} \ln \left(\frac{z+1}{z-1} \right) \quad (24.13)$$

$$\operatorname{csch}^{-1} z = \ln \left(\frac{1 \pm \sqrt{z^2 + 1}}{z} \right) \quad (24.14)$$

$$\operatorname{sech}^{-1} z = \ln \left(\frac{1 \pm \sqrt{1 - z^2}}{2} \right) \quad (24.15)$$

24.4 Relation to Circular Trigonometric Functions

$$\sinh(z) = -i \sin(iz) \quad (24.16)$$

$$\cosh(z) = \cos(iz) \quad (24.17)$$

$$\tanh(z) = -i \tan(iz) \quad (24.18)$$

$$\operatorname{csch}(z) = i \csc(iz) \quad (24.19)$$

$$\operatorname{sech}(z) = \sec(iz) \quad (24.20)$$

$$\coth(z) = i \cot(iz) \quad (24.21)$$

Part V

Calculus

Chapter 25

Limits

$$\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1 \quad (25.1)$$

$$\lim_{x \rightarrow 0} \frac{\tan x}{x} = 1 \quad (25.2)$$

$$\lim_{x \rightarrow 0} \cos x = 1 \quad (25.3)$$

$$\lim_{x \rightarrow a} \frac{x^n - a^n}{x - a} = na^{n-1} \quad (25.4)$$

$$\lim_{x \rightarrow a} f(x)g(x) = \lim_{x \rightarrow a} f(x) \lim_{x \rightarrow a} g(x) \quad (25.5)$$

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow a} f(x)}{\lim_{x \rightarrow a} g(x)}, \lim_{x \rightarrow a} g(x) \neq 0 \quad (25.6)$$

$$\lim_{x \rightarrow 0} \exp(x) = 1 \quad (25.7)$$

$$\lim_{x \rightarrow a} \exp(x) = \exp(c) \quad (25.8)$$

$$\lim_{x \rightarrow 0} \frac{\exp(x) - 1}{x} = 1 \quad (25.9)$$

$$\lim_{x \rightarrow a} c^x = c^a \quad (25.10)$$

$$\lim_{x \rightarrow a} \ln x = \ln a \quad (25.11)$$

$$\lim_{x \rightarrow 0} (1+x)^{\frac{1}{x}} = e \quad (25.12)$$

$$\lim_{x \rightarrow 0} \frac{\ln(1+x)}{x} = 1 \quad (25.13)$$

$$\lim_{x \rightarrow 0} \frac{a^x - 1}{x} = \ln a \quad (25.14)$$

$$\lim_{n \rightarrow \infty} \frac{x^n}{n!} = 0, \forall x \in \mathbb{R} \quad (25.15)$$

25.1 L'Hospital Rule

If:

$$L = \lim_{x \rightarrow a} \frac{f(x)}{g(x)}$$

is such that $f(a) = 0$ and $g(a) = 0$, or $f(a) = \infty$ and $g(a) = \infty$, then:

$$L = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$$

Chapter 26

Differentiation

26.1 Differentiation by First Principle

$$\frac{df(x)}{dx} = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \quad (26.1)$$

26.2 Standard Differentiation Formulae

$$\frac{dk}{dx} = 0 \quad (26.2)$$

$$\frac{dx^n}{dx} = nx^{n-1} \quad (26.3)$$

$$\frac{da^x}{dx} = \ln a \cdot a^x \quad (26.4)$$

$$\frac{d \exp(x)}{dx} = \exp(x) \quad (26.5)$$

$$\frac{d \ln x}{dx} = \frac{1}{x} \quad (26.6)$$

$$\frac{d\sqrt{x}}{dx} = \frac{1}{2\sqrt{x}} \quad (26.7)$$

$$(26.8)$$

26.2.1 Circular Trigonometric Functions

$$\frac{d \sin x}{dx} = \cos x \quad (26.9)$$

$$\frac{d \cos x}{dx} = -\sin x \quad (26.10)$$

$$\frac{d \tan x}{dx} = \sec^2 x \quad (26.11)$$

$$\frac{d \sec x}{dx} = \sec x \tan x \quad (26.12)$$

$$\frac{d \csc x}{dx} = -\csc x \cot x \quad (26.13)$$

$$\frac{d \cot x}{dx} = -\csc^2 x \quad (26.14)$$

26.2.2 Inverse Circular Trigonometric Functions

$$\frac{d \arcsin x}{dx} = \frac{1}{\sqrt{1-x^2}}, |x| \leq 1 \quad (26.15)$$

$$\frac{d \arccos x}{dx} = -\frac{1}{\sqrt{1-x^2}}, |x| \leq 1 \quad (26.16)$$

$$\frac{d \arctan x}{dx} = \frac{1}{1+x^2} \quad (26.17)$$

$$\frac{d \cot^{-1} x}{dx} = -\frac{1}{1+x^2} \quad (26.18)$$

$$\frac{d \sec^{-1} x}{dx} = \frac{1}{x\sqrt{x^2-1}}, |x| \geq 1 \quad (26.19)$$

$$\frac{d \csc^{-1} x}{dx} = -\frac{1}{x\sqrt{x^2-1}}, |x| \geq 1 \quad (26.20)$$

26.2.3 Hyperbolic Trigonometric Function

$$\frac{d \sinh x}{dx} = \cosh x \quad (26.21)$$

$$\frac{d \cosh x}{dx} = \sinh x \quad (26.22)$$

$$\frac{d \tanh x}{dx} = 1 - \tanh^2 x = \operatorname{sech}^2(x) \quad (26.23)$$

$$\frac{d \coth x}{dx} = 1 - \coth^2 x = -\operatorname{csch}^2(x) \quad (26.24)$$

$$\frac{d[\operatorname{sech}(x)]}{dx} = -\tanh x \operatorname{sech} x \quad (26.25)$$

$$\frac{d[\operatorname{csch}(x)]}{dx} = -\coth x \operatorname{csch} x \quad (26.26)$$

26.2.4 Inverse Hyperbolic Trigonometric Function

$$\frac{d \sinh x}{dx} = \frac{1}{\sqrt{x^2 + 1}} \quad (26.27)$$

$$\frac{d \cosh x}{dx} = \frac{1}{\sqrt{x^2 - 1}} \quad (26.28)$$

$$\frac{d \tanh x}{dx} = \frac{1}{1 - x^2} \quad (26.29)$$

$$\frac{d \coth x}{dx} = \frac{1}{1 - x^2} \quad (26.30)$$

$$\frac{d[\operatorname{sech}(x)]}{dx} = \frac{1}{x\sqrt{1 - x^2}} \quad (26.31)$$

$$\frac{d[\operatorname{csch}(x)]}{dx} = \frac{1}{|x|\sqrt{1 + x^2}} \quad (26.32)$$

26.3 Rules of Differentiation

$$\frac{d[cf(x)]}{dx} = c \frac{df(x)}{dx} \quad (26.33)$$

$$\frac{d[f_1(x) + f_2(x)]}{dx} = \frac{d[f_1(x)]}{dx} + \frac{d[f_2(x)]}{dx} \quad (26.34)$$

$$\frac{d[f_1 f_2]}{dx} = f_1 f_2' + f_2 f_1' \quad (26.35)$$

$$\frac{d\left(\frac{f_1}{f_2}\right)}{dx} = \frac{f_2 f_1' - f_1 f_2'}{f_2^2} \quad (26.36)$$

26.4 Chain Rule

If two functions are defined as $z = f(y)$ and $y = g(x)$:

$$\frac{dz}{dx} = \frac{dz}{dy} \cdot \frac{dy}{dx} \quad (26.37)$$

If two functions are defined as $x = f(\theta)$ and $y = g(\theta)$:

$$\frac{d^2 y}{dx^2} = \left[\frac{d}{d\theta} \left(\frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}} \right) \right] \frac{d\theta}{dx} \quad (26.38)$$

Chapter 27

Successive Differentiation

$$D^n(ax+b)^m = m(m-1)\cdots(m-n+1)a^n(ax+b)^{m-n} \quad (27.1)$$

$$D^n\left(\frac{1}{ax+b}\right) = \frac{(-1)^n n! a^n}{(ax+b)^{n+1}} \quad (27.2)$$

$$D^n \ln(ax+b) = \frac{(-1)^{n-1} (n-1)! a^n}{(ax+b)^n}, n \geq 2 \quad (27.3)$$

$$D^n(a^{mx}) = m^n (\ln a)^n a^{mx} \quad (27.4)$$

$$D^n(e^{mx}) = m^n e^{mx} \quad (27.5)$$

$$D^n \sin(ax+b) = a^n \sin(ax+b+n\frac{\pi}{2}) \quad (27.6)$$

$$D^n \cos(ax+b) = a^n \cos(ax+b+n\frac{\pi}{2}) \quad (27.7)$$

$$D^n[e^{ax} \sin(bx+c)] = (a^2+b^2)^{\frac{n}{2}} e^{ax} \sin(bx+c+n \arctan \frac{b}{a}) \quad (27.8)$$

$$D^n[e^{ax} \cos(bx+c)] = (a^2+b^2)^{\frac{n}{2}} e^{ax} \cos(bx+c+n \arctan \frac{b}{a}) \quad (27.9)$$

27.1 Leibnitz's Theorem

For two functions u and v of x , the successive differentiation of their product is defined as:

$$\begin{aligned} (uv)_n &= {}^nC_0 u_n v + {}^nC_1 u_{n-1} v_1 + \cdots + {}^nC_n u v_n \\ &= \sum_{i=0}^n {}^nC_i u_{n-i} v_i \end{aligned} \quad (27.10)$$

Chapter 28

Partial Derivative

If $f(x, y)$ is a function of (x, y) , then $\frac{\delta f(x, y)}{\delta x}$ is the differentiation of $f(x, y)$ w.r.t. x , keeping all other parameters constant.

28.1 Chain Rule

If f is a function of u and v , which are functions of x and y , then:

$$\frac{\delta f}{\delta x} = \frac{\delta f}{\delta u} \frac{\delta u}{\delta x} + \frac{\delta f}{\delta v} \frac{\delta v}{\delta x} \quad (28.1)$$

$$\frac{\delta f}{\delta y} = \frac{\delta f}{\delta u} \frac{\delta u}{\delta y} + \frac{\delta f}{\delta v} \frac{\delta v}{\delta y} \quad (28.2)$$

If f is a function of x and y , which are functions of t , then:

$$\frac{df}{dt} = \frac{\delta f}{\delta x} \frac{dx}{dt} + \frac{\delta f}{\delta y} \frac{dy}{dt} \quad (28.3)$$

28.2 Euler's Theorem

For a homogeneous function ¹, $f(x_i)$ of degree n :

$$\sum x_i \frac{\delta f}{\delta x_i} = n f(x_i) \quad (28.4)$$

¹Homogeneous functions are defined as $f(ax, ay) = a^\kappa f(x, y)$, where κ is the degree of homogeneity. E.g. $f(x, y) = x^2 + y^2$, then $f(tx, ty) = t^2(x^2 + y^2)$, and the degree of homogeneity is 2.

Chapter 29

Application of Differentiation

29.1 Rolle's Theorem

For a function $f(x)$:

1. is continuous in $[a, b]$
2. is differentiable in (a, b)
3. $f(a) = f(b)$,

then there exists a point $x = c$ such that $f'(c) = 0$, $c \in (a, b)$

29.2 Mean Value Theorem or LaGrange's Theorem

For a function $f(x)$:

1. is continuous in $[a, b]$
2. is differentiable in (a, b) ,

then there exists a point $x = c$ such that $f'(c) = \frac{f(b) - f(a)}{b - a}$, $c \in (a, b)$,

i.e., the tangent is parallel to the line joining the the points $(a, f(a))$ and $(b, f(b))$.

29.3 Cauchy's Mean Value Theorem

For a function $f(x)$ and $g(x)$:

1. are continuous in $[a, b]$
2. are differentiable in (a, b)
3. $g'(x) \neq 0$ in (a, b) ,

then there exists a point $c \in (a, b)$, such that $\frac{f'(c)}{g'(c)} = \frac{f(b) - f(a)}{g(b) - g(a)}$.

29.4 Maxima and Minima

29.4.1 Maxima

For the local maxima of a function $f(x)$:

1. $f'(c) = 0$ and

$$\lim_{\epsilon \rightarrow c^-} f'(\epsilon) > 0$$

$$\lim_{\epsilon \rightarrow c^+} f'(\epsilon) < 0$$

OR

2. $f'(c) = 0$ and $f''(x) < 0$,

then $f(c)$ is the local maxima point of the function $f(x)$.

29.4.2 Minima

For the local minima of a function $f(x)$:

1. $f'(c) = 0$ and

$$\lim_{\epsilon \rightarrow c^-} f'(\epsilon) < 0$$

$$\lim_{\epsilon \rightarrow c^+} f'(\epsilon) > 0$$

OR

2. $f'(c) = 0$ and $f''(x) > 0$,

then $f(c)$ is the local minima point of the function $f(x)$.

29.5 Taylor's Theorem

For a function which is differentiable n times:

$$f(a+h) = f(a) + hf'(a) + \frac{h^2}{2!}f''(a) + \cdots + \frac{h^{n-1}}{(n-1)!}f^{n-1}(a) + \frac{h^n}{x!}R_n \quad (29.1)$$

where R_n is the remainder term.

29.5.1 Remainder Term

LeGrange's Form

$$R_n = f^n(a + \theta h), \theta \in (0, 1) \quad (29.2)$$

Cauchy's Form

$$R_n = n(1 - \theta)^{n-1} f^n(a + \theta h), \theta \in (0, 1) \quad (29.3)$$

29.5.2 Conditions for Validity of Expansion

For validity of Taylor Expansion, the condition

$$\lim_{n \rightarrow \infty} R_n = 0 \quad (29.4)$$

needs to be satisfied either where R_n is the remainder term in either LeGrange's Form or Cauchy's Form. If the condition is satisfied in a certain domain, then the expansion is valid within that domain only.

29.5.3 Taylor's Theorem for Two Variables

$$\begin{aligned} f(a + x, b + y) = & f(x, y) + \left(a \frac{\delta}{\delta x} + b \frac{\delta}{\delta y} \right) f(x, y) + \\ & \frac{1}{2!} \left(a^2 \frac{\delta^2}{\delta x^2} + b^2 \frac{\delta^2}{\delta y^2} \right) f(x, y) + \dots + \\ & \frac{1}{n!} \left(a^n \frac{\delta^n}{\delta x^n} + b^n \frac{\delta^n}{\delta y^n} \right) f(x + \theta a, y + \theta b), \theta \in (0, 1) \end{aligned} \quad (29.5)$$

29.6 Maclaurin's Series

$$\begin{aligned} f(x) = & f(0) + x f'(0) + \frac{1}{2!} x^2 f''(0) + \frac{1}{3!} x^3 f'''(0) + \dots \infty \\ = & \sum_{i=0}^{\infty} \frac{1}{i!} x^i f^i(0) \end{aligned} \quad (29.6)$$

29.6.1 Maclaurin's Series with Two Variables

$$\begin{aligned} f(a, b) = & f(0, 0) + \left(a \frac{\delta}{\delta x} + b \frac{\delta}{\delta y} \right) f(0, 0) + \\ & \frac{1}{2!} \left(a^2 \frac{\delta^2}{\delta x^2} + b^2 \frac{\delta^2}{\delta y^2} \right) f(0, 0) + \dots \infty \\ = & \sum_{i=0}^{\infty} \frac{1}{n!} \left(a^i \frac{\delta^i}{\delta x^i} + b^i \frac{\delta^i}{\delta y^i} \right) f(0, 0) \end{aligned} \quad (29.7)$$

29.7 Curvature

Curvature is the rate of change of direction w.r.t. arc. Mathematically:

$$\text{Curvature} = \frac{d(\text{direction})}{d(\text{arc})}$$

$$\lim_{\Delta s \rightarrow 0} \frac{\Delta \psi}{\Delta s} = \frac{d\psi}{ds} \quad (29.8)$$

29.7.1 Radius of Curvature

Cartesian Form

For a curve $y = f(x)$:

$$\rho = \frac{(1 + y'^2)^{\frac{3}{2}}}{y''} \quad (29.9)$$

However, this formula fails for $y' \rightarrow \infty$.

Parametric Form

For a curve defined as $x = \phi(t)$ and $y = \psi(t)$:

$$\rho = \frac{(\ddot{x}^2 + \ddot{y}^2)^{\frac{3}{2}}}{x\ddot{y} - y\ddot{x}} \quad (29.10)$$

29.7.2 Newton's Formula

1. If the curve passes through origin, and the tangent at origin is the x-axis:

$$\rho = \lim_{\substack{x \rightarrow 0 \\ y \rightarrow 0}} \frac{x^2}{2y} \quad (29.11)$$

2. If the curve passes through origin, and the tangent at origin is the y-axis:

$$\rho = \lim_{\substack{x \rightarrow 0 \\ y \rightarrow 0}} \frac{y^2}{2x} \quad (29.12)$$

3. If the curve passes through origin and $ax + by + c = 0$ is the tangent at origin:

$$\rho(0,0) = \frac{1}{2} \sqrt{a^2 + b^2} \lim_{\substack{x \rightarrow 0 \\ y \rightarrow 0}} \frac{a^2 + y^2}{ax + by} \quad (29.13)$$

29.7.3 Tangent at Origin

For a curve

$$\sum c_i x^j y^k = 0, i \in \mathbb{N} \text{ and } j, k \in \mathbb{Z} - \{0\} \quad (29.14)$$

The curve passes through origin $\because c = 0$. Then the lowest degree term equated to x gives the tangent at origin.

29.8 Asymptotes

If the distance between a line P and a curve $f(x)$, s is such that $s \rightarrow 0$, as $x \rightarrow \infty$, then P is the asymptote of $f(x)$. For asymptotes not parallel to x-axis:

Let $y = mx + c$ be the asymptote of the function $y = f(x)$, then:

$$m = \lim_{x \rightarrow \infty} \frac{y}{x} \quad (29.15)$$

$$c = \lim_{x \rightarrow \infty} (y - mx) \quad (29.16)$$

29.8.1 Asymptote of Algebraic Curves

For an algebraic curve, passing through origin, defined as:

$$\begin{aligned} & (a_0 x^n + a_1 x^{n-1} y^1 + a_2 x^{n-2} y^2 + \cdots + a_{n-1} x y^{n-1} + a_n y^n) + \\ & (b_0 x^{n-1} + b_1 x^{n-2} y^1 + b_2 x^{n-3} y^2 + \cdots + b_{n-1} x y^{n-2} + a_n y^{n-1}) + \\ & \quad \quad \quad \dots = 0 \end{aligned}$$

$$\Rightarrow x^n \phi_n \left(\frac{y}{x} \right) + x^{n-1} \phi_{n-1} \left(\frac{y}{x} \right) + \cdots + x \phi_1 \left(\frac{y}{x} \right) = 0$$

The asymptote(s) defined as $y = mx + c$,

1. m is the solution for the equation

$$\phi_n(m) = 0 \quad (29.17)$$

- 2.

$$c = -\frac{\phi_{n-1}(m)}{\phi_n(m)} \quad (29.18)$$

where c is a finite value.

Chapter 30

Integration

30.1 General Formulae

¹

$$\int nx^{n-1}dx = x^n + A \quad (30.1)$$

$$\int x^n dx = \frac{1}{n+1}x^{n+1} + A \quad (30.2)$$

$$\int e^x dx = e^x + A \quad (30.3)$$

$$\int \frac{1}{x} dx = \ln x + A \quad (30.4)$$

$$\int \ln x dx = x(\ln x - 1) + A \quad (30.5)$$

¹A is the constant of integration in all cases

30.2 Circular Trigonometric Functions

$$\int \sin x dx = -\cos x + A \quad (30.6)$$

$$\int \cos x dx = \sin x + A \quad (30.7)$$

$$\int \sec^2 x dx = \tan x + A \quad (30.8)$$

$$\int \csc^2 x dx = -\cot x + A \quad (30.9)$$

$$\int \sec x \tan x dx = \sec x + A \quad (30.10)$$

$$\int \csc x \cot x dx = -\csc x + A \quad (30.11)$$

$$\int \sec x dx = \ln(\sec x + \tan x) + A \quad (30.12)$$

$$\int \csc x dx = -\ln(\csc x + \cot x) + A \quad (30.13)$$

$$\begin{aligned} \int \tan x dx &= -\ln(\cos x) + A \\ &= \ln(\sec x) + A \end{aligned} \quad (30.14)$$

$$\int \cot x dx = \ln(\sin x) + A \quad (30.15)$$

30.3 Inverse Circular Trigonometric Function

$$\int \frac{1}{\sqrt{1-x^2}} dx = \sin^{-1} x + A \quad (30.16)$$

$$\int \frac{-1}{\sqrt{1-x^2}} dx = \cos^{-1} x + A \quad (30.17)$$

$$\int \frac{1}{1+x^2} dx = \tan^{-1} x + A \quad (30.18)$$

$$\int \frac{-1}{1+x^2} dx = \cot^{-1} x + A = -\tan^{-1} x + A \quad (30.19)$$

$$\int \frac{1}{x\sqrt{x^2-1}} dx = \sec^{-1} x + A = -\csc^{-1} x + A \quad (30.20)$$

$$\int \frac{-1}{x\sqrt{x^2-1}} dx = \csc^{-1} x + A = -\sec^{-1} x + A \quad (30.21)$$

30.4 Standard Integrals

$$\int \frac{dx}{\sqrt{a^2 - x^2}} = \sin^{-1} \frac{x}{a} + A \quad (30.22)$$

$$\int \frac{dx}{\sqrt{x^2 + a^2}} = \ln(x + \sqrt{x^2 + a^2}) + A \quad (30.23)$$

$$\int \frac{dx}{a^2 + x^2} = \frac{1}{a} \tan^{-1} \frac{x}{a} + A \quad (30.24)$$

$$\int \frac{dx}{\sqrt{x^2 - a^2}} = \ln(x + \sqrt{x^2 - a^2}) + A \quad (30.25)$$

$$\int \frac{dx}{x\sqrt{x^2 - a^2}} = \frac{1}{a} \sec^{-1} \frac{x}{a} + A \quad (30.26)$$

$$\int \sqrt{a^2 - x^2} dx = \frac{x\sqrt{a^2 - x^2}}{2} + \frac{a^2}{2} \sin^{-1} \frac{x}{a} + A \quad (30.27)$$

$$\int \sqrt{a^2 + x^2} dx = \frac{x\sqrt{a^2 + x^2}}{2} + \frac{a^2}{2} \ln(x + \sqrt{x^2 + a^2}) + A \quad (30.28)$$

$$\int \sqrt{x^2 - a^2} dx = \frac{x\sqrt{x^2 - a^2}}{2} - \frac{a^2}{2} \ln(x + \sqrt{a^2 - x^2}) + A \quad (30.29)$$

$$\int \frac{dx}{x^2 - a^2} = \frac{1}{2a} \ln \left| \frac{x - a}{x + a} \right| + A \quad (30.30)$$

$$\int \frac{dx}{a^2 - x^2} = \frac{1}{2a} \ln \left| \frac{a + x}{a - x} \right| + A \quad (30.31)$$

2

30.5 Special Forms

For a function $f(x)$:

$$\int [f(x)]^n f'(x) dx = \begin{cases} \frac{[f(x)]^{n+1}}{n+1} + A, n \neq -1 \\ \ln|f(x)| + A, n = -1 \end{cases} \quad (30.32)$$

² a is a constant $\in \mathbb{R}$

30.5.1 Integration by Part

For two functions $u(x)$ and $v(x)$:

$$\int u(x)v(x)dx = u(x) \left[\int v(x)dx \right] - \int \left[\frac{du(x)}{dx} \left(\int v(x)dx \right) dx \right] \quad (30.33)$$

Chapter 31

Definite Integral

31.1 Definition

For a function $f(x)$ for which $\int f(x)dx = F(x) + A$,

$$\int_a^b f(x)dx = F(b) - F(a) \quad (31.1)$$

31.2 Properties of Definite Integration

$$\int_a^b f(x)dx = \int_a^b f(t)dt \quad (31.2)$$

$$\int_a^b f(x)dx = - \int_b^a f(x)dx \quad (31.3)$$

$$\int_a^b f(x)dx = \int_a^c f(x)dx + \int_c^b f(x)dx \quad (31.4)$$

$$\int_a^b f(x)dx = \int_a^b f(a+b-x)dx \quad (31.5)$$

$$\int_0^a f(x)dx = \int_0^a f(a-x)dx \quad (31.6)$$

$$\int_0^{2a} f(x)dx = \begin{cases} 2 \int_0^a f(x)dx, & f(2a-x) = f(x) \\ 0, & f(2a-x) = f-(x) \end{cases} \quad (31.7)$$

$$\int_{-a}^a f(x)dx = \begin{cases} 2 \int_0^a f(x)dx, & f(x) \text{ is even} \\ 0, & f(x) \text{ is odd} \end{cases} \quad (31.8)$$

31.3 Approximation

$$f(a)(b-a) \leq \int_a^b f(x)dx \leq f(b)(b-a) \quad (31.9)$$

31.4 Sum of Infinite Series as a Definite Integral

Refer to 3.5.2.

Chapter 32

Reduction Formulae

$$\int \sin^n x dx = -\frac{1}{n} \sin^{n-1} x \cos x + \frac{n-1}{n} \int \sin^{n-2} x dx \quad (32.1)$$

$$\int \cos^n x dx = \frac{1}{n} \cos^{n-1} x \sin x + \frac{n-1}{n} \int \cos^{n-2} x dx \quad (32.2)$$

$$\int_0^{\frac{\pi}{2}} \sin^n x dx = \int_0^{\frac{\pi}{2}} \cos^n x dx = \begin{cases} \frac{(n-1) \cdot (n-3) \cdots 3 \cdot 1}{n \cdot (n-2) \cdots 4 \cdot 2} \left(\frac{\pi}{2}\right), n \rightarrow \text{even} \\ \frac{(n-1) \cdot (n-3) \cdots 4 \cdot 2}{n \cdot (n-2) \cdots 3 \cdot 1}, n \rightarrow \text{odd} \end{cases} \quad (32.3)$$

$$\int \sin^m x \cos^n x dx = \frac{-\sin^{m-1} x \cos^{n+1} x}{m+n} + \frac{m-1}{m+n} \int \sin^{m-2} x \cos^n x dx \quad (32.4)$$

For $I(m, n) = \int_0^{\frac{\pi}{2}} \sin^m x \cos^n x dx$:

When m and n are both even:

$$I(m, n) = \frac{[(m-1).(m-3) \cdots 3.1][(n-1).(n-3) \cdots 3.1]}{(m+n).(m+n-1) \cdots (4).(2)} \cdot \frac{\pi}{2} \quad (32.5)$$

Otherwise:

$$I(m, n) = \frac{[(m-1).(m-3) \cdots (2 \text{ or } 1)][(n-1).(n-3) \cdots (2 \text{ or } 1)]}{(m+n).(m+n-1) \cdots (2 \text{ or } 1)} \quad (32.6)$$

$$\begin{aligned} I_n &= \int \tan^n x dx \\ \Rightarrow I_n &= \frac{\tan^{n-2} x}{n-1} - I_{n-2} \end{aligned} \quad (32.7)$$

$$\begin{aligned} I_n &= \int \cot^n x dx \\ \Rightarrow I_n &= -\frac{\cot^{n-2} x}{n-1} - I_{n-2} \end{aligned} \quad (32.8)$$

$$\begin{aligned} I_n &= \int \sec^n x dx \\ \Rightarrow I_n &= \frac{\sec^{n-2} x \tan x}{n-1} + \frac{n-2}{n-1} I_{n-2} \end{aligned} \quad (32.9)$$

$$\begin{aligned} I_n &= \int \csc^n x dx \\ \Rightarrow I_n &= -\frac{\csc^{n-2} x \cot x}{n-1} + \frac{n-2}{n-1} I_{n-2} \end{aligned} \quad (32.10)$$

$$I_n = \int x^n e^{ax} dx \quad (32.11)$$

$$\int x^n e^{ax} dx = \frac{x^n e^{ax}}{a} - \frac{n}{a} I_{n-2} \quad (32.12)$$

$$I(m, n) = \int x^m (\ln x)^n dx \quad (32.13)$$

$$\int x^m (\ln x)^n dx = \frac{x^{m+1}}{m+1} (\ln x)^n - \frac{n}{m+1} I_{m, n-1} \quad (32.14)$$

Chapter 33

Multiple Integrals

33.1 Two Variables

For

$$I = \iint_R f(x, y) dx dy \quad (33.1)$$

The following substitution are made:

$$x = g(r, \theta) \quad (33.2)$$

$$y = h(r, \theta) \quad (33.3)$$

$$\therefore dx dy = |J| dr d\theta \quad (33.4)$$

Where J is the Jacobian, defined as:

$$J = \begin{vmatrix} \frac{\delta x}{\delta r} & \frac{\delta y}{\delta r} \\ \frac{\delta x}{\delta \theta} & \frac{\delta y}{\delta \theta} \end{vmatrix} \quad (33.5)$$

The equivalent integral is:

$$I = \iint_{R_1} f(g(r, \theta), h(r, \theta)) |J| dr d\theta \quad (33.6)$$

33.2 Three Variables

For

$$I = \iiint_R f(x, y, z) dx dy dz \quad (33.7)$$

The following substitution are made:

$$x = g(r, \theta, \phi) \quad (33.8)$$

$$y = h(r, \theta, \phi) \quad (33.9)$$

$$z = k(r, \theta, \phi) \quad (33.10)$$

$$\therefore dx dy dz = |J| dr d\theta d\phi \quad (33.11)$$

Where J is the Jacobian, defined as:

$$J = \begin{vmatrix} \frac{\delta x}{\delta r} & \frac{\delta y}{\delta r} & \frac{\delta z}{\delta r} \\ \frac{\delta x}{\delta \theta} & \frac{\delta y}{\delta \theta} & \frac{\delta z}{\delta \theta} \\ \frac{\delta x}{\delta \phi} & \frac{\delta y}{\delta \phi} & \frac{\delta z}{\delta \phi} \end{vmatrix} \quad (33.12)$$

The equivalent integral is:

$$I = \iiint_{R_1} f(g(r, \theta, \phi), h(r, \theta, \phi), k(r, \theta, \phi)) |J| dr d\theta d\phi \quad (33.13)$$

Chapter 34

Differential Equation

34.1 1st Order, 1st Degree Differential Equation

For the equation:

$$\frac{dy}{dx} + P(x)y = Q(x) \quad (34.1)$$

Then an Integral Function (I.F.) is defined as:

$$I.F. = e^{\int P(x)dx} \quad (34.2)$$

Then the solution of the equation 34.1 is given by:

$$y(I.F.) = \int Q(I.F.)dx \quad (34.3)$$

34.2 2nd Order, 1st Degree Differential Equation

For the equation:

$$\frac{d^2y}{dx^2} + a\frac{dy}{dx} + by = 0 \quad (34.4)$$

OR

$$y'' + ay' + by = 0 \quad (34.5)$$

By substituting $y = e^{\lambda x}$, the equation obtained is:

$$\begin{aligned} \lambda^2 e^{\lambda x} + \lambda e^{\lambda x} + b e^{\lambda x} &= 0 \\ \therefore e^{\lambda x} &\neq 0 \\ \Rightarrow \lambda^2 + a\lambda + b &= 0 \end{aligned} \quad (34.6)$$

If α and β are the solutions of the equation 34.6, then the solution of 34.4 can be:

1. If $\alpha = \beta$ and $\alpha, \beta \in \mathbb{R}$:

$$y = (c_1 + c_2x)e^{\alpha x} \quad (34.7)$$

2. If $\alpha \neq \beta$ and $\alpha, \beta \in \mathbb{R}$:

$$y = c_1e^{\alpha x} + c_2e^{\beta x} \quad (34.8)$$

3. If $\lambda = \alpha + i\beta$:

$$y = e^{\alpha x} [A \cos(\beta x) + B \sin(\beta x)] \quad (34.9)$$

34.3 Special Cases of Differential Equation

34.3.1 Definition of Inverse Operator

The operator D is equivalent to $\frac{d}{dx}$. If $Df(x) = X$, then $f(x) = \frac{1}{D}X = \int X dx$.

34.3.2 Special Cases

- 1.

$$f(x) = \frac{1}{D-a}X = e^{ax} \int X e^{-ax} dx \quad (34.10)$$

- 2.

$$\frac{1}{f(D)}e^{ax} = \begin{cases} \frac{e^{ax}}{f(a)}, f(a) \neq 0 \\ x \frac{e^{ax}}{f'(a)}, f(x) = 0 \text{ and } f'(a) \neq 0 \\ x^2 \frac{e^{ax}}{f''(a)}, f(x) = 0 \text{ and } f'(a) = 0 \end{cases} \quad (34.11)$$

- 3.

$$\frac{1}{f(D)}x^m = [f(D)]^{-1}x^m \quad (34.12)$$

$[f(D)]^{-1}$ is expanded and arranged in terms of ascending powers of D and operated on x^m .

4. (a)

$$\begin{aligned} \frac{1}{f(D)} \sin(ax) &= \frac{1}{\phi(D^2)} \sin(ax) \\ &= \frac{1}{\phi(-a^2)} \sin(ax) \end{aligned} \quad (34.13)$$

(b)

$$\begin{aligned}
\frac{1}{f(D)} \cos(ax) &= \frac{1}{\phi(D^2)} \cos(ax) \\
&= \frac{1}{\phi(-a^2)} \cos(ax)
\end{aligned} \tag{34.14}$$

5. (a)

$$\begin{aligned}
\frac{1}{f(D)} \sin(ax) &= \frac{1}{\phi(D^2, D)} \sin(ax) \\
&= \frac{1}{\phi(-a^2, D)} \sin(ax)
\end{aligned} \tag{34.15}$$

(b)

$$\begin{aligned}
\frac{1}{f(D)} \cos(ax) &= \frac{1}{\phi(D^2, D)} \cos(ax) \\
&= \frac{1}{\phi(-a^2, D)} \cos(ax)
\end{aligned} \tag{34.16}$$

6. (a)

$$\begin{aligned}
\frac{1}{f(D)} \sin(ax) &= \frac{\psi(D)}{\phi(D^2)} \sin(ax) \\
&= \frac{\psi(D)}{\phi(-a^2)} \sin(ax)
\end{aligned} \tag{34.17}$$

(b)

$$\begin{aligned}
\frac{1}{f(D)} \cos(ax) &= \frac{\psi(D)}{\phi(D^2)} \cos(ax) \\
&= \frac{\psi(D)}{\phi(-a^2)} \cos(ax)
\end{aligned} \tag{34.18}$$

7. (a)

$$\frac{1}{f(D)} \sin(ax) = x \frac{1}{f'(D)} \sin(ax) \tag{34.19}$$

(b)

$$\frac{1}{f(D)} \cos(ax) = x \frac{1}{f'(D)} \cos(ax) \tag{34.20}$$

34.4 Method of Variation of Parameters

If the equation is of the form:

$$\frac{d^2y}{dx^2} + a \frac{dy}{dx} + by = f \tag{34.21}$$

where a, b, f are functions of x . The solution for 34.21 is obtained by solving for:

$$\frac{d^2y}{dx^2} + a\frac{dy}{dx} + by = 0 \quad (34.22)$$

If y_1 and y_2 are the two independent solution of equation 34.22. Then the general solution of the equation is:

$$y = c_1y_1 + c_2y_2 \quad (34.23)$$

where c_1 and c_2 are the constants.

The particular solution of equation 34.22 will be:

$$y = y_1 \left(\int \frac{y_2(-f)}{W} dx \right) + y_2 \left(\int \frac{y_1 f}{W} dx \right) \quad (34.24)$$

W is the Wronskian, which is defined by:

$$W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} \quad (34.25)$$

34.5 Singular and Ordinary Point

For a differential equation:

$$P_0 \frac{d^n y}{dx^n} + P_1 \frac{d^{n-1} y}{dx^{n-1}} + \cdots + P_{n-1} \frac{dy}{dx} + P_n y = R(x) \quad (34.26)$$

where $P_0 \cdots P_n$ are functions of x .

If at a point $x = x_0$:

1. $P_0(x_0) \neq 0$, x_0 is an ordinary point.
2. $P_0(x_0) = 0$, x_0 is an singular point:

(a)

$$\lim_{x \rightarrow x_0} (x - x_0) P_1(x) = c_1 \quad (34.27)$$

$$\lim_{x \rightarrow x_0} (x - x_0)^2 P_2(x) = c_2 \quad (34.28)$$

$$(34.29)$$

where c_1 and c_2 are both finite quantities x_0 is a regular singular point.

- (b) otherwise it is an irregular singular point.

Chapter 35

Beta and Gamma Functions

For $m, n > 0$:

$$\begin{aligned}\beta(m, n) &= \int_0^1 x^{m-1} (1-x)^{n-1} dx \\ &= 2 \int_0^{\frac{\pi}{2}} \sin^{2m-1} x \cos^{2n-1} x dx\end{aligned}\tag{35.1}$$

$$\Gamma(n) = \int_0^\infty e^{-x} x^{n-1} dx\tag{35.2}$$

35.1 Important Relations between $\beta(m, n)$ and $\Gamma(n)$ Functions

$$\Gamma(n) = \frac{\Gamma(n+1)}{n}\tag{35.3}$$

$$\Gamma(1) = 1\tag{35.4}$$

$$\Gamma\left(\frac{1}{2}\right) = \sqrt{\pi} \approx 1.772\tag{35.5}$$

$$\Gamma(n+1) = n!, n \in \mathbb{N}\tag{35.6}$$

$$\Gamma(m)\Gamma\left(m + \frac{1}{2}\right) = \sqrt{\pi}\Gamma(2m)\tag{35.7}$$

$$\Gamma(m)\Gamma(m-1) = \pi \csc(m\pi)\tag{35.8}$$

$$\beta(m, n) = \beta(n, m) \quad (35.9)$$

$$\beta(m, n) = \frac{\Gamma(m)\Gamma(n)}{\Gamma(m+n)} \quad (35.10)$$

$$\beta\left(\frac{1}{2}, \frac{1}{2}\right) = \pi \quad (35.11)$$

$$\int_0^{\frac{\pi}{2}} \sin^p x \cos^q x = \frac{1}{2} \frac{\Gamma(\frac{p+1}{2})\Gamma(\frac{q+1}{2})}{\Gamma(\frac{p+q}{2})} \quad (35.12)$$

$$(35.13)$$

Chapter 36

Laplace Transformations

The Laplace Transformation of a function $f(t)$ is defined as:

$$F(s) = \mathcal{L}\{f(t)\} = \lim_{x \rightarrow \infty} \int_0^x e^{-st} f(t) dt \tag{36.1}$$

36.1 Basic Transformations

$f(t)$	$F(s)$
$af(t) + bg(t)$	$aF(s) + bG(s)$
1	$\frac{1}{s}$
t	$\frac{1}{s^2}$
t^n	$\frac{n!}{s^{n+1}}$
e^{at}	$\frac{1}{s - a}$
$\cos(\omega t)$	$\frac{s}{s^2 + \omega^2}$
$\sin(\omega t)$	$\frac{\omega}{s^2 + \omega^2}$
$\sinh at$	$\frac{s}{s^2 - a^2}$
$\cosh at$	$\frac{s}{s^2 - a^2}$

Table 36.1: Table of Laplace Transformations

36.2 Important Relations

$$\mathcal{L}\{e^{at}f(t)\} = F(s-a) \quad (36.2)$$

$$\mathcal{L}\{tf(t)\} = -F'(s) \quad (36.3)$$

$$\mathcal{L}\{t^n f(t)\} = (-1)^n F^{(n)}(s) \quad (36.4)$$

$$\mathcal{L}\left\{\frac{f(t)}{t}\right\} = \lim_{x \rightarrow \infty} \int_s^x F(u) du \quad (36.5)$$

$$\mathcal{L}\left\{\frac{f(t)}{t^n}\right\} = \lim_{x \rightarrow \infty} \int_1 \int_2 \cdots \int_s^x F(u) du \cdots du \quad (36.6)$$

36.3 Convolution

For two functions $f(t)$ and $g(t)$ be given such that their Laplace transforms are $F(s)$ and $G(s)$, then:

$$\mathcal{L}\{f(t) \star g(t)\} = F(s)G(s) \quad (36.7)$$

where $f(t) \star g(t)$ is defined as:

$$\int_0^t f(u)g(t-u)du \quad (36.8)$$

36.4 Laplace Transforms of Differentials

If the Laplace Transform of $f(t)$ is $F(s)$ ¹:

$$\mathcal{L}\{f'(t)\} = sF(s) - y(0) \quad (36.9)$$

$$\mathcal{L}\{f''(t)\} = s^2F(s) - [sy(0) + y'(0)] \quad (36.10)$$

$$\vdots$$

$$\mathcal{L}\{f^n(t)\} = s^n F(s) - \left[\sum_{i=0}^{n-1} s^{n-i} y^{(i)}(0) \right] \quad (36.11)$$

¹Used in initial value problems