

Diferenças espectrais entre genótipos e condições, dia a dia

A. ANOVA entre genótipos dentro de cada condição

IRRIG	1.814 88.4%	1.871 91.2%	2.035 99.2%	2.046 99.8%	1.823 88.9%	1.555 75.8%	2.019 98.4%
NIRRIG	606 29.5%	1.331 64.9%	895 43.6%	2.040 99.5%	1.229 59.9%	1.967 95.9%	1.661 81.0%
	D02	D03	D04	D05	D06	D09	D10

B. Tukey HSD: genótipos comparados dentro da mesma condição

IRRIG BR16 × CD202	822 40.1%	1.242 60.6%	1.864 90.9%	678 33.1%	271 13.2%	1.546 75.4%	1.815 88.5%
IRRIG BR16 × EMB48	482 23.5%	716 34.9%	521 25.4%	1.812 88.3%	1.652 80.5%	58 2.8%	285 13.9%
IRRIG CD202 × EMB48	1.650 80.4%	1.709 83.3%	1.900 92.6%	1.889 92.1%	1.507 73.5%	1.028 50.1%	1.909 93.1%
NIRRIG BR16 × CD202	412 20.1%	712 34.7%	508 24.8%	506 24.7%	684 33.3%	1.557 75.9%	1.527 74.5%
NIRRIG BR16 × EMB48	259 12.6%	340 16.6%	651 31.7%	1.877 91.5%	737 35.9%	580 28.3%	349 17.0%
NIRRIG CD202 × EMB48	409 19.9%	1.064 51.9%	522 25.5%	1.792 87.4%	429 20.9%	1.911 93.2%	1.029 50.2%
	D02	D03	D04	D05	D06	D09	D10



C. Tukey HSD: irrigado × não irrigado dentro de cada genótipo

BR16: Irrigado × Não irrigado	124 6.0%	1.734 84.5%	368 17.9%	2.007 97.9%	2.022 98.6%	995 48.5%	917 44.7%
CD202: Irrigado × Não irrigado	1.767 86.2%	1.767 86.2%	1.895 92.4%	1.958 95.5%	1.986 96.8%	1.917 93.5%	1.758 85.7%
EMB48: Irrigado × Não irrigado	79 3.9%	1.625 79.2%	1.134 55.3%	1.838 89.6%	1.796 87.6%	1.011 49.3%	972 47.4%
	D02	D03	D04	D05	D06	D09	D10