NUTRITION GUIDE

Eating well, from first milk to family table

A guide. Tap a card to explore, then take the quiz.

Start exploring

Take the quiz



Pick a topic to explore

Take the quiz ›



Breastfeeding

Duration and benefits



Starting solids

Principles and textures



Iron

Who needs it, how much



Vitamin A and zinc

Doses and why



Deworming

Drugs, timing, cautions



Iron deficiency anemia

Who to test, and when

Breastfeeding: how long and why

Menu

6 months

Exclusive breastfeeding

2 years+

Continue breastfeeding alongside food



Easy to digest and absorb



Contains antibodies



DHA fats support brain development and intelligence



Promotes emotional bonding



Protects mom from breast, uterine and ovarian cancer, obesity and bleeding



Earlier return to pre-pregnancy weight, and big family savings

Starting solids at 6 months

Menu



Timely

Introduce complementary foods at 6 months.

Adequate

Plan with the PSRGN Dietary Prescription Food Guide Pyramid.

Safe

Do not add salt before one year of age.

Properly fed

Practice responsive feeding.

Textures and how often

Menu

6 — 8 — 10 — 12

Pureed food
at 6 months

Finger foods
at 8 months

Lumpy or chopped
at 10 months

Table food
at 12 months

Go slowly

One new food at a time, for 3 days. Start with 1 to 2 teaspoons a day of lugaw or cereals, fruits or vegetables.

How often

2 to 3 times a day at 6 to 8 months. 3 to 4 times a day from 9 to 24 months.



Iron supplementation

Menu

Who	Preparation	Dose and duration
Low birth weight	Drops: 15 mg iron per 0.6 mL	0.3 mL once a day, from 2 to 6 months of age
Infants 6–11 months	Drops: 15 mg iron per 0.6 mL	0.6 mL once a day for 3 months
Children 1–5 years	Syrup: 30 mg iron per 5 mL	1 tsp once a day for 3 months, or 30 mg once a week for 6 months
Girls 10–19 years	Coated tablet: 60 mg iron with 400 mcg folic acid	One tablet once a day



Why it matters

Low iron is linked to problems with thinking and movement skills.

Vitamin A and zinc

Menu

100,000 IU

Infants 6 to 11 months. One dose only, usually given at 9 months.

200,000 IU

Children 12 to 59 months. One capsule every 6 months.

Zinc

Zinc supplements help prevent pneumonia and diarrhea.

Deworming every 6 months

Menu

Albendazole

200 mg for ages 12 to 23 months. 400 mg for 24 months and above.

Mebendazole

500 mg for ages 12 months and above.

Who and when

All children aged 1 to 12 years (DOH). A single dose every 6 months, taken on a full stomach.

Do not deworm if

The child has high-grade fever, profuse diarrhea or abdominal pain.

Iron deficiency anemia

Menu

It is associated with cognitive and psychomotor problems in children.

At-risk children should have a complete blood count (CBC) at least once at:

6–24 months

2–6 years

10–19 years

Also at risk: actively menstruating adolescent girls and fad dieters. A CBC is recommended at each stage of adolescence.

QUICK QUIZ

Test what you learned

4 questions. Tap an answer to see if you got it right. Wrong answer? You can try again.

Start quiz

Menu



Question 1 of 4

Menu

BREASTFEEDING

How long is exclusive breastfeeding recommended?

A Up to 6 months

B Up to 3 months

C Up to 12 months

Correct!



Up to 6 months

Exclusive breastfeeding up to 6 months, then continued up to 2 years and beyond alongside food.

Menu

Next question ›

Not quite



Hint

Think about the age when solid foods begin.

Menu

Try again

Question 2 of 4

Menu

STARTING SOLIDS

How should new foods be introduced to a 6-month-old?

A Several new foods at the same time

B A new food at every meal, every day

C One new food at a time, for 3 days

Correct!



One new food at a time, for 3 days

Introduce one new food at a time for 3 days, starting with 1 to 2 teaspoons a day.

Menu

Next question ›

Not quite



Hint

Go slowly so you can tell how each food agrees with the baby.

Menu

Try again

Question 3 of 4

Menu

TEXTURES

At what age is table food introduced?

A 6 months

B 12 months

C 8 months

Correct!



12 months

Pureed at 6 months, finger foods around 8 months, lumpy or chopped at 10 months, table food at 12 months.

Menu

Next question ›

Not quite



Hint

It is the last step on the texture timeline.

Menu

Try again

Question 4 of 4

Menu

DEWORMING

When should deworming be skipped?

A When the child has just eaten

B When the child is 5 years old

C When there is high-grade fever, profuse diarrhea or abdominal pain

Correct!



When there is high-grade fever, profuse diarrhea or abdominal pain

Do not deworm with high-grade fever, profuse diarrhea or abdominal pain. Otherwise, dose every 6 months on a full stomach.

Menu

Finish ›

Not quite



Hint

Think of signs that a child is already unwell.

Menu

Try again

WELL DONE

Good food, good start

- Breast milk only to 6 months, then keep going
- One new food at a time, for 3 days
- No salt before age 1
- Keep iron, vitamin A and deworming on schedule

Back to menu

Retake quiz

