CEGEP Linear Algebra Problems

AN OPEN SOURCE COLLECTION OF CEGEP LEVEL LINEAR ALGEBRA PROBLEMS

EDITED BY

YANN LAMONTAGNE, ADD YOUR NAME HERE

Contents

1	\mathbf{Sys}	tems of Linear Equations	1
	1.1	Introduction to Systems of Linear Equations	1
	1.2	Gaussian and Gauss-Jordan Elimination	1
	1.3	Applications of Linear Systems	1
2	Ma	trix Algebra	3
4	2.1	Introduction to Matrices and Matrix Operations	3
	$\frac{2.1}{2.2}$	•	ა 3
		Matrix Inverses and Algebraic Properties	3
	2.3	Elementary Matrices	3
	2.4	Linear Systems and Matrices	3
3	Det	erminants	5
	3.1	The Laplace Expansion	5
	3.2	Determinants and Elementary Operations	5
	3.3	Determinants and Elementary Matrices	5
	3.4	Applications of the Determinant	5
4	Vec	etor Geometry	7
	4.1	Introduction to Vectors and Lines	7
	4.2	Dot Product and Projections	7
	4.3	Cross Product and Planes	7
	4.4	Areas, Volumes and Distances	7
	4.5	Geometry of Solutions of Linear Systems	7
	1.0	domestry of polations of Emocal Systems 1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.	•
5	Vec	etor Spaces	9
	5.1	Introduction to Vector Spaces	9
	5.2	Subspaces	10
	5.3	Spanning Sets	10
	5.4	Linear Independence	10
	5.5	Basis	10
	5.6	Dimension	11
A	nswe	ers to Exercises	13
\mathbf{R}	efere	ences	16
T.	dor		17

Systems of Linear Equations

1.1 Introduction to Systems of Linear Equations

1.1.1 Place Holder

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam tincidunt cursus volutpat. Quisque non congue sem. Vivamus nec nibh sed est dapibus auctor eu sed nulla. Praesent ornare eleifend nibh a finibus. Proin rutrum neque nec massa tincidunt, non malesuada dolor interdum. Nam a massa sit amet diam efficitur pharetra. Nulla interdum efficitur sem, sit amet commodo orci mattis non. Duis tortor ex, maximus a sapien id, molestie maximus risus.

1.2 Gaussian and Gauss-Jordan Elimination

1.2.1 [JH] Use Gauss's Method to find the unique solution for each system.

a.
$$2x+3y=13 \\ x-y=-1$$
 b.
$$x-z=0 \\ 3x+y=1 \\ -x+y+z=4$$

1.3 Applications of Linear Systems

1.3.1 Place Holder

Matrix Algebra

2.1 Introduction to Matrices and Matrix Operations

2.1.1 [**HE**] Let

$$A = \begin{bmatrix} 1 & -2 & 3 \\ 1 & -1 & 0 \end{bmatrix}, \ B = \begin{bmatrix} 3 & 4 \\ 5 & -1 \\ 1 & -1 \end{bmatrix}, \ C = \begin{bmatrix} 4 & -1 & 2 \\ -1 & 5 & 1 \end{bmatrix},$$

$$D = \begin{bmatrix} -1 & 0 & 1 \\ 0 & 2 & 1 \end{bmatrix}, E = \begin{bmatrix} 3 & 4 \\ -2 & 3 \\ 0 & 1 \end{bmatrix},$$

$$F = \begin{bmatrix} 2 \\ -3 \end{bmatrix}, \ G = \begin{bmatrix} 2 & -1 \end{bmatrix}.$$

Compute each of the following and simplify, whenever possible. If a computation is not possible, state why.

- a. 3C 4D
- b. A (D + 2C)
- c. A-E
- d. AE
- e. 3BC 4BD
- f. CB + D
- g. GC
- h. FG
- i. Illustrate the associativity of matrix multiplication by multiplying (AB)C and A(BC) where A, B, and C are matrices above.

2.2 Matrix Inverses and Algebraic Properties

2.2.1 [YL] Prove: If A and B are square matrices satisfying AB = I, then $A = B^{-1}$.

2.3 Elementary Matrices

2.3.1 [YL] Express

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 0 & 1 \\ 0 & 2 & 1 \end{bmatrix}$$

as a product of 4 elementary matrices.

2.4 Linear Systems and Matrices

2.4.1 [YL] Consider

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 2 \\ 0 & 0 & 3 \end{bmatrix}.$$

- a. Find A^{-1} .
- b. Using A^{-1} solve Ax = b where

$$x = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$
 and $b = \begin{bmatrix} 1 \\ -2 \\ 1 \end{bmatrix}$.

Determinants

3.1 The Laplace Expansion

3.1.1 Place Holder

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam tincidunt cursus volutpat. Quisque non congue sem. Vivamus nec nibh sed est dapibus auctor eu sed nulla. Praesent ornare eleifend nibh a finibus. Proin rutrum neque nec massa tincidunt, non malesuada dolor interdum. Nam a massa sit amet diam efficitur pharetra. Nulla interdum efficitur sem, sit amet commodo orci mattis non. Duis tortor ex, maximus a sapien id, molestie maximus risus.

3.2 Determinants and Elementary Operations

3.2.1 Place Holder

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam tincidunt cursus volutpat. Quisque non congue sem. Vivamus nec nibh sed est dapibus auctor eu sed nulla. Praesent ornare eleifend nibh a finibus. Proin rutrum neque nec massa tincidunt, non malesuada dolor interdum. Nam a massa sit amet diam efficitur pharetra. Nulla interdum efficitur sem, sit amet commodo orci mattis non. Duis tortor ex, maximus a sapien id, molestie maximus risus.

3.3 Determinants and Elementary Matrices

3.3.1 Place Holder

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam tincidunt cursus volutpat. Quisque non congue sem. Vivamus nec nibh sed est dapibus auctor eu sed nulla. Praesent ornare eleifend nibh a finibus. Proin rutrum neque nec massa tincidunt, non malesuada dolor interdum. Nam a massa sit amet diam efficitur pharetra. Nulla interdum efficitur sem, sit amet commodo orci mattis non. Duis tortor ex, maximus a sapien id, molestie maximus risus.

3.4 Applications of the Determinant

3.4.1 Place Holder

Vector Geometry

4.1 Introduction to Vectors and Lines

4.1.1 Place Holder

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam tincidunt cursus volutpat. Quisque non congue sem. Vivamus nec nibh sed est dapibus auctor eu sed nulla. Praesent ornare eleifend nibh a finibus. Proin rutrum neque nec massa tincidunt, non malesuada dolor interdum. Nam a massa sit amet diam efficitur pharetra. Nulla interdum efficitur sem, sit amet commodo orci mattis non. Duis tortor ex, maximus a sapien id, molestie maximus risus.

4.2 Dot Product and Projections

4.2.1 Cauchy-Schwartz Inequality [YL] Prove without assuming that the law of cosine holds in \mathbb{R}^n : If $\vec{u}, \vec{v} \in \mathbb{R}^n$ then $|\vec{u} \cdot \vec{v}| \leq ||\vec{u}|| ||\vec{v}||$.

4.3 Cross Product and Planes

4.3.1 Place Holder

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam tincidunt cursus volutpat. Quisque non congue sem. Vivamus nec nibh sed est dapibus auctor eu sed nulla. Praesent ornare eleifend nibh a finibus. Proin rutrum neque nec massa tincidunt, non malesuada dolor interdum. Nam a massa sit amet diam efficitur pharetra. Nulla interdum efficitur sem, sit amet commodo orci mattis non. Duis tortor ex, maximus a sapien id, molestie maximus risus.

4.4 Areas, Volumes and Distances

4.4.1 Place Holder

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam tincidunt cursus volutpat. Quisque non congue sem. Vivamus nec nibh sed est dapibus auctor eu sed nulla. Praesent ornare eleifend nibh a finibus. Proin rutrum neque nec massa tincidunt, non malesuada dolor interdum. Nam a

and massa sit amet diam efficitur pharetra. Nulla interdum efficitur sem, sit amet commodo orci mattis non. Duis tortor ex, maximus a sapien id, molestie maximus risus.

4.5 Geometry of Solutions of Linear Systems

4.5.1 Place Holder

Vector Spaces

5.1 Introduction to Vector Spaces

5.1.1 [JH] Name the zero vector for each of these vector spaces.

- a. The space of degree three polynomials under the natural operations.
- b. The space of 2×3 matrices.
- c. The space $\{f: [0,1] \to \mathbb{R} \mid f \text{ is continuous}\}.$
- d. The space of real-valued functions of one natural number variable.

5.1.2 [JH] Find the additive inverse, in the vector space, of the vector.

- a. In \mathcal{P}_3 , the vector $-3 2x + x^2$.
- b. In the space $\mathcal{M}_{2\times 2}$,

$$\begin{bmatrix} 1 & -1 \\ 0 & 3 \end{bmatrix}.$$

- c. In $\{ae^x + be^{-x} \mid a, b \in \mathbb{R}\}$, the space of functions of the real variable x under the natural operations, the vector $3e^x 2e^{-x}$.
- **5.1.3** [JH] For each, list three elements and then show it is a vector space.
 - a. The set of linear polynomials $\mathcal{P}_1 = \{a_0 + a_1x \mid a_0, a_1 \in \mathbb{R}\}$ under the usual polynomial addition and scalar multiplication operations.
 - b. The set of linear polynomials $\{a_0 + a_1x \mid a_0 2a_1 = 0\}$, under the usual polynomial addition and scalar multiplication operations.

5.1.4 [JH] For each, list three elements and then show it is a vector space.

- a. The set of 2×2 matrices with real entries under the usual matrix operations.
- b. The set of 2×2 matrices with real entries where the 2, 1 entry is zero, under the usual matrix operations.
- **5.1.5** by Jim Hefferon [JH] For each, list three elements and then show it is a vector space.
 - a. The set of three-component row vectors with their

usual operations.

b. The set

$$\{(x, y, z, w) \in \mathbb{R}^4 \mid x + y - z + w = 0\}$$

under the operations inherited from \mathbb{R}^4 .

5.1.6 [JH] Show that the following are not vector spaces.

a. Under the operations inherited from \mathbb{R}^3 , this set

$$\{(x, y, z) \in \mathbb{R}^3 \mid x+y+z=1\}$$

b. Under the operations inherited from \mathbb{R}^3 , this set

$$\{(x, y, z) \in \mathbb{R}^3 \mid x^2 + y^2 + z^2 = 1\}$$

c. Under the usual matrix operations,

$$\left\{ \begin{bmatrix} a & 1 \\ b & c \end{bmatrix} \mid a, b, c \in \mathbb{R} \right\}$$

d. Under the usual polynomial operations,

$$\{a_0 + a_1 x + a_2 x^2 \mid a_0, a_1, a_2 \in \mathbb{R}^+ \}$$

where \mathbb{R}^+ is the set of reals greater than zero

e. Under the inherited operations,

$$\left\{ (x,\ y) \in \mathbb{R}^2 \ \middle| \ x + 3y = 4, \, 2x - y = 3 \text{ and } 6x + 4y = 10 \right\}$$

- **5.1.7** [JH] Is the set of rational numbers a vector space over \mathbb{R} under the usual addition and scalar multiplication operations?
- **5.1.8** [JH] Prove that the following is not a vector space: the set of two-tall column vectors with real entries subject to these operations.

$$\begin{pmatrix} x_1 \\ y_1 \end{pmatrix} + \begin{pmatrix} x_2 \\ y_2 \end{pmatrix} = \begin{pmatrix} x_1 - x_2 \\ y_1 - y_2 \end{pmatrix} \qquad r \cdot \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} rx \\ ry \end{pmatrix}$$

5.1.9 [JH] Prove or disprove that \mathbb{R}^3 is a vector space under these operations.

a.
$$\begin{pmatrix} x_1 \\ y_1 \\ z_1 \end{pmatrix} + \begin{pmatrix} x_2 \\ y_2 \\ z_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$
 and $r \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} rx \\ ry \\ rz \end{pmatrix}$
b. $\begin{pmatrix} x_1 \\ y_1 \\ z_1 \end{pmatrix} + \begin{pmatrix} x_2 \\ y_2 \\ z_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$ and $r \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$

5.1.10 [JH] For each, decide if it is a vector space; the intended operations are the natural ones.

a. The set of diagonal 2×2 matrices

$$\left\{ \begin{bmatrix} a & 0 \\ 0 & b \end{bmatrix} \mid a, b \in \mathbb{R} \right\}$$

b. The set of 2×2 matrices

$$\left\{ \begin{bmatrix} x & x+y \\ x+y & y \end{bmatrix} \mid x,y \in \mathbb{R} \right\}$$

- c. $\{(x, y, z, w) \in \mathbb{R}^4 \mid x+y+w=1\}$
- d. The set of functions $\{f: \mathbb{R} \to \mathbb{R} \mid df/dx + 2f = 0\}$
- e. The set of functions $\{f: \mathbb{R} \to \mathbb{R} \mid df/dx + 2f = 1\}$

5.1.11 [JH] Show that the set \mathbb{R}^+ of positive reals is a vector space when we interpret 'x+y' to mean the product of x and y (so that 2+3 is 6), and we interpret ' $r \cdot x$ ' as the r-th power of x.

5.1.12 [JH] Prove or disprove that the following is a vector space: the set of polynomials of degree greater than or equal to two, along with the zero polynomial.

5.1.13 [JH]

Is $\{(x, y) \mid x, y \in \mathbb{R}\}$ a vector space under these operations?

- a. $(x_1, y_1)+(x_2, y_2)=(x_1+x_2, y_1+y_2)$ and $r\cdot(x,y)=(rx,y)$
- b. $(x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2)$ and $r \cdot (x, y) = (rx, 0)$

5.1.14 [JH]

Prove the following:

- a. For any $\vec{v} \in V$, if $\vec{w} \in V$ is an additive inverse of \vec{v} , then \vec{v} is an additive inverse of \vec{w} . So a vector is an additive inverse of any additive inverse of itself.
- b. Vector addition left-cancels: if $\vec{v}, \vec{s}, \vec{t} \in V$ then $\vec{v} + \vec{s} = \vec{v} + \vec{t}$ implies that $\vec{s} = \vec{t}$.

5.1.15 [JH]

The definition of vector spaces does not explicitly say that $\vec{0} + \vec{v} = \vec{v}$ (it instead says that $\vec{v} + \vec{0} = \vec{v}$). Show that it must nonetheless hold in any vector space.

5.1.16 [JH]

Prove or disprove that the following is a vector space: the set of all matrices, under the usual operations.

5.1.17 [JH]

In a vector space every element has an additive inverse. Is the additive inverse unique (Can some elements have two or more)?

5.1.18 [JH]

Assume that $\vec{v} \in V$ is not $\vec{0}$.

- a. Prove that $r \cdot \vec{v} = \vec{0}$ if and only if r = 0.
- b. Prove that $r_1 \cdot \vec{v} = r_2 \cdot \vec{v}$ if and only if $r_1 = r_2$.
- c. Prove that any nontrivial vector space is infinite.

5.2 Subspaces

5.2.1 [JH]

- a. Prove that every point, line, or plane thru the origin in \mathbb{R}^3 is a subspace of \mathbb{R}^3 under the inherited operations.
- b. What if it doesn't contain the origin?

5.3 Spanning Sets

5.3.1 Place Holder

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam tincidunt cursus volutpat. Quisque non congue sem. Vivamus nec nibh sed est dapibus auctor eu sed nulla. Praesent ornare eleifend nibh a finibus. Proin rutrum neque nec massa tincidunt, non malesuada dolor interdum. Nam a massa sit amet diam efficitur pharetra. Nulla interdum efficitur sem, sit amet commodo orci mattis non. Duis tortor ex, maximus a sapien id, molestie maximus risus.

5.4 Linear Independence

5.4.1 Place Holder

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam tincidunt cursus volutpat. Quisque non congue sem. Vivamus nec nibh sed est dapibus auctor eu sed nulla. Praesent ornare eleifend nibh a finibus. Proin rutrum neque nec massa tincidunt, non malesuada dolor interdum. Nam a massa sit amet diam efficitur pharetra. Nulla interdum efficitur sem, sit amet commodo orci mattis non. Duis tortor ex, maximus a sapien id, molestie maximus risus.

5.5 Basis

5.5.1 Place Holder

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam tincidunt cursus volutpat. Quisque non congue sem. Vivamus nec nibh sed est dapibus auctor eu sed nulla. Praesent ornare eleifend nibh a finibus. Proin rutrum neque nec massa tincidunt, non malesuada dolor interdum. Nam a

massa sit amet diam efficitur pharetra. Nulla interdum efficitur sem, sit amet commodo orci mattis non. Duis tortor ex, maximus a sapien id, molestie maximus risus.

5.6 Dimension

5.6.1 Place Holder

Answers to Exercises

1.1.1 Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam tincidunt cursus volutpat. Quisque non congue sem. Vivamus nec nibh sed est dapibus auctor eu sed nulla. Praesent ornare eleifend nibh a finibus. Proin rutrum neque nec massa tincidunt, non malesuada dolor interdum. Nam a massa sit amet diam efficitur pharetra. Nulla interdum efficitur sem, sit amet commodo orci mattis non. Duis tortor ex, maximus a sapien id, molestie maximus risus.

1.2.1

a.
$$x = 2$$
, $y = 3$
b. $x = -1$, $y = 4$, and $z = -1$.

1.3.1 Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam tincidunt cursus volutpat. Quisque non congue sem. Vivamus nec nibh sed est dapibus auctor eu sed nulla. Praesent ornare eleifend nibh a finibus. Proin rutrum neque nec massa tincidunt, non malesuada dolor interdum. Nam a massa sit amet diam efficitur pharetra. Nulla interdum efficitur sem, sit amet commodo orci mattis non. Duis tortor ex, maximus a sapien id, molestie maximus risus.

2.1.1

a.
$$\begin{bmatrix} 16 & -3 & 2 \\ -3 & 7 & -1 \end{bmatrix}$$

b.
$$\begin{bmatrix} -2 & 0 & -2 \\ 3 & -13 & -3 \end{bmatrix}$$

b.
$$\begin{bmatrix} -2 & 0 & -2 \\ 3 & -13 & -3 \end{bmatrix}$$

c. Not possible, since dimension of A and E are not the same.

d.
$$\begin{bmatrix} 7 & 1 \\ 5 & 1 \end{bmatrix}$$

e.
$$\begin{bmatrix} 36 & 19 & 2 \\ 83 & -22 & 11 \\ 19 & -10 & 3 \end{bmatrix}$$

f. Not possible, since the dimension of CD is 2×2 and is not equal to the dimension of D.

g.
$$[9 -7 3]$$

h.
$$\begin{vmatrix} 4 & -2 \\ -6 & 3 \end{vmatrix}$$

2.2.1 Hint: Show that the homogeneous system Ax = 0 has only the trivial solution.

2.3.1

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Note: The answer is not unique.

a.
$$A^{-1} = \begin{bmatrix} 1 & -2 & \frac{1}{3} \\ 0 & 1 & -\frac{2}{3} \\ 0 & 0 & \frac{1}{3} \end{bmatrix}$$

b.
$$x = \begin{bmatrix} \frac{16}{3} \\ \frac{8}{3} \\ \frac{1}{3} \end{bmatrix}$$

3.1.1 Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam tincidunt cursus volutpat. Quisque non congue sem. Vivamus nec nibh sed est dapibus auctor eu sed nulla. Praesent ornare eleifend nibh a finibus. Proin rutrum neque nec massa tincidunt, non malesuada dolor interdum. Nam a massa sit amet diam efficitur pharetra. Nulla interdum efficitur sem, sit amet commodo orci mattis non. Duis tortor ex, maximus a sapien id, molestie maximus risus.

3.2.1 Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam tincidunt cursus volutpat. Quisque non congue sem. Vivamus nec nibh sed est dapibus auctor eu sed nulla. Praesent ornare eleifend nibh a finibus. Proin rutrum neque nec massa tincidunt, non malesuada dolor interdum. Nam a massa sit amet diam efficitur pharetra. Nulla interdum efficitur sem, sit amet commodo orci mattis non. Duis tortor ex, maximus a sapien id, molestie maximus risus.

3.3.1 Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam tincidunt cursus volutpat. Quisque non congue sem. Vivamus nec nibh sed est dapibus auctor eu sed nulla. Praesent ornare eleifend nibh a finibus. Proin rutrum neque nec massa tincidunt, non malesuada dolor interdum. Nam a massa sit amet diam efficitur pharetra. Nulla interdum efficitur sem, sit amet commodo orci mattis non. Duis tortor ex, maximus a sapien id, molestie maximus risus.

3.4.1 Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam tincidunt cursus volutpat. Quisque non congue sem. Vivamus nec nibh sed est dapibus auctor eu sed nulla. Praesent ornare eleifend nibh a finibus. Proin rutrum neque nec massa tincidunt, non malesuada dolor interdum. Nam a massa sit amet diam efficitur pharetra. Nulla interdum efficitur sem, sit amet commodo orci mattis non. Duis tortor ex, maximus a sapien id, molestie maximus risus.

4.1.1 Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam tincidunt cursus volutpat. Quisque non congue sem. Vivamus nec nibh sed est dapibus auctor eu sed nulla.

Praesent ornare eleifend nibh a finibus. Proin rutrum neque nec massa tincidunt, non malesuada dolor interdum. Nam a massa sit amet diam efficitur pharetra. Nulla interdum efficitur sem, sit amet commodo orci mattis non. Duis tortor ex, maximus a sapien id, molestie maximus risus.

- **4.2.1** Analyse the squared norm of $\|\vec{u}\|\vec{v} \|\vec{v}\|\vec{u}$ and $\|\vec{u}\|\vec{v} + \vec{v}\|\vec{u}$
- **4.3.1** Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam tincidunt cursus volutpat. Quisque non congue sem. Vivamus nec nibh sed est dapibus auctor eu sed nulla. Praesent ornare eleifend nibh a finibus. Proin rutrum neque nec massa tincidunt, non malesuada dolor interdum. Nam a massa sit amet diam efficitur pharetra. Nulla interdum efficitur sem, sit amet commodo orci mattis non. Duis tortor ex, maximus a sapien id, molestie maximus risus.
- **4.4.1** Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam tincidunt cursus volutpat. Quisque non congue sem. Vivamus nec nibh sed est dapibus auctor eu sed nulla. Praesent ornare eleifend nibh a finibus. Proin rutrum neque nec massa tincidunt, non malesuada dolor interdum. Nam a massa sit amet diam efficitur pharetra. Nulla interdum efficitur sem, sit amet commodo orci mattis non. Duis tortor ex, maximus a sapien id, molestie maximus risus.
- **4.5.1** Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam tincidunt cursus volutpat. Quisque non congue sem. Vivamus nec nibh sed est dapibus auctor eu sed nulla. Praesent ornare eleifend nibh a finibus. Proin rutrum neque nec massa tincidunt, non malesuada dolor interdum. Nam a massa sit amet diam efficitur pharetra. Nulla interdum efficitur sem, sit amet commodo orci mattis non. Duis tortor ex, maximus a sapien id, molestie maximus risus.

5.1.1

- a. $0 + 0x + 0x^2 + 0x^3$ b. $\begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$
- c. The constant function f(x) = 0
- d. The constant function f(n) = 0

5.1.2

- a. $3 + 2x x^2$ b. $\begin{bmatrix} -1 & +1 \\ 0 & -3 \end{bmatrix}$
- c. $-3e^x + 2e^{-x}$

5.1.3

- a. 1 + 2x, 2 1x, and x.
- b. 2 + 1x, 6 + 3x, and -4 2x.

5.1.4

a.

$$\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}, \begin{bmatrix} -1 & -2 \\ -3 & -4 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

b.

$$\begin{bmatrix} 1 & 2 \\ 0 & 4 \end{bmatrix}, \begin{bmatrix} -1 & -2 \\ 0 & -4 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

5.1.5

- a. (1, 2, 3), (2, 1, 3), and (0, 0, 0).
- b. (1, 1, 1, -1), (1, 0, 1, 0) and (0, 0, 0, 0).

5.1.6

For each part the set is called Q. For some parts, there are more than one correct way to show that Q is not a vector

a. It is not closed under addition.

$$(1, 0, 0), (0, 1, 0) \in Q$$
 $(1, 1, 0) \notin Q$

b. It is not closed under addition.

$$(1, 0, 0), (0, 1, 0) \in Q$$
 $(1, 1, 0) \notin Q$

c. It is not closed under addition.

$$\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} \in Q \qquad \begin{bmatrix} 1 & 2 \\ 0 & 0 \end{bmatrix} \not\in Q$$

d. It is not closed under scalar multiplication.

$$1 + 1x + 1x^2 \in Q$$
 $-1 \cdot (1 + 1x + 1x^2) \notin Q$

- e. The set is empty, violating the existence of the zero vector.
- **5.1.7** No, it is not closed under scalar multiplication since, e.g., $\pi \cdot (1)$ is not a rational number.
- **5.1.8** The '+' operation is not commutative; producing two members of the set witnessing this assertion is easy.

5.1.9

a. It is not a vector space.

$$(1+1) \cdot \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \neq \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} + \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$$

b. It is not a vector space.

$$1 \cdot \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \neq \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$$

- 5.1.10 For each "yes" answer, you must give a check of all the conditions given in the definition of a vector space. For each "no" answer, give a specific example of the failure of one of the conditions.
 - a. Yes.
 - b. Yes.

- c. No, this set is not closed under the natural addition operation. The vector of all 1/4's is an element of this set but when added to itself the result, the vector of all 1/2's, is not an element of the set.
- d. Yes.
- e. No, $f(x) = e^{-2x} + (1/2)$ is in the set but $2 \cdot f$ is not (that is, closure under scalar multiplication fails).
- **5.1.11** Check all 10 conditions of the definition of a vector space.
- **5.1.12** It is not a vector space since it is not closed under addition, as $(x^2) + (1 + x x^2)$ is not in the set.

5.1.13

- a. No since $1 \cdot (0, 1) + 1 \cdot (0, 1) \neq (1+1) \cdot (0, 1)$.
- b. No since the same calculation as the prior part shows a condition in the definition of a vector space that is violated. Another example of a violation of the conditions for a vector space is that $1 \cdot (0, 1) \neq (0, 1)$.

5.1.14

- a. Let V be a vector space, let $\vec{v} \in V$, and assume that $\vec{w} \in V$ is an additive inverse of \vec{v} so that $\vec{w} + \vec{v} = \vec{0}$. Because addition is commutative, $\vec{0} = \vec{w} + \vec{v} = \vec{v} + \vec{w}$, so therefore \vec{v} is also the additive inverse of \vec{w} .
- b. Let V be a vector space and suppose $\vec{v}, \vec{s}, \vec{t} \in V$. The additive inverse of \vec{v} is $-\vec{v}$ so $\vec{v} + \vec{s} = \vec{v} + \vec{t}$ gives that $-\vec{v} + \vec{v} + \vec{s} = -\vec{v} + \vec{v} + \vec{t}$, which implies that $\vec{0} + \vec{s} = \vec{0} + \vec{t}$ and so $\vec{s} = \vec{t}$.

5.1.15

Addition is commutative, so in any vector space, for any vector \vec{v} we have that $\vec{v} = \vec{v} + \vec{0} = \vec{0} + \vec{v}$.

5.1.16

It is not a vector space since addition of two matrices of unequal sizes is not defined, and thus the set fails to satisfy the closure condition.

5.1.17

Each element of a vector space has one and only one additive inverse.

For, let V be a vector space and suppose that $\vec{v} \in V$. If $\vec{w}_1, \vec{w}_2 \in V$ are both additive inverses of \vec{v} then consider $\vec{w}_1 + \vec{v} + \vec{w}_2$. On the one hand, we have that it equals $\vec{w}_1 + (\vec{v} + \vec{w}_2) = \vec{w}_1 + \vec{0} = \vec{w}_1$. On the other hand we have that it equals $(\vec{w}_1 + \vec{v}) + \vec{w}_2 = \vec{0} + \vec{w}_2 = \vec{w}_2$. Therefore, $\vec{w}_1 = \vec{w}_2$.

5.1.18

Assume that $\vec{v} \in V$ is not $\vec{0}$.

- a. One direction of the if and only if is clear: if r=0 then $r \cdot \vec{v} = \vec{0}$. For the other way, let r be a nonzero scalar. If $r\vec{v} = \vec{0}$ then $(1/r) \cdot r\vec{v} = (1/r) \cdot \vec{0}$ shows that $\vec{v} = \vec{0}$, contrary to the assumption.
- b. Where r_1, r_2 are scalars, $r_1 \vec{v} = r_2 \vec{v}$ holds if and only if $(r_1 r_2)\vec{v} = \vec{0}$. By the prior item, then $r_1 r_2 = 0$.

c. A nontrivial space has a vector $\vec{v} \neq \vec{0}$. Consider the set $\{k \cdot \vec{v} \mid k \in \mathbb{R}\}$. By the prior item this set is infinite.

5.2.1

- a. Every such set has the form $\{r \cdot \vec{v} + s \cdot \vec{w} \mid r, s \in \mathbb{R}\}$ where either or both of \vec{v}, \vec{w} may be $\vec{0}$. With the inherited operations, closure of addition $(r_1\vec{v} + s_1\vec{w}) + (r_2\vec{v} + s_2\vec{w}) = (r_1 + r_2)\vec{v} + (s_1 + s_2)\vec{w}$ and scalar multiplication $c(r\vec{v} + s\vec{w}) = (cr)\vec{v} + (cs)\vec{w}$ is clear.
- b. No such set can be a vector space under the inherited operations because it does not have a zero element.
- **5.3.1** Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam tincidunt cursus volutpat. Quisque non congue sem. Vivamus nec nibh sed est dapibus auctor eu sed nulla. Praesent ornare eleifend nibh a finibus. Proin rutrum neque nec massa tincidunt, non malesuada dolor interdum. Nam a massa sit amet diam efficitur pharetra. Nulla interdum efficitur sem, sit amet commodo orci mattis non. Duis tortor ex, maximus a sapien id, molestie maximus risus.
- **5.4.1** Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam tincidunt cursus volutpat. Quisque non congue sem. Vivamus nec nibh sed est dapibus auctor eu sed nulla. Praesent ornare eleifend nibh a finibus. Proin rutrum neque nec massa tincidunt, non malesuada dolor interdum. Nam a massa sit amet diam efficitur pharetra. Nulla interdum efficitur sem, sit amet commodo orci mattis non. Duis tortor ex, maximus a sapien id, molestie maximus risus.
- **5.5.1** Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam tincidunt cursus volutpat. Quisque non congue sem. Vivamus nec nibh sed est dapibus auctor eu sed nulla. Praesent ornare eleifend nibh a finibus. Proin rutrum neque nec massa tincidunt, non malesuada dolor interdum. Nam a massa sit amet diam efficitur pharetra. Nulla interdum efficitur sem, sit amet commodo orci mattis non. Duis tortor ex, maximus a sapien id, molestie maximus risus.
- **5.6.1** Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam tincidunt cursus volutpat. Quisque non congue sem. Vivamus nec nibh sed est dapibus auctor eu sed nulla. Praesent ornare eleifend nibh a finibus. Proin rutrum neque nec massa tincidunt, non malesuada dolor interdum. Nam a massa sit amet diam efficitur pharetra. Nulla interdum efficitur sem, sit amet commodo orci mattis non. Duis tortor ex, maximus a sapien id, molestie maximus risus.

References

- [HE] Harold W. Ellingsen Jr., Matrix Arithmetic, Licensed under the Creative Commons Attribution-ShareAlike 2.5 License.
- [JH] Jim Hefferon, *Linear Algebra*, http://joshua.smcvt.edu/linearalgebra, Licensed under the GNU Free Documentation License or the Creative Commons Attribution-ShareAlike 2.5 License, 2014.
- [YL] Yann Lamontagne, http://obeymath.org, Licensed under the GNU Free Documentation License or the Creative Commons License Creative Commons Attribution-ShareAlike License.

Index

```
additive inverse, 9, 10
Cauchy-Schwartz Inequality, 7
diagonal matrix, 10
function space, 9, 10
Gaussian Elimination, 1
line, plane, 10
matrix space, 9, 10
ph, 1, 3, 5, 7, 10, 11
place holder, 1, 3, 5, 7, 10, 11
polynomial space, 9
positive real numbers, 10
rational numbers, 9
subspace, 10
trivial subspace, 10
vector space, 9, 10
zero vector, 9, 10
```