
a Calls before and after full composition adjustment
Covariates (p: family-wise, max |t|)
batch, menopause, epithelial score
+ all composition scores and age
earlier calls surviving the full model: 0 of 120
RBP calls (BH q < 0.1)
5
p = 0.016
0
p = 0.242
66
p = 0.003
0
p = 0.273
0
p = 0.053
1
p = 0.017
49
p = 0.001
0
p = 0.137
fallopian splicing
fallopian expression
breast splicing
breast expression
b Fallopian splicing-layer calls under graded adjustment
t of carrier status on RBP activity
RRP9
EIF3D
EIF2S1
SRSF3
SUGP2
TAF15
none (p = 0.01)
batch + menopause (p = 0.08)
+ epithelial (p = 0.02)
+ 4 cell-type scores (p = 0.54)
all 10 scores + age (p = 0.30)
Covariates in the model (family-wise p over 478 RBPs)
c Matched pairs, no regression (sign-flip p)
Paired t, carrier minus matched non-carrier (41 composition-matched pairs)
diff +0.0003
p = 0.921
diff +0.0019
p = 0.610
diff -0.0053
p = 0.159
diff -0.0198
p = 0.001
diff -0.0010
p = 0.777
diff +0.0059
p = 0.117
largest |t| of 478
p = 0.001
RRP9
EIF3D
EIF2S1
SRSF3
SUGP2
TAF15
CSNK1E (|t|)
d RRP9 (call lost)
non-carrier (n = 53)
BRCA carrier (n = 41)
RRP9 activity score
r = -0.72 (all)
r = -0.70 (non-carriers)
Ciliated-cell score (log2 CPM)
SRSF3 (kept by matching)
SRSF3 activity score
r = -0.17 (all)
r = -0.27 (non-carriers)
Ciliated-cell score (log2 CPM)