Climate data in Python

A hands-on notebook: test a claim about CO_2 and warming from public data, step by step

Ahmad Ammar

13 November 2026

Contents

1 Before you start **2**

1.1 What you will learn . . . 2

1.2 Getting Python and the notebook . . . 2

1.3 How to read the cells . . . 3

1.4 The question and the libraries . . . 3

2 The data **4**

2.1 Download the data . . . 4

2.2 Look inside each file . . . 5

2.3 Prepare: units, baselines and one joined table . . . 6

3 Trends and their uncertainty **8**

3.1 A trend line, by hand and for every window . . . 8

3.2 Neighbouring years are not independent . . . 9

4 Forcing and a simple climate model **11**

4.1 Forcing and a simple energy-balance model . . . 11

4.2 What drove each period . . . 12

5 Figures and records **13**

5.1 Making and reading the figures . . . 13

5.2 Recording what the run used . . . 14

5.3 Where to go next . . . 15

1 Before you start

This guide is a notebook you run yourself. It tests a claim made at a conference in November 2026, that the Earth warmed fastest in 1899–1940 when CO_2 emissions were small, using four public datasets and a few dozen lines of Python. The essay Reading climate data tells the story and the conclusions; this guide shows every step that produced them, so you can repeat, change and extend it.

1.1 What you will learn

- **Data:** where climate records come from, what a temperature anomaly is, and how to download, inspect and join tables with `pandas`.
- **Statistics:** how a trend line is fitted, by hand and with a library; what a standard error is; why neighbouring years are not independent, and how to allow for it.
- **Physics:** what radiative forcing is, and how a two-reservoir energy-balance model turns forcing into temperature.
- **Figures:** how to read any chart by naming its parts, and how each line and bar maps to a column of a table.
- **Practice:** keeping a record of what each run used, so others can repeat it.

1.2 Getting Python and the notebook

You need Python 3.10 or newer with `numpy`, `pandas`, `matplotlib` and `scipy`. Three ways, from simplest to most flexible:

1. **Google Colab**, in a web browser, with nothing to install: open the *Notebooks* page of this guide and choose *Open in Colab*. Colab already has every library used here.
2. **Anaconda or Miniforge** on your own computer: install it, then in a terminal run the commands below, and open the downloaded `climate_data.ipynb` in the Jupyter window that appears.
3. **VS Code** with its Python and Jupyter extensions: open the `.ipynb` file and choose a Python environment that has the four libraries.

```
conda create -n climate python=3.12 numpy pandas matplotlib scipy jupyterlab
conda activate climate
jupyter lab
```

1.3 How to read the cells

Each step below is a cell of the notebook. A box labelled *In [n]* is the code; the box labelled *Out [n]* under it is what that code printed when the notebook was run, and a figure under *Out [n]* is what it drew. Run the cells in order from the top: later cells use tables made by earlier ones. If you change something and want to start again, use *Restart and run all*.

1.4 The question and the libraries

A working paper presented at a conference in November 2026 argued that the Earth warmed fastest in 1899–1940, when carbon dioxide (CO_2) emissions were small, and concluded that CO_2 is not the main driver of warming. In this notebook we test that claim ourselves, from public data, one step at a time. By the end you will have downloaded four datasets, joined them into one table, fitted trend lines by hand and with a library, measured how uncertain those trends are, built a small physical model of the climate, and made the figures that answer the question.

Each code cell is followed by its output, exactly as it appeared when this notebook was run. Run the cells in order from the top: later cells use what earlier cells made.

We use four Python libraries. **pandas** holds tables (rows and columns, like a spreadsheet); **numpy** does arithmetic on whole columns at once; **matplotlib** draws figures; and **scipy** provides statistical tests.

In [1]

```
import json
import urllib.request
from pathlib import Path

import numpy as np
import pandas as pd
import matplotlib.pyplot as plt
from scipy import stats

pd.set_option("display.width", 110)
plt.rcParams.update({"figure.dpi": 110, "axes.spines.top": False, "axes.spines.right": False})
print("numpy", np.__version__, "| pandas", pd.__version__)
```

Out [1]

```
numpy 2.0.1 | pandas 2.2.3
```

2 The data

Four files hold everything we need. This chapter downloads them, looks inside each one, and joins them into a single table with one row per year.

2.1 Download the data

Every number in this notebook comes from four public files. Each is downloaded once into a folder called `data` next to the notebook, and reused after that. Writing the download as code, rather than clicking in a browser, means anyone can repeat it exactly.

File	What it holds	Who publishes it
`hadcrut5.csv`	Global temperature anomaly, one row per year since 1850	Met Office Hadley Centre and University of East Anglia
`conus.csv`	Temperature of the contiguous United States, one row per year since 1895	NOAA
`owid_co2.csv`	CO_2 emitted by every country, every year	Our World in Data
`forcing.csv`	Radiative forcing by cause, one row per year since 1750	Indicators of Global Climate Change

In [2]

```
DATA = Path("data")
DATA.mkdir(exist_ok=True)
SOURCES = {
    "hadcrut5.csv": "https://www.metoffice.gov.uk/hadobs/hadcrut5/data/HadCRUT.5.0.2.0/analysis/diagnostics/"
                    "HadCRUT.5.0.2.0.analysis.summary_series.global.annual.csv",
    "conus.csv": "https://www.ncei.noaa.gov/access/monitoring/climate-at-a-glance/national/time-series/110/tavg/12/12/1895-2024/data.csv",
    "owid_co2.csv": "https://raw.githubusercontent.com/owid/co2-data/master/owid-co2-data.csv",
    "forcing.csv": "https://raw.githubusercontent.com/ClimateIndicator/forcing-timeseries/main/output/ERF_best_aggregates_1750-2024.csv",
}
for name, url in SOURCES.items():
    target = DATA / name
    if not target.exists():
        req = urllib.request.Request(url, headers={"User-Agent": "climate-data-notebook/1.0"})
        target.write_bytes(urllib.request.urlopen(req, timeout=120).read())
    print(f"{name:14s} {target.stat().st_size / 1e6:6.2f} MB")
```

Out [2]

```
hadcrut5.csv     0.01 MB
conus.csv        0.00 MB
owid_co2.csv    14.38 MB
forcing.csv      0.10 MB
```

2.2 Look inside each file

Before using any data, look at it: how many rows and columns, what the columns are called, and what a few rows contain. `pd.read_csv` reads a file into a table called a DataFrame; `.shape` gives (rows, columns); `.head()` shows the first rows.

HadCRUT5. The anomaly is the temperature minus the average of 1961–1990, so 0 means "normal", positive means warmer than normal. The two limits give the range within which the true value lies with 95% probability.

In [3]

```
g = pd.read_csv(DATA / "hadcrut5.csv")
print(g.shape)
print(g.head(3).to_string())
print("...")
print(g.tail(2).to_string())
```

Out [3]

```
(176, 4)
   Time  Anomaly (deg C)  Lower confidence limit (2.5%)  Upper confidence limit (97.5%)
0  1850        -0.417711                      -0.589256                       -0.246166
1  1851        -0.233350                      -0.411868                       -0.054832
2  1852        -0.229399                      -0.409382                       -0.049416
...
     Time  Anomaly (deg C)  Lower confidence limit (2.5%)  Upper confidence limit (97.5%)
174  2024         1.180800                       1.141589                        1.220011
175  2025         1.190283                       1.048175                        1.332391
```

The last row is the current year, which is not finished, so we will stop at 2024.

NOAA contiguous US. The file starts with lines beginning #, which are notes; `comment="#"` skips them. The `Date` 189512 means the twelve months ending December 1895. Values are in degrees Fahrenheit.

In [4]

```
us_raw = pd.read_csv(DATA / "conus.csv", comment="#")
print(us_raw.shape)
print(us_raw.head(3).to_string())
```

Out [4]

```
(130, 2)
     Date  Value
0  189512  50.33
1  189612  51.98
2  189712  51.54
```

Our World in Data. One row per country per year, with 79 columns. We keep only the rows for the whole World and two columns: `co2`, the CO_2 emitted that year, and `cumulative_co2`, the running total since records began. Both are in millions of tonnes; dividing by 1,000 gives gigatonnes (Gt).

In [5]

```
owid = pd.read_csv(DATA / "owid_co2.csv", usecols=["country", "year", "co2", "cumulative_co2"])
print(owid.shape, "rows and columns;", owid.country.nunique(), "countries and regions")
world = owid[owid.country == "World"].drop(columns="country")
print(world[world.year.isin([1900, 1950, 2000, 2024])].to_string(index=False))
```

Out [5]

```
(50411, 4) rows and columns; 254 countries and regions
 year       co2  cumulative_co2
 1900  1962.884       45105.176
 1950  5930.406      231890.109
 2000 25511.482     1045316.812
 2024 38598.578     1849123.875
```

Radiative forcing. Radiative forcing is the extra energy, in watts per square metre of the Earth's surface (W/m^2), flowing into the climate because something changed: more greenhouse gases (positive, warming), more sulfate pollution (negative, cooling), a brighter Sun, a volcanic eruption. Each column is one cause; `anthro` is all human causes together and `total` adds the natural ones.

In [6]

```
f = pd.read_csv(DATA / "forcing.csv").rename(columns={"Unnamed: 0": "year"})
f["year"] = f["year"].astype(int)
print(f.shape)
print(f[["year", "CO2", "aerosol", "solar", "volcanic", "anthro", "total"]].iloc[[100, 200, 274]].round
    (3).to_string(index=False))
```

Out [6]

```
(275, 19)
 year   CO2  aerosol  solar  volcanic  anthro  total
 1850 0.140   -0.083  0.008     0.244   0.114  0.365
 1950 0.648   -0.714  0.057     0.218   0.344  0.618
 2024 2.332   -1.066  0.091     0.238   2.974  3.303
```

2.3 Prepare: units, baselines and one joined table

Three preparation steps. **Units:** convert NOAA's Fahrenheit to Celsius with °C = (°F − 32) × 5/9. **Baseline:** turn the US temperatures into anomalies by subtracting their 1901–2000 mean. **Join:** put everything into one table with one row per year, matching rows on the `year` column with `merge`.

A worked example of the conversion for 2024: NOAA gives 55.48 °F, so (55.48 − 32) × 5/9 = 13.04 °C.

In [7]

```
us = pd.DataFrame({"year": us_raw["Date"] // 100, "conus": (us_raw["Value"] - 32) * 5 / 9})
base = us.loc[us.year.between(1901, 2000), "conus"].mean()
print(f"US 1901-2000 mean: {base:.2f} °C; 2024: {us.conus.iloc[-1]:.2f} °C, anomaly {us.conus.iloc[-1] -
     base:+.2f} °C")
us["conus"] -= base

d = (g.rename(columns={"Time": "year", "Anomaly (deg C)": "global"})[["year", "global"]]
       .merge(us, on="year", how="left")
       .merge(world, on="year", how="left")
       .merge(f[["year", "anthro", "aerosol", "solar", "volcanic", "total"]], on="year", how="left"))
d = d[d.year <= 2024].reset_index(drop=True)
print(d.shape)
print(d[d.year.isin([1900, 1940, 1982, 2024])].round(3).to_string(index=False))
```

Out [7]

```
US 1901-2000 mean: 11.12 °C; 2024: 13.04 °C, anomaly +1.93 °C
(175, 10)
```

year	global	conus	co2	cumulative_co2	anthro	aerosol	solar	volcanic	total
1900	-0.234	0.417	1962.884	45105.176	0.124	-0.366	-0.037	0.179	0.265
1940	0.076	-0.072	4884.662	181025.359	0.304	-0.658	0.040	0.227	0.571
1982	0.034	-0.378	18729.742	640191.688	0.809	-1.393	0.067	-0.244	0.632
2024	1.181	1.928	38598.578	1849123.875	2.974	-1.066	0.091	0.238	3.303

`how="left"` keeps every year of the first table; where another table has no value for that year (the US series starts in 1895) the cell is `NaN`, which means ”no value”. Missing values are skipped by the calculations below.

3 Trends and their uncertainty

A trend is a slope; the question is how fast temperature rose in each window and how sure we can be. We fit slopes by hand and with a library, then allow for the fact that one year is a lot like the next.

3.1 A trend line, by hand and for every window

A trend is the slope of the straight line that best fits the points. ”Best” means the line that makes the sum of the squared vertical distances from the points to the line as small as possible: ordinary least squares (OLS). For points (x, y), with $\bar{x}$ and $\bar{y}$ their means, the slope is

$$b = \frac{\sum (x - \bar{x})(y - \bar{y})}{\sum (x - \bar{x})^2}$$

It is in degrees per year; multiplying by ten gives degrees per decade. Here it is by hand, for five years:

In [8]

```
small = d[d.year.between(2010, 2014)][["year", "global"]].copy()
small["x - xbar"] = small.year - small.year.mean()
small["y - ybar"] = small["global"] - small["global"].mean()
small["product"] = small["x - xbar"] * small["y - ybar"]
small["(x - xbar)^2"] = small["x - xbar"] ** 2
print(small.round(4).to_string(index=False))
b = small["product"].sum() / small["(x - xbar)^2"].sum()
print(f"\nslope = {small['product'].sum():.4f} / {small['(x - xbar)^2'].sum():.0f} = {b:.4f} °C per year
    ")
print(f"numpy gives {np.polyfit(small.year, small['global'], 1)[0]:.4f}")
```

Out [8]

```
 year  global  x - xbar  y - ybar  product  (x - xbar)^2
 2010  0.6804      -2.0    0.0619  -0.1239           4.0
 2011  0.5377      -1.0   -0.0807   0.0807           1.0
 2012  0.5776       0.0   -0.0408  -0.0000           0.0
 2013  0.6236       1.0    0.0052   0.0052           1.0
 2014  0.6729       2.0    0.0544   0.1089           4.0

slope = 0.0709 / 10 = 0.0071 °C per year
numpy gives 0.0071
```

Five years is far too short for climate: a single El Niño year can tip the slope either way. The paper used 42-year windows, so we do the same, for the global record and for the United States.

In [9]

```
WINDOWS = [(1899, 1940), (1941, 1982), (1983, 2024)]
rows = []
for a, e in WINDOWS:
    w = d[d.year.between(a, e)]
    for col in ["global", "conus"]:
        slope = np.polyfit(w.year, w[col], 1)[0] * 10
        rows.append({"window": f"{a}-{e}", "series": col, "trend per decade": round(slope, 3),
```

```
                    "CO2 emitted (Gt)": round(w.co2.sum() / 1000), "change in human forcing (W/m2)":
    round(w.anthro.iloc[-1] - w.anthro.iloc[0], 2)})
trends = pd.DataFrame(rows)
print(trends.to_string(index=False))
gl = trends[trends.series == "global"].set_index("window")["trend per decade"]
print(f"\nRead across the global rows: the recent window warmed {gl['1983-2024'] / gl['1899-1940']:.1f}
    times as fast as 1899-1940,")
print(f"while {trends['CO2 emitted (Gt)'].iloc[4]:,} Gt of CO2 were emitted in it against {trends['CO2
    emitted (Gt)'].iloc[0]:,} Gt in the first window.")
```

Out [9]

```
   window series  trend per decade  CO2 emitted (Gt)  change in human forcing (W/m2)
1899-1940 global             0.101               140                            0.18
1899-1940  conus             0.168               140                            0.18
1941-1982 global             0.012               459                            0.51
1941-1982  conus            -0.059               459                            0.51
1983-2024 global             0.216              1209                            2.05
1983-2024  conus             0.280              1209                            2.05

Read across the global rows: the recent window warmed 2.1 times as fast as 1899-1940,
while 1,209 Gt of CO2 were emitted in it against 140 Gt in the first window.
```

How to read the table. One row per window and series. ”trend per decade” is the slope in °C per decade: 0.1 means the line rose a tenth of a degree every ten years. ”CO2 emitted” is the total released in that window, and ”change in human forcing” is how much the human push on the climate grew across the window, in W/m^2.

What it says. Both the globe and the United States warmed fastest in the most recent window, 1983–2024, not in 1899–1940. In the middle window the human forcing grew, yet warming nearly stopped: something cancelled it, which the physics section identifies as sulfate aerosols. Comparing trends with emissions in the same window, as the paper did, hides that.

3.2 Neighbouring years are not independent

How sure are we of each slope? The standard error (SE) of a slope measures how much it would vary if the ”noise” around the line were drawn again. The usual formula assumes every point is an independent piece of information. Climate years are not: a warm year tends to follow a warm year, because the oceans store heat and El Niño lasts more than one year. This is **autocorrelation**. We measure it with the lag-1 correlation r_1 of the leftovers (residuals) after removing the trend, and then count how many independent years we really have:

$$n_{\text{eff}} = n\,\frac{1 - r_1}{1 + r_1}$$

The SE grows accordingly. `scipy.stats.linregress` gives the slope and the naive SE; we then correct it.

In [10]

```
def trend_with_uncertainty(years, values):
    fit = stats.linregress(years, values)
    resid = values - (fit.intercept + fit.slope * years)
    r1 = np.corrcoef(resid[:-1], resid[1:])[0, 1]
    n = len(values)
    n_eff = n * (1 - r1) / (1 + r1) if r1 > 0 else n
```

```
    se = fit.stderr * np.sqrt((n - 2) / max(n_eff - 2, 1))
    return {"trend": fit.slope * 10, "se_naive": fit.stderr * 10, "r1": r1, "n_eff": n_eff, "
     se_corrected": se * 10}

rows = []
for a, e in WINDOWS:
    w = d[d.year.between(a, e)]
    t = trend_with_uncertainty(w.year.to_numpy(float), w["global"].to_numpy())
    rows.append({"window": f"{a}-{e}", **{k: round(v, 3) for k, v in t.items()}})
unc = pd.DataFrame(rows)
print(unc.to_string(index=False))
```

Out [10]

```
   window  trend  se_naive    r1  n_eff  se_corrected
1899-1940  0.101     0.014 0.483 14.648         0.026
1941-1982  0.012     0.016 0.475 14.935         0.028
1983-2024  0.216     0.013 0.217 27.023         0.016
```

How to read the table. ”trend” is the slope (°C per decade); ”se_naive” its standard error if every year were independent; ”r1” how alike neighbouring years are after removing the trend (0 = not at all, 1 = identical); ”n_eff” the number of independent years that leaves; ”se_corrected” the standard error allowing for it.

Read the table: in 1983–2024 the global trend is about 0.22 °C per decade with a corrected SE of about 0.02. A 95% range is roughly the trend ± 2 SE, so 0.18 to 0.25. In 1899–1940 it is about 0.10 ± 0.05. The ranges do not overlap: the recent warming really is faster.

4 Forcing and a simple climate model

Temperature responds to the total energy pushed into the climate, from all causes together. This chapter builds the smallest useful model of that response and uses it to split each period's warming into its causes.

4.1 Forcing and a simple energy-balance model

Temperature responds to the total forcing, not to the CO_2 emitted in a window. To see this we build the simplest useful climate model: two reservoirs of heat. The first (land, air, upper ocean) responds within a few years; the second (deep ocean) over centuries. Each year, each reservoir keeps most of last year's warmth and moves a little towards the level the current forcing would eventually produce:

$$T_j(t) = T_j(t-1)\,e^{-1/\tau_j} + q_j\,F(t)\,(1 - e^{-1/\tau_j})$$

with τ the response time (4 and 250 years) and q the sensitivity (0.33 and 0.41 °C per W/m^2), values typical of the calibrations in the IPCC's Sixth Assessment Report. The temperature is the sum of the two reservoirs.

In [11]

```
def respond(forcing, tau=(4.0, 250.0), q=(0.33, 0.41)):
    out = np.zeros(len(forcing))
    for t_, q_ in zip(tau, q):
        a, box = np.exp(-1 / t_), 0.0
        for i, F in enumerate(forcing):
            box = box * a + q_ * F * (1 - a)
            out[i] += box
    return out

# a step of 1 W/m2 switched on in year 2, to see how the model behaves
step = respond(np.r_[0, 0, np.ones(8)])
print("years 0-9 after a 1 W/m2 step:", np.round(step, 3))
```

Out [11]

```
years 0-9 after a 1 W/m2 step: [0.    0.    0.075 0.133 0.179 0.215 0.244 0.266 0.284 0.298]
```

The fast reservoir warms by about 0.07 °C in the first year and keeps rising; the slow one barely moves in ten years. Now we feed in each real cause of forcing separately, at its published size, and add them up.

In [12]

```
f24 = f[f.year <= 2024].reset_index(drop=True)
parts = pd.DataFrame({"year": f24.year,
                      "greenhouse and other": respond((f24.anthro - f24.aerosol).to_numpy()),
                      "aerosol": respond(f24.aerosol.to_numpy()),
                      "solar": respond(f24.solar.to_numpy()),
                      "volcanic": respond(f24.volcanic.to_numpy())})
m = d[["year", "global"]].merge(parts, on="year")
causes = ["greenhouse and other", "aerosol", "solar", "volcanic"]
```

```
offset = (m["global"] - m[causes].sum(axis=1)).mean()        # the only thing fitted: the baseline
m["model"] = m[causes].sum(axis=1) + offset
m["unexplained"] = m["global"] - m["model"]
r2 = 1 - (m.unexplained ** 2).sum() / ((m["global"] - m["global"].mean()) ** 2).sum()
print(f"share of year-to-year variation the model explains: {r2:.0%}")
```

Out [12]

```
share of year-to-year variation the model explains: 87%
```

4.2 What drove each period

Because the model is a sum of causes, each period's trend splits into the trend of each part. The parts add up to the observed trend exactly, with the ”unexplained” column holding what forcing does not account for: natural swings in the oceans and measurement errors.

In [13]

```
rows = []
for a, e in [(1910, 1945), (1946, 1975), (1976, 2024)]:
    w = m[m.year.between(a, e)]
    rows.append({"period": f"{a}-{e}", **{c: round(np.polyfit(w.year, w[c], 1)[0] * 10, 3) for c in ["
     global", *causes, "unexplained"]}})
split = pd.DataFrame(rows)
print(split.to_string(index=False))
for _, row in split.iterrows():
    forced = row[causes].sum()
    print(f"{row.period}: observed {row['global']:+.3f}, forcing explains {forced:+.3f} (greenhouse {row
     ['greenhouse and other']:+.3f}, "
          f"aerosol {row.aerosol:+.3f}); unexplained {row.unexplained:+.3f} °C/decade")
```

Out [13]

```
   period  global  greenhouse and other  aerosol  solar  volcanic  unexplained
1910-1945   0.156                 0.048   -0.021  0.004     0.009        0.116
1946-1975  -0.004                 0.094   -0.088  0.001    -0.010       -0.001
1976-2024   0.204                 0.183   -0.001 -0.003     0.018        0.007
1910-1945: observed +0.156, forcing explains +0.040 (greenhouse +0.048, aerosol -0.021); unexplained
     +0.116 °C/decade
1946-1975: observed -0.004, forcing explains -0.003 (greenhouse +0.094, aerosol -0.088); unexplained
     -0.001 °C/decade
1976-2024: observed +0.204, forcing explains +0.197 (greenhouse +0.183, aerosol -0.001); unexplained
     +0.007 °C/decade
```

How to read the table. One row per period; each column after ”global” is the trend of one cause's part of the model, in °C per decade, and they add up to ”global”. A positive number warmed the period, a negative one cooled it. The printed lines under the table group them: what forcing explains in total, and what is left over.

What it says. Since the mid-1970s greenhouse gases account for almost all of the warming. In 1946–1975 their warming was cancelled by aerosol cooling. In 1910–1945 forcing explains only part of the warming; the rest is natural variability, which climate scientists already knew.

5 Figures and records


5.1 Making and reading the figures

A figure answers a question only if you can say what each part shows. Before reading any chart, name its parts: what one mark is (a year, a bar), which column of the table goes on the x axis and which on the y axis, and in what units; what colour encodes; and whether uncertainty is shown.

In [14]

```
fig, ax = plt.subplots(figsize=(8, 3.6))
ax.plot(d.year, d["global"], color="0.45", lw=1, label="Observed, global (HadCRUT5)")      # x: column '
    year', y: column 'global'
ax.plot(m.year, m.model, color="#1f5c7a", lw=2, label="Model: response to all forcings")    # y: column
    'model'
for (a, e), col in zip(WINDOWS, ["#ca8a04", "#64748b", "#b91c1c"]):                          # one trend
     line per window
    w = d[d.year.between(a, e)]
    k = np.polyfit(w.year, w["global"], 1)
    ax.plot(w.year, np.polyval(k, w.year), color=col, lw=2.5)
ax.set(xlabel="Year", ylabel="Anomaly (°C, vs −19611990)", title="Global temperature, the model, and
    three window trends")
ax.legend(frameon=False, fontsize=8)
plt.tight_layout()
plt.show()
```

Out [14]




Reading it: the x axis is the `year` column; the y axis is the anomaly in °C. Each point on the grey line is one row of the table. The coloured segments are the three window trends from the trend table: the red one, 1983–2024, is the steepest. The blue line is the model, which knows nothing of the observed temperatures except their average level, yet follows the long rise, the dips after volcanic eruptions and the mid-century flat period.

In [15]

```
fig, ax = plt.subplots(figsize=(7, 3.4))
```

```
split.set_index("period")[[*causes, "unexplained"]].plot.bar(ax=ax, width=0.8,
    color=["#b91c1c", "#64748b", "#ca8a04", "#0f766e", "#cbd5e1"])
ax.scatter(range(len(split)), split["global"], color="k", zorder=3, label="observed trend")
ax.axhline(0, color="0.5", lw=0.8)
ax.set(xlabel="", ylabel="Trend (°C per decade)", title="What drove each period's trend")
ax.legend(frameon=False, fontsize=8)
plt.xticks(rotation=0)
plt.tight_layout()
plt.show()
```

Out [15]



Reading it: one group of bars per row of the `split` table; one bar per cause column; the height is that cause's trend in °C per decade, below zero for cooling. The black dot is the observed trend; the bars in each group add up to it.

5.2 Recording what the run used

Good practice is to record what each run used: the versions of the libraries, the size of the data, and the time it took. Then anyone can see what they need to repeat it, and you can compare when the work grows.

In [16]

```
record = {"python_packages": {"numpy": np.__version__, "pandas": pd.__version__},
          "data_files_mb": {p.name: round(p.stat().st_size / 1e6, 2) for p in sorted(DATA.glob("*.csv"))
    },
          "rows_in_joined_table": len(d),
          "result": {"global_trend_1983_2024": round(float(unc.trend.iloc[2]), 3), "
    global_trend_1899_1940": round(float(unc.trend.iloc[0]), 3)}}
print(json.dumps(record, indent=1))
```

Out [16]

```
{
 "python_packages": {
  "numpy": "2.0.1",
  "pandas": "2.2.3"
 },
```

```
 "data_files_mb": {
  "conus.csv": 0.0,
  "forcing.csv": 0.1,
  "hadcrut5.csv": 0.01,
  "owid_co2.csv": 14.38
 },
 "rows_in_joined_table": 175,
 "result": {
  "global_trend_1983_2024": 0.216,
  "global_trend_1899_1940": 0.101
 }
}
```


5.3 Where to go next

- Change a window in `WINDOWS` and rerun: does the ordering of trends change?
- Replace the global series with the US one in the energy-balance section: how much more does a single region vary?
- Read the essay Reading climate data for what these results mean, and the guide Hypothesis testing with climate data for the statistics in depth.