
EGM305 The Atmosphere

Practical Companion

Practicals 1, 2 and 3, worked step by step

The vertical structure of the atmosphere ▪ solar radiation ▪ moisture, motion and weather maps



Practical 1 Practical 2 Practical 3

Part I Practical 1: Vertical structure of the atmosphere

Part II Practical 2: Solar radiation

Part III Practical 3: Moisture, motion and weather maps

This companion works through every question on the three EGM305 practical sheets, in the order of the sheets. Each idea is built from first principles, every formula is derived before it is used, every number is shown and checked, and each chapter ends with practice problems and their answers. All figures were drawn for it from the equations in it and the data on the sheets.

Ahmad Ammar ▪ School of Geography and Environmental Sciences, Ulster University

Contents

1 How to use this companion **1**

1.1 What is in it . . . 1

1.2 How each chapter is laid out . . . 1

1.3 Conventions used throughout . . . 2

1.4 What to have beside you . . . 3

1.5 The printed answer keys . . . 3

1.6 Reading . . . 3

I Practical 1: Vertical structure of the atmosphere **4**

2 Practical 1 at a glance **5**

2.1 What the practical asks of you . . . 5

2.2 The data on the sheet . . . 5

2.3 A word on the printed answer key . . . 6

2.4 Before the practical . . . 6

3 How much air is above you: the halving rule **7**

3.1 Air has weight . . . 7

3.2 Air is squeezed by its own weight . . . 7

3.3 Why the fall is exponential . . . 8

3.4 Question 1: the percentage of the atmosphere above you . . . 9

3.5 Question 2: adding a pressure axis . . . 10

3.6 How good is the halving rule? . . . 11

3.7 Try it yourself: Practice Set 3 . . . 12

4 Reading the graph, and oxygen at altitude **13**

4.1 Reading values from a graph, and checking them . . . 13

4.2 Question 3: five heights . . . 13

4.3 What “21 % oxygen” means . . . 14

4.4 Partial pressure . . . 14

4.5 Question 4: oxygen at five heights . . . 15

4.6 Why it matters . . . 16

4.7 Try it yourself: Practice Set 4 . . . 17

5 Temperature with height: the standard atmosphere **18**

5.1 Temperature and the lapse rate . . . 18

5.2 The standard atmosphere . . . 18

5.3 Question 5: Table 2 . . . 19

5.4 The layers of the atmosphere . . . 19

5.5 Try it yourself: Practice Set 5 . . . 21

6 Two soundings: inversions and the tropopause **22**

6.1 Measuring the atmosphere in the vertical . . . 22

6.2 Question 6: plotting the soundings . . . 22

6.3 Inversions . . . 23

6.4 Question 7: finding the inversions . . . 24

6.5 The tropopause . . . 24

6.6 Question 8: marking the tropopause . . . 25

6.7 Try it yourself: Practice Set 6 . . . 25

7 Lapse rates and the height of the troposphere **26**

7.1 Average and local lapse rates . . . 26

7.2 Question 9: the lapse rate over Hillsborough . . . 26

7.3 Question 10: a tropical profile . . . 27

7.4 Question 11: is the troposphere the same thickness everywhere? . . . 28

7.5 Try it yourself: Practice Set 7 . . . 29

8 Pressure aloft, buoyancy and stability **30**

8.1 Question 12: equal steps, unequal drops . . . 30

8.2 “Cold air sinks”? Density depends on pressure too . . . 31

8.3 What happens to air that moves up or down . . . 31

8.4 Question 13: why the cold air aloft does not sink . . . 33

8.5 Try it yourself: Practice Set 8 . . . 34

9 Two soundings, two climates **35**

9.1 The puzzle . . . 35

9.2 What the soundings show . . . 35

9.3 The ocean and the winds . . . 35

9.4 Question 14: Hillsborough and St John’s . . . 36

9.5 Try it yourself: Practice Set 9 . . . 37

10 Putting Practical 1 together **38**

10.1 One chain of ideas . . . 38
10.2 All the answers in one table . . . 38
10.3 Quick sheet for the lab . . . 39
10.4 Mistakes to avoid . . . 39
10.5 Self-test . . . 40

II Practical 2: Solar radiation **41**

11 Practical 2 at a glance **42**

11.1 What the practical asks of you . . . 42
11.2 Conventions for this practical . . . 43
11.3 A word on the printed answer key . . . 43
11.4 Before the practical . . . 43

12 What radiation is **44**

12.1 Energy that needs no material to travel . . . 44
12.2 Wavelength, frequency and speed . . . 44
12.3 Photons: radiation comes in packets . . . 45
12.4 The electromagnetic spectrum . . . 46
12.5 Why everything above absolute zero radiates . . . 46
12.6 Black bodies . . . 47
12.7 The Planck curve . . . 47
12.8 Questions that come up . . . 48
12.9 Try it yourself: Practice Set 12 . . . 49

13 Wien's law and the temperature of the Sun **50**

13.1 Hotter bodies peak at shorter wavelengths . . . 50
13.2 Wien's displacement law . . . 50
13.3 Question 1: the temperature of the Sun . . . 51
13.4 Wien's law in practice . . . 52
13.5 Try it yourself: Practice Set 13 . . . 52

14 The Stefan–Boltzmann law: how much the Sun and the Earth emit **54**

14.1 The total emission of a black body . . . 54
14.2 What a fourth power means . . . 54
14.3 Question 2a: the emission of the Sun . . . 55
14.4 Question 2b: the emission of the Earth . . . 56

14.5 Question 2c: how many times hotter is the Sun? . . . 56
14.6 Question 2d: how many times more does the Sun emit? . . . 57
14.7 Question 2e: why the two ratios are so different . . . 57
14.8 From the Sun's surface to the Earth . . . 58
14.9 Questions that come up . . . 58
14.10Try it yourself: Practice Set 14 . . . 59

15 Plotting three planetary spectra **60**
15.1 The data . . . 60
15.2 How to plot the graph . . . 60
15.3 Question 3a: the wavelength of the peak . . . 61
15.4 Question 3b: the temperatures of the three planets . . . 62
15.5 What the graph shows . . . 62
15.6 Try it yourself: Practice Set 15 . . . 63

16 The temperatures of the planets **64**
16.1 A planet in balance . . . 64
16.2 Sunlight at each planet: the inverse-square law . . . 64
16.3 Albedo . . . 64
16.4 The black-body temperature of a planet . . . 65
16.5 Question 3c: calculating the temperatures . . . 65
16.6 Question 3d: which spectrum belongs to which planet? . . . 66
16.7 Question 3e: why the real temperatures differ . . . 67
16.8 Try it yourself: Practice Set 16 . . . 69

17 The Earth's energy balance and the greenhouse effect **71**
17.1 What Question 4 sets out to do . . . 71
17.2 Question 4a: how much is absorbed? . . . 71
17.3 Question 4b: sunlight facing the Sun . . . 71
17.4 Question 4b′: why the average is only 341 $\mathrm{W\,m^{-2}}$ . . . 72
17.5 Question 4c: the global average . . . 73
17.6 Question 4d: the temperature of an airless Earth . . . 74
17.7 Question 4e: what the atmosphere does . . . 74
17.8 Question 4f: day and night without an atmosphere . . . 76
17.9 Question 4g: the assumptions . . . 77
17.10Try it yourself: Practice Set 17 . . . 77

18 Sun angle, solar intensity and the seasons **79**

18.1 A spread beam is a weaker beam . . . 79
18.2 Measuring the spreading . . . 79
18.3 Where the noon Sun is overhead . . . 80
18.4 Question 5a: sketches of the noon Sun . . . 80
18.5 Question 5b: why the sun angle makes the seasons . . . 82
18.6 Question 5c: comparing solar intensities . . . 83
18.7 Try it yourself: Practice Set 18 . . . 84

19 Putting Practical 2 together **86**

19.1 One chain of ideas . . . 86
19.2 All the answers in one table . . . 86
19.3 Mistakes to avoid . . . 87
19.4 Self-test . . . 88

III Practical 3: Moisture, motion and weather maps **89**

20 Practical 3 at a glance **90**

20.1 What the practical asks of you . . . 90
20.2 What to have beside you . . . 91
20.3 Symbols and units . . . 91
20.4 A word on the printed answer key . . . 91
20.5 About the weather map . . . 91
20.6 Before the practical . . . 92

21 Moisture in the air: mixing ratio and relative humidity **93**

21.1 Why water vapour matters . . . 93
21.2 Vapour pressure and saturation . . . 93
21.3 Mixing ratio . . . 94
21.4 Relative humidity . . . 95
21.5 A picture to hold on to: the glass . . . 96
21.6 Reading Table 1 . . . 96
21.7 Question 1: relative humidity of five samples . . . 97
21.8 Question 2: cold air heated in a house . . . 98
21.9 Question 3: who holds the most water? . . . 98
21.10 Question 4: the disadvantage of relative humidity . . . 100
21.11 The two slips in Table 1 . . . 100
21.12 Try it yourself: Practice Set 21 . . . 101

22 The dew point **102**

22.1 Cooling the glass . . . 102
22.2 Three ways to find a dew point . . . 102
22.3 Question 5: 20 °C and 10 g kg^{-1} . . . 102
22.4 Question 6: 34 °C and 62 % humidity . . . 104
22.5 A quick check by rule of thumb . . . 104
22.6 What happens below the dew point . . . 105
22.7 Why meteorologists prefer the dew point . . . 106
22.8 The dew point and the height of cloud base . . . 106
22.9 Questions that come up . . . 107
22.10 Try it yourself: Practice Set 22 . . . 107

23 Why air moves: the three forces **109**

23.1 Newton's law for a parcel of air . . . 109
23.2 Pressure, and why it falls with height . . . 109
23.3 The pressure gradient force . . . 109
23.4 The Coriolis force . . . 111
23.5 Friction . . . 112
23.6 The sheet's table, completed . . . 112
23.7 Three situations to hold in your head . . . 112
23.8 Try it yourself: Practice Set 23 . . . 113

24 Pressure surfaces and the pressure gradient **114**

24.1 What a pressure surface is . . . 114
24.2 Why cold air gives low pressure surfaces . . . 114
24.3 The data of Figure 1 . . . 115
24.4 Question 1: plot the pressure surfaces . . . 116
24.5 Questions 2 and 3: where does pressure fall fastest? . . . 117
24.6 Question 4: which point has the higher pressure? . . . 117
24.7 Question 5: where is the pressure gradient greater? . . . 118
24.8 Question 6: what this means for the wind . . . 119
24.9 An experiment: change the temperatures . . . 120
24.10 Try it yourself: Practice Set 24 . . . 120

25 The geostrophic wind **122**

25.1 Two forces in balance . . . 122
25.2 How the balance comes about: the four stages of Figure 3 . . . 122
25.3 Question 7 (Figure 3): the Northern Hemisphere . . . 124

25.4 Question 7 (Figure 4): the Southern Hemisphere . . . 124
25.5 Buys-Ballot's law . . . 125
25.6 How fast is the geostrophic wind? . . . 126
25.7 When is the wind not geostrophic? . . . 127
25.8 Questions that come up . . . 127
25.9 Try it yourself: Practice Set 25 . . . 127

26 Wind near the ground: friction **129**
26.1 Question 8: what does a slower wind do to the Coriolis force? . . . 129
26.2 The balance of three forces . . . 129
26.3 Question 9: the forces in Figure 5 . . . 129
26.4 The numbers: how much does friction turn the wind? . . . 131
26.5 Question 10: which system is which? . . . 132
26.6 Question 11: the forces on one box in each system . . . 133
26.7 Why this matters for weather . . . 133
26.8 Questions that come up . . . 134
26.9 Try it yourself: Practice Set 26 . . . 135

27 Reading a station model **136**
27.1 Why a station model? . . . 136
27.2 The anatomy of a station model . . . 136
27.3 Decoding the pressure . . . 136
27.4 Temperature and dew point . . . 137
27.5 Sky cover . . . 137
27.6 Wind: the barbs . . . 138
27.7 Present weather and pressure tendency . . . 139
27.8 Question 12: the four station models of the sheet . . . 140
27.9 What the models tell you . . . 142
27.10Try it yourself: Practice Set 27 . . . 142

28 Streamlines **144**
28.1 What a streamline is . . . 144
28.2 The method . . . 144
28.3 Figure 8: the stations . . . 145
28.4 Question 13: the streamlines . . . 146
28.5 What the streamlines show . . . 147
28.6 Common mistakes . . . 147
28.7 Try it yourself: Practice Set 28 . . . 148

29 Fronts and the weather map **149**

29.1 Air masses and fronts . . . 149

29.2 Figure 9: a low with a warm front and a cold front . . . 149

29.3 Question 14: the full analysis of Figure 10 . . . 151

29.3.1 Step 0: read the pressures . . . 151

29.3.2 Step 1: isobars every 4 mb . . . 151

29.3.3 Step 2: the L . . . 153

29.3.4 Step 3: the fronts . . . 154

29.3.5 Step 4: cloud cover above 75 % . . . 155

29.3.6 Step 5: the precipitation . . . 155

29.4 Reading the finished picture . . . 159

29.5 Questions that come up . . . 159

29.6 Try it yourself: Practice Set 29 . . . 160

30 Putting Practical 3 together **161**

30.1 One chain of ideas . . . 161

30.2 All the answers in one table . . . 161

30.3 Mistakes to avoid . . . 161

30.4 Self-test . . . 162

30.5 Before the practical: a checklist . . . 164

A Practical 1: formulas, derivations, data and corrections **165**

A.1 Formula sheet . . . 165

A.2 Derivations for the curious . . . 165

A.2.1 Pressure falls exponentially with height . . . 165

A.2.2 The dry adiabatic lapse rate and potential temperature . . . 166

A.3 The data of the sheet . . . 166

A.4 Corrections to the printed key, at a glance . . . 167

A.5 Symbols . . . 168

A.6 About the figures and the data . . . 168

B Practical 2: formulas, derivations, data and corrections **170**

B.1 Formula sheet . . . 170

B.2 Derivations for the curious . . . 171

B.2.1 Wien's law from the Planck curve . . . 172

B.2.2 The Stefan–Boltzmann law from the Planck curve . . . 172

B.2.3 The black-body temperature of a planet . . . 172

B.2.4 Beam spreading . . . 173

B.2.5 The noon zenith angle . . . 173
B.3 The spectra of Question 3, recomputed . . . 173
B.4 Corrections to the printed key, at a glance . . . 173
B.5 Symbols and constants . . . 175
B.6 About the figures and the data . . . 176
B.6.1 How the figures were made . . . 176
B.6.2 Limits to bear in mind . . . 176

C Practical 3: formulas, data, derivations and corrections 178
C.1 Formula sheet . . . 178
C.2 Table 1, with the two corrections . . . 179
C.3 The 34 stations of Figures 8 and 10 . . . 180
C.4 Derivations for the curious . . . 181
C.4.1 Mixing ratio and vapour pressure . . . 181
C.4.2 Saturation vapour pressure . . . 181
C.4.3 The dew point . . . 182
C.4.4 Cloud base from the dew point . . . 182
C.4.5 The hypsometric equation . . . 182
C.4.6 The pressure gradient force on a pressure surface . . . 182
C.4.7 The geostrophic wind . . . 182
C.4.8 The steady wind with friction . . . 183
C.4.9 Pressure at a height between two reported levels . . . 183
C.5 Weather symbols: quick reference . . . 183
C.6 Corrections to the printed key, at a glance . . . 183
C.7 About the figures, the data and further reading . . . 185
C.7.1 How the figures were made . . . 185
C.7.2 Limits to bear in mind . . . 185
C.7.3 Further reading . . . 186

D Constants, units and conversions 187
D.1 Constants . . . 187
D.2 Units and conversions . . . 188

References 189

Chapter 1

How to use this companion

1.1 What is in it

This companion works through the three practicals of EGM305 The Atmosphere, in the order of the module. Each practical is one part, and each part follows on from the one before: Practical 1 describes the structure of the atmosphere, Practical 2 explains how the Sun heats it, and Practical 3 how that uneven heating fills it with moisture and sets it in motion.

Table 1.1. The three parts of the companion.

Part	Practical and what it covers	Chapters	Appendix
I	**Practical 1: Vertical structure of the atmosphere.** Pressure and the halving rule, oxygen at altitude, the standard atmosphere, soundings, inversions and the tropopause, lapse rates, stability, climates.	2–10	A
II	**Practical 2: Solar radiation.** The laws of radiation, black bodies and planetary temperatures, the Earth's energy balance and the greenhouse effect, sun angle and the seasons.	11–19	B
III	**Practical 3: Moisture, motion and weather maps.** Humidity and the dew point, the forces that move the air, the geostrophic wind and friction, station models, streamlines, fronts and a full weather-map analysis.	20–30	C

Every part opens with a chapter that maps each question of the sheet to the chapter that answers it, and closes with a chapter that gathers all the answers in one table, a list of mistakes to avoid and a self-test. Its appendix holds the formulas, derivations, data, corrections to the printed key and symbols for that practical. Appendix D lists the constants and units shared by all three, and the references for the whole companion are at the end.

The companion is published as one book and as three separate booklets, one per practical, with the same numbering: Chapter 14 is the same chapter, with the same numbered equations, figures and tables, in the book and in the Practical 2 booklet. A reference in a booklet to another part opens the full companion at that place.

1.2 How each chapter is laid out

Every chapter follows the same order, so you always know where to look.

1. **The ideas first,** in plain language, before any symbol appears. Every important equation is written on its own line and numbered, for example (3.4), and is followed by a *where* list that says what each symbol stands for, its unit and the value used. Formulas are derived before they are used: short derivations sit in the text, longer ones in the part's appendix.
2. **The question**, as printed on the sheet, in a box like this:

The question on the sheet

The wording of the sheet.

3. **The method**, one step per line, with every number substituted and every unit carried.
4. **The answer**, in a green box, followed by a **check** that it is sensible.

Answer to Question n

The answer, with the key number in bold.

5. **Worked examples** in blue-headed boxes, in which every number is shown, and **key ideas** in blue boxes.
6. **Corrections to the printed key**, where the official answer sheet has a slip or a gap, in a red box. Each part collects its corrections in its appendix.
7. **Questions that come up**: the doubts that most often arise at that point, answered (orange boxes).
8. **Try it yourself**: a practice set at the end of the chapter, named after the chapter (Practice Set 3 belongs to Chapter 3), with answers in the green box that follows.

How to get the most out of the companion

Cover the green answer box, work the question on paper, then uncover it. If your answer differs, find the step where you diverged before reading on. That step is where the learning is. The PDF is fully hyperlinked: every equation, figure, table, chapter, section and reference number can be clicked, and the back button of your PDF reader returns you to where you were.

1.3 Conventions used throughout

- **Numbering.** "Eq. (4.2)" is equation 2 of Chapter 4; figures and tables are numbered the same way. "§ 4.3" is section 3 of Chapter 4. Appendices are lettered A to D, one per practical and one for shared constants.
- **Units** are SI, written as powers: $\mathrm{W\,m^{-2}}$ means watts per square metre, $\mathrm{m\,s^{-1}}$ metres per second. Pressure is in hectopascals, which are the same as millibars ($1\ \mathrm{hPa} = 1\ \mathrm{mb} = 100\ \mathrm{Pa}$). Heights are in metres or kilometres, as the sheets give them.
- **Temperature.** Temperatures read from or written on the sheets are in degrees Celsius. Any formula in which temperature is multiplied, divided or raised to a power (the gas law, the radiation laws, the potential temperature) needs kelvin: $T_{\mathrm{K}} = T_{^\circ\mathrm{C}} + 273.15$. A *difference* of temperature is the same in both scales, so lapse rates can be written in $^\circ\mathrm{C\,km^{-1}}$ or $\mathrm{K\,km^{-1}}$.
- **Constants.** Each part uses the values printed on its sheet, so that your answers match the key; Appendix D gives precise values and sources.
- **Rounding.** Intermediate results keep at least four significant figures; answers are rounded at the end, usually to three or four.
- **Cross-references between parts** are written in the same way as within a part, with the part named: "Part II, § 17.7".
- **References** to the literature are given as author and year, for example Stull (2017), and listed in full at the end. The lectures are cited by slide number, for example EGM305 module team (2020, slide 14).

1.4 What to have beside you

- The practical sheet itself, with its tables and figures; the companion prints the data again (in the part's appendix) wherever a question needs them.
- A calculator with powers, logarithms and trigonometric functions, set to **degrees**.
- Graph paper or the graphs on the sheet, a ruler, a soft pencil, and three or four coloured pencils so that each line on a graph has its own colour.

1.5 The printed answer keys

Every answer in the companion was recomputed independently from the data on the sheets and checked against the printed answer keys. Almost everything in the keys is right. Where a key has a slip (a decimal point, a typing error, a unit), a gap, or an explanation that can be sharpened, the companion says so in a red box at the place where it arises, and each part lists all of its corrections in its appendix: Practical 1 in § A.4, Practical 2 in § B.4, Practical 3 in § C.6. None of the corrections changes the physics a practical is teaching.

1.6 Reading

The module's set texts are Lutgens et al. (2019) and Ahrens and Henson (2022); each part names the chapters to read. A free textbook that covers all three practicals with worked examples is Stull (2017). For more depth, Wallace and Hobbs (2006) treats the physics of every chapter of this companion at the next level up.

Part I

Practical 1: Vertical structure of the atmosphere

Chapter 2

Practical 1 at a glance

2.1 What the practical asks of you

Practical 1 is about the *vertical structure of the atmosphere*: how the amount of air, its pressure, its oxygen and its temperature change as you go up (EGM305 module team, 2025b). The fourteen questions fall into four groups, and each group needs the one before it.

How much air is above you (Questions 1–4). A simple rule, "half the atmosphere is above you for every 5.6 km you climb", gives the fraction of the atmosphere and the pressure at any height. You draw that relationship, read values from it, and use it to find how much oxygen there is on Everest or in an aircraft cabin.

How the temperature changes (Questions 5–8). You build the standard atmosphere, plot two real weather-balloon soundings beside it, and find the inversions and the tropopause on each.

Lapse rates and the shape of the troposphere (Questions 9–11). You measure a lapse rate from data, add a tropical sounding, and explain why the troposphere is thickest over the tropics.

Why the structure holds together (Questions 12–14). You explain why pressure falls fast near the ground and slowly high up, why the cold air aloft does not sink, and why Northern Ireland is milder than Newfoundland in March.

Table 2.1. Where each question of the Practical 1 sheet is answered, and the equations it needs.

Question	What it asks	Chapter	Equations
1, 2	percentage of the atmosphere above, and pressure, at each height; Table 1 and the graph	3	(3.5), (3.6)
3	read five heights off the graph	4	(3.4)
4	partial pressure of oxygen at those heights	4	(4.1)
5	the standard atmosphere, Table 2	5	(5.1), (5.2)
6, 7, 8	plot the soundings; inversions; tropopause	6	(5.1)
9, 10, 11	lapse rate over Hillsborough; Florida; thickness of the troposphere	7	(5.1)
12, 13	why pressure falls more slowly aloft; why cold air aloft does not sink	8	(3.2)–(8.3)
14	Hillsborough against St John's	9	–

Chapter 10 gathers every answer in one table. The formulas, derivations, data, corrections to the key and symbols for this practical are in Appendix A.

2.2 The data on the sheet

The sheet gives four sets of numbers. All are reproduced in Appendix A.3.

- **Table 1** is yours to complete: height, percentage of the atmosphere above, and pressure, in steps of 5.6 km.
- **Table 2** is the *standard atmosphere*: the temperature at ten heights from sea level to 16 km, three of them filled in.

- **Table 3** gives two radiosonde (weather-balloon) soundings made on 21 March 1990: **3a** at Hillsborough, County Down, Northern Ireland, and **3b** at St John's, Newfoundland, Canada. Each line is a pressure, a height and a temperature, from near the ground to about 30 km.
- **Table 4** is a temperature profile for Florida, every 2 km from sea level to 18 km.

A note on the printed Table 3

On the sheet the two soundings are printed side by side in one table, and when the sheet is copied or opened on a screen the columns can run together. The two stations are told apart most easily by the first line: Hillsborough starts at 1001 hPa, 38 m and 11.05 °C; St John's at 1005 hPa, 140 m and −1.65 °C. Table A.1 (Appendix A.3) prints them separately.

2.3 A word on the printed answer key

The answer key was checked against independent calculation and against the data, one answer at a time. Most of it is right. Question 9 has no answer in the key; the tropopause temperature for Florida is mistyped; the oxygen percentages of Question 4 answer one of two possible readings of the question; and two of the written explanations (Questions 12 and 13) are worth sharpening. Each is explained where it arises, and § A.4 lists them all.

2.4 Before the practical

The Week 1 lecture introduces the atmosphere as a system and its vertical structure (EGM305 module team, 2026, slides 35–50), and the first half of the Week 2 lecture covers its composition, the gas law, pressure and density with height, and the layers of temperature (EGM305 module team, 2020, slides 5–26). The lecture asks you to read Chapter 1 of Lutgens et al. (2019) before this practical; Chapter 1 of Ahrens and Henson (2022) covers the same ground. A free alternative is Chapter 1 of Stull (2017), which develops pressure, density and temperature with height with worked examples.

Chapter 3

How much air is above you: the halving rule

This chapter answers Questions 1 and 2 of the sheet.

3.1 Air has weight

Air is matter: every cubic metre near the ground has a mass of about 1.2 kg. Gravity pulls all of it towards the Earth, so every layer of the atmosphere rests on the layers below. **Atmospheric pressure** at any height is the weight of all the air above, pressing on each square metre (EGM305 module team, 2020, slides 7 and 14). At sea level it averages 1013.25 hPa (NOAA et al., 1976); the sheet rounds this to 1000 hPa.

Units of pressure

Pressure is force per unit area. The SI unit is the pascal, 1 Pa = 1 $\mathrm{N\,m^{-2}}$. Meteorologists use the **hectopascal**, 1 hPa = 100 Pa, which is exactly the same as the older **millibar**: 1 hPa = 1 mb. Sea-level pressure is about 1000 hPa = 100 000 Pa.

How heavy is the air above you? A pressure of 100 000 Pa means a force of 100 000 N on every square metre. Dividing by the acceleration due to gravity, $g = 9.81\ \mathrm{m\,s^{-2}}$, gives the mass that exerts it:

$$\frac{100\,000\ \mathrm{N\,m^{-2}}}{9.81\ \mathrm{m\,s^{-2}}} = 10\,200\ \mathrm{kg\,m^{-2}}.$$

About ten tonnes of air sit above every square metre of the ground. We do not feel it because the pressure acts equally in all directions and the air inside our bodies pushes back.

3.2 Air is squeezed by its own weight

Water is almost incompressible: the bottom of the ocean is hardly denser than the top. Air is very compressible, and the air near the ground carries the weight of all the air above it, so it is squeezed the most. The lowest layers are therefore the densest, and density (and pressure) fall with height (EGM305 module team, 2020, slides 8–9). How fast? Two physical laws answer that.

The ideal gas law links the pressure, density and temperature of a gas (EGM305 module team, 2020, slide 12):

$$p = \rho R_d T. \tag{3.1}$$

where

p	pressure (Pa)	
ρ	density of the air ($\mathrm{kg\,m^{-3}}$)	about 1.2 near the ground
R_d	gas constant for dry air ($\mathrm{J\,kg^{-1}\,K^{-1}}$)	287, Table D.1
T	absolute temperature (K)	

At a fixed temperature, pressure and density are proportional: squeeze the air to twice the density and its pressure doubles.

Hydrostatic balance. Picture a thin horizontal slab of air, one square metre in area and Δz thick. Its weight is its mass times g, that is $\rho\,\Delta z\,g$. The slab stays where it is (it does not fall or fly upwards) because the pressure pushing up on its underside exceeds the pressure pushing down on its top by exactly its weight. So the pressure must fall by $\rho g\,\Delta z$ across the slab (Wallace and Hobbs, 2006, ch. 1):

$$\frac{\Delta p}{\Delta z} = -\rho g. \tag{3.2}$$

where

Δp	change in pressure across the slab (Pa)	negative going up
Δz	thickness of the slab (m)	
ρ	density of the air in the slab ($\mathrm{kg\,m^{-3}}$)	
g	acceleration due to gravity ($\mathrm{m\,s^{-2}}$)	9.81




Figure 3.1. Hydrostatic balance. The slab neither rises nor falls, so the upward push of the pressure on its underside, p, exceeds the downward push on its top, $p + \Delta p$ (with Δp negative), by exactly its weight: $-\Delta p = \rho g\,\Delta z$, Eq. (3.2).

Near the ground $\rho g = 1.2 \times 9.81 = 11.8\,\mathrm{Pa}$ per metre, so the pressure falls by about 1 hPa for every 8.5 m you climb. You can feel this in your ears in a lift or on a hill.

3.3 Why the fall is exponential

Put the two laws together. By Eq. (3.2) the pressure falls by an amount proportional to the density, and by Eq. (3.1) the density is proportional to the pressure. So *the pressure falls by an amount proportional to the pressure itself*: where there is a lot of air, it falls fast; where there is little, it falls slowly. A quantity that shrinks in proportion to itself decays **exponentially**, like a radioactive substance or money losing value at a fixed rate of inflation. If the temperature is taken as constant, the result is (Appendix A.2.1)

$$p(z) = p_0\, e^{-z/H}, \qquad H = \frac{R_d T}{g}. \tag{3.3}$$

where

$p(z)$	pressure at height z (hPa)	calculated
p_0	pressure at sea level (hPa)	1000, on the sheet
z	height above sea level (km)	
H	scale height: the height over which the pressure falls by a factor $e = 2.718$ (km)	8.1, for the sheet's rule
T	a representative temperature of the air column (K)	

The **scale height** H is about 7 to 8.5 km for the temperatures of the lower atmosphere (EGM305 module team, 2020, slide 9).

The halving rule is the same law

Any exponential decay has a fixed *halving distance*: the height over which the pressure halves is $z_{1/2} = H \ln 2 = 0.693\,H$. The sheet's rule, "for every 5.6 km you go up, there is half the atmosphere above you", is Eq. (3.3) with $z_{1/2} = 5.6$ km, that is $H = 5.6/0.693 = 8.08$ km. Written with halvings instead of powers of e, the fraction of the atmosphere above height z is

$$f(z) = \left(\tfrac{1}{2}\right)^{z/5.6} = 2^{-z/5.6}, \tag{3.4}$$

where

$f(z)$	fraction of the whole atmosphere that lies above height z (0 to 1; multiply by 100 for %)	calculated
z	height above sea level (km)	
5.6	the halving height (km)	on the sheet

and after n whole steps of 5.6 km (that is, at $z = 5.6\,n$ km) it is simply

$$f_n = \frac{1}{2^n}, \qquad \text{percentage above} = \frac{100\,\%}{2^n}. \tag{3.5}$$

3.4 Question 1: the percentage of the atmosphere above you

The question on the sheet

"For every 5.6 km you go up, there is half the atmosphere above you as when you started." Keeping this rule in mind, calculate and draw the percentage of atmosphere that is above each point (x-axis: percentage of the atmosphere above (%); y-axis: height above sea level (km)).

Method.

Step 1. Start at sea level with 100 % of the atmosphere above you.

Step 2. Each row of Table 1 is 5.6 km higher than the last, so add 5.6 km to the height and halve the percentage, Eq. (3.5): $100 \to 50 \to 25 \to 12.5 \to \ldots$

Step 3. Continue until the percentage is well below 1 %: ten rows take you to 50.4 km.

Step 4. Plot each row as a point, percentage across and height up, and join the points with a *smooth* curve (Figure 3.2).

Answer to Question 1

100, 50, 25, 12.5, 6.25, 3.13, 1.56, 0.78, 0.39 and 0.20 % of the atmosphere is above sea level, 5.6, 11.2, 16.8, 22.4, 28.0, 33.6, 39.2, 44.8 and 50.4 km (Table 3.1), drawn as the curve of Figure 3.2.

Check. Half the air is below 5.6 km and three-quarters below 11.2 km, which matches the lecture's statement that half the mass of the atmosphere is within about 5 km of the surface (EGM305 module team, 2020, slide 14). The curve never reaches zero: each step halves what is left, so there is always a little air above you. That is why there is no sharp "top" to the atmosphere.

Table 3.1. Table 1 of the sheet, completed (Questions 1 and 2). The last column is the U.S. Standard Atmosphere (NOAA et al., 1976), for comparison only.

Step n	Height (km)	% above	Pressure (hPa)	Standard atmosphere (hPa)
0	sea level	100	1000	1013
1	5.6	50	500	498
2	11.2	25	250	219
3	16.8	12.5	125	91
4	22.4	6.25	62.5	38
5	28.0	3.13	31.3	16
6	33.6	1.56	15.6	6.8
7	39.2	0.78	7.8	3.0
8	44.8	0.39	3.9	1.4
9	50.4	0.20	2.0	0.6

A question that comes up

Why do I get a curve and not a straight line?
Because equal steps up remove equal *fractions*, not equal *amounts*. The first 5.6 km removes 50 percentage points; the next removes 25; the next 12.5. Plotted against height, that is a curve that is steep near the ground and flattens higher up, the shape of every exponential decay. Plotted with a logarithmic pressure axis it would become a straight line (Figure 8.1).

3.5 Question 2: adding a pressure axis

The question on the sheet

As the density of the atmosphere drops by 50 % when we go up 5.6 km, the pressure also drops by 50 %. At sea level the pressure is approximately 1000 hPa. Add the pressure for each of your heights in the third column of Table 1 and add a second x-axis to your graph to show atmospheric pressure.

Why pressure follows the percentage. Pressure is the weight of the air above (§ 3.1). If only half the atmosphere is above you, only half its weight presses down, so the pressure is half the sea-level pressure. In general

$$p(z) = p_0 \times f(z) = 1000 \text{ hPa} \times f(z). \tag{3.6}$$

where

$p(z)$	pressure at height z (hPa)	calculated
p_0	sea-level pressure (hPa)	1000, on the sheet
$f(z)$	fraction of the atmosphere above, Eq. (3.4)	Question 1

With $p_0 = 1000$ hPa the pressure in hectopascals is simply ten times the percentage.

The second axis. Because the pressure is exactly ten times the percentage, the pressure axis is the percentage axis relabelled: 0 to 1000 hPa along the top where 0 to 100 % runs along the bottom (Figure 3.2). The same curve then answers both questions.




Figure 3.2. Questions 1 and 2. The percentage of the atmosphere above each height (bottom axis) and the pressure (top axis) by the halving rule. Each step of 5.6 km up halves both. The dashed line is the U.S. Standard Atmosphere: the rule is very close near the ground and overestimates the pressure higher up (§ 3.6).

Answer to Question 2

Pressure = 10× the percentage above: **1000, 500, 250, 125, 62.5, 31.3, 15.6, 7.8, 3.9 and 2.0 hPa** at sea level, 5.6, 11.2, …, 50.4 km (Table 3.1). The second axis runs from 0 to 1000 hPa above the 0 to 100 % axis.

3.6 How good is the halving rule?

The last column of Table 3.1 compares the rule with the U.S. Standard Atmosphere, an idealised but realistic average atmosphere (NOAA et al., 1976). Up to about 6 km the two agree within 1 %. Higher up the rule gives too much pressure: 250 against 219 hPa at 11.2 km, 125 against 91 hPa at 16.8 km.

The reason is the temperature in the scale height, $H = R_d T/g$, Eq. (3.3). The rule's $H = 8.08$ km corresponds to a constant temperature of

$$T = \frac{gH}{R_d} = \frac{9.81 \times 8080}{287} = 276 \text{ K } (3\ ^\circ\text{C}),$$

a reasonable average for the lowest few kilometres. But above about 8 km the air is far colder (around 217 K, −56 °C, Chapter 5), its scale height is only about 6.3 km, and the pressure falls faster than the rule assumes. Cold air is denser at the same pressure, Eq. (3.1), so a given weight of it is packed into a thinner layer. Question 12 lets you measure this from the soundings (§ 8.1).

A question that comes up

The sheet says the *density* halves every 5.6 km. Is that the same as the pressure halving?
Only if the temperature does not change with height, because by Eq. (3.1) $\rho = p/(R_d T)$. In the real troposphere the temperature falls as you go up, so the density falls a little more slowly than the pressure (the cooler air is relatively denser). In the standard atmosphere the density at 5.6 km is 56 % of its sea-level value while the pressure is 49 %. The rule is about pressure, the weight of the air above; it is the right one to use for Questions 2 to 4.

A question that comes up

Where does the atmosphere end?
It thins out gradually and has no top. By convention the edge of space is put at 100 km, the Kármán line, where by the rule only about $2^{-100/5.6} \approx 0.0004$ % of the atmosphere remains (the real figure is smaller still). The sheet notes that there are very few gas molecules left beyond 100 km.

3.7 Try it yourself: Practice Set 3

Practice Set 3: pressure and height

Use the halving rule unless told otherwise; $g = 9.81\,\mathrm{m\,s^{-2}}$, $R_d = 287\,\mathrm{J\,kg^{-1}\,K^{-1}}$.

3.1 What percentage of the atmosphere is above you at 16.8 km, and what is the pressure there?

3.2 What fraction of the whole atmosphere lies between 5.6 and 11.2 km?

3.3 At what height does the pressure fall to 100 hPa?

3.4 What mass of air lies above each square metre at sea level, where the pressure is 1013 hPa?

3.5 What would the halving height be in an atmosphere at a constant 250 K?

Answers to Practice Set 3

3.1 Three halvings: 12.5 % and 125 hPa, Eqs. (3.5) and (3.6).

3.2 50 % is above 5.6 km and 25 % above 11.2 km, so $50 - 25 = 25$ % lies between them.

3.3 100 hPa is a tenth of 1000 hPa: $2^{-z/5.6} = 0.1$, so $z/5.6 = \log_2 10 = 3.32$ and $z = 18.6$ km, Eq. (3.4).

3.4 $101\,300/9.81 = 10\,330$ kg: about 10.3 tonnes.

3.5 $H = 287 \times 250/9.81 = 7.31$ km, so $z_{1/2} = 0.693 \times 7.31 = 5.07$ km. Colder air halves its pressure over a shorter distance.

Chapter 4

Reading the graph, and oxygen at altitude

This chapter answers Questions 3 and 4 of the sheet.

4.1 Reading values from a graph, and checking them

A graph turns a rule into a tool: once the curve of Figure 3.2 is drawn, any height can be looked up without calculation. To read a value:

Step 1. Find the height on the vertical axis (convert metres to kilometres first: 8848 m = 8.848 km).

Step 2. Draw a horizontal line across to the curve.

Step 3. From the meeting point, draw a vertical line down to the percentage axis and up to the pressure axis, and read both.

A hand-drawn curve and a ruler give two significant figures at best. The exact value comes from Eq. (3.4): divide the height by 5.6 km to get the number of halvings x, then $f = 2^{-x}$. On a calculator, 2^{-x} is entered as `2 ^ (-x)` or computed as $e^{-0.6931\,x}$.

Worked example: the summit of Everest, 8848 m

Number of halvings: $x = 8.848/5.6 = 1.580$. Fraction above: $f = 2^{-1.580} = 0.3345$, so **33.4 %** of the atmosphere is above the summit. Pressure: $1000 \times 0.3345 =$ **334 hPa**, Eq. (3.6). Graph check: 8.8 km lies between 5.6 km (50 %) and 11.2 km (25 %), closer to 11.2, so the answer must be between 25 and 50 %, nearer 25. It is.

4.2 Question 3: five heights

The question on the sheet

Using the graph you just drew, estimate the percentage of atmosphere above and the total pressure for the following heights: Portstewart strand (0 m); the top of Mount Everest (8848 m); the cruising altitude of a jumbo jet (12 km); Machu Picchu, Peru (2430 m); the world-record hot-air balloon (21 027 m).

Answer to Question 3

Portstewart strand: **100 %, 1000 hPa**. Everest: **33.4 %, 334 hPa**. Jumbo jet at 12 km: **22.6 %, 226 hPa**. Machu Picchu: **74.0 %, 740 hPa**. Record balloon: **7.4 %, 74 hPa**. (Values read from a hand-drawn graph within about ±2 percentage points are fine.)

Check against measurement. On the summit of Everest the barometric pressure has been measured: 253 mmHg, which is 337 hPa (West et al., 1983), within 1 % of the rule's 334 hPa. The standard atmosphere gives only 314 hPa there, because it describes a cooler, mid-latitude column;

Table 4.1. Question 3 by the halving rule, Eqs. (3.4) and (3.6). The last column gives the U.S. Standard Atmosphere for comparison (NOAA et al., 1976).

Place	Height (km)	Halvings x	% above	Pressure (hPa)	Std. atm. (hPa)
Portstewart strand	0	0	100	1000	1013
Machu Picchu	2.430	0.434	74.0	740	753
Everest summit	8.848	1.580	33.4	334	314
Jumbo jet	12.000	2.143	22.6	226	193
Record balloon	21.027	3.755	7.4	74	47




Figure 4.1. Question 3 read from the graph. For each place, go across from its height to the curve and down to the percentage axis; the pressure is ten times the percentage.

Everest stands at 28° N, where the warm air column holds the pressure up at height (West, 1996). For the jet and the balloon, high in the cold upper air, the rule overestimates the pressure, for the reason given in § 3.6.

4.3 What "21 % oxygen" means

Dry air is a mixture of gases: by volume about 78 % nitrogen, 21 % oxygen and 0.9 % argon, with carbon dioxide and other trace gases making up the rest (EGM305 module team, 2020, slide 6); precise values are 78.08, 20.95 and 0.93 % (Wallace and Hobbs, 2006, ch. 1). "By volume" means by number of molecules: of every hundred molecules of dry air, about 21 are oxygen. Up to about 80 km the air is stirred so thoroughly that these proportions stay the same at every height (EGM305 module team, 2020, slide 24). So the air on Everest is still 21 % oxygen. What changes is how many molecules there are in each breath.

4.4 Partial pressure

In a mixture of gases each gas contributes to the total pressure in proportion to its share of the molecules. The contribution of one gas is its **partial pressure**, and the partial pressures add up

to the total (Dalton's law). For oxygen

$$p_{O_2} = x_{O_2}\, p = 0.21\, p. \tag{4.1}$$

where

p_{O_2}	partial pressure of oxygen (hPa)	calculated
x_{O_2}	fraction of the molecules that are oxygen	0.21, on the sheet
p	total air pressure (hPa)	from Question 3

At sea level $p_{O_2} = 0.21 \times 1000 = 210\,\text{hPa}$ and $p_{N_2} = 0.78 \times 1000 = 780\,\text{hPa}$, as the sheet says. The partial pressure of oxygen is what matters to the body: it pushes oxygen from the lungs into the blood. Halve the total pressure and you halve the oxygen partial pressure, even though the air is still 21 % oxygen. The same idea of a partial pressure returns for water vapour in Practical 3, as the vapour pressure e (§ 21.2).

4.5 Question 4: oxygen at five heights

The question on the sheet

The partial pressure of each gas drops at the same rate as the total atmospheric pressure. Use your answers to Question 3 to first calculate the partial pressure of oxygen at different elevations. Then use your answer to calculate the percentage of oxygen available compared to the atmosphere at sea level.

Method.

Step 1. Partial pressure: multiply each pressure from Question 3 by 0.21, Eq. (4.1).

Step 2. Percentage: the second part of the question can be read in two ways, and the two readings give different numbers (Table 4.2).

- *(a) Oxygen as a percentage of the sea-level atmosphere:* divide the oxygen partial pressure by the sea-level *total* pressure, 1000 hPa. This is the reading used in the answer key.
- *(b) Oxygen available compared with sea level:* divide the oxygen partial pressure by the sea-level *oxygen* partial pressure, 210 hPa. This says how much oxygen each breath brings compared with a breath at the coast.

Table 4.2. Question 4. Partial pressure of oxygen, Eq. (4.1), and the two percentages.

Place	Pressure (hPa)	p_{O_2} (hPa)	(a) $p_{O_2}/1000$	(b) $p_{O_2}/210$
Portstewart strand	1000	210	21.0 %	100 %
Machu Picchu	740	155.5	15.5 %	74 %
Everest summit	334	70.2	7.0 %	33 %
Jumbo jet	226	47.6	4.8 %	23 %
Record balloon	74	15.6	1.6 %	7.4 %




Figure 4.2. Question 4: the partial pressure of oxygen at the five places. The air is 21 % oxygen at all of them; what falls is the number of molecules in each breath, in step with the total pressure (Table 4.2).

Answer to Question 4

Partial pressure of oxygen: Portstewart **210 hPa**; Machu Picchu **155 hPa**; Everest **70 hPa**; jumbo jet **48 hPa**; record balloon **16 hPa**. As a percentage of the sea-level atmosphere (the key's reading): 21, 15.5, 7.0, 4.8 and 1.6 %. Compared with the oxygen available at sea level: 100, 74, 33, 23 and 7 %. On Everest each breath brings only a third of the oxygen it brings at sea level.

A note on the printed key: two readings, and one rounding

The key gives the percentages of reading (a): 21, 7, 4.7, 15.5 and 1.6 %. Reading (b) is the one that answers "how much oxygen is available compared with sea level" and is the more useful for explaining breathlessness, so both are given above. Notice that reading (b) is simply the percentage of the atmosphere above you from Question 3: since the oxygen fraction does not change with height, the available oxygen falls in exactly the same proportion as the total pressure. The key's 4.7 % for the jet comes from rounding 47.5 down; $0.21 \times 226.4 = 47.6$ hPa, which is 4.8 %.

4.6 Why it matters

- **Machu Picchu** (about three-quarters of sea-level oxygen): most visitors notice breathlessness on climbing stairs; the body adjusts within days by breathing faster and making more red blood cells.
- **Everest** (a third): close to the limit of human survival without bottled oxygen. Climbers acclimatise for weeks, and most use supplementary oxygen near the summit (West et al., 1983).
- **A jet at 12 km** (under a quarter): a person would lose consciousness within minutes. Airliner cabins are therefore pressurised, to roughly the pressure found at about 2 km, and oxygen masks drop if the pressure is lost.
- **The record balloon at 21 km** (7 %): no human could breathe this air; balloonists wear pressure suits or ride in sealed gondolas.

A question that comes up

If the air is still 21 % oxygen at the top of Everest, why is it harder to breathe? Because "21 %" is a proportion, not an amount. At the summit each lungful of air contains only a third as many molecules of every gas as at sea level, so it delivers a third as much oxygen. The lungs move oxygen into the blood in proportion to its partial pressure, 70 hPa instead of 210.

4.7 Try it yourself: Practice Set 4

Practice Set 4: reading heights and partial pressures

Use the halving rule and an oxygen fraction of 0.21.

4.1 Ben Nevis is 1345 m high. What are the pressure and the oxygen partial pressure at the summit?

4.2 Kilimanjaro is 5895 m high. Find the pressure, the oxygen partial pressure, and the oxygen available compared with sea level.

4.3 On a summit the pressure is 600 hPa. What percentage of the air molecules there are oxygen? What is the oxygen partial pressure? At what height is the summit, by the rule?

4.4 What is the partial pressure of nitrogen on the summit of Everest?

Answers to Practice Set 4

4.1 $x = 1.345/5.6 = 0.240$; $f = 2^{-0.240} = 0.847$; pressure 847 hPa; $p_{O_2} = 0.21 \times 847 = 178$ hPa.

4.2 $x = 5.895/5.6 = 1.053$; $f = 0.482$; pressure 482 hPa; $p_{O_2} = 101$ hPa; 48 % of the sea-level oxygen.

4.3 Still 21 % (the proportion does not change). $p_{O_2} = 0.21 \times 600 = 126$ hPa. Height: $2^{-z/5.6} = 0.6$, so $z = 5.6 \times \log_2(1/0.6) = 5.6 \times 0.737 = 4.1$ km.

4.4 $0.78 \times 334 = 261$ hPa.

Chapter 5

Temperature with height: the standard atmosphere

This chapter answers Question 5 of the sheet and introduces the layers of the atmosphere used in Questions 6 to 11.

5.1 Temperature and the lapse rate

Temperature measures how fast the molecules of the air are moving: the warmer the air, the faster they move (EGM305 module team, 2020, slide 16). Unlike pressure, temperature does not fall steadily all the way up. It falls, then rises, then falls and rises again, and those reversals divide the atmosphere into layers (§ 5.4).

The rate at which the temperature falls with height is the **lapse rate**:

$$\Gamma = -\frac{\Delta T}{\Delta z} = \frac{T_{\text{lower}} - T_{\text{upper}}}{z_{\text{upper}} - z_{\text{lower}}}. \tag{5.1}$$

where

Γ	lapse rate (°C km^{-1}, the same as K km^{-1})	calculated
ΔT	change in temperature going up (°C)	from a table or graph
Δz	change in height (km)	from a table or graph

The minus sign makes the lapse rate *positive* when the temperature falls with height, which is the usual case in the lowest layer. A negative lapse rate means the temperature *rises* with height: an **inversion** (§ 6.3). A lapse rate of zero means the temperature does not change: an **isothermal** layer.

5.2 The standard atmosphere

Real temperature profiles change from day to day and from place to place. For aircraft instruments, engineering and comparison, meteorologists use an agreed average: the **standard atmosphere**. Near the ground it is defined by (NOAA et al., 1976):

$$T(z) = \begin{cases} 15 - 6.5\,z & 0 \le z \le 11 \text{ km}, \\ -56.5 & 11 \le z \le 20 \text{ km}, \end{cases} \tag{5.2}$$

where

$T(z)$	temperature at height z (°C)	calculated
z	height above sea level (km)	
15	sea-level temperature (°C)	288.15 K
6.5	the standard (“normal”) lapse rate (°C km^{-1})	on the sheet

At sea level the standard pressure is 1013.25 hPa and the standard temperature 15 °C. The temperature falls by 6.5 °C for every kilometre up to 11 km, the standard **tropopause**, where it reaches −56.5 °C, and then stays constant.

5.3 Question 5: Table 2

The question on the sheet

In the troposphere temperature decreases by 6.5 °C per km in the lowest 11 km; this is called the average or normal lapse rate. Based on this average lapse rate, complete the drop in temperature for the "standard atmosphere" in Table 2.

Method. Use Eq. (5.2). For each height up to 11 km, convert to kilometres and subtract $6.5 \times z$ from 15 °C. Above 11 km the temperature stays at −56.5 °C.

$$T(2\text{ km}) = 15 - 6.5 \times 2 = 15 - 13 = 2\ ^\circ\text{C}, \qquad T(8\text{ km}) = 15 - 6.5 \times 8 = 15 - 52 = -37\ ^\circ\text{C}.$$

Table 5.1. Table 2 of the sheet, completed. Values in bold are those to be filled in; the others are printed on the sheet.

Height (m)	0	1000	2000	4000	6000	8000	10 000	11 000	14 000	16 000
T (°C)	15	8.5	**2**	**−11**	**−24**	**−37**	**−50**	−56.5	−56.5	−56.5

Answer to Question 5

At 2000, 4000, 6000, 8000 and 10 000 m the standard temperatures are **2, −11, −24, −37 and −50** °C. At 11 000 m the temperature reaches −56.5 °C and stays there to 16 000 m (Table 5.1).

Check. The printed values agree with the rule: $15 - 6.5 \times 1 = 8.5$ °C at 1000 m and $15 - 6.5 \times 11 = -56.5$ °C at 11 000 m. Each 2 km step lowers the temperature by 13 °C.

5.4 The layers of the atmosphere

Figure 5.1 extends the standard atmosphere to 100 km. The turning points of the temperature (the *pauses*) divide it into four layers (EGM305 module team, 2020, slides 16–22).

Troposphere (surface to about 11 km). Temperature falls with height, on average by 6.5 °C per km. The air is heated mainly from below: sunlight passes largely through it and warms the ground, which then warms the air (Part II, § 17.7). Rising air also cools as it expands (Chapter 8). About three-quarters of the air and nearly all the water vapour are here, and almost all weather happens here.

Stratosphere (about 11 to 50 km). Temperature is constant at first, then rises with height. The warming comes from ozone, which absorbs the Sun's ultraviolet radiation (Part II, § 12.4). Warm air sitting on cooler air is very stable, so the stratosphere is calm and layered.

Mesosphere (about 50 to 85 km). Temperature falls again, to the coldest values in the atmosphere, below −85 °C at the mesopause.

Thermosphere (above about 90 km). Temperature rises steeply, as the few molecules there absorb the most energetic solar radiation (EGM305 module team, 2020, slide 21).




Figure 5.1. The U.S. Standard Atmosphere from the ground to 100 km (NOAA et al., 1976). Temperature falls through the troposphere, is constant and then rises through the stratosphere, falls again through the mesosphere, and rises in the thermosphere. The shaded bands are the layers; the dashed lines are the pauses between them.

A question that comes up

Is the thermosphere really hot?
Its molecules move very fast, which is what a high temperature means, and in sunlight the upper thermosphere reaches several hundred degrees. But they are so few that they would hardly heat anything: an unprotected thermometer there would lose more by radiating than it gained from the rare collisions, and would read very cold. Temperature and heat are not the same thing.

A question that comes up

Why does the troposphere get colder going up, when it is closer to the Sun?
The extra closeness is negligible (a few kilometres in 150 million). What matters is where the heating happens. Sunlight passes mostly through the air and is absorbed by the ground, so the troposphere is heated from below, like a pan of water on a hob, and is warmest at the bottom.

5.5 Try it yourself: Practice Set 5

Practice Set 5: lapse rates and the standard atmosphere

5.1 The temperature falls from 20 °C at the ground to −10 °C at 5 km. What is the lapse rate?

5.2 What is the standard-atmosphere temperature at 3 km? At the summit of Everest (8848 m)?

5.3 At what height does the standard atmosphere reach 0 °C?

5.4 Convert −56.5 °C to kelvin.

5.5 Over a layer from 1.2 to 1.6 km the temperature rises from −2 to 0 °C. What is its lapse rate, and what is such a layer called?

Answers to Practice Set 5

5.1 $\Gamma = (20 - (-10))/5 = 6\,^\circ\mathrm{C\,km^{-1}}$, Eq. (5.1).

5.2 $15 - 6.5 \times 3 = -4.5\,^\circ\mathrm{C}$; $15 - 6.5 \times 8.848 = -42.5\,^\circ\mathrm{C}$, Eq. (5.2).

5.3 $15 - 6.5\,z = 0$ gives $z = 15/6.5 = 2.31$ km.

5.4 $-56.5 + 273.15 = 216.65$ K.

5.5 $\Gamma = (-2 - 0)/(1.6 - 1.2) = -5\,^\circ\mathrm{C\,km^{-1}}$: negative, so the temperature rises with height. It is an inversion.

Chapter 6

Two soundings: inversions and the tropopause

This chapter answers Questions 6, 7 and 8 of the sheet.

6.1 Measuring the atmosphere in the vertical

A **radiosonde** is a small instrument package carried up by a helium or hydrogen balloon. As it rises at about $5\,\mathrm{m\,s^{-1}}$ it measures pressure, temperature and humidity and sends them to the ground by radio; tracking its drift gives the wind. The balloon expands as the pressure falls and bursts at around 30 km, which is why the soundings on the sheet stop near that height. Radiosondes are launched at hundreds of stations worldwide, at the same times each day, so that the measurements can be compared and fed into weather forecasts (World Meteorological Organization, 2023). The record of one ascent is called a **sounding**.

6.2 Question 6: plotting the soundings

The question on the sheet

Table 3 gives the pressure, height and temperature from a radiosonde measurement taken on 21 March 1990 for Hillsborough (Northern Ireland) and St John's, Newfoundland (Canada). Plot the temperature values against height for these stations and for the standard atmosphere (Table 2) on the graph on the final page, using different colours for the different plots.

Method.

Step 1. **Axes.** Height (km) up the side, from 0 to 31 km; temperature (°C) along the bottom, from −80 to +30 °C. Height goes on the vertical axis even though it is the "input": a profile drawn this way looks like the atmosphere it describes, with the ground at the bottom.

Step 2. **Convert heights to kilometres** (divide the metres by 1000): 1330 m is 1.33 km.

Step 3. **Plot each sounding with its own colour and symbol** (Table A.1 in Appendix A.3 lists the values), and the standard atmosphere from Table 5.1.

Step 4. **Join the points of each sounding with straight lines.** A sounding is a set of measurements at separate levels; straight lines are the honest way to connect them.

Step 5. **Label** each line and give the graph a title and a key.




Figure 6.1. Questions 6 to 8: the two soundings of Table 3 and the standard atmosphere (Table 2), with the inversions in the troposphere circled (Question 7) and the tropopause of each profile marked (Question 8).

Answer to Question 6

The completed graph is Figure 6.1. Both soundings cool with height up to a turning point near 9 to 11 km, above which the temperature stops falling and stays between about −50 and −62 °C to 30 km. Near the ground Hillsborough is 12.7 °C warmer than St John's, but the two profiles cross at about 2 km, and between 3 and 7 km St John's is the warmer, by about 7 °C: the contrast at the surface is confined to the lowest layer (Question 14, Chapter 9).

6.3 Inversions

An **inversion** is a layer in which the temperature *rises* with height: the lapse rate, Eq. (5.1), is negative. It is called an inversion because it inverts the normal pattern of the troposphere. Inversions form in three common ways (Ahrens and Henson, 2022, ch. 3):

- **Radiation inversions:** on clear, calm nights the ground cools by radiating to the sky (Part II, § 17.8), and the air next to it cools most.
- **Subsidence inversions:** in an anticyclone, air sinking from above is compressed and warms, forming a warm layer above cooler air near the ground.
- **Frontal inversions:** warm air rides up over a wedge of colder air at a front (Part III,

Chapter 29).

Inversions matter because they are very stable: they suppress rising air, trap pollution and fog beneath them, and cap the growth of clouds (Chapter 8).

6.4 Question 7: finding the inversions

The question on the sheet

You will see that in some situations the temperature increases with height in the troposphere. Such layers are called inversion layers. Circle these on the graph.

Method. Go up each sounding below its tropopause and look for any step in which the temperature at the higher level is warmer than at the level below. Table A.3 in Appendix A.3 lists the lapse rate of every layer of both soundings, Eq. (5.1).

- **Hillsborough:** from 1358 m (−1.25 °C) to 1503 m (−1.05 °C) the temperature rises by 0.2 °C: a weak inversion, sitting on top of an isothermal step from 1330 to 1358 m. Below it the air cools fast, by 9.5 °C per km.
- **St John's:** no level in the troposphere is warmer than the one below, but from 140 m to 3006 m the temperature falls by only 1.2 °C in almost 3 km (0.3 to 0.7 °C per km). This deep, nearly isothermal layer is very stable and behaves like an inversion.

Answer to Question 7

Hillsborough: an inversion between about **1360 and 1500 m** (−1.25 to −1.05 °C), just above the steep cooling of the lowest 1.3 km. **St John's:** no true inversion in the troposphere, but a deep, almost **isothermal layer from the ground to about 3 km**. Both are circled in Figure 6.1.

A question that comes up

Above the tropopause the temperature rises with height too. Are those inversions? Strictly, yes: Hillsborough warms from −51 to −45 °C between 9.4 and 11.1 km, and St John's from −62 to −55 °C between 11.5 and 12.1 km. But they are in the stratosphere, where warming with height is the normal state (Figure 5.1). The question asks only about the troposphere.

6.5 The tropopause

The sheet defines the **tropopause** as the top of the troposphere, where the temperature ceases to decrease with height. The World Meteorological Organization makes this precise: it is the lowest level at which the lapse rate falls to 2 °C km^{-1} or less, provided the average lapse rate over the 2 km above does not exceed 2 °C km^{-1} (Reichler et al., 2003). The second condition stops a thin inversion in the troposphere, like the one at 1.4 km over Hillsborough, from being mistaken for the tropopause. Above the tropopause the stable stratosphere begins; weather, and the tops of most clouds, stop below it.

6.6 Question 8: marking the tropopause

The question on the sheet

The tropopause is defined as the top boundary of the troposphere where the temperature ceases to decrease with height. Mark this boundary on the graph and note the temperature and the height for the standard atmosphere, Hillsborough and St John's.

Method. For each profile, follow the temperature up through the troposphere and find the level above which it no longer falls. Check the WMO condition: above that level the temperature must not keep falling by more than 2 °C per km over the next 2 km.

- **Standard atmosphere:** by definition, 11 000 m and −56.5 °C (Table 5.1).
- **Hillsborough:** the temperature falls to −51.05 °C at 9383 m and then *rises* to −49.05 °C at 10 164 m and −44.65 °C at 11 116 m.
- **St John's:** the temperature falls to −62.05 °C at 11 482 m and then rises to −58.05 °C at 11 843 m.

Answer to Question 8

Standard atmosphere: **11 000 m, −56.5 °C.** Hillsborough: **9383 m, −51.05 °C.** St John's: **11 482 m, −62.05 °C.** Marked in Figure 6.1.

Check. All three lie between 9 and 12 km, the usual range for the middle latitudes (Hoinka, 1998). The higher tropopause (St John's) is also the colder one: a tropopause that sits higher has had more kilometres of cooling below it. Chapter 7 uses this to explain why the tropical tropopause is both the highest and the coldest of all.

6.7 Try it yourself: Practice Set 6

Practice Set 6: reading soundings

Use the soundings of Table A.1.

6.1 What is the lapse rate over Hillsborough between 38 m and 1330 m? How does it compare with the standard 6.5 °C km^{-1}?

6.2 Is the layer over St John's from 9103 to 10 446 m an inversion? What is its lapse rate?

6.3 At what pressure is the tropopause over Hillsborough? Over St John's?

6.4 A sounding shows −40 °C at 9 km, −55 °C at 11 km, −54 °C at 12 km and −53 °C at 13 km. Where is the tropopause?

Answers to Practice Set 6

6.1 $(11.05 - (-1.25))/(1.330 - 0.038) = 12.30/1.292 = 9.5$ °C km^{-1}: much steeper than the standard rate, and close to the rate at which rising dry air cools (§ 8.3).

6.2 No: the temperature falls from −41.05 to −53.45 °C, a lapse rate of $12.4/1.343 = 9.2$ °C km^{-1}.

6.3 Hillsborough 282 hPa; St John's 212 hPa (Table A.1).

6.4 At 11 km (−55 °C): above it the temperature rises, so the lapse rate is negative, below 2 °C km^{-1}, over the 2 km above.

Chapter 7

Lapse rates and the height of the troposphere

This chapter answers Questions 9, 10 and 11 of the sheet.

7.1 Average and local lapse rates

The lapse rate of Eq. (5.1) can be taken over any layer. Taken between the ground and the tropopause it gives the **average lapse rate** of the troposphere, one number for the whole layer. Taken between neighbouring levels of a sounding it gives the **local lapse rate**, which varies a great deal: in the Hillsborough sounding it ranges from $-1.4\,^\circ\mathrm{C\,km^{-1}}$ (the inversion at 1.4 km) to $+9.5\,^\circ\mathrm{C\,km^{-1}}$ (the lowest 1.3 km), as Table A.3 shows. The standard $6.5\,^\circ\mathrm{C\,km^{-1}}$ is an average of averages: over many places and many days.

7.2 Question 9: the lapse rate over Hillsborough

The question on the sheet

Based on your graph and Table 3a, calculate the lapse rate for the part of the atmosphere below the tropopause for Hillsborough (have another look at Question 5 to work out how to calculate the lapse rate).

Method. Take the lowest level of the sounding and the tropopause found in Question 8 (§ 6.6), and apply Eq. (5.1).

Step 1. Lowest level: 38 m, 11.05 °C. Tropopause: 9383 m, −51.05 °C.

Step 2. Fall in temperature: $11.05 - (-51.05) = 62.10\,^\circ\mathrm{C}$.

Step 3. Rise in height: $9383 - 38 = 9345\,\mathrm{m} = 9.345\,\mathrm{km}$.

Step 4. Lapse rate: $\Gamma = 62.10/9.345 = 6.65\,^\circ\mathrm{C\,km^{-1}}$.

Answer to Question 9

$\Gamma = \dfrac{11.05 - (-51.05)}{9.383 - 0.038} = \dfrac{62.10\ ^\circ\mathrm{C}}{9.345\ \mathrm{km}} = \mathbf{6.65}\,^\circ\mathrm{C}$ **per km**, very close to the standard lapse rate of 6.5 °C per km.

Check. The value lies between the extremes of the local lapse rates in Table A.3, and the straight line from the surface point to the tropopause point in Figure 6.1 runs close to the dashed standard-atmosphere line. For comparison, the same calculation for St John's gives $(-1.65 - (-62.05))/(11.482 - 0.140) = 5.33\,^\circ\mathrm{C\,km^{-1}}$: smaller, because its surface layer is so cold.

A gap in the printed key

The answer key leaves Question 9 blank. The answer above, 6.65 °C per km, follows from the data and the definition on the sheet.

7.3 Question 10: a tropical profile

The question on the sheet

Add the temperature curve for Florida to your graph (Table 4). Where does the tropopause occur?




Figure 7.1. Question 10: the Florida profile (Table 4) added to the graph, shown here to 20 km. Florida is warmest at the ground and coldest at its tropopause, which is 5 to 7 km higher than the others.

Method. Plot Table 4 on the same axes (Figure 7.1) and look for the level above which the temperature stops falling: it falls all the way to −75.6 °C at 16 000 m and then rises to −73.3 °C at 18 000 m.

Answer to Question 10

Florida: the tropopause is at about **16 000 m**, at **−75.6** °C. Its average lapse rate below the tropopause is $(27.2 - (-75.6))/16 = 6.4\,^{\circ}\mathrm{C\,km^{-1}}$.

A correction to the printed key: -75.6, not -75.4 °C

The key gives the Florida tropopause temperature as -75.4 °C. Table 4 prints -75.6 °C at 16 000 m, the coldest value of the profile; -75.4 is a typing slip.

Check. The data are only every 2 km, so the true tropopause could be anywhere between about 15 and 17 km. That range is typical of the tropics and subtropics (Hoinka, 1998).

7.4 Question 11: is the troposphere the same thickness everywhere?

The question on the sheet

Examining all your graphs, do you think the troposphere is the same thickness across the globe? Where do you think it is thickest and why is that?

The evidence. The four tropopauses are at 9.4 km (Hillsborough, 54° N), 11.0 km (the standard atmosphere, which represents the middle latitudes), 11.5 km (St John's, 48° N) and 16 km (Florida, about 25 to 31° N). The further from the pole, the higher the tropopause (Figure 7.2).




Figure 7.2. Question 11: the tropopause heights of the four profiles against latitude. The grey band shows the typical pattern (Hoinka, 1998; Wallace and Hobbs, 2006): about 16 to 17 km over the tropics, falling steeply across the subtropics and middle latitudes, to about 8 to 9 km near the poles. Florida is plotted at its middle latitude, 28° N; the standard atmosphere at 45° N.

The reasons. Three effects work together.

1. **More heating at the surface.** The tropics receive the most solar energy per square metre, because the Sun is high in the sky all year (Part II, Chapter 18). The ground and the air above it are warm.
2. **Warm air columns are taller.** Warm air is less dense than cold air at the same pressure, Eq. (3.1), so the same weight of air (the same drop in pressure) occupies a greater depth: the scale height $H = R_d T/g$ is larger, Eq. (3.3). Every pressure level, and with them the whole troposphere, sits higher above a warm surface. The same effect makes pressure surfaces slope from the warm tropics down to the cold poles, which drives the winds of Practical 3 (Part III, Chapter 24).
3. **Deep convection.** Strong heating makes air rise in towering thunderstorm clouds that carry heat and mix the troposphere up to 15 to 18 km. The tropopause sits where this stirring

from below gives way to the stable, radiatively heated stratosphere, so where convection reaches higher, the tropopause is higher.

A taller troposphere also has a *colder* top: with a similar lapse rate, the temperature keeps falling over more kilometres. That is why Florida, the warmest at the ground (27 °C), has the coldest tropopause (−75.6 °C).

Answer to Question 11

No. The troposphere is **thickest over the tropics** (about 16 to 17 km; 16 km in the Florida profile) and **thinnest near the poles** (about 8 to 9 km), with middle latitudes in between (9.4 to 11.5 km in the soundings). The tropics receive the most solar heating; warm air is less dense, so a warm air column is taller; and strong convection mixes warm air to great heights before the stable stratosphere begins. The tropopause is also higher in summer than in winter, for the same reasons.

A question that comes up

The coldest air in the troposphere is over the equator? That sounds backwards. At the ground the tropics are the warmest places on Earth. But high up, at 16 to 17 km, the tropical tropopause is colder than the tropopause anywhere else, often below −75 °C, because it is so far above the surface. Near the poles the tropopause is low and so less cold: around −50 to −60 °C.

7.5 Try it yourself: Practice Set 7

Practice Set 7: lapse rates

7.1 What is the lapse rate in the lowest 2 km of the Florida profile? Between 10 and 12 km?

7.2 A tropical sounding has 28 °C at the ground and an average lapse rate of 6.5 °C km^{-1} up to its tropopause at 16 km. What is the tropopause temperature?

7.3 A polar sounding has −20 °C at the ground and the tropopause at 8.5 km with −55 °C. What is its average lapse rate?

7.4 By how much is the scale height over Florida (mean tropospheric temperature about −24 °C) larger than over a polar station (about −40 °C)?

Answers to Practice Set 7

7.1 $(27.2 - 15.2)/2 = 6.0$ °C km^{-1}; $(-32.6 - (-50.5))/2 = 8.95$ °C km^{-1}.

7.2 $28 - 6.5 \times 16 = 28 - 104 = -76$ °C.

7.3 $(-20 - (-55))/8.5 = 35/8.5 = 4.1$ °C km^{-1}.

7.4 $H = R_d T/g$: $287 \times 249/9.81 = 7.29$ km against $287 \times 233/9.81 = 6.82$ km, about 0.47 km (7 %) larger. Every pressure level therefore sits about 7 % higher above the ground in the warm column: 500 hPa, for example, at about 5.1 km instead of 4.7 km.

Chapter 8

Pressure aloft, buoyancy and stability

This chapter answers Questions 12 and 13 of the sheet.

8.1 Question 12: equal steps, unequal drops

The question on the sheet

In St John's, how much is the pressure drop between 140 m and 3006 m? How much is the pressure drop between 23 834 m and 26 404 m? Seeing that the difference in altitude is almost the same, why is the difference in pressure so dissimilar?

The two drops. From Table A.1 (St John's):

$$\begin{aligned} \text{lower layer:} &\quad 1005 - 700 = 305 \text{ hPa} \quad \text{over} \quad 3006 - 140 = 2866 \text{ m}, \\ \text{upper layer:} &\quad 30 - 20 = 10 \text{ hPa} \quad \text{over} \quad 26\,404 - 23\,834 = 2570 \text{ m}. \end{aligned}$$

Look at fractions, not amounts. In the lower layer the pressure falls to $700/1005 = 0.70$ of its starting value, a fall of 30 %; in the upper layer it falls to $20/30 = 0.67$, a fall of 33 %. The *fractional* drops over these similar depths are almost the same. That is exactly what exponential decay means (§ 3.3): each kilometre removes the same *fraction* of the air above, so where there is a lot of air (near the ground) the same fraction is a large amount, and where there is little (at 24 km, only 3 % of the atmosphere is above) it is a tiny amount. Plotted on a logarithmic pressure axis, an exponential becomes a straight line (Figure 8.1).

The data even measure the scale height. Rearranging Eq. (3.3) for two levels gives the scale height of the layer between them (Appendix A.2.1):

$$H = \frac{z_2 - z_1}{\ln(p_1/p_2)}. \tag{8.1}$$

where

H	scale height of the layer (km)	calculated
$z_1,\ z_2$	heights of the bottom and top of the layer (km)	from the sounding
$p_1,\ p_2$	pressures at those heights (hPa)	from the sounding

For the lower layer $H = 2.866/\ln(1005/700) = 2.866/0.362 = 7.9$ km; for the upper layer $H = 2.570/\ln(30/20) = 2.570/0.405 = 6.3$ km. By Eq. (3.3), $H = R_d T/g$, so these correspond to temperatures of $gH/R_d = 271$ K (-2 °C) and 218 K (-55 °C): exactly the temperatures measured in those layers. The upper layer is colder, so its air is denser for its pressure and its pressure falls off over a shorter distance.




Figure 8.1. Question 12: the St John’s sounding with pressure on a logarithmic axis (decreasing to the right, so that height and falling pressure go the same way). An exponential decay is a straight line here; the slope is steeper (shorter scale height) where the air is colder.

Answer to Question 12

Between 140 and 3006 m the pressure drops by **305 hPa**; between 23 834 and 26 404 m by only **10 hPa**. Pressure decreases **exponentially** with height: each step up removes about the same *fraction* of the air above (30 % and 33 % here), not the same amount. Near the ground there is a great deal of air above, compressed by its own weight, so the same fraction is a large drop; at 24 km only about 3 % of the atmosphere remains above, so the drop is tiny.

8.2 “Cold air sinks”? Density depends on pressure too

Everyday experience says cold air sinks: cold draughts collect at the floor, frost forms in hollows. That is true when cold and warm air are side by side at the *same pressure*, because by the gas law, Eq. (3.1), $\rho = p/(R_d T)$ and lower T means higher ρ. But in the vertical the pressure changes enormously, and pressure wins. Using the Hillsborough sounding:

$$\text{surface } (1001\,\text{hPa},\ 11.05\,^\circ\text{C}): \quad \rho = \frac{100\,100}{287 \times 284.2} = 1.23\ \text{kg m}^{-3},$$

$$\text{tropopause } (282\,\text{hPa},\ -51.05\,^\circ\text{C}): \quad \rho = \frac{28\,200}{287 \times 222.1} = 0.44\ \text{kg m}^{-3}.$$

The cold air at the tropopause is only a third as dense as the warm air at the ground: it is lighter, not heavier, and it does not “fall through” the air below it.

8.3 What happens to air that moves up or down

The full answer needs one more idea. Air that moves vertically changes its pressure, and so its temperature. A parcel of air brought *down* is squeezed by the rising pressure, and the work done in compressing it warms it; a parcel that rises expands and cools. If no heat is exchanged with its

surroundings (the process is *adiabatic*), dry air warms or cools at a fixed rate, the **dry adiabatic lapse rate** (Wallace and Hobbs, 2006, ch. 3):

$$\Gamma_d = \frac{g}{c_p} = \frac{9.81}{1004}\ \mathrm{K\,m^{-1}} = 9.8\ ^\circ\mathrm{C\,km^{-1}}. \tag{8.2}$$

where

Γ_d	dry adiabatic lapse rate ($^\circ$C km^{-1})	9.8
g	acceleration due to gravity (m s^{-2})	9.81
c_p	specific heat of dry air at constant pressure (J kg^{-1} K^{-1})	1004, Table D.1

The derivation is in Appendix A.2.2. The same rate decides the height of cloud base in Practical 3: rising air cools at 9.8 °C km^{-1} until it reaches its dew point (Part III, § 22.8).

A convenient way to compare air at different heights is to ask what temperature each parcel would have if brought down adiabatically to the standard level of 1000 hPa. That temperature is the **potential temperature**:

$$\theta = T\left(\frac{1000\ \mathrm{hPa}}{p}\right)^{R_d/c_p} = T\left(\frac{1000}{p}\right)^{0.286}. \tag{8.3}$$

where

θ	potential temperature (K)	calculated
T	temperature of the parcel (K)	from the sounding
p	pressure of the parcel (hPa)	from the sounding
R_d/c_p	ratio of the gas constant to the specific heat	$287/1004 = 0.286$

Potential temperature does not change when air moves up or down adiabatically, so it labels a parcel the way a name does: two parcels with the same θ would have the same temperature if brought to the same level.




Figure 8.2. Question 13: potential temperature, Eq. (8.3), for both soundings up to 16 km. It increases with height everywhere, which means the atmosphere is stable: any parcel moved down arrives warmer, and so lighter, than the air around it. The increase is slow in the troposphere and fast in the stratosphere.

8.4 Question 13: why the cold air aloft does not sink

The question on the sheet

Thinking of your previous answer, if cold air is supposed to sink, why is the colder air of the upper troposphere not flowing down to the surface?

Follow a parcel down. Take the air at the Hillsborough tropopause, $-51.05\,^{\circ}\mathrm{C}$ (222.1 K) at 282 hPa, and bring it down to the ground. As it sinks it is compressed and warms at $9.8\,^{\circ}\mathrm{C\,km^{-1}}$. Its potential temperature, Eq. (8.3), tells us where it would end up:

$$\theta = 222.1 \times \left(\frac{1000}{282}\right)^{0.286} = 222.1 \times 1.436 = 318.9\ \mathrm{K} = 45.8\ ^{\circ}\mathrm{C}.$$

It would arrive at the surface at about 46 °C, some 35 °C *warmer* than the 11 °C air already there. Warmer air at the same pressure is less dense, so the parcel would be buoyant and would be pushed straight back up. The cold air aloft is not "waiting to fall": it is held up because it would become warm, and light, on the way down.

This is general. Whenever the temperature in the atmosphere falls with height more slowly than $9.8\,^{\circ}\mathrm{C\,km^{-1}}$ (the standard $6.5\,^{\circ}\mathrm{C\,km^{-1}}$, or the $6.65\,^{\circ}\mathrm{C\,km^{-1}}$ of Question 9), the potential temperature increases with height (Figure 8.2) and the air is **stably stratified**: air displaced downwards comes back up, and air displaced upwards sinks back. The atmosphere is layered like oil on water, but the layers are sorted by potential temperature, not by temperature.

Answer to Question 13

The upper air is cold but it is at very low pressure, so it is **much less dense** than the air near the ground (0.44 against $1.23\,\mathrm{kg\,m^{-3}}$ at Hillsborough): "cold air sinks" compares air at the same pressure. If the upper air did sink, it would be **compressed and warm up** at about $9.8\,^{\circ}\mathrm{C}$ per km; the tropopause air over Hillsborough would reach the ground at about **46** °C, far warmer and lighter than the surface air, and would rise again. Because the temperature falls with height more slowly than air warms on descent, the atmosphere is **stably layered**: its potential temperature increases with height.

Sharpening the printed key for Questions 12 and 13

The key's explanation of Question 12 (exponential decay, compressibility) is right; the fractional drops and scale heights above add the evidence from the data. For Question 13 the key says that the temperature decreases at a constant rate while pressure and density decay exponentially, so that the cold air aloft "floats" on the surface air like oil on water. The comparison of densities is right, but the reason the layering is *stable* is the warming of sinking air by compression: without it, Question 13 has no complete answer. Use the potential temperature argument above.

A question that comes up

Then when does air overturn?
When the temperature falls with height *faster* than $9.8\,^{\circ}\mathrm{C\,km^{-1}}$, a rising parcel stays warmer than its surroundings and keeps rising: the air is unstable, and convection overturns it. This happens over ground heated strongly by the Sun, where the lowest layer can exceed the dry adiabatic rate. In the Hillsborough sounding the lowest 1.3 km cools at $9.5\,^{\circ}\mathrm{C\,km^{-1}}$, almost exactly the dry adiabatic rate: a well-mixed layer stirred by the wind and by heating from the ground.

8.5 Try it yourself: Practice Set 8

Practice Set 8: pressure and stability

Use $R_d = 287\,\mathrm{J\,kg^{-1}\,K^{-1}}$, $c_p = 1004\,\mathrm{J\,kg^{-1}\,K^{-1}}$, $g = 9.81\,\mathrm{m\,s^{-2}}$ and kelvin $=^\circ\mathrm{C} + 273.15$.

8.1 What is the density of air at 500 hPa and $-20\,^\circ$C?

8.2 What is the potential temperature of the tropopause air over St John's (212 hPa, $-62.05\,^\circ$C)?

8.3 From the Hillsborough sounding, find the scale height of the layer from 1001 hPa (38 m) to 500 hPa (5414 m), and the temperature it corresponds to.

8.4 Air at 5414 m over Hillsborough ($-22.45\,^\circ$C) is brought down to the ground (38 m) without exchanging heat. Roughly what temperature would it have?

Answers to Practice Set 8

8.1 $\rho = 50\,000/(287 \times 253.15) = 0.688\,\mathrm{kg\,m^{-3}}$, Eq. (3.1): just over half the surface density.

8.2 $\theta = 211.1 \times (1000/212)^{0.286} = 328.8\,\mathrm{K}$ (55.7 °C), Eq. (8.3).

8.3 $H = 5.376/\ln(1001/500) = 5.376/0.694 = 7.74\,\mathrm{km}$, Eq. (8.1); $T = gH/R_d = 9.81 \times 7744/287 = 265\,\mathrm{K}$ ($-8\,^\circ$C), close to the average temperature of that layer.

8.4 It warms by about $9.8 \times 5.4 = 53\,^\circ$C, to about 30 °C, Eq. (8.2): 19 °C warmer than the surface air, so it would rise again.

Chapter 9

Two soundings, two climates

This chapter answers Question 14 of the sheet.

9.1 The puzzle

On 21 March 1990 the air near the ground was at 11.05 °C over Hillsborough and −1.65 °C over St John's (Table A.1). Hillsborough is at about 54.4° N; St John's at about 47.6° N, about 750 km closer to the equator. If latitude alone set the temperature, St John's should be the warmer.

The Sun does not explain it either. The soundings were made at the March equinox, when the noon Sun is overhead at the equator and its angle above the horizon is 90° minus the latitude (Part II, § 18.3). By Eq. (18.3), the noon solar intensity is $100\,\% \times \sin\beta$:

$$\text{Hillsborough: } 100\,\% \times \sin(35.6^\circ) = 58\,\%; \qquad \text{St John's: } 100\,\% \times \sin(42.4^\circ) = 67\,\%.$$

St John's received *more* solar energy per square metre at noon, yet was 12.7 °C colder. Something other than sunshine controls the temperature of the air near the ground.

9.2 What the soundings show

The difference is confined to the lowest layer. The two profiles cross at about 2 km, and from 3 to 7 km St John's is the warmer by about 7 °C (Figure 6.1). Over St John's the lowest 3 km is cold and almost isothermal (Question 7, § 6.4); over Hillsborough the lowest 1.3 km is warm and well mixed, cooling at nearly the dry adiabatic rate (§ 8.3). The warmth of Hillsborough, and the cold of St John's, come from the surface beneath and the air that has flowed across it.

9.3 The ocean and the winds

Two things decide the temperature of the air near the ground in winter and early spring: the temperature of the surface it lies over, and the surface it has just crossed (Lutgens et al., 2019, ch. 3).

1. **Ocean currents.** The Gulf Stream carries warm water from the Gulf of Mexico along the coast of North America and, as the North Atlantic Current, across to north-west Europe. The seas around Ireland are therefore far warmer in winter than their latitude would suggest. Newfoundland is washed instead by the Labrador Current, which brings cold water (and, in late winter and spring, sea ice and icebergs) south from the Arctic.
2. **Prevailing winds.** In the middle latitudes the winds blow mostly from the west. Air reaching Northern Ireland has crossed thousands of kilometres of relatively warm ocean and arrives mild and moist (maritime air). Air reaching St John's has usually crossed the North American continent, which in winter is snow-covered and very cold (continental polar air).
3. **Water stores heat.** Water heats and cools far more slowly than land, so the ocean keeps the air over it mild in winter and cool in summer (EGM305 module team, 2020, slide 60). Downwind of an ocean, winters are mild; downwind of a continent, they are harsh.




Figure 9.1. Why Hillsborough is milder than St John's in late winter (schematic). Westerly winds reach Northern Ireland after crossing ocean warmed by the Gulf Stream and North Atlantic Current; they reach Newfoundland after crossing the cold continent, and its coast is bathed by the cold Labrador Current.

9.4 Question 14: Hillsborough and St John's

The question on the sheet

Compare the surface temperatures of St John's and Hillsborough. Although Northern Ireland is located at a higher latitude, why do you think the surface temperature is higher? Do you think that this difference is the same throughout the year?

Answer to Question 14

On 21 March 1990 Hillsborough was at **11.05** °C and St John's at **−1.65** °C: **12.7** °C **warmer** in Northern Ireland, although it is about 7° of latitude further north and received less sunshine at noon. Northern Ireland lies downwind of the North Atlantic, whose surface is kept warm by the **Gulf Stream and North Atlantic Current**, so the prevailing westerly winds bring it mild maritime air. St John's is bathed by the cold **Labrador Current** and its westerly winds arrive from the frozen North American continent. **The difference is not the same all year:** it is largest in winter and early spring, when the ocean is much warmer than the land and the continent is coldest, and much smaller in summer, when the continent warms up (summer days in St John's can be warmer than in Northern Ireland).

Check. The soundings agree with this explanation: the contrast is confined to the lowest kilometre or two, the layer in contact with the surface, while higher up St John's is the warmer (§ 9.2). A difference caused by latitude or by the Sun would extend through the whole troposphere.

A question that comes up

Is one day's sounding enough to say anything about climate?
No. One day shows weather; climate is the average over many years (EGM305 module team, 2026). But this day is typical of the pattern: north-west Europe has some of the mildest winters for its latitude anywhere in the world, and eastern Canada some of the coldest. The sounding illustrates a climatic contrast; the climate itself is established from long records.

9.5 Try it yourself: Practice Set 9

Practice Set 9: places and seasons

9.1 At the June solstice, what are the noon solar intensities at Hillsborough (54.4° N) and St John's (47.6° N)? Use Part II, Eqs. (18.4) and (18.3).

9.2 Bergen in Norway (60.4° N) has milder winters than many places 15° further south in eastern North America. Give two reasons.

9.3 Which of the two soundings has the more stable lowest kilometre, and why does that fit the explanation of Question 14?

Answers to Practice Set 9

9.1 Zenith angles $54.4 - 23.5 = 30.9°$ and $47.6 - 23.5 = 24.1°$; sun angles $59.1°$ and $65.9°$; intensities $\sin 59.1° = 86\,\%$ and $\sin 65.9° = 91\,\%$ per unit.

9.2 The North Atlantic Current keeps the Norwegian Sea warm and ice-free in winter, and the prevailing westerly winds bring mild maritime air from the ocean, whereas eastern North America receives air from the cold continent.

9.3 St John's: its lowest 3 km is almost isothermal (a lapse rate under $1\,^\circ\mathrm{C\,km^{-1}}$), so cold air lies trapped beneath warmer air, as over a cold surface. Hillsborough's lowest 1.3 km cools at $9.5\,^\circ\mathrm{C\,km^{-1}}$, a mixed layer over a relatively warm surface.

Chapter 10

Putting Practical 1 together

10.1 One chain of ideas

The fourteen questions are one argument, each step resting on the one before:

1. Air has weight, and pressure is the weight of the air above. Because air is compressible, the air near the ground is squeezed by everything above it: pressure and density fall **exponentially** with height, halving roughly every 5.6 km (Questions 1–2, Chapter 3).
2. Reading that curve gives the pressure anywhere; because the proportion of oxygen is the same at all heights, its **partial pressure** falls in step, to a third on Everest (Questions 3–4, Chapter 4).
3. Temperature falls through the troposphere, on average by 6.5 °C per km, then stops falling at the **tropopause**, above which the ozone-heated stratosphere begins (Question 5, Chapter 5).
4. Real soundings follow the standard atmosphere only roughly: they contain **inversions**, and their tropopauses lie at different heights (Questions 6–8, Chapter 6).
5. The troposphere is **thickest and its top coldest over the tropics**, where heating and convection are strongest and warm air columns are tallest (Questions 9–11, Chapter 7).
6. Equal height steps remove equal *fractions* of the air, so pressure falls fast low down and slowly high up; and air that sinks warms by compression, so the cold air aloft is held up: the atmosphere is **stably layered** (Questions 12–13, Chapter 8).
7. Near the ground, the surface and the winds over it control the temperature: an ocean warmed by the Gulf Stream keeps Northern Ireland mild (Question 14, Chapter 9).

Practical 2 (Part II) explains where the heat comes from in the first place, and Practical 3 (Part III) how the differences of pressure set up here make the air move.

10.2 All the answers in one table

Table 10.1. Every answer of Practical 1, with the section where it is worked out.

Q	Answer	Where
1	100, 50, 25, 12.5, 6.25, 3.13, 1.56, 0.78, 0.39, 0.20 % above 0, 5.6, . . . , 50.4 km	§ 3.4
2	1000, 500, 250, 125, 62.5, 31.3, 15.6, 7.8, 3.9, 2.0 hPa; second axis = 10× the percentage	§ 3.5
3	Portstewart 100 %, 1000 hPa; Everest 33.4 %, 334 hPa; jet 22.6 %, 226 hPa; Machu Picchu 74.0 %, 740 hPa; balloon 7.4 %, 74 hPa	§ 4.2
4	p_{O_2}: 210, 70, 48, 155, 16 hPa; as % of the sea-level atmosphere 21, 7.0, 4.8, 15.5, 1.6 %; compared with sea-level oxygen 100, 33, 23, 74, 7 %	§ 4.5

Q	Answer	Where
5	2, −11, −24, −37, −50 °C at 2, 4, 6, 8, 10 km; −56.5 °C from 11 to 16 km	§ 5.3
6	graph of the two soundings and the standard atmosphere (Figure 6.1)	§ 6.2
7	Hillsborough inversion about 1360–1500 m; St John's nearly isothermal from the ground to about 3 km	§ 6.4
8	standard 11 000 m, −56.5 °C; Hillsborough 9383 m, −51.05 °C; St John's 11 482 m, −62.05 °C	§ 6.6
9	62.10/9.345 = 6.65 °C per km	§ 7.2
10	Florida tropopause about 16 000 m, −75.6 °C	§ 7.3
11	no: thickest over the tropics (16–17 km), thinnest at the poles (8–9 km); heating, warm tall columns, deep convection	§ 7.4
12	305 hPa against 10 hPa; exponential decay: equal fractions (30 and 33 %), not equal amounts	§ 8.1
13	upper air is far less dense (low pressure); sinking air warms by compression (9.8 °C per km): tropopause air would arrive at 46 °C; stable layering	§ 8.4
14	Hillsborough 12.7 °C warmer: Gulf Stream and North Atlantic Current, westerly maritime air; Labrador Current and continental air at St John's; largest contrast in winter and spring	§ 9.4

10.3 Quick sheet for the lab

Practical 1 on one page

Halving rule: $f = 2^{-z/5.6}$; % above = $100f$; pressure = $1000f$ hPa — Eqs. (3.4), (3.6)
Steps of 5.6 km: 100, 50, 25, 12.5, 6.25 … % — Table 3.1
Oxygen: $p_{O_2} = 0.21\,p$; available oxygen falls in step with p — Eq. (4.1)
Lapse rate: $\Gamma = (T_{\text{lower}} - T_{\text{upper}})/(z_{\text{upper}} - z_{\text{lower}})$; negative means inversion — Eq. (5.1)
Standard atmosphere: $T = 15 - 6.5\,z$ to 11 km, then −56.5 °C — Eq. (5.2)
Tropopause: lowest level where the temperature stops falling (and stays within 2 °C km^{-1} for the 2 km above) — § 6.5
Scale height: $H = R_d T/g \approx 8$ km; from data $H = \Delta z/\ln(p_1/p_2)$ — Eqs. (3.3), (8.1)
Sinking air warms: 9.8 °C per km; $\theta = T(1000/p)^{0.286}$ — Eqs. (8.2), (8.3)

Graph habits: height always on the vertical axis; one colour and symbol per profile; straight lines between sounding points, a smooth curve for the halving rule; convert metres to kilometres before plotting.

10.4 Mistakes to avoid

- **Halving the height instead of the air.** Each 5.6 km step halves the *percentage*; the heights go up in equal steps.
- **Joining the halving-rule points with straight lines.** The rule describes a smooth exponential curve; straight segments make the readings of Question 3 wrong between the points.

- **Mixing metres and kilometres** in the lapse rate or in $z/5.6$.
- **Getting the sign of the lapse rate wrong.** Temperature falling with height is a *positive* lapse rate.
- **Taking any warm step as the tropopause.** A thin inversion low in the troposphere (Hillsborough, 1.4 km) is not the tropopause: the temperature falls again above it.
- **Saying the air aloft is "heavier" because it is colder.** At the same pressure, yes; at its own low pressure it is far lighter.

10.5 Self-test

Self-test: one line each

S1. Why does pressure fall faster near the ground than high up?

S2. Why does the air on Everest have less oxygen, though it is still 21 % oxygen?

S3. What is the difference between an inversion and the tropopause?

S4. Why is the tropical tropopause the coldest?

S5. Why does the cold air of the upper troposphere not sink?

S6. Why is Northern Ireland milder in winter than Newfoundland?

Answers to the self-test

S1. Each height step removes the same fraction of the air above, and near the ground there is far more air above (§ 8.1).

S2. Each breath holds a third as many molecules; the oxygen partial pressure is a third of that at sea level (§ 4.5).

S3. An inversion is any layer where the temperature rises with height; the tropopause is the top of the troposphere, above which the temperature stops falling for good (§ 6.5).

S4. It is the highest, so the temperature falls over the most kilometres before it stops (§ 7.4).

S5. It is far less dense than the air below, and if it sank it would warm by compression and become lighter still (§ 8.4).

S6. Westerly winds off an ocean warmed by the Gulf Stream, against a cold current and a frozen continent upwind of Newfoundland (§ 9.4).

Part II

Practical 2: Solar radiation

Chapter 11

Practical 2 at a glance

11.1 What the practical asks of you

Practical 2 has three learning goals, printed at the top of the sheet: the *laws of radiation*, *black bodies and radiant energy*, and the *determination of solar intensity* (EGM305 module team, 2025a). The five questions build on each other, so it pays to work through them in order. They also build on Practical 1: the same atmosphere whose layers you plotted there (Chapter 5) is now heated by the Sun and cooled by its own radiation.

The laws of radiation. Every object gives off radiation, and its temperature decides both *what kind* (the wavelength of its peak, Wien's law, Question 1) and *how much* (the total, the Stefan–Boltzmann law, Question 2).

Black bodies and planets. You plot the spectra of three planets, read their temperatures from the peaks (Questions 3a and 3b), calculate the temperature each planet *should* have from its distance to the Sun and its reflectivity (Question 3c), match the spectra to the planets (Question 3d) and explain why the real temperatures differ (Question 3e).

The Earth's energy balance and solar intensity. You calculate the temperature the Earth would have without an atmosphere, which measures the greenhouse effect (Question 4), and then show why the angle of the Sun makes the tropics warm, the poles cold and the seasons change (Question 5).

The sheet numbers two different parts of Question 3 as "3d". In this companion the first ("link the data sets with the planets") is **3d** and the second ("using the table you constructed, explain…") is **3e**. The unnumbered question in Question 4 ("why is the average incoming radiation only $341\,\mathrm{W\,m^{-2}}$?") is called **4b′**.

Table 11.1. Where each question of the Practical 2 sheet is answered, which equations it needs, and where the same material is in the Week 2 lecture (EGM305 module team, 2020) and the set texts (Lutgens et al., 2019; Ahrens and Henson, 2022).

Question	What it asks	Chapter	Equations	L2 slides	Texts
1	the Sun's temperature from its peak wavelength	13	(13.1)	39–42	ch. 2
2a–2e	emission of the Sun and the Earth; why T and E ratios differ	14	(14.1), (14.2)	39–42	ch. 2
3 (graph), 3a–3b	plot three spectra; peaks and temperatures	15	(13.2)	40–41	ch. 2
3c–3e	planetary temperatures and why they differ	16	(16.2), (16.1)	43–44, 52, 73	ch. 2
4a–4g	albedo, energy balance, greenhouse effect	17	(17.1)–(17.3)	36, 43–54	ch. 2
5a–5c	sun angle, beam spreading, seasons	18	(18.2)–(18.3)	55–59	ch. 2–3

Chapter 12 builds the physics that every question uses; Chapter 19 gathers all the answers in one table. The formulas, derivations, recomputed data, corrections to the key, symbols and constants for this practical are in Appendix B.

11.2 Conventions for this practical

The conventions of the whole companion are in § 1.3. Three more apply here.

- **Temperatures in the radiation laws are always in kelvin.** The sheet converts with $T_{^\circ\mathrm{C}} = T_{\mathrm{K}} - 273.16$; the exact offset is 273.15 (Tiesinga et al., 2021). The difference (0.01 K) never changes an answer at the precision of the sheet, so this part uses the sheet's 273.16 to keep your numbers identical to the key.
- **Constants.** The sheet's values are used for every answer: Wien's constant $C = 2898\,\mu\mathrm{m\,K}$ and the Stefan–Boltzmann constant $\sigma = 5.67 \times 10^{-8}\,\mathrm{W\,m^{-2}\ K^{-4}}$. Precise values and sources are listed in Table B.4.
- **Units** are written as powers: $\mathrm{W\,m^{-2}}$ means watts per square metre (the sheet writes $\mathrm{W/m^2}$). Wavelengths are in micrometres, $1\,\mu\mathrm{m} = 10^{-6}\,\mathrm{m}$.

11.3 A word on the printed answer key

The answer sheet was checked against independent calculation, one answer at a time. Most of it is right. A few slips need correcting: one misplaced decimal point (Question 4b), a mix of Celsius and kelvin (2c), one physical description that is not quite right (2e), and some rounded or simplified values that are worth knowing about (3c, 3e, 4b′, 5c). Each is explained where it arises, and § B.4 lists them all. None of them changes the physics the practical is teaching.

11.4 Before the practical

The lecture asks you to read Chapters 2 and 3 of Lutgens et al. (2019) or of Ahrens and Henson (2022) before this practical. Chapter 2 of each covers the heating of the Earth's surface and atmosphere (the radiation laws, the energy balance and the greenhouse effect) and Chapter 3 covers temperature and its controls, including latitude and the seasons. A free alternative is Chapter 2 of Stull (2017), which treats solar and infrared radiation with worked examples at the level of this practical.

Chapter 12

What radiation is

This chapter builds the ideas that every question of the sheet uses. It answers no question directly; Chapters 13 to 18 refer back to it.

12.1 Energy that needs no material to travel

Energy moves from place to place in three ways. *Conduction* passes it from molecule to molecule through contact (a spoon warming in hot tea). *Convection* carries it in a moving fluid (warm air rising from a radiator). *Radiation* needs no material at all: it is energy travelling as an electromagnetic wave, a ripple of electric and magnetic fields (Lutgens et al., 2019, ch. 2). Between the Sun and the Earth there are 150 million kilometres of almost empty space, so radiation is the only way the Sun's energy can reach us.

Two properties of radiation matter for this practical:

- **It carries energy.** The body that emits it loses energy; the body that absorbs it gains energy and warms.
- **It comes in many wavelengths.** Radio waves, microwaves, infrared, visible light, ultraviolet and X-rays are all the same kind of thing. They differ only in wavelength.

A question that comes up

Is this the same as radioactivity?
No. In meteorology "radiation" always means electromagnetic radiation: sunlight, the infrared glow of the ground, and so on. Radioactivity is the decay of unstable atomic nuclei, an entirely different process. Only the word is shared.

12.2 Wavelength, frequency and speed

A wave is a repeating pattern of crests and troughs. The **wavelength** λ (the Greek letter lambda) is the distance from one crest to the next (Figure 12.1). The **frequency** f is the number of crests that pass a point each second. Shake one end of a rope slowly and the crests are far apart; shake it quickly and they bunch up.

All electromagnetic waves travel through space at the same speed, the speed of light c. In one second a source sends out f crests, each λ apart, and together they cover the distance the wave travels in one second:

$$c = f\lambda. \tag{12.1}$$

where

c	speed of light ($\mathrm{m\,s^{-1}}$)	3.00×10^8, Table B.4
f	frequency: crests per second ($\mathrm{s^{-1}}$, or hertz, Hz)	calculated
λ	wavelength (m)	given or calculated

Because c is fixed, frequency and wavelength are tied together: a short wavelength means a high frequency, and a long wavelength a low one.




Figure 12.1. Wavelength and frequency. Both waves move at the speed of light, so the one whose source oscillates faster (blue) has its crests closer together, Eq. (12.1). Not to scale: the crests of the Earth's infrared are about 10 μm apart, a seventh of the width of a hair.

Worked example: the frequencies of sunlight and of the Earth's glow

Rearranging Eq. (12.1), $f = c/\lambda$. For the Sun's peak wavelength of $0.5\,\mu\text{m} = 0.5 \times 10^{-6}$ m:

$$f_{\text{Sun}} = \frac{3.00 \times 10^{8}}{0.5 \times 10^{-6}} = 6.0 \times 10^{14}\ \text{Hz}.$$

For the Earth's peak near $10\,\mu\text{m} = 10 \times 10^{-6}$ m:

$$f_{\text{Earth}} = \frac{3.00 \times 10^{8}}{10 \times 10^{-6}} = 3.0 \times 10^{13}\ \text{Hz}.$$

Sunlight oscillates twenty times faster than the Earth's infrared, because its wavelength is twenty times shorter.

12.3 Photons: radiation comes in packets

Radiation is emitted and absorbed in separate packets of energy called **photons**, much as rain arrives as separate drops. The energy of one photon depends only on its frequency (Planck, 1901):

$$\varepsilon = hf = \frac{hc}{\lambda}, \tag{12.2}$$

where

ε	energy of one photon (J)	calculated
h	Planck's constant (J s)	6.626×10^{-34}, Table B.4
$f,\ c,\ \lambda$	as in Eq. (12.1)	

where the second form follows from substituting $f = c/\lambda$ from Eq. (12.1). So **the shorter the wavelength, the more energy each photon carries**:

$$\varepsilon_{0.5\,\mu\text{m}} = \frac{(6.626 \times 10^{-34})(3.00 \times 10^{8})}{0.5 \times 10^{-6}} = 3.98 \times 10^{-19}\ \text{J},$$
$$\varepsilon_{10\,\mu\text{m}} = \frac{(6.626 \times 10^{-34})(3.00 \times 10^{8})}{10 \times 10^{-6}} = 1.99 \times 10^{-20}\ \text{J}.$$

A photon of sunlight carries twenty times the energy of a photon of the Earth's infrared. This is why ultraviolet light, with even shorter wavelengths, can break chemical bonds and burn skin, while infrared only warms.

12.4 The electromagnetic spectrum

Table 12.1 and Figure 12.2 arrange the kinds of radiation by wavelength. Two ranges matter for the atmosphere. The Sun emits mostly between about 0.2 and 4 μm (ultraviolet, visible and near infrared): this is called **shortwave** radiation. The Earth and its atmosphere emit mostly between about 4 and 100 μm (thermal infrared): this is **longwave** radiation (EGM305 module team, 2020, slide 38). Most of the Sun's ultraviolet never reaches the ground: ozone absorbs it, and that absorption is what warms the stratosphere in the temperature profiles of Practical 1 (Part I, § 5.4).

Table 12.1. The electromagnetic spectrum. All of these are photons; they differ only in wavelength and so in energy per photon, Eq. (12.2).

Type	Wavelength	Energy per photon	Example source
Radio	metres to kilometres	smallest	transmitters
Microwave	1 mm to 1 m	very small	ovens; radar
Infrared	0.75 to 1000 μm	small	the Earth, people, warm objects (peak near 10 μm)
Visible	0.4 to 0.75 μm	moderate	the Sun (peak near 0.5 μm)
Ultraviolet	0.01 to 0.4 μm	large	the Sun; mostly absorbed by ozone
X-ray, gamma	shorter still	largest	medical X-rays; radioactive decay




Figure 12.2. The part of the spectrum that matters for the atmosphere. Solar (shortwave) and terrestrial (longwave) radiation meet near 4 μm.

12.5 Why everything above absolute zero radiates

The first law on the sheet says that *all objects continually radiate energy.* Why?

Absolute zero (0 K, or −273.15 °C) is the lowest possible temperature, at which the particles of a substance have the least possible motion. The kelvin scale starts there; that is why the radiation laws need temperatures in kelvin (§ 11.2).

All matter (rock, soil, sea, ice, cloud, air, people) is made of atoms that contain negative electrons and positive nuclei. Above absolute zero these particles are always vibrating and colliding, so their electric charges are always being jostled. Moving charges send out electromagnetic waves, just as a radio transmitter drives electrons up and down an aerial. So

$$\text{above 0 K} \;\Rightarrow\; \text{particles move} \;\Rightarrow\; \text{charges are jostled} \;\Rightarrow\; \text{radiation is emitted.}$$

Cold things radiate too, only less. An ice cube at −10 °C gives off about 270 W m^{-2}, and your skin about 500 W m^{-2} (you will be able to check both with Eq. (14.1) in Chapter 14). A thermal camera sees people in the dark for exactly this reason. Ordinary objects look dark only because almost all their emission is infrared, which our eyes cannot see.

12.6 Black bodies

Definition

A **black body** is an ideal object that absorbs *all* the radiation falling on it, at every wavelength, and reflects none. At any given temperature it also *emits* the most radiation that any object at that temperature can emit, at every wavelength (EGM305 module team, 2020, slide 37).

The last law on the sheet, *good absorbers are good emitters*, is Kirchhoff's law: at each wavelength, the fraction of incoming radiation a surface absorbs equals the fraction of black-body radiation it emits. That fraction is the **emissivity**, ϵ, between 0 and 1; a black body has $\epsilon = 1$ at all wavelengths (Wallace and Hobbs, 2006, ch. 4).

No real object is a perfect black body, but many come close at some wavelengths. Snow is white (a poor absorber) in visible light but almost black in the thermal infrared, where its emissivity is close to 1; most natural surfaces, water and thick cloud behave nearly as black bodies in the infrared. The Sun is close to a black body over the whole spectrum. This is why the black-body laws work so well for this practical.

12.7 The Planck curve

How much energy does a black body emit at each wavelength? Imagine sorting all the radiation leaving a hot plate into wavelength "bins" and plotting how much energy lands in each bin. The result is a **spectrum**, and for a black body its shape was found by Planck (1901):

$$B_\lambda(T) = \frac{2hc^2}{\lambda^5} \frac{1}{\exp\left(\dfrac{hc}{\lambda k_B T}\right) - 1}. \tag{12.3}$$

where

B_λ	spectral radiance: power per unit area, per unit solid angle, per unit wavelength ($\mathrm{W\,m^{-2}\,sr^{-1}\,m^{-1}}$)	calculated
h, c	as in Eqs. (12.1) and (12.2)	Table B.4
k_B	Boltzmann's constant ($\mathrm{J\,K^{-1}}$)	1.381×10^{-23}, Table B.4
λ	wavelength (m)	the horizontal axis
T	absolute temperature of the body (K)	e.g. 5796 K for the Sun (Question 1)

You will not need to calculate with Eq. (12.3). What matters is the shape it draws (Figure 12.3) and the two laws that come out of it.

- **The curve rises steeply, peaks, then tails off slowly.** At short wavelengths each photon is expensive in energy, Eq. (12.2), and only the most energetic particles can make it, so emission falls off a cliff. At long wavelengths photons are cheap but each carries little, so the curve tails off gradually.
- **The position of the peak** says *what kind* of radiation the body mainly emits. Hotter bodies peak at shorter wavelengths. This is Wien's law (Chapter 13).
- **The area under the curve** is the *total* radiation emitted. Hotter bodies have far larger areas. This is the Stefan–Boltzmann law (Chapter 14).

A geographer's analogy is the storm hydrograph. Its peak tells you *when* discharge was greatest; the area under it tells you the total *volume* of water. The Planck curve is read in the same way, with energy in place of water and wavelength in place of time.




Figure 12.3. Black-body spectra of the Sun and the Earth from Eq. (12.3), each divided by its own peak value so that both can be seen on one graph. The two curves hardly overlap: the Sun's radiation is shortwave and the Earth's longwave. On a common scale the Earth curve would be invisible, because per square metre the Sun emits about 166 000 times more energy (Chapter 14).

The four laws on the sheet, and where each is used

1. *All objects continually radiate energy over a range of wavelengths* (§ 12.5; the spread is the Planck curve, Eq. (12.3)).
2. *Hotter objects radiate more total energy per unit area*: the Stefan–Boltzmann law, Eq. (14.1), used in Questions 2 and 4.
3. *Hotter objects radiate more at short wavelengths*: Wien's displacement law, Eq. (13.1), used in Questions 1 and 3.
4. *Good absorbers are good emitters*; the perfect one is a black body (§ 12.6), the assumption behind Questions 3 and 4.

12.8 Questions that come up

A question that comes up

If the air radiates too, why do we not feel it?
We do. On a clear night the ground cools quickly because it radiates to a cold sky; under cloud it stays warmer because the cloud radiates back to it. That downward infrared from cloud and air is part of the greenhouse effect of Chapter 17.

A question that comes up

When does an object start to glow visibly?
When enough of its Planck curve reaches into the visible. For a black body this begins at roughly 800 K, with a dull red glow, as in an electric hob ring. At the Sun's temperature the peak is in the middle of the visible.

12.9 Try it yourself: Practice Set 12

Practice Set 12: waves and photons

Use $c = 3.00 \times 10^{8}\,\mathrm{m\,s^{-1}}$ and $h = 6.626 \times 10^{-34}\,\mathrm{J\,s}$.

12.1 What is the frequency of light with a wavelength of $0.5\,\mu$m?

12.2 What is the energy of one photon of wavelength $10\,\mu$m?

12.3 A microwave oven works at a wavelength of 12.2 cm. What is its frequency?

12.4 How many times more energy does a $0.3\,\mu$m ultraviolet photon carry than a $3\,\mu$m infrared photon?

12.5 Name the part of the spectrum, and say whether it is shortwave or longwave, for radiation of 0.3, 0.55, 2 and $15\,\mu$m.

Answers to Practice Set 12

12.1 $f = c/\lambda = 3.00 \times 10^{8}/0.5 \times 10^{-6} = 6.0 \times 10^{14}\,\mathrm{Hz}$, Eq. (12.1).

12.2 $\varepsilon = hc/\lambda = (6.626 \times 10^{-34})(3.00 \times 10^{8})/(10 \times 10^{-6}) = 1.99 \times 10^{-20}\,\mathrm{J}$, Eq. (12.2).

12.3 $f = 3.00 \times 10^{8}/0.122 = 2.46 \times 10^{9}\,\mathrm{Hz}$ (2.46 GHz).

12.4 The energy is inversely proportional to the wavelength, so the ratio is $3/0.3 = 10$ times.

12.5 $0.3\,\mu$m: ultraviolet, shortwave. $0.55\,\mu$m: visible (green), shortwave. $2\,\mu$m: near infrared, shortwave (still part of the solar spectrum). $15\,\mu$m: thermal infrared, longwave.

Chapter 13

Wien's law and the temperature of the Sun

This chapter answers Question 1 of the sheet.

13.1 Hotter bodies peak at shorter wavelengths

Heat a piece of iron in a forge. At first it gives off only invisible infrared; then it glows dull red, then orange, then yellow-white. As it gets hotter, the colour of its glow moves towards shorter wavelengths. Figure 13.1 shows the same thing with Planck curves, Eq. (12.3): as the temperature rises, every curve gets taller, and its peak slides to the left.




Figure 13.1. Planck curves, Eq. (12.3), for four temperatures, drawn as the power emitted per square metre per micrometre of wavelength. The dots mark the peaks and the grey line joins them: the hotter the body, the shorter the wavelength of its peak (Wien's law, Eq. (13.1)) and the larger the area under its curve (the Stefan–Boltzmann law, Eq. (14.1)).

13.2 Wien's displacement law

The wavelength of the peak is inversely proportional to the temperature (EGM305 module team, 2020, slides 39–41):

$$\lambda_{\max} = \frac{C}{T}. \tag{13.1}$$

where

$\lambda_{\max}$	wavelength of peak emission (μm)	read from a spectrum, or calculated
C	Wien's constant (μm K)	2898, given on the sheet
T	absolute temperature of the body (K)	calculated, or given

"Inversely proportional" means that when one doubles the other halves: a body twice as hot peaks at half the wavelength. The constant C is not arbitrary. It follows from finding the peak of the Planck curve, which is done step by step in Appendix B.2.1; the result is $C = hc/(4.965\,k_B) = 2897.8\,\mu\text{m K}$.

To find a temperature from a peak wavelength, multiply both sides of Eq. (13.1) by T and divide both sides by $\lambda_{\max}$:

$$T = \frac{C}{\lambda_{\max}}. \tag{13.2}$$

where

T, C, $\lambda_{\max}$ as in Eq. (13.1)

The units work out: μm K divided by μm leaves K. This is why $\lambda_{\max}$ must be in micrometres when $C = 2898\,\mu\text{m K}$.

13.3 Question 1: the temperature of the Sun

The question on the sheet

If the wavelength emitted more than any other by the Sun is $0.5\,\mu$m, what is the black-body temperature of the Sun in Celsius? Note the conversion from kelvin to Celsius is $^\circ\text{C} = \text{K} - 273.16$.

What the question asks, in plain words. We know where the Sun's spectrum peaks ($0.5\,\mu$m, in the blue-green part of the visible). Assuming the Sun is a black body, Wien's law tells us what temperature produces a peak there.

Method.

Step 1. Write down the rearranged law, Eq. (13.2): $T = C/\lambda_{\max}$.

Step 2. Substitute $C = 2898\,\mu\text{m K}$ and $\lambda_{\max} = 0.5\,\mu$m:

$$T_{\text{Sun}} = \frac{2898\ \mu\text{m K}}{0.5\ \mu\text{m}} = 5796\ \text{K}.$$

Step 3. Convert to Celsius with the sheet's offset:

$$T_{\text{Sun}} = 5796 - 273.16 = 5522.84\ ^\circ\text{C} \approx 5523\ ^\circ\text{C}.$$

Answer to Question 1

$T_{\text{Sun}} = 2898/0.5 = \mathbf{5796\,K}$, which is $5796 - 273.16 = \mathbf{5522.8}\,^\circ$C (about $5500\,^\circ$C).

Check. Measurements give the Sun an effective temperature of 5772 K (Prša et al., 2016), within half a percent of our 5796 K: the small difference comes from rounding the peak to $0.5\,\mu$m (the exact value for 5772 K is $0.502\,\mu$m). The order of magnitude is also sensible: a body of a few thousand kelvin glows white-hot, and the Sun's peak lies in the visible, which is the light our eyes evolved to use.

The 5796 K found here is used again in Question 2a (§ 14.3) and Question 2c (§ 14.5).

A question that comes up

Why does the answer say 5522.8 °C and not 5522.85 °C?
The sheet subtracts 273.16; the exact offset between the kelvin and Celsius scales is 273.15 (Tiesinga et al., 2021). Either way the answer rounds to 5523 °C, and at the precision of a peak read as "0.5 μm" the last digits mean nothing. Quote about 5500 °C when you describe it in words.

A question that comes up

If the Sun peaks in blue-green light, why does it look white, or yellow?
Because its spectrum is broad (Figure 12.3): it emits strongly at all visible wavelengths, and the mixture looks white. Near the horizon, the long path through the air scatters blue light out of the beam, so the Sun looks yellow or red. The position of the peak also depends on how the spectrum is drawn (per unit wavelength, as here, or per unit frequency), so "the colour of the peak" is not the colour we see.

13.4 Wien's law in practice

- **Satellites.** A satellite cannot put a thermometer on the ground. It measures radiation in chosen wavelength bands and converts it to temperature with the radiation laws (EGM305 module team, 2020, slide 69). Wien's law tells the designer which band suits which target: about 10–12 μm for the land, sea and cloud tops (near 300 K and below), and about 4 μm for fires, which are far hotter.
- **Stars.** Astronomers read the temperature of a star from its colour: red stars are cool (around 3000 K) and blue-white stars hot (10 000 K or more).
- **Shortwave and longwave.** Because the Sun is about twenty times hotter than the Earth, Wien's law puts its peak at a wavelength about twenty times shorter. This single fact separates solar from terrestrial radiation (Figure 12.3) and makes the greenhouse effect possible (Chapter 17).

13.5 Try it yourself: Practice Set 13

Practice Set 13: Wien's law

Use $C = 2898\,\mu\text{m K}$ and $°\text{C} = \text{K} - 273.16$.

13.1 The star Sirius has a surface temperature of about 9940 K. At what wavelength does it emit most, and in which part of the spectrum is that?

13.2 Human skin is at about 33 °C. At what wavelength does it emit most?

13.3 A star's spectrum peaks at 1.0 μm. What is its temperature in kelvin and in Celsius?

13.4 A forest fire burns at about 1000 K. Where is its peak? Why do satellites look for fires near 4 μm rather than at 10 μm?

13.5 An ice cube at −10 °C emits radiation. At what wavelength does it emit most?

Answers to Practice Set 13

13.1 $\lambda_{\max} = 2898/9940 = 0.29\,\mu$m: ultraviolet, Eq. (13.1). Sirius looks blue-white.

13.2 $T = 33 + 273.16 = 306.16\,$K, so $\lambda_{\max} = 2898/306.16 = 9.47\,\mu$m, in the thermal infrared. Thermal cameras are built to see near this wavelength.

13.3 $T = 2898/1.0 = 2898\,$K $= 2624.8\,^{\circ}$C, Eq. (13.2). A cool, red star.

13.4 $\lambda_{\max} = 2898/1000 = 2.9\,\mu$m. Near $4\,\mu$m a fire emits far more than the cool ground around it (whose peak is near $10\,\mu$m), so fires stand out clearly; at $10\,\mu$m the contrast is much smaller.

13.5 $T = 263.16\,$K, so $\lambda_{\max} = 2898/263.16 = 11.0\,\mu$m. Cold objects emit at longer wavelengths than warm ones.

Chapter 14

The Stefan–Boltzmann law: how much the Sun and the Earth emit

This chapter answers Questions 2a to 2e of the sheet.

14.1 The total emission of a black body

Wien's law (Chapter 13) told us *where* a spectrum peaks. The Stefan–Boltzmann law tells us *how much* a body emits in total, adding up every wavelength: it is the area under the Planck curve (EGM305 module team, 2020, slides 39–42):

$$E = \sigma T^4. \tag{14.1}$$

where

E	total radiation emitted per square metre of surface, the "intensity of radiation" ($\mathrm{W\,m^{-2}}$)	calculated
σ	Stefan–Boltzmann constant ($\mathrm{W\,m^{-2}\ K^{-4}}$)	5.67×10^{-8}, given on the sheet
T	absolute temperature of the surface (K)	from Question 1, or given

The law is derived from the Planck curve in Appendix B.2.2: integrating Eq. (12.3) over all wavelengths gives exactly T^4 times a combination of constants, and that combination is σ.

It applies to a black body. A real surface emits a fraction ϵ (its emissivity, § 12.6) of this: $E = \epsilon\sigma T^4$. For the Sun, and for most natural surfaces in the infrared, ϵ is close to 1, so Eq. (14.1) is a good approximation.

14.2 What a fourth power means

T^4 means $T \times T \times T \times T$. A fourth power grows very fast:

Temperature multiplied by	1.1	1.5	2	3	10	20
Emission multiplied by	1.46	5.06	16	81	10 000	160 000

A body only 10 % hotter emits 46 % more; a body twice as hot emits sixteen times as much. Figure 14.1 shows this on a scale wide enough to hold both the Earth and the Sun.




Figure 14.1. The Stefan–Boltzmann law, Eq. (14.1), on logarithmic axes, where a power law becomes a straight line of slope 4. Going from the Earth to the Sun the temperature rises twenty-fold (Question 2c) but the emission rises 166 000-fold (Question 2d).

14.3 Question 2a: the emission of the Sun

The question on the sheet

Using this equation, calculate how much radiation is emitted at the surface of the Sun.

Method. Use Eq. (14.1) with the Sun's temperature from Question 1, $T = 5796\,\mathrm{K}$ (§ 13.3). Raise to the fourth power by squaring twice; this avoids slips on a calculator:

Step 1. $T^2 = 5796^2 = 3.3594 \times 10^7$.

Step 2. $T^4 = (3.3594 \times 10^7)^2 = 1.1285 \times 10^{15}$.

Step 3. $E_{\mathrm{Sun}} = \sigma T^4 = (5.67 \times 10^{-8})(1.1285 \times 10^{15}) = 6.399 \times 10^7\,\mathrm{W\,m^{-2}}$.

When multiplying powers of ten, add the exponents: $10^{-8} \times 10^{15} = 10^7$.

Answer to Question 2a

$E_{\mathrm{Sun}} = 5.67 \times 10^{-8} \times 5796^4 = \mathbf{6.40 \times 10^7}\,\mathrm{W\,m^{-2}}$: about 64 million watts from every square metre of the Sun's surface.

Check. With the measured effective temperature of 5772 K (Prša et al., 2016) the same calculation gives $6.29 \times 10^7\,\mathrm{W\,m^{-2}}$, within 2 %. Spread over the whole Sun, this emission is the Sun's total power output, and diluted over the distance to the Earth it becomes the solar constant of Question 3c (see the worked example in § 14.8).

14.4 Question 2b: the emission of the Earth

The question on the sheet

If the Earth has a black-body temperature of 14 °C, what is its intensity of radiation?

Method.

Step 1. Convert to kelvin: $T = 14 + 273.16 = 287.16\,\mathrm{K}$. The radiation laws need kelvin (§ 11.2).
Step 2. $T^2 = 287.16^2 = 82\,460.9$, and $T^4 = 82\,460.9^2 = 6.7998 \times 10^9$.
Step 3. $E_{\mathrm{Earth}} = (5.67 \times 10^{-8})(6.7998 \times 10^9) = 385.5\,\mathrm{W\,m^{-2}}$.

Answer to Question 2b

$E_{\mathrm{Earth}} = 5.67 \times 10^{-8} \times 287.16^4 = \mathbf{385.5}\,\mathrm{W\,m^{-2}}$.

Check. Observed energy budgets put the emission of the Earth's surface at about $396\,\mathrm{W\,m^{-2}}$ (Trenberth et al., 2009), close to our value (the real surface is a little warmer in places and the fourth power weights warm places heavily). Notice that $385.5\,\mathrm{W\,m^{-2}}$ is *more* than the Earth receives from the Sun on average (about $240\,\mathrm{W\,m^{-2}}$ absorbed, Question 4c). That surplus is supplied by the atmosphere radiating back down to the surface: the greenhouse effect of Chapter 17.

A correction to the printed key: how the formula is written

The key writes the calculation as "$E = (5.67 \times 10^{-8} \times 287.16)^4$" with the unit "Wm2". Read literally, that raises σ to the fourth power as well, which would give about 7×10^{-20}, not 385.5. The number in the key (385.5) is right; the bracket is in the wrong place. Write $E = 5.67 \times 10^{-8} \times (287.16)^4$, and the unit as $\mathrm{W\,m^{-2}}$ (watts per square metre).

14.5 Question 2c: how many times hotter is the Sun?

The question on the sheet

How many times warmer is the Sun than the Earth (solar temperature in kelvin / Earth temperature in kelvin)?

Method. Divide the two temperatures, both in kelvin, from Questions 1 and 2b:

$$\frac{T_{\mathrm{Sun}}}{T_{\mathrm{Earth}}} = \frac{5796\ \mathrm{K}}{287.16\ \mathrm{K}} = 20.18.$$

Answer to Question 2c

The Sun is $5796/287.16 \approx$ **20 times** hotter than the Earth (20.18), measured on the kelvin scale.

A correction to the printed key: kelvin, not Celsius

The key writes "$5522.8/287.16 = 20$", dividing the Sun's temperature in *Celsius* by the Earth's in *kelvin*. That division gives 19.2, not 20; the key's 20 comes from the correct kelvin ratio, $5796/287.16 = 20.2$. Ratios of temperatures only make sense on the kelvin scale, which starts at absolute zero. On the Celsius scale they are meaningless: 20 °C is not "twice as hot" as

10 °C (in kelvin it is 293.16/283.16 = 1.035 times as hot), and nothing is "−2 times" as hot as −10 °C.

14.6 Question 2d: how many times more does the Sun emit?

The question on the sheet

How many more times is the Sun's radiation more intense than the Earth's?

Method. Divide the emissions from Questions 2a and 2b:

$$\frac{E_{\text{Sun}}}{E_{\text{Earth}}} = \frac{6.399 \times 10^{7}\ \text{W m}^{-2}}{385.5\ \text{W m}^{-2}} = 1.660 \times 10^{5}.$$

Answer to Question 2d

Each square metre of the Sun emits about **166 000 times** as much radiation as a square metre of the Earth.

14.7 Question 2e: why the two ratios are so different

The question on the sheet

How do you explain the difference between your answer to Question 2c and 2d?

The idea. Write Eq. (14.1) for both bodies and divide. The constant σ cancels, and only the ratio of temperatures remains, raised to the fourth power:

$$\frac{E_{\text{Sun}}}{E_{\text{Earth}}} = \frac{\sigma T_{\text{Sun}}^{4}}{\sigma T_{\text{Earth}}^{4}} = \left(\frac{T_{\text{Sun}}}{T_{\text{Earth}}}\right)^{4}. \tag{14.2}$$

where

E_{Sun}, E_{Earth}	emissions per square metre (W m^{-2})	Questions 2a and 2b
T_{Sun}, T_{Earth}	temperatures (K)	5796 and 287.16

So the emission ratio is the temperature ratio of Question 2c raised to the fourth power: $20.18^4 = 20.18 \times 20.18 \times 20.18 \times 20.18 = 1.660 \times 10^5$, exactly the answer to Question 2d.

Answer to Question 2e

Emission depends on the **fourth power** of temperature, Eq. (14.1). The Sun is about 20 times hotter, so it emits about $20^4 = 160\,000$ times more per square metre (precisely $20.18^4 = 166\,000$). A modest difference in temperature becomes an enormous difference in emission.

A correction to the printed key: a power law, not an exponential

The key says that "E changes exponentially, i.e. if T doubles, E becomes 16 times greater". The example is right, but the word is not. In an *exponential* relationship, such as $E \propto e^{kT}$, the temperature sits in the exponent and every equal step in T multiplies E by the same factor. In the Stefan–Boltzmann law the exponent is fixed at 4 and the temperature is the base: this is a **power law**, $E \propto T^4$. The difference matters: an exponential eventually outgrows any power law, and on the log–log axes of Figure 14.1 a power law is a straight line while an exponential is not. Say "E is proportional to the fourth power of T".

14.8 From the Sun's surface to the Earth

Questions 2a and 3c are linked. The Sun's surface emits σT^4 per square metre; this energy spreads out over ever larger spheres as it travels, so by the time it reaches the Earth it has been diluted to the solar constant S.

Worked example: the solar constant from the Sun's temperature

The Sun's radius is $R_\odot = 6.957 \times 10^8$ m and the Earth–Sun distance is $d = 1.496 \times 10^{11}$ m (Prša et al., 2016). The Sun's total power is its emission per square metre times its surface area $4\pi R_\odot^2$:

$$P = \sigma T^4 \times 4\pi R_\odot^2 = (6.399 \times 10^7) \times 4\pi (6.957 \times 10^8)^2 = 3.89 \times 10^{26}\ \text{W}.$$

At the Earth this power is spread over a sphere of radius d, so each square metre facing the Sun receives

$$S = \frac{P}{4\pi d^2} = \frac{3.89 \times 10^{26}}{4\pi (1.496 \times 10^{11})^2} = 1384\ \text{W m}^{-2}.$$

With the measured 5772 K the same steps give 1361 W m^{-2}, the value measured by satellites (Kopp and Lean, 2011). The sheet's 1368 W m^{-2} is an older measurement of the same quantity. The dilution with distance, $S \propto 1/d^2$, is the inverse-square law used for the planets in Chapter 16, Eq. (16.1).

14.9 Questions that come up

A question that comes up

The sheet calls 14 °C the Earth's "black-body temperature". Later the Earth's black-body temperature is 255 K (−18 °C). Which is right?
They are two different things. 14 °C (287 K) is the average temperature of the Earth's *surface*, and Question 2b treats that surface as a black body. 255 K is the temperature the *planet as a whole* must have to send back to space as much energy as it absorbs (Questions 3c and 4d). The 33 K difference between them is the greenhouse effect (§ 17.7).

A question that comes up

My calculator shows 6.3988E7. How do I write that?
"E7" means "times ten to the power 7": 6.3988×10^7 W m^{-2}, or 63 988 000 W m^{-2}. Round to three or four significant figures, 6.40×10^7 W m^{-2}.

14.10 Try it yourself: Practice Set 14

Practice Set 14: the Stefan–Boltzmann law

Use $\sigma = 5.67 \times 10^{-8}\,\mathrm{W\,m^{-2}\ K^{-4}}$ and $^\circ\mathrm{C} = \mathrm{K} - 273.16$.

14.1 How much does a black body at 0 °C emit?

14.2 By what percentage does the emission rise if the temperature rises by 10 %?

14.3 What temperature must a black body have to emit $1000\,\mathrm{W\,m^{-2}}$? Give kelvin and Celsius.

14.4 The Earth's surface warms from 287.16 K to 288.16 K. By how much does its emission increase?

14.5 Using the worked example in § 14.8, explain in one sentence why the Earth receives only about $1368\,\mathrm{W\,m^{-2}}$ from a Sun that emits $6.4 \times 10^{7}\,\mathrm{W\,m^{-2}}$.

Answers to Practice Set 14

14.1 $E = 5.67 \times 10^{-8} \times 273.16^4 = 315.7\,\mathrm{W\,m^{-2}}$, Eq. (14.1).

14.2 $1.1^4 = 1.464$, so the emission rises by 46 %, Eq. (14.2).

14.3 $T = (E/\sigma)^{1/4} = (1000/5.67 \times 10^{-8})^{1/4} = 364.4\,\mathrm{K} = 91.3\,^\circ\mathrm{C}$. The rearrangement is Eq. (17.3).

14.4 $5.67 \times 10^{-8} \times (288.16^4 - 287.16^4) = 5.40\,\mathrm{W\,m^{-2}}$. One kelvin of warming adds about $5.4\,\mathrm{W\,m^{-2}}$ of emission; this is why warming is a strong brake on further warming.

14.5 The Sun's energy spreads over a sphere whose area grows with the square of the distance; at 150 million kilometres each square metre receives only $(R_\odot/d)^2 = 1/46\,000$ of what each square metre of the Sun emits.

Chapter 15

Plotting three planetary spectra

This chapter answers the graph of Question 3 and Questions 3a and 3b of the sheet.

15.1 The data

The sheet gives the spectral emission of three black bodies, labelled Data A, B and C, at fourteen wavelengths from 4 to 30 μm (Table 15.1). Each column is one planet; the task is to plot all three on one graph, read off where each peaks, and use Wien's law to find each planet's temperature.

Table 15.1. The data of Question 3, as printed on the sheet (EGM305 module team, 2025a). The largest value in each column is in bold. The same table, with the values computed from the Planck curve, is in Appendix B.3.

Wavelength (μm)	Data A	Data B	Data C
4	26.199	1.3338	0.11992
6	386.28	53.056	10.648
8	970.65	218.86	65.619
10	1313.3	398.14	151.8
12	**1363.5**	502.49	224.7
14	1248.6	**527.86**	264.23
16	1074.3	502.41	**273.39**
18	896.25	452.58	262.51
20	737.1	395.22	241.03
22	603.1	339.25	215.38
24	493.45	288.62	189.36
26	404.91	244.61	164.92
28	333.76	207.16	142.9
30	276.58	175.67	123.56

A note on the units of the data

The sheet heads the columns "Irradiance, W m^{-2}". Strictly these are *spectral* values: emission per square metre *per micrometre of wavelength* (W m^{-2} μm^{-1}), on an arbitrary scale. Recomputing them shows that each column is a Planck curve, Eq. (12.3), multiplied by the same constant (about 316) for all three (Appendix B.3). The units do not affect any answer, because the questions only use *where* each curve peaks and how the curves compare.

15.2 How to plot the graph

Step 1. Axes. Wavelength (μm) along the bottom, from 0 to 30. Emission up the side, from 0 to at least 1400, since the largest value is 1363.5.

Step 2. Scale. Choose round steps that use most of the paper: for example 2 μm per major division across and 100 or 200 per division up.

Step 3. Plot each data set with its own symbol (for example crosses for A, circles for B and triangles for C), so that the graph still reads correctly in black and white.

Step 4. Join the points with a smooth curve, not with straight lines. The underlying spectrum is smooth (§ 12.7); straight segments would put the peak exactly on a data point, which hides the fact that the true peak may lie between two points.

Step 5. Label each curve (A, B, C), the axes with quantity and unit, and give the graph a title.




Figure 15.1. The graph of Question 3. Symbols are the values on the sheet; the smooth lines are the Planck curves that pass through them (254.1, 210.0 and 184.1 K; Appendix B.3). The dotted lines mark the highest data point of each set, which is the answer to Question 3a.

15.3 Question 3a: the wavelength of the peak

The question on the sheet

At what wavelength does the maximum intensity occur for each data set?

Method. Find the largest number in each column of Table 15.1 (in bold), or the highest point of each curve in Figure 15.1, and read its wavelength.

Answer to Question 3a

Data A: $\lambda_{\max} = \mathbf{12}\,\mu$m (1363.5). Data B: $\lambda_{\max} = \mathbf{14}\,\mu$m (527.86). Data C: $\lambda_{\max} = \mathbf{16}\,\mu$m (273.39).

Check. The curve with the highest peak (A) peaks furthest to the left, and the lowest (C) furthest to the right. That is what Wien's law and the Stefan–Boltzmann law together predict (Figure 13.1): the hottest body is both the brightest and the shortest in wavelength. All three peaks lie in the thermal infrared, as they must for planets far cooler than the Sun.

15.4 Question 3b: the temperatures of the three planets

The question on the sheet

Using Wien's law, calculate the black-body temperature for the three planets.

Method. Use the rearranged Wien's law, Eq. (13.2), $T = C/\lambda_{\max}$, with $C = 2898\,\mu\text{m K}$ and the peaks from Question 3a:

$$T_A = \frac{2898}{12} = 241.5\text{ K},$$
$$T_B = \frac{2898}{14} = 207.0\text{ K},$$
$$T_C = \frac{2898}{16} = 181.1\text{ K}.$$

Answer to Question 3b

$T_A = \mathbf{241.5\,K}$ ($-31.7\,^\circ$C), $T_B = \mathbf{207.0\,K}$ ($-66.2\,^\circ$C), $T_C = \mathbf{181.1\,K}$ ($-92.0\,^\circ$C).

Check. The order is right: A is the warmest and C the coldest, matching the heights of the curves. The values are cold, which is reasonable for planets heated by a distant Sun; Chapter 16 identifies which planet each curve belongs to.

Why these temperatures are a little too low

The data are given only every $2\,\mu$m, so the highest *data point* is not exactly the *peak* of the curve. Fitting the Planck curve to each column (Appendix B.3) shows that the true peaks are at 11.4, 13.8 and 15.7 μm, and the true temperatures are **254.1, 210.0 and 184.1 K**. Reading the nearest point puts every peak slightly too far to the right, and so every temperature slightly too low; the error is largest for A (12 instead of 11.4 μm, so 241.5 instead of 254 K). The answers above are the right answers to the question as set; the true values matter in Chapter 16, where they match the calculated planetary temperatures almost exactly.

A question that comes up

Could I find the peak more accurately from the data?
Partly. Fitting a parabola through the three highest points of a column and taking its top moves the A peak from 12 to 11.6 μm (249.7 K), closer to the truth. It does much less for B and C, because a Planck curve is lopsided (steep on the short-wave side, with a long tail) while a parabola is symmetric. The reliable way is to fit the Planck curve itself, as in Appendix B.3, or to take readings closer together near the peak.

15.5 What the graph shows

- **The curves never cross.** At every wavelength the warmer body emits more. This is a property of all Planck curves: heating a black body raises its emission at every wavelength.
- **The area under each curve** is the total emission, which grows as T^4, Eq. (14.1). With the true temperatures, A emits $(254.1/184.1)^4 = 3.6$ times as much in total as C.
- **The gap between the curves is largest on the short-wave side.** At 10 μm A emits 8.7 times as much as C; at 30 μm only 2.2 times. Short wavelengths are the most sensitive to temperature, which is why the peak moves (Wien's law) and why satellites measuring fires use short infrared bands (§ 13.4).

15.6 Try it yourself: Practice Set 15

Practice Set 15: reading spectra

15.1 If you had readings every 0.2 μm near the top of curve A, you would find its peak at 11.4 μm. What temperature does that give?

15.2 Using Table 15.1, how many times more does A emit than C at 10 μm? At 30 μm? Why are the two ratios different?

15.3 A fourth planet's spectrum peaks near 25 μm. What is its temperature, and which planet of Table 16.1 might it be?

15.4 Explain in one sentence why two Planck curves for different temperatures can never cross.

Answers to Practice Set 15

15.1 $T = 2898/11.4 = 254.2$ K, Eq. (13.2). That is the Earth's calculated black-body temperature (Question 3c).

15.2 $1313.3/151.8 = 8.7$ at 10 μm; $276.58/123.56 = 2.2$ at 30 μm. Emission on the short-wave side of the peak depends far more strongly on temperature than on the long-wave tail, Eq. (12.3).

15.3 $T = 2898/25 = 116$ K: Jupiter, whose calculated black-body temperature is 110 K (Question 3c).

15.4 Raising T in Eq. (12.3) makes the exponential in the denominator smaller at every wavelength, so the emission rises at every wavelength and the warmer curve lies above the cooler one everywhere.

Chapter 16

The temperatures of the planets

This chapter answers Questions 3c, 3d and 3e of the sheet.

16.1 A planet in balance

A planet is heated by the Sun and cools by radiating to space. If it absorbed more than it emitted it would warm up; warming raises its emission as T^4, Eq. (14.1), until the two are equal. A planet that has settled is therefore **in radiative balance**:

$$\text{solar energy absorbed} = \text{infrared energy emitted}.$$

The temperature at which this happens is the planet's **equilibrium** or **black-body temperature** (EGM305 module team, 2020, slides 43–44). To calculate it we need three things: how much sunlight arrives (the solar constant S), how much of it is reflected (the albedo a), and the geometry of a sphere.

16.2 Sunlight at each planet: the inverse-square law

Sunlight spreads out over ever larger spheres as it travels from the Sun (§ 14.8). The area of a sphere grows with the square of its radius, so the energy reaching each square metre falls with the square of the distance:

$$S = \frac{S_E}{d^2}. \tag{16.1}$$

where

S	solar constant at the planet: sunlight per square metre facing the Sun, at the top of the atmosphere ($\mathrm{W\,m^{-2}}$)	calculated; listed on the sheet
S_E	solar constant at the Earth ($\mathrm{W\,m^{-2}}$)	1368, on the sheet
d	planet's distance from the Sun, in astronomical units (AU; Earth–Sun = 1)	listed on the sheet

The sheet's values follow this law exactly. For Mars, $d = 1.52\,\mathrm{AU}$, so $S = 1368/1.52^2 = 1368/2.310 = 592\,\mathrm{W\,m^{-2}}$: at one and a half times the distance, Mars receives only 43 % of the Earth's sunlight. For Mercury, $1368/0.39^2 = 8994\,\mathrm{W\,m^{-2}}$, six and a half times the Earth's.

16.3 Albedo

Definition

The **albedo** a of a surface or a planet is the fraction of the sunlight falling on it that is reflected, from 0 (absorbs everything) to 1 (reflects everything). The fraction absorbed is $1 - a$ (EGM305 module team, 2020, slides 36 and 67).

Fresh snow and thick cloud reflect most sunlight (albedo up to 0.8–0.9); forests, soils and especially open water reflect little (water below about 0.1 with the Sun high in the sky) (Ahrens and Henson, 2022, ch. 2). A planet's overall albedo, called its *Bond albedo*, is the average over its whole sunlit side, cloud included: about 0.3 for the Earth (Forster et al., 2021).

16.4 The black-body temperature of a planet




Figure 16.1. Why the solar constant is divided by four. A planet intercepts sunlight over a disc of area πR^2 (its shadow) but radiates from its whole surface, $4\pi R^2$, day and night side alike. Averaged over the sphere, each square metre receives $S/4$.

Set the energy absorbed equal to the energy emitted, for a planet of radius R (Figure 16.1):

$$\begin{aligned} \text{absorbed} &= S\,(1-a) \times \pi R^2, && \text{(sunlight on the disc, minus what is reflected)} \\ \text{emitted} &= \sigma T^4 \times 4\pi R^2. && \text{(Stefan–Boltzmann law, Eq. (14.1), over the whole sphere)} \end{aligned}$$

Setting them equal and cancelling πR^2 from both sides, the size of the planet drops out:

$$S\,(1-a) = 4\sigma T^4 \quad \Longrightarrow \quad T^4 = \frac{S\,(1-a)}{4\sigma},$$

and taking the fourth root of both sides gives the formula printed as a hint on the sheet:

$$T = \left[\frac{S\,(1-a)}{4\sigma}\right]^{1/4}. \tag{16.2}$$

where

T	black-body (equilibrium) temperature of the planet (K)	calculated
S	solar constant at the planet (W m^{-2})	on the sheet; Eq. (16.1)
a	planet's albedo (0 to 1)	on the sheet
σ	Stefan–Boltzmann constant	5.67×10^{-8} W m^{-2} K^{-4}
4	ratio of the sphere's area to the disc's, Figure 16.1	geometry

A fourth root is a square root taken twice; on a calculator use $x^{0.25}$ or press $\sqrt{\ }$ twice.

16.5 Question 3c: calculating the temperatures

The question on the sheet

Using the table below, calculate the black-body temperatures T for the following planets (hint: $T = [S(1-a)/(4\sigma)]^{1/4}$).

Worked example: the Earth, step by step

Step 1. Absorbed sunlight per square metre of disc: $S(1-a) = 1368 \times (1-0.3) = 1368 \times 0.7 = 957.6\,\mathrm{W\,m^{-2}}$.

Step 2. Divide by $4\sigma = 4 \times 5.67 \times 10^{-8} = 2.268 \times 10^{-7}$: $957.6 / 2.268 \times 10^{-7} = 4.2222 \times 10^{9}\,\mathrm{K^4}$.

Step 3. Square root: $\sqrt{4.2222 \times 10^{9}} = 64\,979$.

Step 4. Square root again: $\sqrt{64\,979} = 254.9\,\mathrm{K}$.

So $T_{\mathrm{Earth}} = 254.9\,\mathrm{K}$, or $254.9 - 273.16 = -18.3\,^{\circ}\mathrm{C}$.

The same four steps for every planet give Table 16.1.

Table 16.1. Question 3c completed. The first four columns and the last but one are from the sheet (EGM305 module team, 2025a); $S(1-a)$ and T are calculated with Eq. (16.2); the last column is the actual minus the calculated temperature.

Planet	d (AU)	S ($\mathrm{W\,m^{-2}}$)	a	$S(1-a)$	T calc. (K)	actual (K)	difference
Mercury	0.39	8994	0.06	8454.4	**439.4**	440	+0.6
Venus	0.72	2639	0.90	263.9	**184.7**	737	+552.3
Earth	1.00	1368	0.30	957.6	**254.9**	288	+33.1
Mars	1.52	592	0.25	444.0	**210.3**	210	−0.3
Jupiter	5.18	51	0.34	33.7	**110.4**	152	+41.6

Answer to Question 3c

Mercury **439 K**, Venus **185 K**, Earth **255 K**, Mars **210 K**, Jupiter **110 K** (to the nearest kelvin; Table 16.1).

Check. The temperatures fall with distance from the Sun, as expected, with one exception: Venus is calculated colder than the Earth although it is closer to the Sun. Question 3e asks why. NASA's fact sheets, which use slightly different solar constants and albedos, give black-body temperatures of 439.6 K for Mercury, 254.0 K for the Earth, 209.8 K for Mars and 109.9 K for Jupiter (Williams, 2024), all within a kelvin or so of ours.

16.6 Question 3d: which spectrum belongs to which planet?

The question on the sheet

Using the temperatures from the table above, link the data sets you plotted previously with the planet.

Method. Compare the temperatures from the spectra (Question 3b, § 15.4) with the calculated temperatures in Table 16.1, and pair each data set with the closest planet.

	From the peak (Q3b)	From the full curve (§ 15.4)	Closest planet (Q3c)
Data A	241.5 K	254.1 K	Earth, 254.9 K
Data B	207.0 K	210.0 K	Mars, 210.3 K
Data C	181.1 K	184.1 K	Venus, 184.7 K

Answer to Question 3d

Data A: **Earth**. Data B: **Mars**. Data C: **Venus**.

Check. No other pairing works: Mercury (439 K) would peak near 6.6 μm and Jupiter (110 K) near 26 μm, Eq. (13.1), both outside the three peaks we found. Fitted to the full curves, the three data sets match the calculated temperatures to within one kelvin: the sheet's spectra were made from exactly these planetary temperatures.

16.7 Question 3e: why the real temperatures differ

The question on the sheet

Using the table you constructed, explain: (i) why Mars is theoretically colder than Earth; (ii) why Venus is theoretically colder than Earth; (iii) what the actual surface temperature of Venus is, how you explain the difference between the theoretical and actual temperature, and which process could be causing this; (iv) of all these planets, Mercury is the only one that has an actual average temperature which is the same as its black-body temperature. Why is that?




Figure 16.2. Calculated and actual temperatures from Table 16.1. Where the bars match (Mercury, Mars) there is little or no atmosphere; where the actual temperature is higher, an atmosphere traps heat (Venus, Earth) or the planet has an internal source of heat (Jupiter).

(i) Why Mars is theoretically colder than the Earth

Mars is 1.52 times as far from the Sun, so by the inverse-square law, Eq. (16.1), it receives only $1/1.52^2 = 43\,\%$ of the Earth's sunlight: 592 against 1368 W m^{-2}. Its albedo is a little lower (0.25 against 0.30), so it absorbs a slightly larger fraction, but that cannot make up for receiving less than half as much. It absorbs 444 W m^{-2} (per square metre of disc) against the Earth's 957.6, and T scales as the fourth root of that: $(444/957.6)^{1/4} = 0.825$, so $0.825 \times 254.9 = 210$ K.

(ii) Why Venus is theoretically colder than the Earth

Venus is closer to the Sun and receives almost twice the Earth's sunlight (2639 W m^{-2}). But it is wrapped in a permanent, bright cloud deck that reflects most of that light: with the sheet's albedo of 0.9 it absorbs only 10 %, which is 263.9 W m^{-2}, less than a third of what the Earth absorbs and even less than Mars. **Albedo, not distance, controls its black-body temperature.**

(iii) The actual surface temperature of Venus

The actual average surface temperature of Venus is about **737 K (464 °C)** (Williams, 2024), hot enough to melt lead, and almost the same by day and by night. That is 552 K hotter than its black-body temperature. The process responsible is the **greenhouse effect**. Venus has a very thick atmosphere (a surface pressure of 92 bar, about ninety times the Earth's), made almost entirely of carbon dioxide, which absorbs the infrared radiation emitted by the surface (Chapter 17). Only the cold upper atmosphere can radiate freely to space, and it does so at roughly the black-body temperature; to push the energy up through such a thick, opaque atmosphere, the surface must be extremely hot (Pierrehumbert, 2011).

(iv) Why Mercury's actual and black-body temperatures agree

Mercury has practically no atmosphere (a surface pressure of less than 10^{-14} bar; Williams, 2024). With no gases to absorb its infrared emission, nothing traps heat, so the surface radiates straight to space exactly as the black-body calculation assumes. Its *average* temperature therefore matches: 440 against 439 K. (The average hides a huge range: with no atmosphere to carry heat around, the sunlit side reaches 590–725 K and the night side is far colder.)

Answer to Question 3e

(i) Mars is further from the Sun (1.52 AU) and receives only 43 % of the Earth's sunlight (inverse-square law), so it absorbs less and is colder, despite its slightly lower albedo.

(ii) Venus has a very high albedo (0.9 on the sheet): its bright clouds reflect most sunlight, so it absorbs less than a third of what the Earth absorbs, although it is closer to the Sun.

(iii) About **737 K (464 °C)**, some 550 K above its black-body temperature. Its dense carbon dioxide atmosphere produces an intense **greenhouse effect**: the surface emission is absorbed by the atmosphere, and the surface must become very hot before the planet can emit to space as much as it absorbs. The black-body calculation leaves the atmosphere out.

(iv) Mercury has almost no atmosphere and so no greenhouse effect: it absorbs and re-radiates directly to space, exactly as Eq. (16.2) assumes.

Corrections to the printed key and sheet for Question 3

1. **Mercury is not the only match.** In the sheet's own table Mars matches too (210 against 210 K). Mars has a very thin atmosphere (about 6 mb, under 1 % of the Earth's) and so only a small greenhouse effect: its measured average is about 214 K (Williams, 2024), a few kelvin above its black-body temperature. Thin or absent atmospheres give a match; thick ones do not.
2. **Venus's albedo.** The sheet's 0.9 is high: the measured Bond albedo is 0.77 (Williams, 2024), which gives a black-body temperature of 227 K. Venus is then still colder than the Earth in theory, and the answers to (ii) and (iii) are unchanged.

3. **"Runaway heating".** The key says Venus's atmosphere "traps heat leading to runaway heating". Venus today is not heating up: it is in balance, absorbing as much as it emits, at a very high surface temperature. The *runaway greenhouse* describes how Venus probably reached that state long ago, as rising temperatures evaporated any oceans and the water vapour trapped still more heat (Kasting, 1988). Use "an intense greenhouse effect" for the present.
4. **Two parts labelled 3d.** The sheet uses "3d" twice; this part calls the second one 3e (§ 11.2).

A question that comes up

Why is Jupiter 42 K warmer than its black-body temperature, if it is so far from the Sun?
Because the Sun is not its only source of heat. Jupiter still releases heat left over from its formation and from slow contraction, and Voyager measurements showed that it emits about 1.7 times as much energy as it absorbs from the Sun (Hanel et al., 1981). Equation (16.2) assumes the Sun is the only heat source, so it underestimates Jupiter. The Earth has an internal source too, but it is tiny: the flow of heat from the Earth's interior averages under $0.1\,\mathrm{W\,m^{-2}}$ (Davies and Davies, 2010), less than one two-thousandth of the $238.7\,\mathrm{W\,m^{-2}}$ of sunlight absorbed (Question 4c).

A question that comes up

The Earth is 33 K warmer than its black-body temperature. Is that also the greenhouse effect?
Yes, and Question 4 measures it directly (§ 17.7).

16.8 Try it yourself: Practice Set 16

Practice Set 16: planetary temperatures

Use $\sigma = 5.67 \times 10^{-8}\,\mathrm{W\,m^{-2}\ K^{-4}}$ and $S_E = 1368\,\mathrm{W\,m^{-2}}$.

16.1 Saturn is 9.54 AU from the Sun and has an albedo of 0.34. Find its solar constant and its black-body temperature.

16.2 What would the Earth's black-body temperature be with an albedo of 0.25? Of 0.35?

16.3 The Moon is at the same distance from the Sun as the Earth and has an albedo of about 0.11. What is its black-body temperature?

16.4 What albedo would give the Earth a black-body temperature of 288 K, its actual average? What does your answer tell you?

16.5 If the Earth were moved to half its present distance from the Sun, keeping its albedo of 0.3, what would its black-body temperature be? Find it two ways.

Answers to Practice Set 16

16.1 $S = 1368/9.54^2 = 15.0\,\mathrm{W\,m^{-2}}$, Eq. (16.1); $T = [15.0 \times 0.66/(4 \times 5.67 \times 10^{-8})]^{1/4} = 81.3\,\mathrm{K}$, Eq. (16.2).

16.2 $a = 0.25$: 259.3 K. $a = 0.35$: 250.2 K. A change of 0.1 in albedo changes the temperature by about 9 K.

16.3 $T = [1368 \times 0.89/(2.268 \times 10^{-7})]^{1/4} = 270.7\,\mathrm{K}$. With no atmosphere, the Moon's average is close to this, but its days are scorching and its nights extremely cold (§ 17.8).

16.4 From Eq. (16.2), $a = 1 - 4\sigma T^4/S = 1 - (2.268 \times 10^{-7} \times 288^4)/1368 = -0.14$. A negative albedo is impossible (no planet can absorb more than all the light falling on it). So no albedo can explain the Earth's 288 K: something other than sunlight absorption, the greenhouse effect, must be keeping the surface warmer.

16.5 At half the distance $S = 4 \times 1368 = 5472\,\mathrm{W\,m^{-2}}$, so $T = [5472 \times 0.7/(2.268 \times 10^{-7})]^{1/4} = 360.5\,\mathrm{K}$. Second way: S is four times larger and $T \propto S^{1/4}$, so T is $4^{1/4} = \sqrt{2} = 1.414$ times larger: $1.414 \times 254.9 = 360.5\,\mathrm{K}$.

Chapter 17

The Earth's energy balance and the greenhouse effect

This chapter answers Questions 4a to 4g of the sheet.

17.1 What Question 4 sets out to do

If the Earth had no atmosphere there would be no greenhouse effect, and the planet would be colder. Question 4 estimates *how much* colder, by working out the temperature an airless Earth would need to send back to space all the sunlight it absorbs, and comparing it with the temperature actually observed. It is the calculation of Question 3c for the Earth (§ 16.5), taken one step at a time, with the energy flows made explicit (Figure 17.1).

17.2 Question 4a: how much is absorbed?

The question on the sheet

If the global albedo of the Earth is 30 %, how much solar radiation is absorbed by the Earth?

Whatever is not reflected is absorbed (§ 16.3). As a fraction of the incoming sunlight,

$$\text{fraction absorbed} = 1 - a, \qquad \text{fraction reflected} = a. \tag{17.1}$$

where

a	albedo, as a fraction	0.30 (30 %), on the sheet

Answer to Question 4a

$1 - 0.30 = 0.70$: the Earth absorbs **70 %** of the solar radiation it receives.

17.3 Question 4b: sunlight facing the Sun

The question on the sheet

If the Earth receives 1368 W m^{-2} solar radiation at the equator, how much is reflected back into space, and how much is absorbed by the Earth?

Method. Multiply the incoming radiation by the fractions of Eq. (17.1):

$$\text{reflected} = a \times S = 0.30 \times 1368 = 410.4 \text{ W m}^{-2},$$
$$\text{absorbed} = (1 - a) \times S = 0.70 \times 1368 = 957.6 \text{ W m}^{-2}.$$

Check: $410.4 + 957.6 = 1368$ W m^{-2}. Every watt is accounted for.

Answer to Question 4b

Reflected: $0.3 \times 1368 = \mathbf{410.4}\,\mathrm{W\,m^{-2}}$. Absorbed: $0.7 \times 1368 = \mathbf{957.6}\,\mathrm{W\,m^{-2}}$.

A correction to the printed key: 410.4, not 41.04

The key gives the reflected radiation as "$0.3 \times 1368 = 41.04\,\mathrm{W/m^2}$". The decimal point has slipped one place: $0.3 \times 1368 = 410.4\,\mathrm{W\,m^{-2}}$. The check above shows why 41.04 cannot be right: $41.04 + 957.6 = 998.6$, not 1368, so $369\,\mathrm{W\,m^{-2}}$ would be unaccounted for.

A question that comes up

Is $1368\,\mathrm{W\,m^{-2}}$ really what arrives "at the equator"?
Not quite. The solar constant is the sunlight falling on a surface *at right angles to the Sun's rays, at the top of the atmosphere*, at the Earth's average distance from the Sun (Kopp and Lean, 2011). It arrives at the equator only at noon when the Sun is overhead there (around the equinoxes), and then only above the atmosphere: some is absorbed and scattered on the way down. Today's satellite value is $1361\,\mathrm{W\,m^{-2}}$; the sheet's 1368 is an older figure, and the difference changes no conclusion.

17.4 Question 4b′: why the average is only 341 $\mathrm{W\,m^{-2}}$

The question on the sheet

The average incoming solar radiation for planet Earth is only $341\,\mathrm{W\,m^{-2}}$. Why do you think that is?

The $1368\,\mathrm{W\,m^{-2}}$ is for a surface squarely facing the Sun. Averaged over the whole Earth and over day and night, each square metre receives a quarter of that, for two reasons that each contribute a factor of two:

1. **Night.** At any moment half the Earth faces away from the Sun and receives nothing.
2. **Slanting rays.** On the day side, only the point directly under the Sun receives the full beam. Everywhere else the Sun is lower in the sky and its rays strike the ground at a slant, spreading the same energy over a larger area (this is Question 5, Chapter 18). Averaged over the sunlit hemisphere, this halves the energy per square metre.

Together: the Earth intercepts sunlight over a disc of area πR^2 but has a surface area of $4\pi R^2$ (Figure 16.1), so

$$\bar{S} = \frac{S}{4} = \frac{1368}{4} = 342\ \mathrm{W\,m^{-2}}. \tag{17.2}$$

where

$\bar{S}$	incoming solar radiation averaged over the whole Earth and over day and night ($\mathrm{W\,m^{-2}}$)	calculated
S	solar constant ($\mathrm{W\,m^{-2}}$)	1368, on the sheet

Answer to Question 4b′

The 1368 $\mathrm{W\,m^{-2}}$ falls only on a surface facing the Sun directly. At any time **half the Earth is in darkness**, and over the sunlit half **the Sun's rays arrive at a slant** except where the Sun is overhead, spreading the energy over a larger area. The Earth catches sunlight on a disc (πR^2) but spreads it over a sphere ($4\pi R^2$), so the global average is a quarter: $1368/4 = 342\,\mathrm{W\,m^{-2}}$, about $341\,\mathrm{W\,m^{-2}}$.

Why 341 and not 342? Global energy budgets built from satellite measurements give $341.3\,\mathrm{W\,m^{-2}}$ (Trenberth et al., 2009), a quarter of the solar constant they used ($1365\,\mathrm{W\,m^{-2}}$). The difference from 342 is only in the value of the solar constant.

17.5 Question 4c: the global average

The question on the sheet

If we take the average incoming solar radiation, then how much is reflected back into space, and how much is absorbed by the Earth?

Method. The same fractions, Eq. (17.1), applied to the average of $341\,\mathrm{W\,m^{-2}}$:

$$\text{reflected} = 0.30 \times 341 = 102.3\ \mathrm{W\,m^{-2}},$$
$$\text{absorbed} = 0.70 \times 341 = 238.7\ \mathrm{W\,m^{-2}}.$$

Answer to Question 4c

Reflected: $0.3 \times 341 = \mathbf{102.3}\,\mathrm{W\,m^{-2}}$. Absorbed: $0.7 \times 341 = \mathbf{238.7}\,\mathrm{W\,m^{-2}}$.

Check. Satellite-based budgets give about $102\,\mathrm{W\,m^{-2}}$ reflected and $239\,\mathrm{W\,m^{-2}}$ absorbed (Trenberth et al., 2009): the sheet's simple numbers reproduce the measured global budget.




Figure 17.1. The energy flows of Question 4, in $\mathrm{W\,m^{-2}}$ averaged over the whole Earth, for an Earth without an atmosphere. Arrow widths are proportional to the flows. In balance, the infrared emitted equals the sunlight absorbed; the temperature needed to emit it is the answer to Question 4d.

17.6 Question 4d: the temperature of an airless Earth

The question on the sheet

Earth must lose exactly as much energy as it gains, therefore the Earth must emit all the solar radiation it absorbs. However, if there is no atmosphere this happens very quickly (there is no greenhouse effect). First reorganise the Stefan–Boltzmann equation so that you can calculate the temperature in terms of the intensity of radiation ($T = \dots$). Now calculate what the temperature of the Earth would be if all the radiation it absorbed from the Sun was released at the same time (use the data you calculated in 4c). Convert your answer from kelvin to Celsius.

Rearranging the law. Start from Eq. (14.1), $E = \sigma T^4$, and isolate T in two moves:

Step 1. Divide both sides by σ: $T^4 = E/\sigma$.

Step 2. Take the fourth root of both sides (the inverse of raising to the fourth power):

$$T = \left(\frac{E}{\sigma}\right)^{1/4}. \tag{17.3}$$

where

T	temperature of the emitting surface (K)	calculated
E	radiation emitted per square metre ($\mathrm{W\,m^{-2}}$)	238.7, from Question 4c
σ	Stefan–Boltzmann constant	$5.67 \times 10^{-8}\,\mathrm{W\,m^{-2}\ K^{-4}}$

Calculation. In balance the Earth emits exactly what it absorbs, so $E = 238.7\,\mathrm{W\,m^{-2}}$:

Step 3. $E/\sigma = 238.7/5.67 \times 10^{-8} = 4.2099 \times 10^9\,\mathrm{K^4}$.

Step 4. First square root: $\sqrt{4.2099 \times 10^9} = 64\,884$; second square root: $\sqrt{64\,884} = 254.7\,\mathrm{K}$.

Step 5. In Celsius: $254.7 - 273.16 = -18.5\,^\circ\mathrm{C}$.

Answer to Question 4d

$T = (E/\sigma)^{1/4}$. $T = (238.7/5.67 \times 10^{-8})^{1/4} = \mathbf{254.7\,K}$, which is $\mathbf{-18.5}\,^\circ\mathrm{C}$.

Check. This is the same calculation as the Earth's row of Question 3c, written differently: there $T = [1368 \times 0.7/(4\sigma)]^{1/4} = 254.9\,\mathrm{K}$. The 0.2 K difference comes only from using 341 instead of $1368/4 = 342\,\mathrm{W\,m^{-2}}$. And it is the temperature of curve A in Question 3 (254.1 K): curve A is the spectrum of an Earth without a greenhouse effect.

17.7 Question 4e: what the atmosphere does

The question on the sheet

The actual average global temperature of the Earth between 1961 and 1990 is 14.0 °C (World Meteorological Organization). What difference does the atmosphere make to the global temperature of the Earth?

The size of the effect. The difference between the actual and the airless temperature is

$$14.0 - (-18.5) = 32.5\ ^\circ\mathrm{C},$$

which is also a difference of 32.5 K, since a step of one degree is the same size on both scales. The atmosphere keeps the Earth's surface about **33** °C **warmer** than it would otherwise be: above the freezing point of water instead of well below it.

How it does it. The atmosphere treats the two kinds of radiation differently (Figure 17.2) (EGM305 module team, 2020, slides 45–54):

1. **Sunlight (shortwave) passes through** largely unabsorbed and warms the surface.
2. **The surface emits infrared (longwave)**, Chapter 13, peaking near 10 μm.
3. **Greenhouse gases absorb much of that infrared**: water vapour, carbon dioxide, methane, nitrous oxide and ozone. These are molecules made of three or more atoms (or of two different atoms) whose vibrations and rotations can absorb and emit infrared; the main gases of the air, nitrogen and oxygen, cannot.
4. **The warmed atmosphere radiates in all directions**, including back down to the surface. The surface therefore receives the sunlight *plus* this downward infrared, about 333 W m^{-2} on average (Trenberth et al., 2009), and settles at a higher temperature.
5. **The planet as a whole still balances.** What escapes to space comes mostly from the cold upper atmosphere, and it still totals about 239 W m^{-2}: the Earth seen from space radiates like a body at 255 K, even though its surface is at 288 K.




Figure 17.2. The greenhouse effect. Sunlight passes through the atmosphere and warms the surface; the surface's infrared is absorbed by greenhouse gases, which radiate both to space and back down. The extra downward infrared keeps the surface warmer than the 255 K the planet radiates at from space.

Answer to Question 4e

Without an atmosphere the Earth would average −18.5 °C; with it, 14.0 °C. The atmosphere's **greenhouse effect** raises the average surface temperature by about **32.5** °C (about 33 K). Greenhouse gases absorb the infrared emitted by the surface and radiate part of it back down, so the surface receives more energy than sunlight alone provides and stays at a habitable temperature, above the freezing point of water.

A correction to the printed key: what greenhouse gases trap

The key says the greenhouse effect "prevents all solar energy escaping to space". Two parts of that need changing. Greenhouse gases act mainly on the *infrared* emitted by the Earth, not on incoming sunlight, which passes through largely unabsorbed. And they do not stop energy escaping: in balance the Earth still sends to space exactly as much as it absorbs (238.7 W m^{-2}). What they change is *where* it is emitted from: from the cold upper atmosphere instead of the warm surface. Say: "greenhouse gases absorb the infrared emitted by the surface and re-emit

part of it back down, warming the surface by about 33 K".

Worked example: a one-layer greenhouse

Model the atmosphere as a single layer that lets all sunlight through and absorbs all the surface's infrared. In balance, the layer must emit $238.7\,\mathrm{W\,m^{-2}}$ to space; it emits the same amount downward, so the surface receives $238.7 + 238.7 = 477.4\,\mathrm{W\,m^{-2}}$ and must emit that:

$$T_{\text{surface}} = \left(\frac{2 \times 238.7}{5.67 \times 10^{-8}}\right)^{1/4} = 2^{1/4} \times 254.7 = 303\ \mathrm{K}\ (30\ ^\circ\mathrm{C}).$$

This overshoots the real 288 K, because the real atmosphere lets some infrared escape directly through the "atmospheric window" near 8–12 μm (EGM305 module team, 2020, slide 53) and because rising air and evaporation carry heat up from the surface. But it shows the mechanism, and why a thicker greenhouse atmosphere, like that of Venus (§ 16.7), makes the surface hotter still.

A question that comes up

Why is it called a "greenhouse" effect?
By analogy, and the analogy is loose (EGM305 module team, 2020, slide 46). A real greenhouse stays warm mainly because its glass stops the warm air inside from rising away and mixing with cooler air. The atmosphere warms the surface by radiation, as described above.

17.8 Question 4f: day and night without an atmosphere

The question on the sheet

What effect would no atmosphere have on the variation in temperature on the Earth over the course of a day, compared with an Earth with an atmosphere?

Without an atmosphere, nothing would slow the loss of heat at night or soften the heating by day.

- **At night** the surface would radiate straight to space, with no downward infrared from greenhouse gases and cloud to replace part of the loss (§ 17.7), so it would cool very quickly and very far.
- **By day** there would be no cloud to reflect sunlight and no air to absorb part of it, and no winds to carry heat away, so the sunlit surface would heat strongly.
- **The atmosphere (and the ocean beneath it) also stores and moves heat**: from day to night and from warm regions to cold ones. Without it, each place would respond only to its own sunshine.

The Moon shows the result: at the same distance from the Sun but with no atmosphere, its equatorial surface swings from near 390 K at midday to below 100 K before dawn (Williams et al., 2017), while the Earth's global average daily range is only about 10 K (Williams, 2024).

Answer to Question 4f

The **daily range of temperature would be far larger**: very hot days and very cold nights. The atmosphere reduces the range by radiating infrared back to the surface at night (and through cloud), by reflecting and absorbing part of the sunlight by day, and by storing and moving heat. The airless Moon swings by about 300 K between day and night.

17.9 Question 4g: the assumptions

The question on the sheet

What assumption have we made about the solid Earth for this calculation?

Every model rests on assumptions; naming them is part of using it well. For Question 4d we assumed that the Earth:

1. **has no source of heat of its own**: all its energy comes from the Sun. The heat flowing from the Earth's interior is real but tiny, under $0.1\,\mathrm{W\,m^{-2}}$ on average (Davies and Davies, 2010), so this assumption costs almost nothing (compare Jupiter, § 16.7);
2. **is a perfect black body**, emitting σT^4 with an emissivity of 1 (§ 12.6);
3. **has one uniform temperature** everywhere, by day and by night (in reality the equator and the poles differ by tens of degrees);
4. **is in equilibrium**, emitting at once all it absorbs and storing nothing (the question's "released at the same time");
5. **keeps an albedo of 0.30** even without an atmosphere. In fact much of that reflection is by cloud, which an airless Earth would not have; with a Moon-like albedo of about 0.1 the airless Earth would be nearer 270 K (Practice Set 16, item 16.3).

Answer to Question 4g

That the solid Earth has **no internal source of heat**, so that all the energy it radiates comes from the Sun; and that it behaves as a **black body at a single, uniform temperature in equilibrium**, radiating away immediately everything it absorbs.

A correction to the printed key: the Earth does radiate

The key's answer is "That Earth does not radiate its own heat. It is assumed that 100 % of energy is provided by the Sun." The second sentence is the assumption; the first, read literally, contradicts the whole calculation, in which the Earth radiates $238.7\,\mathrm{W\,m^{-2}}$. What is meant is that the Earth has no internal heat *source*: it radiates only energy that came from the Sun.

17.10 Try it yourself: Practice Set 17

Practice Set 17: the energy balance

Use $\sigma = 5.67 \times 10^{-8}\,\mathrm{W\,m^{-2}\ K^{-4}}$ and $°\mathrm{C} = \mathrm{K} - 273.16$.

17.1 Repeat Questions 4b′ to 4d with today's solar constant, $1361\,\mathrm{W\,m^{-2}}$, and an albedo of 0.30.

17.2 If the Earth's albedo rose from 0.30 to 0.32 (more cloud, or more ice), what would its black-body temperature become?

17.3 The Earth's surface at 14 °C emits $385.5\,\mathrm{W\,m^{-2}}$ (Question 2b), but only $238.7\,\mathrm{W\,m^{-2}}$ leaves the planet. How large is the difference, and what is it a measure of?

17.4 Using the one-layer greenhouse of § 17.7, what would the surface temperature be? Why is the real surface cooler than this?

Answers to Practice Set 17

17.1 $\bar{S} = 1361/4 = 340.25\,\mathrm{W\,m^{-2}}$; absorbed $0.7 \times 340.25 = 238.2\,\mathrm{W\,m^{-2}}$; $T = (238.2/5.67 \times 10^{-8})^{1/4} = 254.6\,\mathrm{K} = -18.6\,^{\circ}\mathrm{C}$, Eqs. (17.2) and (17.3).

17.2 $T = [1368 \times 0.68/(4 \times 5.67 \times 10^{-8})]^{1/4} = 253.1\,\mathrm{K}$, about 1.8 K colder. Small changes in albedo matter, which is why cloud and ice feedbacks are important in climate change.

17.3 $385.5 - 238.7 = 146.8\,\mathrm{W\,m^{-2}}$. This is the infrared emitted by the surface that does not escape to space: a measure of the greenhouse effect in energy terms.

17.4 $2^{1/4} \times 254.7 = 303\,\mathrm{K}$ (30 °C). The real atmosphere is not opaque at all infrared wavelengths (the atmospheric window lets some escape directly), and convection and evaporation carry heat upwards from the surface.

Chapter 18

Sun angle, solar intensity and the seasons

This chapter answers Questions 5a, 5b and 5c of the sheet.

18.1 A spread beam is a weaker beam

Shine a torch straight down at a table: it makes a small, bright circle. Tilt it, and the same light spreads into a long, dim oval. The torch gives out the same energy; it is spread over more table, so each square centimetre receives less. Sunlight does the same. When the Sun is overhead its rays strike the ground squarely and their energy is concentrated; when it is low they strike at a slant and the same energy is spread over a larger area (EGM305 module team, 2020, slides 56–57). The **solar intensity**, the energy received per unit area of ground, therefore depends on the angle of the Sun.

18.2 Measuring the spreading

Two angles describe the Sun's position (Figure 18.1):

- the **sun angle** β (also called the solar elevation or altitude), measured up from the horizon: 90° when the Sun is overhead, 0° when it is on the horizon;
- the **zenith angle** θ_z, measured from the vertical (the zenith, the point straight overhead): 0° when the Sun is overhead.

They add up to a right angle:

$$\beta = 90^\circ - \theta_z. \tag{18.1}$$

where

β	sun angle: height of the Sun above the horizon (degrees)	calculated
θ_z	zenith angle: angle of the Sun from the vertical (degrees)	Eq. (18.4)




Figure 18.1. Beam spreading. A beam one unit wide, arriving at sun angle β, covers a strip of ground of length $1/\sin\beta$: the hypotenuse of a right-angled triangle whose side opposite the angle β is the beam width (Appendix B.2.4). Drawn here for $\beta = 35^\circ$, where the beam covers 1.74 units.

The sheet calls the length of ground covered by a beam one unit wide the "surface area" (EGM305 module team, 2025a). From the right-angled triangle in Figure 18.1 (derived in Appendix B.2.4):

$$A = \frac{1}{\sin\beta}. \tag{18.2}$$

where

A	area over which a beam of unit width spreads ("surface area", "beam spreading"; units)	calculated
β	sun angle (degrees)	Eq. (18.1)

When the Sun is overhead, $\beta = 90°$, $\sin 90° = 1$ and $A = 1$ unit: the beam's full energy, 100 %, falls on one unit. In general the energy falling on each unit of ground is

$$I = \frac{100\,\%}{A} = 100\,\% \times \sin\beta. \tag{18.3}$$

where

I	solar intensity, as a percentage of the overhead value, per unit of ground (% per unit)	calculated
A	beam spreading from Eq. (18.2)	

Make sure your calculator is in **degree** mode: sin(83.5) should give 0.9936.

18.3 Where the noon Sun is overhead

Because the Earth's axis is tilted by 23.5° to its orbit, the latitude at which the noon Sun is directly overhead moves through the year (Figure 18.2). That latitude is the Sun's **declination** δ: +23.5° (the Tropic of Cancer) at the June solstice, 0° (the equator) at the equinoxes, and −23.5° (the Tropic of Capricorn) at the December solstice (EGM305 module team, 2020, slide 59). At solar noon, the Sun's zenith angle at latitude φ is the difference between the two latitudes:

$$\theta_z = |\varphi - \delta|. \tag{18.4}$$

where

θ_z	noon zenith angle (degrees)	calculated
φ	latitude of the site (degrees, positive north)	30 or 60, on the sheet
δ	declination: latitude where the noon Sun is overhead (degrees)	+23.5 in June, −23.5 in December

For a site north of the Tropic of Cancer this gives exactly the two rules on the sheet:


June 21 (summer solstice): $\theta_z = \varphi - 23.5°$; December 22 (winter solstice): $\theta_z = \varphi + 23.5°$.

The reason is geometric: walking north by one degree of latitude tilts your local vertical one degree further from the Sun's direction (Appendix B.2.5).

18.4 Question 5a: sketches of the noon Sun

The question on the sheet

Using this information, draw a simple series of sketches on the figure below for solar noon on June 21st at a site at 30° N, and on December 22nd and June 21st for a site at 60° N.



21 June: $\delta = +23.5°$ **22 December:** $\delta = -23.5°$

Figure 18.2. The Earth at the two solstices, with sunlight arriving from the left. The short coloured lines are the local verticals at the Tropic of Cancer, 30° N and 60° N. In June (left) the northern hemisphere is tilted towards the Sun, the Sun is overhead at the Tropic, and the verticals at 30° N and 60° N are only 6.5° and 36.5° from the Sun's direction. In December (right) they are 53.5° and 83.5° from it, Eq. (18.4).

Indicate the zenith angle, sun angle and beam spreading in each diagram.

Method. For each site and date, apply Eqs. (18.4), (18.1), (18.2) and (18.3) in turn. The calculations are collected in Table 18.1; the 30° N December case is not asked for in 5a but is needed for 5c, so it is included.

Table 18.1. Question 5 for the four cases. Zenith angle from Eq. (18.4), sun angle from Eq. (18.1), beam spreading from Eq. (18.2) and intensity from Eq. (18.3).


Site	Date	Zenith θ_z	Sun angle β	Spread $1/\sin\beta$	Intensity
30° N	21 June	$30 - 23.5 = 6.5°$	83.5°	$1/0.9936 = 1.006$	99.4 %
60° N	21 June	$60 - 23.5 = 36.5°$	53.5°	$1/0.8039 = 1.244$	80.4 %
60° N	22 Dec	$60 + 23.5 = 83.5°$	6.5°	$1/0.1132 = 8.834$	11.3 %
30° N	22 Dec	$30 + 23.5 = 53.5°$	36.5°	$1/0.5948 = 1.681$	59.5 %

Spreading in units of ground per unit of beam width; intensity in % of the overhead value per unit of ground.


Answer to Question 5a

30° N, 21 June: zenith angle 6.5°, sun angle 83.5°, beam spreading $1/\sin 83.5° =$ **1.006 units**.
60° N, 21 June: zenith angle 36.5°, sun angle 53.5°, beam spreading $1/\sin 53.5° =$ **1.24 units**.
60° N, 22 December: zenith angle 83.5°, sun angle 6.5°, beam spreading $1/\sin 6.5° =$ **8.83 units**.
Sketches as in Figure 18.3.


Check. The higher the Sun, the closer the spreading is to 1 unit; with the Sun just 6.5° above the horizon, a beam one unit wide is smeared across almost nine units of ground. In each case $\theta_z + \beta = 90°$.




30° N, 21 June
zenith angle $\theta_z = 6.5°$ (green arc)
sun angle $\beta = 83.5°$ (blue arc)
spreading = 1.01 units (red strip)
intensity 99.4 % per unit



60° N, 21 June
zenith angle $\theta_z = 36.5°$ (green arc)
sun angle $\beta = 53.5°$ (blue arc)
spreading = 1.24 units (red strip)
intensity 80.4 % per unit



60° N, 22 December
zenith angle $\theta_z = 83.5°$ (green arc)
sun angle $\beta = 6.5°$ (blue arc)
spreading = 8.83 units (red strip)
intensity 11.3 % per unit



30° N, 22 December
zenith angle $\theta_z = 53.5°$ (green arc)
sun angle $\beta = 36.5°$ (blue arc)
spreading = 1.68 units (red strip)
intensity 59.5 % per unit (needed for 5c)

Figure 18.3. Question 5a: the noon Sun at each site, drawn to the correct angles and to one scale. Each beam (orange) is one unit wide and the red strip is the ground it covers. The sun angle β (blue arc) is measured from the ground, and the zenith angle θ_z (green arc) from the vertical.

18.5 Question 5b: why the sun angle makes the seasons

The question on the sheet

Given these sketches, explain why the sun angle causes seasonal temperature changes in the mid-latitudes.

As the Earth orbits the Sun with its axis tilted, the latitude where the noon Sun is overhead moves between the tropics (Figure 18.2). At a mid-latitude site the noon Sun is therefore high in summer and low in winter: at 60° N, 53.5° in June and only 6.5° in December. A high Sun concentrates its beam on a small area and a low Sun spreads it over a large one, so each square metre of ground receives far more energy in summer (80 % per unit) than in winter (11 % per unit). More energy absorbed means warmer ground and air.

Answer to Question 5b

Because the Earth's axis is tilted, the noon **sun angle** at a mid-latitude site **rises in spring and summer and falls in autumn and winter**. A high sun angle concentrates the beam on a small area (little beam spreading, high intensity); a low sun angle spreads the same energy over a large area (much spreading, low intensity). So the ground receives more energy per square metre in summer and warms, and less in winter and cools. At 60° N the noon

intensity in June is about seven times that in December.

A question that comes up

Is the sun angle the only reason summers are warmer?
No; it is the main one, and two others work with it. **Day length:** at 60° N the Sun is up for about 18 hours at the June solstice and about 6 in December, so summer days add up much more energy. **Path through the atmosphere:** a low Sun's rays cross far more air, so more is scattered and absorbed before reaching the ground. All three effects point the same way. The distance from the Sun is *not* a reason: the Earth is closest to the Sun in early January, in the northern winter (Lutgens et al., 2019, ch. 2).




Figure 18.4. Noon solar intensity, Eq. (18.3), against latitude at the solstices and equinoxes. The dots are the four cases of Table 18.1. The peak follows the overhead Sun between the tropics; north of the Arctic Circle (66.5° N) the December curve reaches zero: the noon Sun does not rise.

18.6 Question 5c: comparing solar intensities

The question on the sheet

Given your answers: (i) what is the percentage difference in solar intensity between a site at 30° N and 60° N during the winter solstice? (Hint: calculate the solar intensity as % per unit for each site first.) (ii) What is the percentage difference in solar intensity between a site at 60° N for the winter solstice compared to the summer solstice?

Step 1: the intensities, from Eq. (18.3) and Table 18.1:

$$I_{30\text{N, Dec}} = 100/1.681 = 59.5 \text{ \% per unit},$$
$$I_{60\text{N, Dec}} = 100/8.834 = 11.3 \text{ \% per unit},$$
$$I_{60\text{N, Jun}} = 100/1.244 = 80.4 \text{ \% per unit}.$$

Step 2: compare them. "Percentage difference" between two percentages can mean two things, and it is worth giving both:

- the difference in **percentage points**: subtract one percentage from the other;
- the **relative** difference: divide one by the other.

(i) 30° N against 60° N in December:

$59.5 - 11.3 = 48.2$ percentage points; $\dfrac{59.5}{11.3} = 5.25$ times; $\dfrac{59.5 - 11.3}{59.5} = 81\,\%$ less at 60°N.

(ii) 60° N, December against June:

$80.4 - 11.3 = 69.1$ percentage points; $\dfrac{80.4}{11.3} = 7.10$ times; $\dfrac{80.4 - 11.3}{11.3} = 610\,\%$ more in June.

Answer to Question 5c

(i) At the winter solstice 30° N receives **59.5 %** per unit and 60° N **11.3 %** per unit: a difference of **48.2 percentage points**. 30° N receives about **5.3 times** as much as 60° N (60° N receives 81 % less).

(ii) At 60° N the winter solstice gives **11.3 %** per unit and the summer solstice **80.4 %** per unit: a difference of **69.1 percentage points**. The summer noon intensity is about **7.1 times** the winter one.

Check. Both comparisons agree with Figure 18.4: the December curve falls steeply between 30° N and 60° N, and at 60° N the June and December curves are far apart.

Corrections to the printed key for Question 5

1. **Rounding.** The key rounds the spreading at 60° N to 8.8 and 1.24 units before dividing, giving 11.4 and 80.6 % per unit. Unrounded, they are 8.834 and 1.244 units, giving **11.3** and **80.4** % per unit. Elsewhere the key gives the summer spreading as "1.2 units"; it is 1.24.
2. **Question 5c(i) has no final answer in the key**, only the two intensities. The difference is 48.2 percentage points (about 5.3 times).
3. **"69.2 % more".** The key subtracts the two intensities and calls the result "69.2 % more per unit". A difference between two percentages is in *percentage points*: 69.1 points (69.2 with the key's rounding). In relative terms the summer intensity is 7.1 times the winter one, which is 610 % more, not 69 %.
4. **A typing slip**: the key writes "Sun angle = 90 − 83.5.5 = 6.5"; it is 90 − 83.5 = 6.5°.

18.7 Try it yourself: Practice Set 18

Practice Set 18: sun angle

18.1 Coleraine is at 55.1° N. Find the noon zenith angle, sun angle, beam spreading and intensity at the June and December solstices.

18.2 What is the noon intensity at the equator at either solstice? And at the equinoxes?

18.3 What is the noon intensity at 60° N at the equinoxes, when the Sun is overhead at the equator?

18.4 What happens at the Arctic Circle (66.5° N) at noon on 22 December?

18.5 At which latitude is the noon intensity 100 % on 21 June, and why?

Answers to Practice Set 18

18.1 June: $\theta_z = 55.1 - 23.5 = 31.6°$, $\beta = 58.4°$, $A = 1/\sin 58.4° = 1.174$ units, $I = 85.2\,\%$ per unit. December: $\theta_z = 55.1 + 23.5 = 78.6°$, $\beta = 11.4°$, $A = 5.06$ units, $I = 19.8\,\%$ per unit. The June noon Sun gives Coleraine 4.3 times the December intensity.

18.2 At either solstice $\theta_z = 23.5°$, $\beta = 66.5°$, $A = 1.090$, $I = 91.7\,\%$ per unit. At the equinoxes the Sun is overhead: $100\,\%$. The equator's noon intensity hardly changes through the year, which is why the tropics have little temperature seasonality.

18.3 $\theta_z = 60°$, $\beta = 30°$, $A = 1/\sin 30° = 2$ units, $I = 50\,\%$ per unit.

18.4 $\theta_z = 66.5 + 23.5 = 90°$, so $\beta = 0°$: the noon Sun is on the horizon, the beam spreading is infinite and the intensity is zero. North of this, the Sun does not rise at all on that day: the polar night.

18.5 At the Tropic of Cancer, 23.5° N, where $\theta_z = 0°$ and the Sun is directly overhead: the declination on 21 June is +23.5°, Eq. (18.4).

Chapter 19

Putting Practical 2 together

19.1 One chain of ideas

The five questions are one argument, each step resting on the one before:

1. Everything above absolute zero radiates, over a spread of wavelengths described by the Planck curve (Chapter 12).
2. The **peak** of that spread moves to shorter wavelengths as temperature rises (Wien's law): the Sun, at about 5800 K, peaks in visible light; the Earth, near 288 K, in the thermal infrared (Question 1, Chapter 13).
3. The **total** emission rises as the fourth power of temperature (Stefan–Boltzmann law): twenty times the temperature means 166 000 times the emission (Question 2, Chapter 14).
4. A planet settles at the temperature at which it emits as much as it absorbs. That temperature depends on its distance from the Sun (inverse-square law) and its albedo, and can be read from its spectrum (Question 3, Chapters 15 and 16).
5. Where the actual temperature is higher than this, an atmosphere is trapping infrared (the greenhouse effect: Venus, the Earth) or the planet has its own heat (Jupiter). For the Earth the greenhouse effect is worth about 33 K (Question 4, Chapter 17).
6. Across the Earth's surface, the energy received depends on the angle of the Sun. The tilt of the axis moves that angle through the year and so makes the seasons (Question 5, Chapter 18).

This heating explains the structure met in Practical 1: a troposphere heated from below and a stratosphere heated by ozone (Part I, § 5.4), thickest over the tropics (Part I, § 7.4). Practical 3 (Part III) shows how the uneven heating from equator to pole fills the air with moisture and sets it in motion.

19.2 All the answers in one table

Table 19.1. Every answer of Practical 2, with the section where it is worked out.

Q	Answer	Where
1	$T_{\text{Sun}} = 2898/0.5 = 5796\,\text{K} = 5522.8\,^{\circ}\text{C}$	§ 13.3
2a	$E_{\text{Sun}} = 6.40 \times 10^{7}\,\text{W m}^{-2}$	§ 14.3
2b	$E_{\text{Earth}} = 385.5\,\text{W m}^{-2}$ (at 287.16 K)	§ 14.4
2c	$5796/287.16 = 20.2$, about 20 times hotter	§ 14.5
2d	$6.40 \times 10^{7}/385.5 = 166\,000$ times more emission	§ 14.6
2e	$E \propto T^{4}$: $20.18^{4} = 166\,000$ (a power law)	§ 14.7
3a	peaks at 12, 14 and 16 μm (A, B, C)	§ 15.3

Q	Answer	Where
3b	$T_A = 241.5\,\mathrm{K}$, $T_B = 207.0\,\mathrm{K}$, $T_C = 181.1\,\mathrm{K}$ (true curves: 254.1, 210.0, 184.1 K)	§ 15.4
3c	Mercury 439, Venus 185, Earth 255, Mars 210, Jupiter 110 K	§ 16.5
3d	A Earth, B Mars, C Venus	§ 16.6
3e	(i) Mars: further away, 43 % of the Earth's sunlight; (ii) Venus: albedo 0.9; (iii) Venus 737 K, intense CO_2 greenhouse; (iv) Mercury: no atmosphere (Mars nearly none)	§ 16.7
4a	70 % absorbed	§ 17.2
4b	reflected 410.4, absorbed $957.6\,\mathrm{W\,m^{-2}}$	§ 17.3
4b′	night and slanting rays: disc πR^2 against sphere $4\pi R^2$, so $1368/4 = 342 \approx 341\,\mathrm{W\,m^{-2}}$	§ 17.4
4c	reflected 102.3, absorbed $238.7\,\mathrm{W\,m^{-2}}$	§ 17.5
4d	$T = (E/\sigma)^{1/4} = 254.7\,\mathrm{K} = -18.5\,^\circ\mathrm{C}$	§ 17.6
4e	greenhouse effect warms the surface by about 32.5 K (33 K)	§ 17.7
4f	much larger day–night range: very hot days, very cold nights	§ 17.8
4g	no internal heat source; black body at one temperature, in equilibrium	§ 17.9
5a	30° N June: $\theta_z = 6.5°$, $\beta = 83.5°$, 1.006 units; 60° N June: 36.5°, 53.5°, 1.24 units; 60° N December: 83.5°, 6.5°, 8.83 units	§ 18.4
5b	tilt: high summer Sun concentrates the beam, low winter Sun spreads it	§ 18.5
5c	(i) 59.5 against 11.3 % per unit: 48.2 points, 5.3 times; (ii) 80.4 against 11.3 % per unit: 69.1 points, 7.1 times	§ 18.6

19.3 Mistakes to avoid

- **Using Celsius in a radiation law.** σT^4 and C/T need kelvin. Convert first, and convert back only at the end (§ 14.5).
- **Forgetting the fourth root.** When solving $E = \sigma T^4$ for T, divide by σ *and then* take the fourth root, Eq. (17.3); a single square root gives a value about 65 000 instead of 255.
- **Mixing up the sun angle and the zenith angle.** The sun angle is from the horizon, the zenith angle from the vertical; they add to 90°, Eq. (18.1). Beam spreading uses the sun angle.
- **Calculator in radians.** sin(83.5) in radian mode gives 0.97 instead of 0.9936.
- **Subtracting percentages and calling the result a percentage.** Two intensities of 80.4 % and 11.3 % differ by 69.1 percentage points; in relative terms one is 7.1 times the other (§ 18.6).
- **Reading the peak of a coarsely sampled spectrum as exact.** The highest data point lies near, not at, the peak (§ 15.4).

19.4 Self-test

Self-test: one line each

S1. Why is the Sun's radiation called shortwave and the Earth's longwave?

S2. If a star is three times as hot as the Sun, how does its total emission per square metre compare?

S3. Why does Venus have a lower black-body temperature than the Earth but a far higher surface temperature?

S4. Why do we divide the solar constant by four?

S5. Without an atmosphere, would the Earth be colder by day, by night, or both?

S6. Why is there little seasonal change in temperature at the equator?

Answers to the self-test

S1. Wien's law: the Sun is about 20 times hotter, so its emission peaks at a wavelength about 20 times shorter (0.5 against 10 μm) (Chapter 13).

S2. $3^4 = 81$ times as much, Eq. (14.2).

S3. Its bright clouds reflect most sunlight (high albedo), so it absorbs little; but its thick CO_2 atmosphere traps the infrared from the surface (§ 16.7).

S4. The Earth catches sunlight on a disc of area πR^2 but spreads it over a sphere of area $4\pi R^2$ (Figure 16.1).

S5. Mostly by night: nights would be far colder. Days could be hotter, as on the Moon (§ 17.8). On average it would be about 33 K colder.

S6. The noon Sun is always high there (at least 66.5°), so the intensity changes little through the year (Practice Set 18, item 18.2).

Part III

Practical 3: Moisture, motion and weather maps

Chapter 20

Practical 3 at a glance

20.1 What the practical asks of you

Practical 3 has three sections, each building a skill that the next one needs, so it pays to work through them in order (EGM305 module team, n.d.). It brings together the two practicals before it: the vertical structure of Practical 1 (Part I) gives the pressure surfaces whose slopes drive the wind, and the uneven solar heating of Practical 2 (Part II) is what makes them slope.

Moisture. How do we say how much water vapour the air holds? You meet four different measures (vapour pressure, mixing ratio, relative humidity and dew point) and learn to move between them with a table and a graph.

Motion. Why does the air move at all, and why does it move the way it does? You study the three forces that matter for large-scale winds (the pressure gradient force, the Coriolis force and friction) and see how their balance gives the geostrophic wind high up and a wind that crosses the isobars near the ground.

Weather maps. You decode the station model that every weather map uses, draw streamlines from wind barbs, and analyse a whole map: isobars, the low, the fronts, the cloud and the rain.

The sheet numbers its questions in a way that can confuse. The moisture questions are numbered 1 to 6. The numbering then starts again at 1 for the motion questions, which run to 11. The weather map questions carry on from 12 to 14. In this companion a question of Practical 3 is always named with its section, for example *Moisture Q3*, *Motion Q7* or *Map Q12*.

Table 20.1. Where each question of the Practical 3 sheet is answered, and the equations it needs.

Question	What it asks	Chapter	Equations	On the sheet
Moisture Q1	relative humidity of five samples	21	(21.3)	Table 1
Moisture Q2	heating cold air indoors in winter	21	(21.3)	
Moisture Q3	ranking five samples by actual vapour	21	(21.4)	Table 1
Moisture Q4	the weakness of relative humidity	21		
Moisture Q5	dew point of 20 °C, 10 g kg^{-1}	22	(21.2)	Figure 3
Moisture Q6	dew point of 34 °C at 62 %	22	(21.4), (22.1)	
Motion Q1–Q6	pressure surfaces, Saskatoon and Monterrey	24	(24.1), (23.1)	Figures 1, 2
Motion Q7	geostrophic wind, north and south	25	(25.1)	Figures 3, 4
Motion Q8–Q9	friction and the forces on a box	26	(26.1)	Figure 5
Motion Q10–Q11	four circulation systems	26	(26.1)	Figure 6
Map Q12	decode four station models	27	(27.1)	Figure 7
Map Q13	streamlines	28		Figure 8
Map Q14	fronts and the full analysis	29	(25.1)	Figures 9, 10

Chapter 23 builds the physics of the motion section before its questions; Chapter 30 gathers every answer in one table. The formula sheet, the corrected Table 1, the 34 stations of the weather map, derivations, weather symbols and the corrections to the key for this practical are in Appendix C.

20.2 What to have beside you

- The sheet itself, including **Table 1** (saturation vapour pressure and mixing ratio). A corrected copy is printed in § C.2.
- A calculator that has `ln` and `exp` keys, though most questions need only multiplication and division.
- Four coloured pencils for the force diagrams: red for the pressure gradient force, blue for the Coriolis force, green for friction, and black for the wind.
- A ruler and a soft pencil for the weather map.

20.3 Symbols and units

Table 20.2. The symbols used in Part III.

Symbol	Meaning	Unit
T, T_d	air temperature, dew-point temperature	°C (°F on the station models)
e, e_s	vapour pressure, saturation vapour pressure	hPa (= mb)
w, w_s	mixing ratio, saturation mixing ratio	$\mathrm{g\,kg^{-1}}$
RH	relative humidity	%
p	air pressure	hPa (= mb)
z	height above the ground	m
ρ	air density (about 1.2 $\mathrm{kg\,m^{-3}}$ near the ground)	$\mathrm{kg\,m^{-3}}$
f	Coriolis parameter, $f = 2\Omega \sin\varphi$	$\mathrm{s^{-1}}$
V, V_g	wind speed, geostrophic wind speed	$\mathrm{m\,s^{-1}}$ (or knots)
Δn	distance across the isobars (or height lines)	km

Millibar and hectopascal. The sheet says *mb*; modern texts say *hPa*. They are the same: 1 mb = 1 hPa = 100 Pa. This part uses both, following the sheet where it asks the question.

Knots. Wind barbs use knots: 1 kt = 0.514 $\mathrm{m\,s^{-1}} \approx 1.85$ $\mathrm{km\,h^{-1}}$.

20.4 A word on the printed answer key

The printed answer key was checked against Table 1 and the figures, one answer at a time. Almost everything in it is right. Three things need correcting: the dew point of Moisture Q6 (26 °C, not 25 °C), the dew point of station model 1 in Map Q12 (21 °F, not 28 °F), and two typing slips in the mixing ratios of Table 1. Each is explained where it arises, and § C.6 lists them all.

20.5 About the weather map

Figures 8 and 10 of the sheet show the same 34 stations in the central United States. The printed figures are pictures, so the station values used here (positions, temperatures, dew points, pressure codes, cloud cover, wind barbs and weather symbols) were **read from the images**. Wind speeds were obtained by counting barbs and are approximate. The fronts were placed by hand from the temperature, dew-point and wind changes, as an analyst would. § C.3 lists every station so that you can check any value against your own copy of the sheet.

20.6 Before the practical

The chapters on atmospheric humidity, on air pressure and winds, and on air masses, fronts and middle-latitude cyclones in Ahrens and Henson (2022) or Lutgens et al. (2019) cover this practical. Stull (2017) treats the hypsometric equation, the geostrophic wind and station models with worked examples, and Holton and Hakim (2013) takes the balance of forces further.

Chapter 21

Moisture in the air: mixing ratio and relative humidity

This chapter answers Moisture Q1 to Q4 of the sheet.

21.1 Why water vapour matters

Water vapour is the invisible, gaseous form of water. It makes up between a fraction of a percent and about four percent of the air by volume, yet it is the ingredient of every cloud, every shower and every fog. It also sets how comfortable a day feels, because sweat evaporates slowly into air that is already moist. To talk about weather precisely we need a way to say *how much* vapour the air contains. The sheet introduces four measures. This chapter covers the first three; the fourth, the dew point, has its own chapter.

21.2 Vapour pressure and saturation

Air is a mixture of gases, and each gas contributes a share of the total pressure, in proportion to how many of its molecules are present: the partial pressure met for oxygen in Practical 1 (Part I, § 4.4). The share due to water vapour is the **vapour pressure**, written e. In the lowest kilometres it is about 1 to 30 hPa, out of a total pressure of roughly 1000 hPa.

Imagine a closed dish of water with air above it. Molecules leave the water surface (evaporation) and return to it (condensation). At first more leave than return, so the vapour pressure above the water rises. It stops rising when the two flows are equal. The air is then **saturated**, and the vapour pressure has reached its upper limit for that temperature: the **saturation vapour pressure**, e_s.

The warmer the water, the more energetic its molecules and the more easily they escape, so e_s rises with temperature, and it rises steeply. Table 1 on the sheet gives it as the "mb" column: 6.1 mb at 0 °C, 23.4 mb at 20 °C and 73.8 mb at 40 °C. Plotted, it is the curve of Figure 21.1 (this is Figure 1 of the sheet, the Clausius–Clapeyron curve).

A question that comes up

Why does warm air "hold" more water than cold air?
Air does not really hold water the way a sponge does. The saturation vapour pressure is a property of water itself, how readily it evaporates at a given temperature, and the other gases hardly affect it. "Holding capacity" is a convenient picture, and this part uses it, but it is the temperature that sets the limit.




Figure 21.1. Saturation vapour pressure against temperature. The dots are the values printed in Table 1. Air whose vapour pressure and temperature fall below the curve is unsaturated; points above it are not possible for long, because the surplus vapour condenses.

21.3 Mixing ratio

Vapour pressure is not the only way to measure moisture. The **mixing ratio**, w, compares the *mass* of vapour with the mass of the *dry air* that goes with it:

$$w = \frac{\text{mass of water vapour}}{\text{mass of dry air}}. \tag{21.1}$$

where

w	mixing ratio ($\mathrm{g\,kg^{-1}}$: grams of vapour per kilogram of dry air)	calculated

Typical values run from $0.1\,\mathrm{g\,kg^{-1}}$ in the polar winter to $20\,\mathrm{g\,kg^{-1}}$ in the humid tropics. When the air is saturated the ratio is the **saturation mixing ratio**, w_s, the second column of numbers in Table 1 (the "g/kg" column).

The two measures are linked. For a total pressure p and a vapour pressure e,

$$w \approx 622\,\frac{e}{p-e}. \tag{21.2}$$

where

w	mixing ratio ($\mathrm{g\,kg^{-1}}$)	calculated
e	vapour pressure (hPa)	
p	total air pressure (hPa)	995.5 for Table 1
622	ratio of the molar masses of water and dry air, 18/29, times 1000 (to give $\mathrm{g\,kg^{-1}}$)	

The factor 0.622 is the ratio of the molar masses of water (18) and dry air (29): a molecule of vapour is lighter than the average molecule of dry air.

A question that comes up

Why use the mass of *dry* air, and why grams per kilogram?
The mass of dry air in a rising or sinking parcel stays fixed, while the total mass of moist air does not, as soon as vapour condenses or evaporates. Dividing by the dry mass gives a number that stays constant for a parcel until water is added or removed. Grams per kilogram




Figure 21.2. Saturation mixing ratio against temperature (Figure 2 of the sheet). The curve is computed from the saturation vapour pressure at a pressure of 995.5 hPa, which is the pressure Table 1 was evidently calculated for. The two red crosses are the printed values at 0 °C and −8 °C; they lie below the curve (see Section 21.11).

simply keeps the numbers handy: a few units to a few tens.

A question that comes up

The sheet says Table 1 is "at sea-level". Which pressure is that?
Working backwards from the table, $p = e_s + 622\, e_s/w_s$ comes out at about 995.5 hPa for every row from −20 to 40 °C (except the two slips). That is a typical low-ish sea-level pressure, not the standard 1013.25 hPa. At 1013.25 hPa the mixing ratios would be about 1.8 % smaller. The difference never matters for the questions on the sheet, but it explains why the guide's own calculations use 995.5 hPa: so that every answer agrees with the table.

21.4 Relative humidity

Relative humidity (RH) says how close the air is to saturation. On the sheet it is defined in two equivalent ways:

$$\mathrm{RH} = \frac{e}{e_s} \times 100\,\% = \frac{w}{w_s} \times 100\,\%. \tag{21.3}$$

where

RH	relative humidity (%)	calculated
e, e_s	vapour pressure and saturation vapour pressure (hPa)	Table 1
w, w_s	mixing ratio and saturation mixing ratio ($\mathrm{g\,kg^{-1}}$)	Table 1

An RH of 50 % means the air holds half of what it could hold at its present temperature; 100 % means saturated. Because e_s and w_s both depend strongly on temperature, *the same air has a different relative humidity at a different temperature.* That fact is behind most of the puzzles in this chapter.

Relative humidity in one line

Relative humidity = (how much vapour is there) ÷ (how much there could be at this temperature) ×100.

21.5 A picture to hold on to: the glass

Picture the air as a glass. The **size of the glass** is the saturation mixing ratio at the air's temperature. The **water in the glass** is the actual mixing ratio. Relative humidity is how full the glass is. Warm air has a big glass, cold air a small one.




Figure 21.3. The glass picture. The same 5 g of vapour per kilogram of air fills the glass of 14 °C air to 49 %, the glass of 24 °C air to 26 %, and the big glass of 34 °C air to only 14 %. The amount of water is identical; only the size of the glass changes.

21.6 Reading Table 1

Three skills with Table 1 carry you through the whole of Part I.

Skill 1. Look up a value at a temperature that is listed, for example $w_s(24\,^\circ\mathrm{C}) = 19.21\,\mathrm{g\,kg^{-1}}$.

Skill 2. Interpolate between two rows for a temperature that is not listed.

Skill 3. Look up backwards: given a mixing ratio, find the temperature at which it is the saturation value. This is how you find a dew point (Chapter 22).

Worked example: interpolating in Table 1

Find w_s at 25 °C. The table lists 24 °C (19.21) and 26 °C (21.734). 25 °C is half way between, so take the half-way value:

$$w_s(25) \approx \frac{19.21 + 21.734}{2} = 20.47\ \mathrm{g\,kg^{-1}}.$$

The exact value from the formula is $20.44\,\mathrm{g\,kg^{-1}}$. Because the curve is convex (it bends upwards), straight-line interpolation overestimates slightly, by about 0.1 % over a 2-degree step. That is far smaller than the precision of the question.

21.7 Question 1: relative humidity of five samples

The question on the sheet

Using Table 1, determine the saturation mixing ratio and calculate the relative humidity of five air samples at the temperatures and mixing ratios shown.

Method.

Step 1. For each sample read the saturation mixing ratio w_s from Table 1 at the sample's temperature.

Step 2. Divide the sample's mixing ratio w by w_s and multiply by 100.

Step 3. Check: more vapour at the same temperature must give a higher RH; the same vapour in warmer air must give a lower RH.

Answer to Moisture Q1

Temp.	Sat. mixing ratio ($\mathrm{g\,kg^{-1}}$)	Mixing ratio ($\mathrm{g\,kg^{-1}}$)	RH (%)
14 °C	10.14	5	$5/10.14 = \mathbf{49.3}$
14 °C	10.14	9	$9/10.14 = \mathbf{88.8}$
24 °C	19.21	5	$5/19.21 = \mathbf{26.0}$
24 °C	19.21	2	$2/19.21 = \mathbf{10.4}$
34 °C	35.134	7	$7/35.134 = \mathbf{19.9}$




Figure 21.4. The five samples of Question 1 drawn as glasses on the same scale. The number at the top of each glass is its capacity (the saturation mixing ratio); the number just above the water is the sample's mixing ratio; the percentage below is the relative humidity.

Check. The two 14 °C samples have 5 and 9 $\mathrm{g\,kg^{-1}}$; the one with more vapour (49.3 % against 88.8 %) has the higher RH. The two 24 °C samples likewise. The 34 °C sample, with 7 $\mathrm{g\,kg^{-1}}$ in a huge glass, has the lowest RH of the five bar the 2 $\mathrm{g\,kg^{-1}}$ sample, as it should.

21.8 Question 2: cold air heated in a house

The question on the sheet

In winter, cold air is brought into homes and heated. How does this change the humidity in the house?

Method. Follow one parcel of air. Heating changes its temperature, so it changes the size of the glass, but it adds no water, so the amount of vapour stays fixed.

Worked example: a cold day in Belfast

Outside, the air is at $0\,^\circ\mathrm{C}$ with a relative humidity of $80\,\%$. From Table 1 (corrected value) $w_s(0\,^\circ\mathrm{C}) = 3.84\,\mathrm{g\,kg^{-1}}$, so the air holds

$$w = 0.80 \times 3.84 = 3.07\ \mathrm{g\,kg^{-1}}.$$

Bring it indoors and heat it to $21\,^\circ\mathrm{C}$. The saturation mixing ratio at $21\,^\circ\mathrm{C}$ is about $15.9\,\mathrm{g\,kg^{-1}}$ (half way between the 16.0 of $20\,^\circ\mathrm{C}$ and 17.0 of $22\,^\circ\mathrm{C}$; the exact value is 15.93). Nothing has been added, so w is still $3.07\,\mathrm{g\,kg^{-1}}$ and

$$\mathrm{RH} = \frac{3.07}{15.9} \times 100 = 19\,\%.$$

To reach a comfortable $40\,\%$ the air would need $0.40 \times 15.9 = 6.4\,\mathrm{g\,kg^{-1}}$, which means adding $6.4 - 3.07 = 3.3\,\mathrm{g\,kg^{-1}}$ of water.




Figure 21.5. Question 2. Heating leaves the amount of vapour untouched but makes the glass four times bigger, so the relative humidity falls from $80\,\%$ to $19\,\%$.

Answer to Moisture Q2

The humidity in the house will **drop significantly**. Cold air holds little vapour, so the outside air has a low mixing ratio. Heating it without adding water leaves the mixing ratio unchanged but increases the saturation mixing ratio a great deal, so the relative humidity becomes very low.

21.9 Question 3: who holds the most water?

The question on the sheet

Five samples are described by temperature and relative humidity. Complete the mixing ratios and rank them from 1 (most vapour) to 5 (least).




Figure 21.6. How dry the indoor air becomes. Outdoor air at −10 °C and 70 % humidity gives only about 6 % when heated to 21 °C with no water added; on a mild 10 °C day the same calculation gives 28 %.

Method. Turn each relative humidity back into an amount of vapour:

$$w = w_s \times \frac{\mathrm{RH}}{100}. \tag{21.4}$$

where

w	actual mixing ratio ($\mathrm{g\,kg^{-1}}$)	calculated
w_s	saturation mixing ratio at the air temperature ($\mathrm{g\,kg^{-1}}$)	Table 1
RH	relative humidity (%)	given

Step 1. Read w_s from Table 1 for each temperature.

Step 2. Multiply by the relative humidity written as a fraction (90 % is 0.90).

Step 3. Rank the results from the largest to the smallest.

Answer to Moisture Q3

Temp.	w_s ($\mathrm{g\,kg^{-1}}$)	RH	$w = w_s \times$ RH ($\mathrm{g\,kg^{-1}}$)	Rank
14 °C	10.14	90 %	$10.14 \times 0.90 = 9.126$	3
20 °C	14.956	60 %	$14.956 \times 0.60 = 8.974$	4
24 °C	19.21	40 %	$19.21 \times 0.40 = 7.684$	5
30 °C	27.694	40 %	$27.694 \times 0.40 = 11.078$	1
34 °C	35.134	30 %	$35.134 \times 0.30 = 10.54$	2

Check. Make sure the ranking depends on the amount of vapour, not on the humidity. If you had ranked by RH you would have put 14 °C first. The glass picture shows why that is wrong: a small glass 90 % full holds less than a big glass 40 % full.




Figure 21.7. Question 3 ranked (rank 1 at the top). The 14 °C sample is the "dampest" at 90 % but is only third in actual water; the 30 °C sample, which feels "dry" at 40 %, holds the most.

21.10 Question 4: the disadvantage of relative humidity

Answer to Moisture Q4

Relative humidity does not tell you how much water vapour is actually in the air.

That is the short answer the sheet wants. The fuller explanation is worth knowing, because it recurs in weather forecasting.

1. **It depends on temperature as well as on the vapour.** The same air has an RH of 49 % at 14 °C and 26 % at 24 °C (Question 1). Nothing was added or removed.
2. **It changes through the day for no reason connected with moisture.** On a calm day the vapour content hardly changes, but the temperature rises by several degrees after dawn, so the RH falls.
3. **It cannot be used to compare two places.** The most humid-feeling sample in Question 3 (90 %) held less vapour than three drier-feeling ones.

Better measures. The mixing ratio and the dew point measure the vapour itself. Neither changes when air is heated or cooled, as long as nothing condenses. For that reason forecasters watch the dew point more closely than the relative humidity, and so do the station models you meet in Chapter 27.

21.11 The two slips in Table 1

Look again at Figure 21.2. Two of the printed points do not lie on the curve. The table prints $3.189\,\mathrm{g\,kg^{-1}}$ at 0 °C, which is *less* than the 3.313 printed at −2 °C, even though 0 °C is warmer; saturation mixing ratio can only increase with temperature. Applying the formula at 995.5 hPa gives 3.840 for 0 °C. Similarly, the entry for −8 °C (2.009) should be 2.099, a transposition of digits. All other rows fit the formula to better than 0.1 % (the rows at −30 and −25 °C to about 2 %, because only three digits are printed). No question on the sheet uses a temperature below 14 °C except Question 2, and this part has used the corrected 0 °C value there.




Figure 21.8. Table 1 near freezing. The printed values at 0 °C and −8 °C (red crosses) fall away from the curve; the corrected values (green circles) lie on it.

21.12 Try it yourself: Practice Set 21

Practice Set 21: relative humidity and vapour amounts

21.1 Air at 28 °C has a mixing ratio of 15 g kg^{-1}. What is its relative humidity?

21.2 Air at 10 °C has a relative humidity of 70 %. How much water vapour does a kilogram of it contain?

21.3 Sample P: 32 °C, RH 25 %. Sample Q: 20 °C, RH 85 %. Which holds more water vapour, and how much more?

21.4 Air at 0 °C and RH 90 % is heated to 20 °C with no water added. What is its relative humidity now?

21.5 Air at 24 °C and RH 50 % is cooled to 14 °C with no water added (and no condensation). What is its RH now?

21.6 Use interpolation to find w_s at 31 °C, and hence the mixing ratio of air at 31 °C with RH 55 %.

Answers to Practice Set 21

21.1 $w_s(28) = 24.557$, so RH $= 15/24.557 = 61\,\%$.

21.2 $w_s(10) = 7.762$, so $w = 0.70 \times 7.762 = 5.43$ g kg^{-1}.

21.3 P: $31.213 \times 0.25 = 7.80$ g kg^{-1}. Q: $14.956 \times 0.85 = 12.71$ g kg^{-1}. Q holds more, by 4.91 g kg^{-1}, although its temperature is 12 °C lower.

21.4 $w = 0.90 \times 3.84 = 3.456$ g kg^{-1}; $w_s(20) = 14.956$; RH $= 3.456/14.956 = 23\,\%$.

21.5 $w = 0.50 \times 19.21 = 9.605$ g kg^{-1}; $w_s(14) = 10.14$; RH $= 9.605/10.14 = 94.7\,\%$. The air is almost saturated: another degree or so of cooling brings the dew point.

21.6 $w_s(31) \approx (27.694 + 31.213)/2 = 29.45$ g kg^{-1}; $w = 0.55 \times 29.45 = 16.2$ g kg^{-1}.

Chapter 22

The dew point

This chapter answers Moisture Q5 and Q6 of the sheet.

22.1 Cooling the glass

Take a sample of air and cool it, taking nothing away and adding nothing. Two things happen. The vapour in the air stays the same, because no water has gone in or out. The glass (the saturation mixing ratio) shrinks, because colder air can hold less. So the glass gets fuller and the relative humidity rises. At one particular temperature the glass is exactly full. That temperature is the **dew point**, T_d.

Definition

The **dew point** is the temperature to which an air sample must be cooled, at constant pressure and without adding or removing water, to reach saturation. At that temperature the saturation mixing ratio equals the sample's actual mixing ratio, $w_s(T_d) = w$.

If you cool the air below its dew point the surplus vapour cannot stay as vapour. It condenses: into dew on cold grass, into fog near the ground, into cloud higher up. A cold can of drink on a summer day is covered with water for exactly this reason: the air against the can is cooled below its dew point.

22.2 Three ways to find a dew point

All three give the same answer. The sheet asks for the first; the others are checks.

Method A. **Graph** (Figure 3 on the sheet). Plot the air as a point (temperature across, mixing ratio up). Cooling moves the point horizontally to the left, because only the temperature changes. Stop where the point meets the saturation curve. Read the temperature underneath.

Method B. **Table 1, backwards.** Find the two rows whose mixing ratios bracket the sample's, and interpolate for the temperature.

Method C. **Formula.** Compute the vapour pressure $e = wp/(622 + w)$ and invert the saturation formula (Appendix C.1).

22.3 Question 5: 20 °C and 10 g kg^{-1}

The question on the sheet

Consider a sample of air with a temperature of 20 °C and a mixing ratio of 10 g kg^{-1}. Plot the point on Figure 3, and show on the graph to which temperature the sample must cool before it reaches saturation.

Method A: the graph.

Step 1. Mark the point: 20 °C on the horizontal axis, 10 g kg^{-1} on the vertical axis.

Step 2. Cooling moves the point to the left along the horizontal line at 10 g kg^{-1}. Draw an arrow in that direction.

Step 3. The arrow meets the curve where the air is saturated. (At that point the curve's height, w_s, equals 10 g kg^{-1}.)

Step 4. Draw a vertical line down from the meeting point to the temperature axis and read the value.




Figure 22.1. Figure 3 of the sheet solved for both questions. Each air sample is a red dot; cooling moves it horizontally to the curve; the dew point is read on the temperature axis. For Question 5 it is 13.8 °C (about 14 °C); for Question 6 it is 26.0 °C.

Method B: Table 1 backwards. The sample holds 10 g kg^{-1}. Table 1 gives 8.882 g kg^{-1} at 12 °C and 10.14 g kg^{-1} at 14 °C, so the dew point is between 12 and 14 °C. The fraction of the way from 12 to 14 is

$$\frac{10 - 8.882}{10.14 - 8.882} = \frac{1.118}{1.258} = 0.89, \qquad \text{so} \qquad T_d \approx 12 + 0.89 \times 2 = 13.8 \text{ °C}.$$

Method C: formula. The vapour pressure is $e = \dfrac{10 \times 995.5}{622 + 10} = 15.75$ hPa, and inverting $e_s(T) = 6.112 \exp[17.67T/(T + 243.5)]$ (Bolton, 1980) gives $T_d = 13.78$ °C.

Answer to Moisture Q5

The sample cools along the horizontal line at 10 g kg^{-1} and meets the curve at about **14** °C (13.8 °C). That is its dew point.

Check. The dew point must be lower than the air temperature (20 °C), and 14 °C is. At 14 °C Table 1 says the air could hold 10.14 g kg^{-1}, only a little more than the 10 g kg^{-1} it has: so the air is just short of saturated at 14 °C, as it should be.

22.4 Question 6: 34 °C and 62 % humidity

The question on the sheet

A sample of air has a temperature of 34 °C and a relative humidity of 62 %. What is its saturation mixing ratio, its mixing ratio and its dew point?

Method.

Step 1. Saturation mixing ratio at 34 °C, from Table 1: 35.134 g kg^{-1}.

Step 2. Mixing ratio: $w = w_s \times \mathrm{RH}/100 = 35.134 \times 0.62 = 21.78$ g kg^{-1}.

Step 3. Dew point: find the temperature at which $w_s = 21.78$ g kg^{-1}. Table 1 has 21.734 at 26 °C and 24.557 at 28 °C. The value 21.78 is just above the 26 °C entry:

$$\frac{21.78 - 21.734}{24.557 - 21.734} = \frac{0.046}{2.823} = 0.016, \qquad T_d \approx 26 + 0.016 \times 2 = 26.03 \text{ °C}.$$

Answer to Moisture Q6

Saturation mixing ratio: **35.134** g kg^{-1}. Mixing ratio: **35.134 × 0.62 = 21.78** g kg^{-1}. Dew point: about **26** °C.

A correction to the printed key: 26 °C, not 25 °C

The printed answer key gives a dew point of 25 °C for this sample. That cannot be right, and there are three independent ways to see it.

1. **Table 1.** At 25 °C air can hold at most about 20.4 g kg^{-1} (half way between 19.21 and 21.734, which is 20.47). The sample holds 21.78 g kg^{-1}, *more* than that. At 25 °C it would be oversaturated: a relative humidity of 21.78/20.44 = 107 %.
2. **The graph.** In Figure 22.1 the horizontal line at 21.78 g kg^{-1} meets the curve at 26 °C. At 25 °C the line lies above the curve, in the supersaturated region.
3. **The rule of thumb** (next section): RH ≈ 100 − 5 × ($T - T_d$). A dew point of 26 °C gives 100 − 5 × 8 = 60 %, close to the 62 % we started with. A dew point of 25 °C would give 55 %.

22.5 A quick check by rule of thumb

Each degree by which the dew point lies below the temperature lowers the relative humidity by about 5 %. Written as a formula,

$$\mathrm{RH} \approx 100 - 5\,(T - T_d), \tag{22.1}$$

where

RH	relative humidity (%)	estimated
T, T_d	air temperature and dew point (°C)	

good to within a few percent for RH above about 50 %. Turned round, $T - T_d \approx (100 - \mathrm{RH})/5$: air at 80 % is about 4 °C above its dew point; at 50 % about 10 °C.




Figure 22.2. The rule of thumb against the exact relationship. For ordinary temperatures the exact curves stay close to the straight line.

22.6 What happens below the dew point

Continue cooling the Question 6 sample below 26 °C. The air can no longer hold all its vapour, so the surplus condenses. At each temperature the vapour left is $w_s(T)$, and the difference is liquid water:

$$\text{condensed water} = w - w_s(T). \tag{22.2}$$

where

w	mixing ratio of the air before cooling ($\mathrm{g\,kg^{-1}}$)	
$w_s(T)$	saturation mixing ratio at the new, lower temperature ($\mathrm{g\,kg^{-1}}$)	Table 1

Cooled to 20 °C, the sample holds only $14.956\,\mathrm{g\,kg^{-1}}$ as vapour, so $21.78 - 14.956 = 6.8\,\mathrm{g\,kg^{-1}}$ has turned to liquid. That is dew on the ground, or fog, or cloud droplets. Condensation also releases latent heat, which is why cloud keeps air warmer than it would otherwise be as it cools.




Figure 22.3. The Question 6 sample cooled from 34 °C. Above 26 °C nothing condenses and the relative humidity rises as the glass shrinks. Below 26 °C the air stays at 100 % and the surplus vapour condenses: $6.8\,\mathrm{g\,kg^{-1}}$ by the time it reaches 20 °C.

22.7 Why meteorologists prefer the dew point

- **It measures the vapour itself.** Heating or cooling the air leaves the dew point unchanged (until condensation), unlike the relative humidity.
- **It is a temperature,** so it is easy to compare with the air temperature: the gap $T - T_d$ tells you at a glance how close the air is to making dew, fog or cloud.
- **It describes comfort.** Most people find a dew point below about 10 °C dry and comfortable and one above about 20 °C oppressive.
- **It appears on every station model** at the bottom left of the circle, as you will see in Chapter 27.

22.8 The dew point and the height of cloud base

Air that rises expands and cools at about 9.8 K per kilometre (the dry adiabatic lapse rate, derived in Part I, Eq. (8.2)). Its dew point falls more slowly, by about 1.8 K per kilometre, because the pressure falls as it rises. The temperature and the dew point therefore approach each other by 8 K per kilometre, and when they meet the air is saturated and cloud forms. Starting with a gap of $T - T_d$ degrees, they meet after rising

$$z \approx \frac{T - T_d}{8} \text{ km} \approx 125\,(T - T_d) \text{ m}. \tag{22.3}$$

where

z	height of cloud base above the ground	calculated
T, T_d	surface temperature and dew point (°C)	
8	rate at which T and T_d approach in rising air (9.8 − 1.8, K km^{-1})	

Worked example: where will cloud form?

The air of Question 6 has $T - T_d = 34 - 26 = 8$ °C. If it is lifted (by a hill or by convection), cloud forms after $125 \times 8 = 1000$ m: the cloud base is about one kilometre above the surface. A day with $T = 30$ °C and $T_d = 18$ °C has a gap of 12 °C and a cloud base near 1500 m.




Figure 22.4. Rising air, starting at 34 °C with a dew point of 26 °C. The temperature falls 9.8 K per kilometre and the dew point 1.8 K per kilometre. They meet at a height of 1.0 km.

22.9 Questions that come up

A question that comes up

Does the dew point change when the day warms up?
Not because of the warming. If no water is added or removed, the mixing ratio is fixed, and so is the dew point. In the afternoon the *relative humidity* falls because the temperature rises. The dew point changes only when the air gains moisture (evaporation from soil and plants, an incoming sea breeze) or loses it (rain, a dry wind).

A question that comes up

Can the dew point be higher than the temperature?
Not in the open air for more than a moment: it would mean the air is oversaturated, and the surplus vapour would condense at once, lowering the dew point. At most the two are equal (RH $= 100\,\%$), as in fog.

A question that comes up

What if the dew point is below freezing?
Then the vapour deposits directly as ice (frost) when the air is cooled to it. Strictly the temperature of deposition is the *frost point*, which is a degree or two above the dew point. Table 1 is for saturation over liquid water; the questions on the sheet never need the ice case.

22.10 Try it yourself: Practice Set 22

Practice Set 22: the dew point

22.1 Air at $30\,^\circ\mathrm{C}$ has a mixing ratio of $12\,\mathrm{g\,kg^{-1}}$. Find its dew point (a) from Table 1 by interpolation and (b) with the rule of thumb after finding its RH.

22.2 Air at $20\,^\circ\mathrm{C}$ has a relative humidity of $50\,\%$. What is its dew point?

22.3 On an afternoon the air is at $28\,^\circ\mathrm{C}$ with a relative humidity of $40\,\%$. Overnight the temperature falls to $12\,^\circ\mathrm{C}$ with no change in the amount of vapour. Will dew form?

22.4 Find the cloud base for air with $T = 30\,^\circ\mathrm{C}$ and $T_d = 18\,^\circ\mathrm{C}$.

22.5 Air at $25\,^\circ\mathrm{C}$ has a dew point of $15\,^\circ\mathrm{C}$. What are its mixing ratio and relative humidity?

22.6 Cool the Question 6 sample ($34\,^\circ\mathrm{C}$, $21.78\,\mathrm{g\,kg^{-1}}$) to $20\,^\circ\mathrm{C}$: how much liquid water forms per kilogram of air?

Answers to Practice Set 22

22.1 (a) Table 1: $16\,^\circ\mathrm{C}$ holds 11.56 and $18\,^\circ\mathrm{C}$ holds 13.162, so $T_d = 16 + 2 \times (12 - 11.56)/(13.162 - 11.56) = 16.55 \approx 16.6\,^\circ\mathrm{C}$. (b) $w_s(30) = 27.694$, RH $= 12/27.694 = 43\,\%$; the rule gives $T - T_d = (100 - 43)/5 = 11.4$, so $T_d \approx 18.6\,^\circ\mathrm{C}$. The rule is only rough at low humidity; the table answer is better.

22.2 $w = 0.50 \times 14.956 = 7.478\,\mathrm{g\,kg^{-1}}$. Table 1: $8\,^\circ\mathrm{C}$ holds 6.771 and $10\,^\circ\mathrm{C}$ holds 7.762, so $T_d = 8 + 2 \times (7.478 - 6.771)/(7.762 - 6.771) = 9.4\,^\circ\mathrm{C}$. (The rule of thumb gives $10\,^\circ\mathrm{C}$.)

22.3 $w = 0.40 \times 24.557 = 9.82\,\mathrm{g\,kg^{-1}}$. At $12\,^\circ\mathrm{C}$ the air can hold only $8.882\,\mathrm{g\,kg^{-1}}$, which is less: so yes, dew (or fog) forms. The dew point is $13.5\,^\circ\mathrm{C}$, so dew begins when the air

cools through 13.5 °C.

22.4 $125 \times (30 - 18) = 1500\,\mathrm{m}$.

22.5 $w = w_s(15\,^\circ\mathrm{C})$ (the air is just saturated at its dew point) $= 10.83\,\mathrm{g\,kg^{-1}}$. $w_s(25) = 20.44$, so RH $= 10.83/20.44 = 53\,\%$. (Rule: $100 - 5 \times 10 = 50\,\%$.)

22.6 $21.78 - 14.956 = 6.8\,\mathrm{g\,kg^{-1}}$ of liquid water.

Chapter 23

Why air moves: the three forces

This chapter is the background for Part II of the sheet. It also completes the small table of forces printed at the start of "Atmospheric motion".

23.1 Newton's law for a parcel of air

Wind is air that is moving, and air moves for the same reason anything moves: forces act on it. Newton's second law says that the acceleration of a parcel equals the total force on it divided by its mass. It is convenient to talk about the force *per unit mass* (that is, as an acceleration, in $\mathrm{m\,s^{-2}}$), so we never need the mass of the parcel.

For the large-scale winds on the sheet only three horizontal forces matter:

1. the **pressure gradient force**, which starts the air moving;
2. the **Coriolis force**, which steers it once it is moving;
3. **friction**, which slows it near the ground.

If the three add up to zero the wind is steady, and the whole of Part II is about what happens when they do.

23.2 Pressure, and why it falls with height

The pressure at any level is the weight of the air above it, spread over one square metre. At sea level that weight is about 10 tonnes per square metre, which is a pressure of about 1013 hPa. Climb, and there is less air above you, so the pressure drops. Half the atmosphere's mass lies below about 5.5 km, so the pressure there is about 500 hPa; at the cruising height of airliners it is about 250 hPa.

23.3 The pressure gradient force

Pressure also varies from place to place at the same height. Air is pushed from where the pressure is higher towards where it is lower, just as water runs downhill. The push is the **pressure gradient force** (PGF). Per unit mass it is

$$\mathrm{PGF} = -\frac{1}{\rho}\,\frac{\Delta p}{\Delta n}, \tag{23.1}$$

where

PGF	pressure gradient force per unit mass ($\mathrm{m\,s^{-2}}$)	calculated
ρ	air density ($\mathrm{kg\,m^{-3}}$)	about 1.2 near the ground
$\Delta p/\Delta n$	pressure change per unit distance measured *across* the isobars ($\mathrm{Pa\,m^{-1}}$)	from the map

The minus sign means that the force points down the gradient, from high to low pressure (Holton and Hakim, 2013, ch. 1).




Figure 23.1. Pressure falls roughly exponentially with height, by a factor of e every 8 km or so: the halving rule of Practical 1 (Part I, Chapter 3). The red dots mark the standard pressure levels used on upper-air charts; those on the sheet (850, 500 and 300 hPa) are at about 1.5, 5.5 and 9 km.

- **Direction:** from higher to lower pressure, at right angles to the isobars.
- **Magnitude:** proportional to the pressure difference between two points, and inversely proportional to the distance between them. Closely spaced isobars mean a strong force and a strong wind.




Figure 23.2. The same pressure difference (here 12 hPa) squeezed into a short distance (left) and spread over a long one (right). The force, shown by the red arrows, is proportional to the gradient.

Worked example: how big is the pressure gradient force?

The isobars of Figure 5 on the sheet are 1012 and 1016 mb, that is 4 hPa = 400 Pa apart. Suppose they are 200 km apart (2×10^5 m). Then

$$\text{PGF} = \frac{1}{1.2} \times \frac{400}{2 \times 10^5} = 1.7 \times 10^{-3}\ \text{m s}^{-2}.$$

That looks tiny, but acting for an hour (3600 s) with nothing to oppose it, it would add 6 m s^{-1} to the wind speed. That is how quickly a pressure difference sets the air going.

23.4 The Coriolis force

The Earth turns once a day, and we live on its surface, so we measure motion from a spinning frame. An object that moves freely in a straight line (as seen from space) appears to us to curve. We describe the curving with a force, the **Coriolis force**, which has three properties you must remember.

1. **Direction:** at right angles to the motion. It acts to the **right** in the Northern Hemisphere and to the **left** in the Southern Hemisphere.
2. **Magnitude:** it increases with the wind speed V and with latitude: Coriolis $= f\,V$, where $f = 2\Omega \sin\varphi$ is the Coriolis parameter and $\Omega = 7.292 \times 10^{-5}\,\mathrm{s}^{-1}$ is the Earth's rotation rate.
3. **It does no work.** Being at right angles to the motion, it changes the direction of the wind but never its speed.



Figure 23.3. The Coriolis force always acts at right angles to the motion: to the right in the north and to the left in the south.




Figure 23.4. The Coriolis parameter against latitude. It is zero at the equator, where the force vanishes, and largest at the poles.

A question that comes up

Does the Coriolis force make water spin the opposite way in a bath in the two hemispheres?
No. For a bath or a sink the force is minute compared with the shape of the basin and the way the water was stirred. The Coriolis force matters for motions that last hours and cover hundreds of kilometres: winds, ocean currents, storms.

Table 23.1. The Coriolis parameter $f = 2\Omega \sin\varphi$ and the acceleration it gives to a wind of $10\,\mathrm{m\,s^{-1}}$.

Latitude	f ($\mathrm{s^{-1}}$)	Coriolis acceleration for $10\,\mathrm{m\,s^{-1}}$ ($\mathrm{m\,s^{-2}}$)
0°	0	0
10°	2.5×10^{-5}	2.5×10^{-4}
30°	7.3×10^{-5}	7.3×10^{-4}
45°	1.03×10^{-4}	1.03×10^{-3}
60°	1.26×10^{-4}	1.26×10^{-3}
90°	1.46×10^{-4}	1.46×10^{-3}

A question that comes up

Why is the Coriolis force zero at the equator?
It depends on the *vertical* component of the Earth's rotation, which is $\Omega \sin\varphi$. At the equator the rotation axis is horizontal, so that component is zero.

23.5 Friction

Near the ground the air rubs against the surface: grass, trees, buildings, hills, waves. The result is a **frictional force** that always acts **opposite to the wind**, slowing it. Its strength depends on how rough the surface is (a forest or a city is much rougher than a calm sea) and increases with the wind speed. Friction is important only in the lowest kilometre or two of the atmosphere, the *boundary layer*. Above that the air is nearly free of it, which is why the winds at 500 or 300 hPa are so much simpler than those at the surface.

23.6 The sheet's table, completed

The sheet starts Part II with a table of the three forces. Here it is with the entries filled in.

Table 23.2. The three forces (the table "Recall the following from the lectures").

Force	Direction	Magnitude
Pressure gradient	From higher to lower pressure, at right angles to the isobars	Proportional to the pressure difference between two points (and inversely to their distance)
Coriolis	At right angles to the airflow: to the **right** in the Northern Hemisphere, to the **left** in the Southern Hemisphere	Increases with wind strength and with latitude (fV with $f = 2\Omega \sin\varphi$)
Friction	Opposite to the direction of the airflow	Depends on the surface characteristics that obstruct the airflow (and increases with speed)

23.7 Three situations to hold in your head




Figure 23.5. (1) At rest only the pressure gradient force acts. (2) High above the ground the Coriolis force balances it and the wind blows along the isobars: the geostrophic wind (Chapter 25). (3) Near the ground friction joins in, the three forces cancel, and the wind crosses the isobars towards low pressure (Chapter 26). Red: pressure gradient force; blue: Coriolis; green: friction; black: the wind.

23.8 Try it yourself: Practice Set 23

Practice Set 23: forces

23.1 Two isobars are 4 hPa apart and 100 km from each other. Find the pressure gradient force per unit mass, taking $\rho = 1.2\,\mathrm{kg\,m^{-3}}$. How does it compare with the 200 km case in the worked example?

23.2 Find the Coriolis parameter at 60° N and the Coriolis acceleration on a wind of $15\,\mathrm{m\,s^{-1}}$.

23.3 A wind is blowing northward in the Southern Hemisphere. Which way does the Coriolis force push it?

23.4 Why are strong winds found where the isobars are close together?

Answers to Practice Set 23

23.1 PGF $= \dfrac{1}{1.2} \times \dfrac{400}{10^5} = 3.3 \times 10^{-3}\ \mathrm{m\,s^{-2}}$: exactly *twice* the 200 km value, because the distance is halved.

23.2 $f = 2 \times 7.292 \times 10^{-5} \times \sin 60^\circ = 1.26 \times 10^{-4}\ \mathrm{s^{-1}}$; $fV = 1.26 \times 10^{-4} \times 15 = 1.9 \times 10^{-3}\ \mathrm{m\,s^{-2}}$.

23.3 To the left of its motion, which is **west**.

23.4 Close isobars mean a steep pressure gradient, so the pressure gradient force is large; the Coriolis force (and, near the ground, friction) must grow to balance it, which needs a faster wind.

Chapter 24

Pressure surfaces and the pressure gradient

This chapter answers Motion Q1 to Q6 of the sheet (Figures 1 and 2).

24.1 What a pressure surface is

Upper-air charts do not show pressure at a fixed height; they show the *height at which the pressure has a chosen value*. The chart for 500 hPa, for example, is a map of the height of the surface on which the pressure is 500 hPa. Such a surface is called a **pressure surface** (or isobaric surface). Where the pressure surface is high, the pressure at a fixed level nearby is high; where it is low, the pressure is low. The height contours on the chart therefore play the part of isobars, and the wind blows along them (Chapter 25).

Pressure surfaces are not flat. They are high over warm air and low over cold air, and that single fact explains Questions 1 to 6.

24.2 Why cold air gives low pressure surfaces

Pressure at a point is the weight of the air above it. A column of cold air is denser, so a given amount of air (a given pressure drop) occupies less height. A column of warm air is less dense, so the same pressure drop is spread over a greater height: the warm column is thicker. Start both columns with the same pressure at the ground and the 500 hPa surface will be lower over the cold one than over the warm one, and the difference grows with height.

The exact relationship is the **hypsometric equation** (Wallace and Hobbs, 2006, ch. 3). It is the exponential law of Practical 1, $p = p_0 e^{-z/H}$ with $H = R_d T/g$ (Part I, Eq. (3.3)), written for a layer: for the layer between pressures p_1 (lower) and p_2 (upper),

$$\Delta z = z_2 - z_1 = \frac{R_d \overline{T}}{g} \ln \frac{p_1}{p_2}, \tag{24.1}$$

where

Δz	thickness of the layer (m)	calculated
$\overline{T}$	mean temperature of the layer (K)	
R_d	gas constant for dry air ($\mathrm{J\,kg^{-1}\,K^{-1}}$)	287
g	acceleration due to gravity ($\mathrm{m\,s^{-2}}$)	9.81
$p_1,\ p_2$	pressures at the bottom and top of the layer (hPa)	

The thickness Δz is directly proportional to the mean temperature. Practical 1 used the same law the other way round, to find the scale height of a layer from a sounding (Part I, Eq. (8.1)), and to explain why the troposphere is thickest over the warm tropics (Part I, § 7.4).

Worked example: thickness of the 1000 to 500 hPa layer

For a cold layer with $\overline{T} = -10\,^{\circ}\mathrm{C} = 263.15\,\mathrm{K}$,

$$\Delta z = \frac{287 \times 263.15}{9.81} \ln \frac{1000}{500} = 7699 \times 0.693 = 5339\ \mathrm{m}.$$

For a warm layer with $\overline{T} = +10\,^{\circ}\mathrm{C} = 283.15\,\mathrm{K}$ the result is 5745 m. A 20-degree difference in mean temperature makes the layer 406 m thicker, about 20 m per degree.




Figure 24.1. Two columns of air that start with the same pressure at the ground (1013 hPa). Each pressure level sits higher over the warm column, and the difference (the slope of the surface) grows with height: 103 m at 850 hPa, 216 m at 700 hPa, 413 m at 500 hPa and 712 m at 300 hPa. (The heights come from the hypsometric equation with a constant mean temperature.)

24.3 The data of Figure 1

Figure 1 of the sheet gives the pressure level, the height and the temperature at two places, Saskatoon in Canada and Monterrey in Mexico (about 2 990 km apart, Monterrey to the south):

Notice that Monterrey is warmer at every level, and every one of its pressure surfaces is higher.

Table 24.1. The data of Figure 1.

	Saskatoon		Monterrey	
Pressure	**Height**	**Temp.**	**Height**	**Temp.**
300 mb	8700 m	−47 °C	9500 m	−38 °C
500 mb	5400 m	−25 °C	5700 m	−12 °C
850 mb	1360 m	−1 °C	1500 m	15 °C

24.4 Question 1: plot the pressure surfaces

The question on the sheet

Following the example of 850 mb, plot the heights of the 500 and 300 mb levels on Figure 2, draw lines connecting the two points for each level, and label each line.

Method.

Step 1. Figure 2 has Saskatoon on the left and Monterrey on the right, with height up the sides (0 to 10 000 m). The 850 mb line, from 1360 m to 1500 m, is already drawn as the example.

Step 2. For 500 mb put a dot at 5400 m above Saskatoon and a dot at 5700 m above Monterrey. Join them with a straight line and label it "500 mb".

Step 3. For 300 mb put dots at 8700 m (Saskatoon) and 9500 m (Monterrey), join them, and label the line "300 mb".




Figure 24.2. Figure 2 completed for Question 1. The 500 and 300 mb lines slope up towards Monterrey, and the slope is greater for the higher level.

Answer to Motion Q1

Two lines are added to Figure 2: 500 mb from 5400 m (Saskatoon) to 5700 m (Monterrey), and 300 mb from 8700 m to 9500 m. Each rises to the right, more steeply than the one below it.

24.5 Questions 2 and 3: where does pressure fall fastest?

Answer to Motion Q2

Saskatoon. Its pressure surfaces are lower: the 500 mb surface is reached at 5400 m instead of 5700 m, the 300 mb surface at 8700 m instead of 9500 m. Less height is needed to use up the same pressure, so pressure falls faster with height.

Answer to Motion Q3

Saskatoon is colder at each height (-1 against $15\,^\circ$C at 850 mb, -25 against $-12\,^\circ$C at 500 mb, -47 against $-38\,^\circ$C at 300 mb). Cold air is denser, so a column of it is thinner: the pressure falls faster with height.

Check with the hypsometric equation. Use the mean of the two temperatures that bound each layer.

$$\text{Saskatoon: } \Delta z_{850\to300} = 29.27\,[\,260.15\ln\tfrac{850}{500} + 237.15\ln\tfrac{500}{300}\,] = 29.27\,[138.0 + 121.1] = 7587\text{ m},$$
$$\text{Monterrey: } \Delta z_{850\to300} = 29.27\,[\,274.65\ln\tfrac{850}{500} + 248.15\ln\tfrac{500}{300}\,] = 29.27\,[145.7 + 126.8] = 7976\text{ m}.$$

The sheet's figures give $8700 - 1360 = 7340$ m and $9500 - 1500 = 8000$ m. Monterrey agrees to 0.3 %; Saskatoon is 3 % off, because the figure's numbers are rounded. The warmer column is thicker, as predicted.

24.6 Question 4: which point has the higher pressure?

The question on the sheet

Both points x and z are at 9000 m. Which has the highest pressure? At 9000 m write H where the pressure is higher and L where it is lower, then draw an arrow between them showing the direction of the horizontal pressure gradient force.

Method.

Step 1. Pressure decreases with height. At a given height, being *below* a pressure surface means a *higher* pressure than the surface's value, and being *above* it means a lower pressure.

Step 2. Point x is above Saskatoon at 9000 m. The 300 mb surface there is at 8700 m, so x is above it: its pressure is *less than* 300 mb.

Step 3. Point z is above Monterrey at 9000 m. The 300 mb surface there is at 9500 m, so z is below it: its pressure is *more than* 300 mb.

Step 4. So z has the higher pressure: write H at z (the Monterrey side) and L at x (the Saskatoon side).

Step 5. The pressure gradient force points from high to low: draw the arrow from z towards x, that is towards Saskatoon.




Figure 24.3. Question 4. At 9000 m x is above the 300 mb line (pressure below 300 mb, L) and z is below it (pressure above 300 mb, H). The pressure gradient force points from z to x: from Monterrey towards Saskatoon.

Answer to Motion Q4

z has the higher pressure, because it lies below the 300 mb line at 9000 m. Write **H at z** and **L at x**, and draw the pressure gradient force from z towards x (towards Saskatoon).

How big are the pressures? The sheet does not ask, but you can estimate them. Above the 300 mb level at Saskatoon, pressure falls about 1 % every 66 m, so 300 m above the 8700 m level it is about $300\,e^{-0.045} = 287$ mb. At Monterrey, 9000 m is 500 m below the 9500 m level of 300 mb, so the pressure is higher, about 321 mb. The difference between the two places is therefore about 34 mb, over 2 990 km.

24.7 Question 5: where is the pressure gradient greater?

Answer to Motion Q5

The pressure gradient is greater at **9000 m**. The pressure surfaces differ in height by $9500 - 8700 = 800$ m between the two places at the 300 mb level, but only by $1500 - 1360 = 140$ m at the 850 mb level (about 1500 m).

The slope of the surfaces is what drives the wind. The slope grows with height: 140 m at 850 mb, 300 m at 500 mb and 800 m at 300 mb. If you compare pressures at a fixed height instead, you find the same thing: about 15 mb between the two places at 1500 m, but about 34 mb at 9000 m.

A question that comes up

The pressure itself is smaller at 9000 m. How can the gradient be greater? Because what drives the wind is the *change* of pressure from place to place, and that depends on the temperature difference accumulated in the whole layer below. On a pressure surface the force per unit mass is $-g\,\Delta z/\Delta n$, and it does not depend on the air density at all. The slope $\Delta z/\Delta n$ grows with height as the colder column's thickness deficit adds up, so the force




Figure 24.4. The slope of the pressure surfaces between Saskatoon and Monterrey increases with height.

does too.

24.8 Question 6: what this means for the wind

Answer to Motion Q6

Wind speeds at **9000 m should be greater**, because the pressure gradient force is greater there.

You can put numbers on this. The geostrophic wind (Chapter 25) is $V_g = (g/f)\,\Delta z/\Delta n$, Eq. (25.1), where Δz is the difference in height of a pressure surface over a distance Δn. The two places are $\Delta n = 2\,990$ km apart, at a mean latitude of 39°, where $f = 2 \times 7.292 \times 10^{-5} \times \sin 39° = 9.16 \times 10^{-5}\,\mathrm{s}^{-1}$.

Worked example: wind estimates from the slopes

$$850\,\mathrm{mb}: \quad V_g = \frac{9.81}{9.16\times10^{-5}} \times \frac{140}{2.99\times10^{6}} = 5.0\ \mathrm{m\,s^{-1}},$$

$$500\,\mathrm{mb}: \quad V_g = \frac{9.81}{9.16\times10^{-5}} \times \frac{300}{2.99\times10^{6}} = 10.7\ \mathrm{m\,s^{-1}},$$

$$300\,\mathrm{mb}: \quad V_g = \frac{9.81}{9.16\times10^{-5}} \times \frac{800}{2.99\times10^{6}} = 28.6\ \mathrm{m\,s^{-1}}.$$

The wind blows from west to east (Monterrey, with the higher surfaces, is to the south, on the right of the wind in the Northern Hemisphere).

These numbers are rough: the sheet's heights are rounded, and only two places are used. Real winds at 300 mb are often 50 to 80 $\mathrm{m\,s^{-1}}$. The lesson is nevertheless exact. **The faster the temperature falls towards the pole, the faster the westerly winds aloft, and they strengthen with height.** This is the **thermal wind**, and it is why the jet streams sit near 10 km above the zone where the temperature contrast is greatest.




Figure 24.5. Geostrophic wind estimated from the slope of the pressure surfaces between the two places. The wind speeds up fivefold from 850 to 300 mb.

24.9 An experiment: change the temperatures

Replace Saskatoon and Monterrey by two columns that both start at 1013 hPa and have constant mean temperatures T_1 and T_2. The hypsometric equation then gives the height of each surface. If $T_1 = T_2$ every surface is flat: there is no horizontal pressure gradient and no wind, however cold the air. Increase the contrast and the surfaces tilt, increasingly steeply with height.

Table 24.2. Surface slopes and geostrophic winds for two columns with mean temperatures of −25 °C and 0 °C, 2 000 km apart at 45° latitude.

Level	Height, cold column	Height, warm column	Slope / wind
850 hPa	1274 m	1403 m	128 m / 6 m s^{-1}
700 hPa	2685 m	2955 m	270 m / 13 m s^{-1}
500 hPa	5129 m	5645 m	517 m / 25 m s^{-1}
300 hPa	8839 m	9729 m	890 m / 42 m s^{-1}

24.10 Try it yourself: Practice Set 24

Practice Set 24: pressure surfaces

24.1 Find the thickness of the 1000 to 500 hPa layer for mean temperatures of −10 °C and +10 °C, and the difference.

24.2 At 40° N the 500 hPa surface is at 5520 m over town A and 5640 m over town B, which lies 800 km due south of A. Estimate the geostrophic wind and say from which direction it blows.

24.3 The 500 mb heights over two stations are 5400 m (P) and 5600 m (Q). Which has the colder air beneath it?

24.4 The 500 hPa surface slopes up from north to south. Which way does the wind blow in the Northern Hemisphere?

Answers to Practice Set 24

24.1 5339 m and 5745 m (see the worked example); the warm layer is 406 m thicker.

24.2 $f = 2 \times 7.292 \times 10^{-5} \sin 40^\circ = 9.37 \times 10^{-5}\,\mathrm{s}^{-1}$; $V_g = \dfrac{9.81}{9.37 \times 10^{-5}} \times \dfrac{120}{8 \times 10^5} = 15.7\ \mathrm{m\,s}^{-1}$ (about 30 kt). B is to the south and has the higher surface, so the wind has high pressure on its right: it is a **westerly** (blowing from west to east).

24.3 P, because a column that is colder is thinner and has lower pressure surfaces.

24.4 With the low heights to the north (on the left of the wind), the wind blows from west to east, a westerly. Pressure surfaces that rise towards the equator give westerly winds, which is the pattern over mid-latitudes.

Chapter 25

The geostrophic wind

This chapter answers Motion Q7 of the sheet (Figures 3 and 4).

25.1 Two forces in balance

High above the ground, friction can be ignored. Only the pressure gradient force and the Coriolis force act. Suppose they are equal in size and opposite in direction. Then the total force is zero, the air does not accelerate, and the wind is steady. That wind is the **geostrophic wind** ("earth-turning" wind).

The geostrophic wind

The geostrophic wind is the wind for which the pressure gradient force and the Coriolis force exactly balance. It blows *along* the isobars (or the height contours on a pressure-surface chart), with low pressure on its left in the Northern Hemisphere and on its right in the Southern Hemisphere.

Writing the two forces per unit mass, the balance is

$$\underbrace{\frac{1}{\rho}\frac{\Delta p}{\Delta n}}_{\text{pressure gradient force}} = \underbrace{f\,V_g}_{\text{Coriolis force}},$$

so that the geostrophic wind speed is (Holton and Hakim, 2013, ch. 3)

$$V_g = \frac{1}{\rho f}\frac{\Delta p}{\Delta n} = \frac{g}{f}\frac{\Delta z}{\Delta n}. \tag{25.1}$$

where

V_g	geostrophic wind speed ($\mathrm{m\,s^{-1}}$)	calculated
ρ	air density ($\mathrm{kg\,m^{-3}}$)	about 1.2 near the ground
f	Coriolis parameter, $2\Omega\sin\varphi$ ($\mathrm{s^{-1}}$)	$\Omega = 7.292\times10^{-5}\,\mathrm{s^{-1}}$
$\Delta p/\Delta n$	pressure gradient across the isobars ($\mathrm{Pa\,m^{-1}}$)	from a surface map
$\Delta z/\Delta n$	slope of a pressure surface ($\mathrm{m\,m^{-1}}$)	from an upper-air chart
g	acceleration due to gravity ($\mathrm{m\,s^{-2}}$)	9.81

The last form is for a pressure surface, where Δz is the difference in height of the surface over the distance Δn. It is the form used with the 500 hPa heights of the sheet.

25.2 How the balance comes about: the four stages of Figure 3

Figure 3 of the sheet follows a parcel of air as the balance is established. Think of a parcel of air that is at rest between a region of higher pressure (south) and lower pressure (north) in the Northern Hemisphere.

Stage 1. At rest. The parcel is on the high-pressure side. No motion means no Coriolis force, so only the pressure gradient force acts, pointing north, towards low pressure. The parcel starts to move northward.

Stage 2. Moving northward. As soon as it moves, the Coriolis force appears, at right angles and to the right of the motion: towards the east. The parcel's path bends to the right.

Stage 3. Moving north-east. The speed has grown, so the Coriolis force is larger, and since it is at right angles to the motion it now points partly south, against the pressure force. The path keeps bending.

Stage 4. Moving east. The parcel is now moving along the isobars. The Coriolis force points exactly south, equal and opposite to the pressure gradient force pointing north. The forces balance, the speed stops growing, and the wind blows steadily to the east.




Figure 25.1. The four stages of Figure 3, computed for two 500 hPa height lines 60 m and 500 km apart at 45° N. The parcel starts at rest on the high side. At each numbered position the red arrow is the pressure gradient force, the blue arrow is the Coriolis force and the black arrow is the wind. The path bends to the right until the wind is along the isobars and the forces balance (stage 4).

A question that comes up

Why does the parcel not just keep accelerating towards low pressure?
Because as soon as it speeds up, the Coriolis force grows in proportion, and it turns the motion aside. Without any damping the parcel would overshoot and loop around the balance state (an *inertial oscillation* with a period of $2\pi/f$, about 17 hours at 45°). In the real atmosphere the oscillations are damped as the adjustment radiates energy away in waves, and the air settles to the balance. The figure above includes a little damping, which pulls the velocity towards the geostrophic wind, so that the settling is visible. Its final state is the geostrophic wind.




Figure 25.2. The two components of the parcel's velocity against time. The eastward component rises to the geostrophic value (dashed) and the northward component dies away to zero: the air ends up blowing along the isobars.

25.3 Question 7 (Figure 3): the Northern Hemisphere

The question on the sheet

Mark the side with the higher pressure with an H and the side with the lower pressure with an L.

Method.

Step 1. The lines are labelled with the *height* of the 500 hPa surface: 5460 m on the upper line and 5520 m on the lower line.

Step 2. A pressure surface is higher where the pressure is higher. So the line with the larger height, 5520 m, is on the high-pressure side. Write **H** at the 5520 m line (bottom).

Step 3. Write **L** at the 5460 m line (top).

Step 4. Check with the arrows: the pressure gradient force F_{pg} is drawn pointing up, from H to L, as it should.

Answer to Motion Q7, part 1 (Figure 3)

H at the 5520 m line (south, bottom); L at the 5460 m line (north, top). The pressure gradient force points from H to L (north); the Coriolis force points to the right of the wind, south; the geostrophic wind blows to the east, with low pressure on its left.

25.4 Question 7 (Figure 4): the Southern Hemisphere

The question on the sheet

Now consider the Southern Hemisphere. Mark H and L, and in Figure 4 draw the pressure gradient force, the Coriolis force and the resulting geostrophic wind.

What changes. In the Southern Hemisphere the Coriolis force acts to the **left** of the motion. Everything else is the same: the pressure gradient force still points from high to low, and the balance must still be between two opposite forces. The result is a mirror image of Figure 3.

Step 1. The two lines are 5640 m (top) and 5580 m (bottom). The larger height is the high-pressure side: **H** at the 5640 m line (top), **L** at the 5580 m line (bottom).

Step 2. Draw the pressure gradient force F_{pg} from H to L: **downwards**.

Step 3. The Coriolis force must be equal and opposite, so it points **upwards**, towards H. (To check: an eastward wind has its left-hand side to the north, which is up the page. The Coriolis force acts to the left, so it points up. ✓)

Step 4. The wind blows between them, to the **east**, with the low pressure on its right.




Figure 25.3. Figure 4: the Southern Hemisphere. The Coriolis force turns the parcel to the left. At the end (stage 4) the pressure gradient force points down towards L, the Coriolis force points up towards H, and the wind blows to the east with the low pressure on its right.

Answer to Motion Q7, part 2 (Figure 4)

H at the 5640 m line (top); L at the 5580 m line (bottom). Draw F_{pg} **down** (towards L) and the Coriolis force **up** (towards H), of equal length; the geostrophic wind **W** points **east** between them.




Figure 25.4. The two answers side by side. In both figures the wind blows east. In the north the low pressure (5460 m) is on its left; in the south the low pressure (5580 m) is on its right.

25.5 Buys-Ballot's law

You can find where the low pressure lies, without a map, from the wind alone:

- **Northern Hemisphere:** stand with your back to the wind; low pressure is on your **left**.
- **Southern Hemisphere:** stand with your back to the wind; low pressure is on your **right**.

This is Buys-Ballot's law. It follows directly from the balance: the pressure gradient force points towards low pressure and the Coriolis force (to the right in the north) points away from it, so low pressure is on the left of the wind.




Figure 25.5. Buys-Ballot's law. With your back to the wind, low pressure is on the left in the north and on the right in the south.

25.6 How fast is the geostrophic wind?

Worked example: the wind implied by the lines of Figure 3

Suppose the two height lines, 60 m apart, are 500 km apart on the chart, at 45° N. Then

$$f = 2 \times 7.292 \times 10^{-5} \sin 45^\circ = 1.03 \times 10^{-4}\ \text{s}^{-1},\quad V_g = \frac{g\,\Delta z}{f\,\Delta n} = \frac{9.81 \times 60}{1.03 \times 10^{-4} \times 5 \times 10^5} = 11.4\ \text{m s}^{-1}$$

which is 22 knots or 41 km h^{-1}. If the lines were closer, 300 km apart, the wind would be 19 m s^{-1}; at the same spacing but 60° N it would be 9.3 m s^{-1}; at 30° N, 16.1 m s^{-1}.




Figure 25.6. The geostrophic wind for a given height difference rises as the lines get closer together and as the latitude decreases (because f falls). Near the equator $f \to 0$ and the formula breaks down.

25.7 When is the wind not geostrophic?

The balance is an excellent approximation for the winds a kilometre or more above the ground at middle and high latitudes, but not always.

- **Near the ground,** friction acts, and the balance changes (Chapter 26).
- **Near the equator,** f is so small that the balance fails.
- **Where the isobars curve strongly,** as round a hurricane or a tight low, the centrifugal force matters; the result is the *gradient wind*.
- **When the pattern changes quickly,** the air accelerates, as in a developing storm.

25.8 Questions that come up

A question that comes up

Is the geostrophic wind along or across the isobars?
Along. Because the two forces are at right angles to the wind, and the pressure gradient force is at right angles to the isobars, the wind must be parallel to the isobars. This is a much better description of upper-air winds than the picture of air flowing "from high to low".

A question that comes up

On Figure 3, why is H at the bottom? Surely high pressure is up?
"Up" on the page is north, not height. The H and L mark the pressure on a horizontal map at the 500 hPa level: higher pressure to the south (bottom of the page), lower to the north (top).

25.9 Try it yourself: Practice Set 25

Practice Set 25: geostrophic wind

25.1 The 500 hPa height lines 5460 m and 5520 m are 400 km apart at 50° N. Find the geostrophic wind.

25.2 In the Southern Hemisphere the 500 hPa heights are 5600 m at 40° S and 5500 m at 50° S. Which way does the wind blow?

25.3 At what latitude would the lines of Figure 3 (60 m in 500 km) give a geostrophic wind of $20\ \mathrm{m\,s^{-1}}$?

25.4 In the Northern Hemisphere a pilot flying with the wind directly behind notes that the low pressure is on the pilot's left. Is this consistent with Buys-Ballot's law?

Answers to Practice Set 25

25.1 $f = 2 \times 7.292 \times 10^{-5} \sin 50^\circ = 1.117 \times 10^{-4}$; $V_g = \dfrac{9.81 \times 60}{1.117 \times 10^{-4} \times 4 \times 10^5} = 13.2\ \mathrm{m\,s^{-1}}$ (26 kt).

25.2 The heights fall towards the pole, so the low is to the south. In the Southern Hemisphere low pressure is on the right of the wind, so the wind blows from **west to east** (a westerly), as in the Southern Ocean.

25.3 $\sin\varphi = \dfrac{g\Delta z}{2\Omega V_g \Delta n} = \dfrac{588.6}{1458} = 0.404$, so $\varphi = 23.8°$.

25.4 Yes: with your back to the wind, low pressure on the left is exactly Buys-Ballot's law for the Northern Hemisphere.

Chapter 26

Wind near the ground: friction

This chapter answers Motion Q8 to Q11 of the sheet (Figures 5 and 6).

26.1 Question 8: what does a slower wind do to the Coriolis force?

Close to the ground friction acts on the air, always against the wind, slowing it. The sheet's first question about it is: *what effect will a reduction in surface wind speed have on the Coriolis force?*

Answer to Motion Q8

The Coriolis force becomes **weaker**, because it depends on the wind speed. A slower wind means a smaller Coriolis force; a faster wind a larger one.

The reason is that the Coriolis force per unit mass is fV, proportional to the wind speed V. If friction cuts the speed to 70 % of the geostrophic value, the Coriolis force falls to 70 % as well. But the pressure gradient force does not depend on the wind at all. So at the surface the Coriolis force can no longer balance the pressure gradient force on its own. Something has to make up the difference, and that is where the three-force balance comes from.

26.2 The balance of three forces

Near the ground three forces act on the air. If the wind is steady they add up to zero.

- the **pressure gradient force**, from high to low pressure;
- the **Coriolis force**, at right angles to the wind (to its right in the Northern Hemisphere);
- **friction**, directly against the wind.

The sheet draws one box of air with these arrows. To see why the balance forces the wind to cross the isobars, imagine that the wind were parallel to the isobars as in Chapter 25. Then the Coriolis force would point directly against the pressure force, and friction (against the wind, along the isobars) would be left with nothing to balance it. The only way out is for the wind to turn a little *towards low pressure*, so that the pressure force has a component along the wind that cancels friction, and the Coriolis force (now pointing partly backwards) cancels the rest.

26.3 Question 9: the forces in Figure 5

The question on the sheet

Indicate with arrows the three forces acting on the Northern Hemisphere box (pressure gradient in red, Coriolis in blue, friction in green); the black arrow shows the result.

Figure 5 shows two isobars, 1012 mb at the top and 1016 mb at the bottom. The pressure is lower at the top, so the pressure gradient force points *up*.

Method: draw the arrows in this order.

Step 1. Red: the pressure gradient force. From the box, straight towards the lower pressure (up the page), at right angles to the isobars.

Step 2. Black: the wind. It blows across the isobars, towards low pressure but turned to the right, so draw it up and to the right (north-east). Its angle to the isobars is typically 20 to 45 degrees; the sheet's example uses 45°.

Step 3. Blue: the Coriolis force. At right angles to the wind, on its right-hand side. For a north-east wind that is to the south-east (down and to the right).

Step 4. Green: friction. Opposite to the wind: pointing south-west (down and to the left).

Step 5. Check. Join the three coloured arrows tip to tail: they should make a closed triangle. If they do, the forces cancel and the wind is steady.




Figure 26.1. Figure 5 completed. Red: pressure gradient force; blue: Coriolis force; green: friction; black: the wind. The three forces, joined tip to tail on the right, close into a triangle: they cancel. The black arrow is the *wind*, the direction in which the air actually moves.

Answer to Motion Q9

Pressure gradient force (red): up, towards 1012 mb. Wind (black): up and to the right. Coriolis force (blue): at right angles to the wind, down and to the right. Friction (green): opposite to the wind, down and to the left.

A question that comes up

The sheet says the black arrow is the "resulting force". Is it a force?
In a steady wind the three forces add up to zero, so there is *no* resultant force: that is exactly why the wind is steady. What the black arrow shows is the resulting wind: the direction the air moves when the forces balance. It is the right arrow to draw; just do not think of it as a fourth force.

26.4 The numbers: how much does friction turn the wind?

Represent friction as a drag proportional to the wind, with coefficient k. The balance of the three forces then gives, for a pressure gradient force P,

$$\alpha = \arctan\frac{k}{f}, \qquad V = V_g\cos\alpha, \qquad V_g = \frac{P}{|f|}. \tag{26.1}$$

where

α	angle at which the wind crosses the isobars, towards low pressure (degrees)	calculated
k	friction coefficient (s^{-1})	larger over land than sea
f	Coriolis parameter (s^{-1})	
$V,\ V_g$	wind speed with friction, and geostrophic wind speed ($\mathrm{m\,s}^{-1}$)	
P	pressure gradient force per unit mass ($\mathrm{m\,s}^{-2}$)	Eq. (23.1)

The ratio k/f measures how strong friction is compared with the Coriolis force. The sheet's Figure 5 corresponds to $k/f = 1$.

Table 26.1. Friction and the surface wind (the ratio k/f is a guide; real values depend on the surface and the weather).

Surface	k/f	Angle across isobars	Speed V/V_g	Coriolis force (relative)
No friction (high above the ground)	0	0°	100 %	100 %
Open sea	0.3	17°	96 %	96 %
Flat grassland	0.6	31°	86 %	86 %
Forest or suburb	1.0	45°	71 %	71 %
Tall city	1.5	56°	55 %	55 %




Figure 26.2. The surface wind for different surfaces, for the same pressure gradient force (in the Northern Hemisphere, with low pressure to the north). The rougher the surface, the slower the wind and the greater its angle across the isobars towards low pressure. The length of each arrow is proportional to the wind speed.




Figure 26.3. The angle across the isobars (blue) and the speed (red) against the friction coefficient. The speed is $V_g \cos\alpha$.

26.5 Question 10: which system is which?

The question on the sheet

Figure 6 shows winds circulating round surface high and low pressure centres in both hemispheres. Label the correct hemisphere and pressure system for each of the four examples.

Two observations identify each system.

1. **Does the air spiral inwards or outwards?** Friction makes it cross the isobars towards low pressure, so it spirals *in* towards a low and *out* from a high.
2. **Which way does it turn?** In the Northern Hemisphere the Coriolis force deflects the air to the right, so the circulation round a low is *anticlockwise* and round a high *clockwise*. In the Southern Hemisphere it is the other way round.

Table 26.2. The four systems of Figure 6.

System	Rotation	Air spirals	Pressure gradient force
Northern low	anticlockwise	in	towards the centre
Northern high	clockwise	out	away from the centre
Southern low	clockwise	in	towards the centre
Southern high	anticlockwise	out	away from the centre

Answer to Motion Q10

In the order of Figure 6: top left **Northern Hemisphere low**; top right **Northern Hemisphere high**; bottom left **Southern Hemisphere low**; bottom right **Southern Hemisphere high**.




Figure 26.4. A two-question key to the four systems: in or out first, then the sense of rotation.

26.6 Question 11: the forces on one box in each system

The question on the sheet

Choose one box on each example and draw the pressure gradient force in red, the Coriolis force in blue, and the frictional force in green. The black arrow shows the result.

Method. For the box you choose:

Step 1. Draw the pressure gradient force (red): towards the centre for a low, away from it for a high.

Step 2. Draw the wind (black): follow the arrows already on the figure, spiralling in for a low and out for a high.

Step 3. Draw the Coriolis force (blue) at right angles to the wind: to its right in the Northern Hemisphere, to its left in the Southern Hemisphere.

Step 4. Draw friction (green) directly opposite to the wind.

Step 5. Check that the three arrows could be joined tip to tail into a triangle.

Answer to Motion Q11

On every box the red arrow points from high to low pressure; the black wind arrow is turned from it, to the right in the north and the left in the south, and pulled towards low pressure by friction; the blue arrow is at right angles to the wind on the same side as the turn; the green arrow is exactly opposite to the wind. The pattern is identical at every position round the system, so any box gives the same picture rotated.

26.7 Why this matters for weather

Because the air spirals in towards a low, more air flows in at the bottom than flows out sideways, so the surplus must rise: surface **convergence** gives rising air, cloud and rain. In a high the air spirals out, so air sinks from above to replace it, which warms and dries it and gives clear skies. This is why lows (depressions) bring cloud and rain and highs bring settled weather, and it is the link between this chapter and the weather map you analyse in Chapter 29.




Figure 26.5. Figure 6 completed. In each panel the grey arrows show the surface wind, spiralling in (low) or out (high), and the circles are isobars. On the box at the south-east of each system: red, the pressure gradient force (always along the line to the centre); blue, the Coriolis force (right of the wind in the north, left of the wind in the south); green, friction (against the wind); black, the wind. Any other box would show the same pattern turned round the system.

26.8 Questions that come up

A question that comes up

Why does the wind cross the isobars *towards* low pressure and not away from it?
Because friction removes some of the Coriolis balance. The pressure gradient force is now larger than the Coriolis force, and the surplus pushes the air down the pressure gradient. The result is a wind with a component towards low pressure.

A question that comes up

Why is the angle bigger over land than over the sea?
Land is rougher (trees, buildings, hills), so friction is stronger and k/f is larger. Over the open sea the angle is only 10 to 20 degrees; over land it can be 30 to 45 degrees or more.




Figure 26.6. Cross-sections. Near the surface the winds converge into a low and diverge from a high; the continuity of air then requires rising motion over the low and sinking over the high.

26.9 Try it yourself: Practice Set 26

Practice Set 26: friction

26.1 Over a flat grassy plain ($k/f = 0.6$) the geostrophic wind is $15\,\mathrm{m\,s^{-1}}$. Find the surface wind speed and its angle to the isobars.

26.2 In a Southern Hemisphere low, in which direction does the air circulate, and does it spiral in or out?

26.3 Describe the winds round a system in which the air spirals outwards in an anticlockwise sense. Which hemisphere and which kind of system is it?

26.4 Over a city the surface wind is $6\,\mathrm{m\,s^{-1}}$ and the geostrophic wind above it is $11\,\mathrm{m\,s^{-1}}$. Roughly what is k/f?

Answers to Practice Set 26

26.1 $\alpha = \arctan 0.6 = 31°$; $V = 15\cos 31° = 12.9\ \mathrm{m\,s^{-1}}$.

26.2 Clockwise, spiralling **in**.

26.3 Anticlockwise and outward is a **Southern Hemisphere high** (Table above).

26.4 $V/V_g = 6/11 = 0.55 = \cos\alpha$, so $\alpha \approx 57°$ and $k/f = \tan 57° \approx 1.5$: a tall-city value.

Chapter 27

Reading a station model

This chapter answers Map Q12 of the sheet (Figure 7 and the four models).

27.1 Why a station model?

About 10 000 weather stations around the world measure as many as 18 quantities at each observation: temperature, dew point, pressure, wind, cloud, weather, and more. To show all that on one map the World Meteorological Organization (WMO) fixed a compact way of drawing an observation, the **station model**. Every number and symbol has a fixed position around a small circle that marks the station. Once you know the positions, a map becomes a table you can read at a glance.

27.2 The anatomy of a station model




Figure 27.1. A station model with every field labelled. This example reads: temperature 45 °F, dew point 38 °F, pressure 1014.7 hPa, rising by 2.8 hPa in the last three hours, three-quarters of the sky covered, continuous rain, and a wind of 20 knots from the north-west.

27.3 Decoding the pressure

Only three digits are printed, in tenths of a hectopascal, and the leading digits are left off. Sea-level pressure almost always lies between about 950 and 1050 hPa, so the missing digits can be restored without doubt.

Table 27.1. Where each quantity sits around the circle.

Position	What it shows
Top left	air temperature, in °F on the sheet
Bottom left	dew-point temperature
Top right	pressure: three digits (tenths of a hectopascal)
Right, below the pressure	pressure change over the last 3 hours (with a symbol for its pattern)
Left of the circle	present weather symbol
The circle itself	sky cover (how much is filled)
A line from the circle	wind: the staff points **from** where the wind blows, barbs give the speed

The pressure rule

1. Put a decimal point before the last digit: 117 → 11.7.
2. If the three-digit number is **less than 500**, put **10** in front: 11.7 → 1011.7 hPa.
3. If it is **500 or more**, put **9** in front: 998 → 99.8 → 999.8 hPa.

Table 27.2. Examples of the pressure rule (the first four are the models of the sheet).

Printed	Decimal point	< 500 or ≥ 500?	Prefix	Pressure
024	02.4	< 500	10	**1002.4 hPa**
998	99.8	≥ 500	9	**999.8 hPa**
117	11.7	< 500	10	**1011.7 hPa**
997	99.7	≥ 500	9	**999.7 hPa**
147	14.7	< 500	10	1014.7 hPa

27.4 Temperature and dew point

Both are in degrees Fahrenheit on the sheet. To convert to Celsius, subtract 32 and multiply by 5/9:

$$T(^\circ\mathrm{C}) = \tfrac{5}{9}\,[T(^\circ\mathrm{F}) - 32]. \tag{27.1}$$

where

$T(^\circ\mathrm{C})$, $T(^\circ\mathrm{F})$ the same temperature in Celsius and in Fahrenheit °F from the station model

So 37 °F is 2.8 °C, 63 °F is 17.2 °C, and 21 °F is −6.1 °C.

The **dew point is the water-vapour measure of Chapter 22**, and the gap between the two, $T - T_d$, tells you how close the air is to saturation. A gap of 3 °F means nearly saturated air (fog, rain or thick cloud likely); a gap of 20 °F means dry air. You can turn the pair into a relative humidity with the formula $\mathrm{RH} = 100\, e_s(T_d)/e_s(T)$.

27.5 Sky cover

The circle is filled in proportion to the cloud cover. The sheet uses tenths. The shading grows clockwise from the top, as in Figure 27.2.




Figure 27.2. The sky-cover symbols of the sheet's Appendix: from clear (empty) through half (right half filled) to overcast (filled), and an obscured sky (a cross). The "9/10" symbol is filled with a vertical gap: overcast with small openings.

27.6 Wind: the barbs

The wind is drawn as a straight line, the **staff**, that starts at the circle and points **towards the direction the wind comes from**, as a weather vane does. At the free end are short lines, the **barbs**, that tell you the speed in knots.

- a **half barb**: 5 knots;
- a **full barb**: 10 knots;
- a **pennant** (a filled triangle): 50 knots;
- no barb but a staff: 1 to 2 knots; a circle round the station and no staff: calm.




Figure 27.3. Wind barbs for the Northern Hemisphere, drawn with the staff pointing up (a north wind) and the barbs on the right. Add the barbs: 10 + 10 + 5 = 25 knots.

To read a barb.

Step 1. Find the staff and follow it from the circle to its free end. The wind comes *from* that direction: a staff pointing to the lower left is a wind from the south-west.

Step 2. Count the barbs at the free end: pennants (50), full barbs (10), half barbs (5). Add them.

Step 3. Convert if you wish: $1\,\text{knot} = 0.514\ \text{m}\,\text{s}^{-1}$, so $25\,\text{knots} = 12.9\ \text{m}\,\text{s}^{-1}$.

A question that comes up

The staff points where the wind comes from, but the air is moving the other way. Which way do I draw a streamline?
Away from the barbs, i.e. in the direction *opposite* to the staff. A staff pointing to the south-west means the air is heading north-east. Keep that in mind in Chapter 28.

27.7 Present weather and pressure tendency




Figure 27.4. The present-weather symbols used on the sheet.




Figure 27.5. The barometric tendency symbols: how the pressure has changed in the last three hours. A line rising to the right means rising pressure, one falling to the right means falling pressure.

The number next to the symbol, for example +17 or −11, is the pressure change in tenths of a hectopascal: +17 is a rise of 1.7 hPa in three hours, −11 a fall of 1.1 hPa. Rapid falls often warn of a storm approaching; rising pressure usually brings improving weather.


27.8 Question 12: the four station models of the sheet

The question on the sheet

Decode the information from each of the following station models.

Method. For each model work down the list the sheet gives: pressure; temperature; dew point; sky cover; and, where shown, current weather, wind speed, wind direction, pressure change and tendency.

Model 1




Model 1

Pressure: 024 → 02.4 → **1002.4 hPa**
Temperature: **37** °F (2.8 °C) **Dew point:** **21** °F (−6.1 °C)
Sky cover: empty circle: **0 %** (clear). No staff, only a ring round the circle: **calm**. RH ≈ 52 %.

A correction to the printed key: dew point of model 1

The key gives 28 for the dew point of the first model. The model printed on the sheet shows **21** at the bottom left. The correct answer is 21 °F.


Model 2



Model 2

Pressure: 998 → 99.8 → **999.8 hPa**
Temperature: 63 °F (17.2 °C) **Dew point: 60** °F (15.6 °C)
Current weather: thunder with intermittent rain. Sky half filled (5/10); no staff: calm. RH ≈ 90 %.


Model 3



Model 3

Pressure: 117 → 11.7 → **1011.7 hPa**
Temperature: 43 °F (6.1 °C) **Dew point: 20** °F (−6.7 °C) **Sky cover:** empty circle, **0 %**
Wind: staff to the lower left: **from the south-west**; two full barbs and a half barb: **25 knots**
Pressure change: +17: +1.7 **mb**, **rising**

Model 4



Model 4

Pressure: 997 → 99.7 → **999.7 mb**
Temperature: 27 °F (−2.8 °C) **Dew point: 20** °F (−6.7 °C)
Sky cover: filled circle, **100 %** **Current weather: continuous snow**
Wind: staff to the upper left: **from the north-west**, **25 knots**
Pressure change: −11: −1.1 **mb**, **falling**

Table 27.3. All four models decoded, in the layout of the sheet.

	Model 1	Model 2	Model 3	Model 4
Barometric pressure (mb)	1002.4	999.8	1011.7	999.7
Temperature (°F)	37	63	43	27
Dew point (°F)	**21**	60	20	20
Sky coverage (%)	0	50	0	100
Current weather	—	thunder, interm. rain	—	continuous snow
Wind speed (knots)	calm	calm	25	25
Wind direction (from)	—	—	south-west	north-west
Pressure change in 3 h (mb)	—	—	+1.7	−1.1
Tendency	—	—	rising	falling

27.9 What the models tell you

Putting the numbers together gives each station a story.

- **Model 1:** cold, dry (the dew point is 16 °F below the temperature), clear and calm: a cold, high-pressure night.
- **Model 2:** warm and very humid (the dew point only 3 °F below the temperature), with thunder and rain: unstable, thundery air.
- **Model 3:** clear and windy with rising pressure: air moving away from a low towards a high, after a front has passed.
- **Model 4:** overcast, snowing, 27 °F, wind 25 knots and falling pressure: a depression is arriving or overhead.

27.10 Try it yourself: Practice Set 27

Decode each of the three models in Figure 27.6 (pressure, temperature and dew point in °F and °C, relative humidity, sky cover, weather, wind speed and direction, pressure change and tendency), then check against the answers below.




Figure 27.6. Three station models to decode.

Answers to Practice Set 27

27.1 **150** $\rightarrow$ 15.0 $\rightarrow$ **1015.0 hPa**. $T = 72\,^{\circ}\text{F}$ (22.2 °C), $T_d = 65\,^{\circ}\text{F}$ (18.3 °C), RH $\approx 79\,\%$. Sky: 2–3/10 (about 25 %). Wind: staff pointing straight down: **from the south**, one full barb and a half barb: **15 knots**. Tendency: +21 $\rightarrow$ up 2.1 hPa, rising then steady. No weather symbol: fair, warm and humid.

27.2 **975** (≥ 500) $\rightarrow$ 97.5 $\rightarrow$ **997.5 hPa**. $T = 48\,^{\circ}\text{F}$ (8.9 °C), $T_d = 47\,^{\circ}\text{F}$ (8.3 °C); RH $\approx 96\,\%$. Overcast (100 %). Continuous rain. Wind from the **north-east**, three full barbs and a half barb: **35 knots**. Pressure −32: down 3.2 hPa, falling steadily.

27.3 **084** (< 500) $\rightarrow$ 08.4 $\rightarrow$ **1008.4 hPa**. $T = 5\,^{\circ}\text{F}$ (−15 °C), $T_d = 3\,^{\circ}\text{F}$; RH $\approx 91\,\%$. Sky: 6/10. Continuous drizzle. Wind from the **west**, one full barb: **10 knots**. Pressure −4: down 0.4 hPa, rising then falling.

Chapter 28

Streamlines

This chapter answers Map Q13 of the sheet (Figure 8).

28.1 What a streamline is

A wind barb tells you the wind at one station. A **streamline** joins the stations: it is a line that is everywhere **parallel to the wind**. If you could photograph a river with a long exposure, the streaks of floating leaves would be streamlines. On a map they show at a glance where the air is coming from, where it is going, and where it swirls, converges or spreads out.

- At every point of a streamline, the line is tangent to the wind there.
- Streamlines **never cross**, because the wind has only one direction at one place.
- They begin at an **upwind** station and carry an **arrow head** downwind, usually where the flow changes direction.
- They show direction only. Their spacing does not show speed unless you choose to draw them that way.

28.2 The method

Step 1. **Pick a starting point** at an upwind station: one where the wind is blowing towards the area you want to follow.

Step 2. **Read the barb correctly.** The staff points to where the wind comes *from*. The air moves the other way, *away* from the staff. A barb with its staff pointing to the lower left is a south-west wind and the streamline there runs towards the upper right.

Step 3. **Draw a short line in that direction** and move downwind until you are near the next station.


Step 4. **Bend the line smoothly** so that it follows the direction of the barb at the next station, then the next.

Step 5. **Continue** until the stations run out or the flow leaves the map, then draw an **arrow head** at the downwind end.

Step 6. **Repeat from another starting point** until you have the five or six streamlines the question asks for. Keep them smooth and make sure no two cross.

Worked example: a small flow

Twelve stations report winds from the south-west on the left of the map and from the west on the right (the staffs point lower-left on the left and to the left on the right). Starting at a left-hand station the air moves towards the north-east. As you move east the barbs rotate clockwise, so the streamline bends to the right and ends up heading east. Each line starts at a station and stays parallel to the barbs it passes.



A question that comes up

How do I know whether to draw the arrow at the start or the end?
At the *downwind* end: the end towards which the air is moving, that is, away from the stations' barbs. A good test: put your pencil on the station, look at the barb, and ask "which way is this air going next?"

28.3 Figure 8: the stations

Figure 8 of the sheet shows the same 34 stations as Figure 10, but plotted with only the sky cover and the wind barb (no numbers). Here they are, on a real map, with each station numbered so that you can refer to it. Look at the barbs before you draw anything: where does the wind blow from in each part of the map?

Table 28.1. The wind by region (the staff points to where the wind comes from).

Region	Stations	Wind from
Dakotas, Nebraska, Kansas, Oklahoma, north Texas	1, 4, 12, 15, 21, 22, 25, 27, 30	north-west (WNW–NW), about 10 knots
Iowa, Missouri (west of the low)	5, 9, 16	north-west to west-north-west, 10–25 knots
Southern Missouri, Arkansas, Texas (east), Mississippi, Alabama	23, 28, 29, 31, 32, 33, 34	south-south-west to west-south-west, about 10 knots
Illinois, Kentucky, Tennessee, Indiana (south)	13, 17, 18, 19, 20, 24, 26	south-south-west to south, 10–25 knots
Wisconsin, Michigan, Ohio, Indiana (north), northern Illinois	2, 3, 6, 7, 8, 10, 11, 14	north-east to south-east, 2–20 knots




Figure 28.1. The stations of Figure 8 (circle for the sky cover, staff and barbs for the wind; station numbers in grey). The state outlines are real boundaries drawn in the same projection as the sheet.

28.4 Question 13: the streamlines

The question on the sheet

Streamlines begin at an upwind location and are drawn as long lines with an arrow where the wind shifts abruptly. Complete the analysis by drawing 5 to 6 streamlines to illustrate the general wind flow.

On the sheet two streamlines are already drawn as examples. One runs south-east from near the Nebraska–Kansas border across Kansas into Oklahoma, in the north-westerly wind. The other starts in north Texas and curves north and then north-east through Arkansas and Missouri into Illinois, in the southerly wind.

Answer to Map Q13

Five or six smooth lines, each starting upwind and ending with an arrow, such as those in Figure 28.2: north-westerly flow in the west curving south-east then east; southerly flow in the south curving north and then north-west; easterly flow in the north curving south-west. They all swirl **anticlockwise** and inwards towards one centre in Iowa and northern Illinois.




Figure 28.2. Figure 8 completed. The two grey streamlines are the examples printed on the sheet. The six green streamlines were drawn through the barbs of all 34 stations, starting upwind in the Dakotas, Kansas, Texas, Mississippi, Indiana and Michigan. They curve round a centre in Iowa and northern Illinois.

28.5 What the streamlines show

The pattern is the signature of a Northern Hemisphere low-pressure system (Chapter 26).

- The wind is **north-westerly** to the west of the centre, **southerly** to the south and east of it, and **easterly** to the north: an anticlockwise circulation.
- The streamlines curve **inwards**, because friction makes the air cross the isobars towards low pressure. They converge near the centre, so air must rise there: this is where the cloud and rain of Question 14 are.
- The streamlines from the south carry warm, humid air northwards; those from the north-west carry cold, dry air southwards. Where the two meet are the fronts.
- The centre of the swirl is where the pressure is lowest. The station reports confirm it: the lowest pressures on the map, 1002.4 and 1002.6 hPa, are at stations 8 and 9, next to the centre of the swirl.

28.6 Common mistakes

- **Drawing the arrow the wrong way round,** along the staff rather than away from it. Always ask where the air is going.

- **Joining stations with straight segments** instead of a smooth curve that follows every barb.
- **Letting streamlines cross.** If two lines are about to cross you have misread a barb.
- **Starting downwind.** A streamline needs a clear upwind start.
- **Forgetting the arrow heads.** The sheet asks for an arrow where the wind shifts abruptly; put one on every line.

28.7 Try it yourself: Practice Set 28

Practice Set 28: reading wind barbs

28.1 A station has its staff pointing to the right (east) with one full barb and a half barb. From which direction does the wind blow, how strong is it, and in which direction does a streamline through the station run?

28.2 The staff points to the top of the page (north) with two full barbs. State the wind and the direction of the streamline.

28.3 A station near the centre of a low shows a wind from the north-east, another to its east shows a wind from the south-east and another to its south a wind from the south-west. Is the circulation clockwise or anticlockwise?

28.4 In Figure 28.2 what is the wind at station 31 (south-west Texas)? What is the direction of the streamline there?

Answers to Practice Set 28

28.1 From the **east**, 15 knots; the streamline runs **towards the west**.

28.2 From the **north**, 20 knots; the streamline runs **towards the south**.

28.3 Going from the north-east side, to the east, to the south the winds are from NE, SE, SW. A wind from the NE blows to the SW, from the SE to the NW, from the SW to the NE: the air circles anticlockwise, as round a Northern Hemisphere low.

28.4 Station 31 shows a wind from the west-south-west (255°) of about 10 knots. The streamline there runs towards the east-north-east (it is the start of the north-easterly curve in the figure).

Chapter 29

Fronts and the weather map

This chapter answers Map Q14 of the sheet (Figures 9 and 10).

29.1 Air masses and fronts

An **air mass** is a very large body of air, hundreds of kilometres across, whose temperature and humidity are much the same throughout. It takes on the character of the surface it has rested on: air that has sat over a cold, dry continent is cold and dry; air from the tropical ocean is warm and humid. A **front** is the boundary between two unlike air masses. You can recognise one by any of the following:

- a sharp change of **temperature**;
- a sharp change of **moisture** (the dew point);
- a sharp change of **wind direction**.

Fronts are classified by how they move.

Warm front A warm air mass advances over a cooler one. Drawn as a red line with semicircles pointing the way the front moves.

Cold front A cold air mass advances under a warmer one. Drawn as a blue line with triangles pointing the way the front moves.




Figure 29.1. Cross-sections through a warm front (left) and a cold front (right), with the vertical scale greatly exaggerated. In both the front is a sloping surface along which the warm air rises over the cold air.

Wind veering. Notice that in both cases the wind *veers*, that is, turns clockwise, as the front passes. This is what the sheet means by "notice also how wind direction shifts at the frontal boundary".

29.2 Figure 9: a low with a warm front and a cold front

Figure 9 of the sheet shows a low-pressure centre (L) with a warm front curving east and south from it and a cold front trailing to the south-west. Four stations sample the air round them, which gives a small case in which to learn how to read a front before tackling the whole map.

Table 29.1. What happens as each kind of front passes a station in the Northern Hemisphere.

	Warm front	**Cold front**
Before	pressure falls; cloud thickens from cirrus to nimbostratus; wind backs to south-east	pressure falls; warm, humid, south-westerly wind
Passage	steady rain or drizzle; wind veers to south-west; temperature rises	heavy showers, perhaps thunder; wind veers sharply to north-west
After	warm, humid air, cloud breaks; pressure steady	colder, drier air; pressure rises; clearing sky




Figure 29.2. Figure 9: station A is north of the warm front, B and C are in the warm sector between the fronts, D is behind the cold front. The dashed orange arrows mark the two crossings studied below: C to D across the cold front, and A to B across the warm front.

Step 1. The warm sector (B and C). The air between the fronts is warm (68 and 70 °F) and humid (dew points 63 and 64 °F, relative humidity above 80 %), and the wind is from the south.

Step 2. Behind the cold front (D). The air is cooler (50 °F), much drier (dew point 38 °F, RH 63 %), the sky is clear and the wind has swung to the west-north-west.

Step 3. Crossing the cold front, C to D. The temperature falls 20 °F, the dew point falls 26 °F, the wind turns from south-south-west (193°) to west-north-west (292°): clockwise by about 100°. The pressure rises and the cloud clears.

Step 4. Crossing the warm front, A to B. The temperature rises 12 °F, the dew point 10 °F, and the wind turns from east-south-east (113°) to south (177°): clockwise by about 64°. The sky changes from overcast and raining to almost clear.

Table 29.2. The four stations of Figure 9, read from the figure.

Stn	Temp. (°F)	Dew pt (°F)	RH (%)	p (hPa)	Wind from	Sky
A	56	53	90	1001.9	ESE 10 kt	overcast, rain
B	68	63	84	1002.5	S 10 kt	2–3/10
C	70	64	81	1003.0	SSW 10 kt	5/10
D	50	38	63	1004.1	WNW 20 kt	clear

What Figure 9 teaches

A front is marked by *simultaneous* changes of temperature, moisture and wind. At **both** fronts the wind **veers** (turns clockwise). The cold front brings the bigger changes in temperature and dew point; the warm front brings the continuous rain.

29.3 Question 14: the full analysis of Figure 10

The question on the sheet

Using the map of Figure 10: draw isobars at 4 mb intervals (e.g. 1004, 1008, 1012 mb); label the low-pressure centre with an "L"; draw the warm and cold fronts; outline the area where cloud cover exceeds 75 %; shade the areas receiving precipitation.

Figure 10 shows the same 34 stations as Figure 8, but with all the numbers: temperature, dew point, pressure and weather, as well as the sky cover and the wind. The recommended order is the order of the bullets; each step helps the next.

29.3.1 Step 0: read the pressures

Decode every pressure with the rule of Chapter 27 (a three-digit number below 500 gets 10 in front, otherwise 9), and write the full value next to each station.

29.3.2 Step 1: isobars every 4 mb

The isobars are 1004, 1008, 1012, 1016 and 1020 mb. (The lowest pressure is 1002.4, so there is no 1000 mb isobar. The highest, 1019.4 in Alabama, falls just short of 1020; the 1020 isobar lies a little beyond it, where the trend continues, and may be left out.)

Method: interpolating.

1. Look at two neighbouring stations and ask whether a multiple of 4 lies between their pressures.
2. If it does, find where it falls on the straight line between them *in proportion*.
3. Mark that point. Do it for every pair of neighbours.
4. Join the points of equal value with a smooth curve. Do the same for the next value.




Figure 29.3. Step 0: the decoded sea-level pressures (hPa) at the 34 stations. The lowest values, 1002.4 and 1002.6, are in Iowa and Illinois (stations 8 and 9); the highest, 1019.4, is in Alabama (station 34).

Worked example: where does 1008 mb cross between stations 18 and 19?

Station 18 reads 1004.4 hPa and its neighbour station 19 reads 1009.9 hPa. The pressure changes by $1009.9 - 1004.4 = 5.5$ hPa between them, and 1008 is $1008 - 1004.4 = 3.6$ hPa above the first station:

$$\text{fraction of the distance} = \frac{3.6}{5.5} = 0.65.$$

The 1008 mb isobar crosses the line joining the stations 65 % of the way from station 18 towards station 19.



Rules for drawing isobars

- Isobars are smooth curves. Each connects points of exactly the same pressure.
- Isobars **never cross** and never branch.
- They never end inside the map: they either close into a loop (round a low or a high) or leave the map edge.

Table 29.3. More interpolations to practise on.

Stations	Pressures (hPa)	Isobar	Calculation	Fraction
9 and 5	1002.6, 1008.8	1004	$(1004 - 1002.6)/6.2$	23 %
9 and 5	1002.6, 1008.8	1008	$(1008 - 1002.6)/6.2$	87 %
16 and 5	1006.9, 1008.8	1008	$(1008 - 1006.9)/1.9$	58 %
20 and 24	1011.1, 1014.9	1012	$(1012 - 1011.1)/3.8$	24 %

- The pressure rises or falls in steps of 4 mb from one isobar to the next; if the next isobar is much further away the gradient is gentle there.
- Label each isobar with its value.



Figure 29.4. Step 1: isobars at 4 mb intervals through the station pressures. The 1004 mb isobar closes round the lowest pressures; the 1008 and 1012 mb isobars follow it outwards; 1016 and 1020 mb lie to the east and south-east. (The curves here were drawn as a smooth surface through all 34 pressures.)

29.3.3 Step 2: the L

The low-pressure centre is where the isobars close round the lowest pressure. The two lowest pressures on the map are 1002.4 hPa (station 8, northern Illinois) and 1002.6 hPa (station 9, Iowa). The 1004 mb isobar encloses them, together with stations 13 and 17 (1002.9 and 1004.0). Write the **L** between stations 8 and 9, near the centre of the closed isobar, at about 1002 hPa.

Answer: the L

The L goes in northern Illinois and eastern Iowa, between stations 8 and 9, in the middle of the innermost isobar. It is also the centre of the swirl of streamlines you drew in Question 13.

29.3.4 Step 3: the fronts

A front lies where the temperature, the dew point and the wind direction change rapidly. Compare pairs of neighbouring stations: if two neighbours differ strongly in all three, a front runs between them.




Figure 29.5. Step 3: air temperature (°F) at each station, coloured from blue (cold, 32 °F) to red (warm, 72 °F). The warm air fills the south-east and a tongue reaches towards the low: the warm sector. Cold air covers the north-west. The cold front (blue, triangles) separates them in the west; the warm front (red, semicircles) in the east.

The cold front. West of the low, the stations behind the front are cool (42 to 52 °F), drier (dew points 36 to 50 °F) and have winds from the west to north-west; those ahead are warm (60 to 66 °F), humid (dew points 55 to 61 °F) and have winds from the south-south-west to south-west. The pairs 16–17, 22–23, 25–28 and 31–32 straddle it, with temperature jumps of 11 to 19 °F. The front runs from the low towards the south-south-west, down through Missouri, Arkansas and east Texas.

The warm front. East of the low, the stations north of the front are cool (46 to 54 °F) with winds from the north-east or east; those south of it are warm (57 to 66 °F) with winds from the south. The pairs 8–13, 14–19 and 10–20 straddle it, with temperature jumps of 9 to 20 °F and the

Table 29.4. Evidence for the fronts: pairs of stations on either side of each front (cold side → warm side).

Front	Stations	Temperature (°F)	Dew point (°F)	Wind from
cold	16 → 17	45 → 60	40 → 55	NW → SSW
cold	22 → 23	42 → 61	36 → 57	NW → SSW
cold	25 → 28	48 → 66	38 → 59	WNW → SW
cold	31 → 32	52 → 63	50 → 61	WSW → S
warm	8 → 13	48 → 57	45 → 54	NE → SSW
warm	14 → 19	54 → 66	45 → 53	E → S
warm	10 → 20	46 → 66	43 → 55	E → S

wind turning from east or north-east to south. The front runs east from the low through Indiana and Ohio.

Answer: the fronts

A **cold front** (blue, with triangles pointing east, the way it moves) from the L towards the south-south-west to Texas; a **warm front** (red, with semicircles pointing north-east) from the L eastwards through Indiana and Ohio. The warm sector, the wedge of warm air between them, lies to the south-east.

29.3.5 Step 4: cloud cover above 75 %

Cloud cover above 75 % means the stations drawn with a three-quarters, nine-tenths or completely filled circle. Looking at the sky circles: stations 2, 5, 6, 8, 9, 10, 11, 14, 17, 23, 31 and 32 are completely overcast; stations 3, 13, 19 and 28 have 7 to 8 tenths. Draw a line round the group, including the area between them. The result is a broad area wrapped round the low and extending along the cold front to the south-west.

29.3.6 Step 5: the precipitation

Shade round every station that reports precipitation: the weather symbol is a dot or two dots (rain), or the thunderstorm symbol with a dot. These are stations 2, 5, 6, 8, 9, 10, 11, 17, 18 and 23: a band to the north and east of the low, along the warm front, and a narrow band along the cold front in Missouri.




Figure 29.6. Step 4: the area where the cloud cover exceeds 75 % (dashed outline and grey tint). It wraps round the low on its north and east sides and tails away to the south-west along the cold front.




Figure 29.7. Step 5: the areas receiving precipitation (blue shading) at the ten stations that report rain or thunderstorms.




Figure 29.8. Question 14 complete: isobars every 4 mb (orange), the low (L), the cold front (blue triangles) and the warm front (red semicircles), the area with more than 75 % cloud (dashed outline) and the precipitation (blue shading), over the 34 stations of Figure 10. The state outlines are real boundaries. The fronts are an interpretation, drawn from the temperature, dew-point and wind contrasts.

29.4 Reading the finished picture

The map is a text-book mid-latitude depression.

- The low is over Iowa and northern Illinois, at about 1002 hPa.
- The **warm sector** is the tongue of warm, humid air in the south and east, between the two fronts: temperatures in the 60s and 70s °F and dew points of 55 to 64 °F.
- The cold air in the north-west is dry (dew points of 22 to 38 °F) and mostly clear.
- The cloud and rain are on the fronts and near the low, where the air is being lifted.
- The winds circle the low anticlockwise (Chapter 28) and cross the isobars towards low pressure (Chapter 26).

Worked example: is the wind consistent with the isobars?

Between stations 9 (1002.6 hPa) and 5 (1008.8 hPa), about 216 km apart, the pressure changes by 6.2 hPa. The geostrophic wind for that gradient at 41.5° N ($f = 9.7 \times 10^{-5}\,\mathrm{s^{-1}}$) is

$$V_g = \frac{1}{\rho f}\frac{\Delta p}{\Delta n} = \frac{620}{1.2 \times 9.7 \times 10^{-5} \times 2.16 \times 10^{5}} = 25\ \mathrm{m\,s^{-1}} \approx 48\ \text{knots}.$$

The observed surface winds are 10 to 25 knots. That is about what friction does: it slows the surface wind to a half or two thirds of the geostrophic value and turns it across the isobars (Chapter 26).

29.5 Questions that come up

A question that comes up

Where exactly should I put the front?
There is no single right answer: a front analysis is an interpretation. Put the line along the zone where the temperature, dew point and wind change fastest, bend it to follow the stations, and begin it at the low. Check that the stations on the cold side are colder and drier than those on the warm side.

A question that comes up

Why do the isobars bend sharply where a front crosses them?
Because the wind changes direction at a front, and the wind follows the isobars, the isobars must change direction there too. A front lies along a trough of low pressure, with the isobars forming a V pointing away from the low.

A question that comes up

Which stations count as "cloud above 75 %"?
Those whose circle is filled to at least three quarters: 7–8 tenths, 9 tenths (filled with a gap) or overcast. Stations with half or less are outside the outline.

29.6 Try it yourself: Practice Set 29

Practice Set 29: fronts and isobars

29.1 Stations X (1011.1 hPa) and Y (1014.9 hPa) are neighbours. Where does the 1012 mb isobar cross the line between them?

29.2 A station reports a temperature of 61 °F, a dew point of 57 °F and a south-south-west wind of 10 knots. A neighbour to its west reports 42 °F, 36 °F and a north-west wind. What kind of front lies between them?

29.3 Station P is 30 °F with dew point 28 °F, east wind, overcast and snow. Station Q to the south is 41 °F with dew point 38 °F and south-east wind. Is a warm or cold front likely to lie between them? Why?

29.4 Why is there no 1000 mb isobar on Figure 10?

Answers to Practice Set 29

29.1 $(1012 - 1011.1)/(1014.9 - 1011.1) = 0.9/3.8 = 0.24$: 24 % of the way from X to Y.

29.2 A **cold front**: the temperature falls 19 °F, the dew point 21 °F, and the wind turns clockwise from south-south-west to north-west as you cross it from east to west.

29.3 A **warm front**: the warmer, moister air (41 °F, 38 °F) lies to the south with a south-east wind, riding over the colder air with east wind to the north, where it is snowing. The wind turns clockwise from east to south-east.

29.4 No station reports a pressure as low as 1000 hPa (the lowest is 1002.4), and an isobar should only be drawn where the data support it.

Chapter 30

Putting Practical 3 together

30.1 One chain of ideas

The three sections of Practical 3 are one argument, and they close the argument begun in Parts I and II:

1. Air holds water vapour up to a limit set by its temperature. Relative humidity compares the vapour with that limit; the mixing ratio and the dew point measure the vapour itself, and cooling air to its dew point makes dew, fog or cloud (Moisture Q1–Q6, Chapters 21 and 22).
2. Warm air columns are taller than cold ones (the scale-height law of Part I, Chapter 3), so pressure surfaces slope down from the warm tropics towards the cold poles, more steeply with height (Motion Q1–Q6, Chapter 24). The uneven warming that causes this is the sun-angle effect of Part II (Chapter 18).
3. The slope produces a pressure gradient force; on a rotating Earth the Coriolis force turns the moving air until the two balance and the wind blows along the isobars, the geostrophic wind (Motion Q7, Chapter 25).
4. Near the ground friction slows the wind and turns it across the isobars towards low pressure, so air spirals into lows, rises, and forms cloud and rain (Motion Q8–Q11, Chapter 26).
5. A weather map shows all of this at once: station models record the observations, streamlines show the flow, and isobars, fronts, cloud and rain show the weather systems it organises (Map Q12–Q14, Chapters 27 to 29).

30.2 All the answers in one table

30.3 Mistakes to avoid

1. **Ranking by relative humidity** instead of the actual mixing ratio (Moisture Q3).
2. **Reading the dew point from the wrong row** of Table 1, or at the wrong temperature: always check that the air could hold *at least* the sample's vapour at the temperature you give (Moisture Q6: 26, not 25).
3. **Putting H and L the wrong way round** on the pressure surfaces: the lower pressure surface (colder column) is the L (Motion Q4).
4. **Drawing the Coriolis force to the wrong side:** right in the north, left in the south, always at right angles to the wind.
5. **Drawing the pressure gradient force away from low pressure.** It points *towards* low pressure.
6. **Forgetting that the staff of a wind barb points to where the wind comes from** (Map Q12 and Q13).
7. **Mis-reading the pressure code:** 998 is 999.8, not 1099.8 or 998.0 (Map Q12).
8. **Isobars that cross,** or that stop in the middle of the map (Map Q14).

Table 30.1. Every answer of Practical 3, with the chapter where it is worked out.

Question	Answer	Ch.
Moisture Q1	RH = 49.3, 88.8, 26.0, 10.4, 19.9 %	21
Moisture Q2	Humidity falls sharply: the vapour content is unchanged but the capacity of the warmer air is much larger	21
Moisture Q3	Mixing ratios 9.126, 8.974, 7.684, 11.078, 10.54 g kg^{-1}; ranks 3, 4, 5, 1, 2	21
Moisture Q4	RH does not say how much vapour is actually in the air	21
Moisture Q5	Dew point $\approx 14\,^\circ$C (13.8)	22
Moisture Q6	$w_s = 35.134$ g kg^{-1}; $w = 21.78$ g kg^{-1}; dew point $\approx 26\,^\circ$C (the printed key says 25)	22
Motion Q1	Lines 5400→5700 m (500 mb) and 8700→9500 m (300 mb)	24
Motion Q2	Saskatoon	24
Motion Q3	Saskatoon is colder at each height; the cold column is thinner	24
Motion Q4	z has the higher pressure: H at z (Monterrey side), L at x; arrow from z to x	24
Motion Q5	At 9000 m (800 m against 140 m of slope)	24
Motion Q6	Wind speeds are greater at 9000 m	24
Motion Q7	North: H at 5520 m, L at 5460 m; south: H at 5640 m, L at 5580 m; force arrow to L, Coriolis arrow to H, wind between them	25
Motion Q8	The Coriolis force is weakened, because it depends on wind speed	26
Motion Q9	Red up (to low pressure), black up-right, blue down-right (right of wind), green down-left (against wind)	26
Motion Q10	NH low, NH high, SH low, SH high (in the order of Figure 6)	26
Motion Q11	On each box: red along the line to or from the centre; blue at right angles to the wind (right in the north, left in the south); green against the wind	26
Map Q12	Model 1: 1002.4, 37, **21**, 0 %. Model 2: 999.8, 63, 60, thunder with intermittent rain. Model 3: 1011.7, 43, 20, 0 %, 25 kt from SW, +1.7, rising. Model 4: 999.7, 27, 20, 100 %, continuous snow, 25 kt from NW, −1.1, falling	27
Map Q13	Five or six smooth streamlines swirling anticlockwise round a centre in Iowa and Illinois	28
Map Q14	Isobars at 4 mb; L in northern Illinois/Iowa; cold front to the south-south-west, warm front to the east; cloud above 75 % wrapped round the low; rain north and east of the low and along the cold front	29

30.4 Self-test

Try these before the practical. The answers follow.

Self-test: one line each

S1. Air at 20 °C has a mixing ratio of 7.5 g kg^{-1}. What is its relative humidity?

S2. That air is cooled to 10 °C. What is its relative humidity now, and what is its dew point?

S3. Which holds more water vapour: air at 35 °C with RH 30 %, or air at 25 °C with RH 60 %?

S4. Find the dew point of air at 30 °C and RH 50 %, and the height at which cloud would form if it were lifted.

S5. True or false: the dew point can be higher than the air temperature. Explain.

S6. Why do pressure surfaces slope down towards cold air?

S7. The 300 mb surface is at 8700 m over place S and 9500 m over place M. In which direction does the pressure gradient force act at 9000 m?

S8. Two 500 hPa height lines 90 m apart are 600 km apart at 35° N. Find the geostrophic

wind.

S9. In the Southern Hemisphere a wind blows towards the west. Where is the low pressure, to its north or south?

S10. What does friction do to (a) the speed and (b) the direction of the surface wind?

S11. Over land the surface wind is $8\,\mathrm{m\,s^{-1}}$ and crosses the isobars at $40°$. Estimate the geostrophic wind.

S12. Air spirals inward and clockwise. Which hemisphere, which pressure system?

S13. Decode the pressure codes 087 and 962.

S14. Convert $50\,°\mathrm{F}$ and $14\,°\mathrm{F}$ to °C.

S15. A wind barb has two full barbs and one pennant. What is the speed?

S16. Behind a front the temperature falls $18\,°\mathrm{F}$ and the wind turns clockwise from south-west to north-west. Which kind of front?

S17. A staff points to the north-east. In which direction does the air move?

S18. The 1012 mb isobar lies between stations at 1009.6 and 1013.2 hPa. Where?

S19. Why does a strong wind go with closely spaced isobars?

S20. What can relative humidity not tell you?

Answers to the self-test

S1. $7.5/14.956 = 50\,\%$.

S2. $w_s(10) = 7.762$: RH $= 7.5/7.762 = 97\,\%$. Dew point: Table 1 gives 6.771 at $8\,°\mathrm{C}$ and 7.762 at $10\,°\mathrm{C}$, so $T_d = 8 + 2(7.5 - 6.771)/(7.762 - 6.771) = 9.5\,°\mathrm{C}$.

S3. $w_s(35) \approx 37.3$ gives $0.30 \times 37.3 = 11.2\,\mathrm{g\,kg^{-1}}$; $w_s(25) \approx 20.5$ gives $0.60 \times 20.5 = 12.3\,\mathrm{g\,kg^{-1}}$. The $25\,°\mathrm{C}$ air holds more.

S4. $w = 0.5 \times 27.694 = 13.85\,\mathrm{g\,kg^{-1}}$: between $18\,°\mathrm{C}$ (13.162) and $20\,°\mathrm{C}$ (14.956), $T_d = 18 + 2(13.85 - 13.162)/1.794 = 18.8\,°\mathrm{C}$. Cloud base $\approx 125 \times (30 - 18.8) = 1400\,\mathrm{m}$.

S5. False (in open air): it would mean the air is oversaturated and the surplus vapour would condense at once. At most $T_d = T$ at saturation.

S6. Cold air is denser, so a column of it is thinner: the same pressure drop uses up less height, so the pressure surfaces are lower over cold air.

S7. From the high (the 300 mb surface is higher over M, so z over M has the higher pressure) to the low: from M towards S.

S8. $f = 2 \times 7.292 \times 10^{-5} \sin 35° = 8.37 \times 10^{-5}$; $V_g = \dfrac{9.81 \times 90}{8.37 \times 10^{-5} \times 6 \times 10^5} = 17.6\,\mathrm{m\,s^{-1}}$ (34 kt).

S9. To the **north**: in the Southern Hemisphere the low is on the right of the wind (Buys-Ballot), and the right of a westward wind is north.

S10. (a) It reduces the speed. (b) It turns the wind across the isobars towards low pressure.

S11. $V = V_g \cos\alpha$, so $V_g = 8/\cos 40° = 10.4\,\mathrm{m\,s^{-1}}$.

S12. Inward and clockwise: Southern Hemisphere low.

S13. 087 → 1008.7 hPa; 962 → 996.2 hPa.

S14. $50\,°\mathrm{F} = 10\,°\mathrm{C}$; $14\,°\mathrm{F} = -10\,°\mathrm{C}$.

S15. $2 \times 10 + 50 = 70$ knots.

S16. A cold front.

S17. The air moves towards the south-west.

S18. $(1012 - 1009.6)/(1013.2 - 1009.6) = 0.67$: two thirds of the way from the lower to the higher station.

S19. Close isobars mean a large pressure gradient force, which needs a large wind to balance it with the Coriolis force (and friction).

S20. How much vapour is actually in the air (or how the air compares with air at another temperature): it only compares the vapour with the capacity at the present temperature.

30.5 Before the practical: a checklist

- ☐ I can look up, and interpolate, in Table 1 (and I know about the two slips).
- ☐ I can find an RH, a mixing ratio and a dew point from any two of T, w, RH.
- ☐ I can explain why pressure surfaces slope and why wind increases with height.
- ☐ I can draw the three forces correctly in both hemispheres.
- ☐ I can name the four systems of Figure 6 from their rotation.
- ☐ I can decode a station model: pressure, temperature, dew point, sky, wind, weather, tendency.
- ☐ I can draw streamlines (arrows downwind) and isobars (interpolating, never crossing).
- ☐ I can say how to recognise a front, and which way the wind turns across it.

Appendix A

Practical 1: formulas, derivations, data and corrections

A.1 Formula sheet

Every formula used in Part I, with the equation number where it is introduced and explained.

Quantity	Formula	Where, and what for
Ideal gas law	$p = \rho R_d T$	Eq. (3.1); density from pressure and temperature (Q13)
Hydrostatic balance	$\Delta p/\Delta z = -\rho g$	Eq. (3.2); why pressure falls with height
Exponential pressure	$p = p_0 e^{-z/H}$, $H = R_d T/g$	Eq. (3.3); scale height (Q1, Q12)
Fraction of the atmosphere above	$f = 2^{-z/5.6}$ (z in km)	Eq. (3.4); Q1, Q3
After n halvings	$f_n = 1/2^n$	Eq. (3.5); Table 1
Pressure from the fraction	$p = 1000\, f$ hPa	Eq. (3.6); Q2, Q3
Partial pressure of oxygen	$p_{O_2} = 0.21\, p$	Eq. (4.1); Q4
Lapse rate	$\Gamma = -\Delta T/\Delta z$	Eq. (5.1); Q5, Q7–Q10
Standard atmosphere	$T = 15 - 6.5\, z$ to 11 km, then $-56.5\,^\circ$C	Eq. (5.2); Q5
Scale height from data	$H = (z_2 - z_1)/\ln(p_1/p_2)$	Eq. (8.1); Q12
Dry adiabatic lapse rate	$\Gamma_d = g/c_p = 9.8\,^\circ\mathrm{C\,km^{-1}}$	Eq. (8.2); Q13
Potential temperature	$\theta = T(1000/p)^{0.286}$	Eq. (8.3); Q13
Kelvin and Celsius	$T_\mathrm{K} = T_{^\circ\mathrm{C}} + 273.15$	§ 1.3

A.2 Derivations for the curious

A.2.1 Pressure falls exponentially with height

Start from hydrostatic balance, Eq. (3.2), written for a very thin layer, $dp/dz = -\rho g$, and replace the density using the gas law, Eq. (3.1), $\rho = p/(R_d T)$:

$$\frac{dp}{dz} = -\frac{g}{R_d T}\, p \qquad \Longrightarrow \qquad \frac{dp}{p} = -\frac{dz}{H}, \qquad H = \frac{R_d T}{g}.$$

The rate of fall of p is proportional to p itself. If T (and so H) is taken as constant, integrating from sea level ($z = 0$, $p = p_0$) to height z gives $\ln(p/p_0) = -z/H$, that is

$$p = p_0\, e^{-z/H},$$

which is Eq. (3.3). The pressure halves when $e^{-z/H} = \frac{1}{2}$, at $z_{1/2} = H \ln 2 = 0.693\, H$; the sheet's 5.6 km corresponds to $H = 8.08$ km. Writing the same law for two levels and dividing, $p_1/p_2 = e^{(z_2 - z_1)/H}$, so $\ln(p_1/p_2) = (z_2 - z_1)/H$: rearranged, this is Eq. (8.1), which gives the scale height of a layer from two lines of a sounding.

Without calculus. Divide the atmosphere into thin slabs, each Δz thick. By Eqs. (3.2) and (3.1), crossing one slab lowers the pressure by $\Delta p = \rho g\, \Delta z = p\, \Delta z/H$: the same *fraction* $\Delta z/H$ of whatever pressure there is. Removing the same fraction again and again is exactly what an exponential does.

A.2.2 The dry adiabatic lapse rate and potential temperature

The first law of thermodynamics for one kilogram of air says that heat added goes into warming the air and into the work it does as it expands: $dq = c_p\, dT - dp/\rho$. In an adiabatic process no heat is added, $dq = 0$, so

$$c_p\, dT = \frac{dp}{\rho}.$$

For air moving vertically through an atmosphere in hydrostatic balance, $dp = -\rho g\, dz$, Eq. (3.2). Substituting,

$$c_p\, dT = -g\, dz \qquad \Longrightarrow \qquad \frac{dT}{dz} = -\frac{g}{c_p} = -\frac{9.81}{1004} = -0.0098 \ \mathrm{K\,m^{-1}},$$

which is Eq. (8.2): dry air cools by 9.8 °C per km as it rises and warms by the same amount as it sinks. Alternatively, dividing $c_p\, dT = dp/\rho$ by T and using $1/\rho = R_d T/p$ gives $c_p\, dT/T = R_d\, dp/p$; integrating from (T, p) to $(\theta, 1000\ \mathrm{hPa})$,

$$c_p \ln\frac{\theta}{T} = R_d \ln\frac{1000}{p} \qquad \Longrightarrow \qquad \theta = T\left(\frac{1000}{p}\right)^{R_d/c_p},$$

which is Eq. (8.3), with $R_d/c_p = 287/1004 = 0.286$ (Wallace and Hobbs, 2006, ch. 3).

A.3 The data of the sheet

Table 1 completed is Table 3.1 and Table 2 is Table 5.1; the soundings and the Florida profile follow.

Table A.3. The local lapse rate, Eq. (5.1), of every layer of the two soundings up to 16 km. Negative lapse rates are inversions.

Layer (m)	T **bottom** (°C)	T **top** (°C)	Γ (°C km^{-1})	
Hillsborough (tropopause 9383 m)				
38–1330	11.05	−1.25	9.52	
1330–1358	−1.25	−1.25	0.00	isothermal
1358–1503	−1.25	−1.05	−1.38	inversion
1503–2504	−1.05	−7.45	6.39	

Layer (m)	T bottom (°C)	T top (°C)	Γ (°C km^{-1})	
2504–4405	−7.45	−15.25	4.10	
4405–4725	−15.25	−18.05	8.75	
4725–5414	−18.05	−22.45	6.39	
5414–6025	−22.45	−25.45	4.91	
6025–7018	−25.45	−32.85	7.45	
7018–9383	−32.85	−51.05	7.70	
9383–10 164	−51.05	−49.05	−2.56	inversion
10 164–11 116	−49.05	−44.65	−4.62	inversion
11 116–13 524	−44.65	−49.65	2.08	
13 524–14 969	−49.65	−53.85	2.91	
14 969–16 126	−53.85	−56.45	2.25	
St John's (tropopause 11 482 m)				
140–687	−1.65	−2.05	0.73	
687–3006	−2.05	−2.85	0.34	
3006–4062	−2.85	−6.65	3.60	
4062–5604	−6.65	−15.85	5.97	
5604–6263	−15.85	−19.85	6.07	
6263–7248	−19.85	−27.05	7.31	
7248–9103	−27.05	−41.05	7.55	
9103–10 446	−41.05	−53.45	9.23	
10 446–11 482	−53.45	−62.05	8.30	
11 482–11 843	−62.05	−58.05	−11.08	inversion
11 843–12 141	−58.05	−54.65	−11.41	inversion
12 141–16 254	−54.65	−56.25	0.39	

A.4 Corrections to the printed key, at a glance

Each item was found by recomputing the answer sheet (EGM305 module team, 2025b) against the data. The last column links to the full explanation.

Table A.4. Corrections and clarifications to the Practical 1 sheet and answer key.

Q	The key or sheet says	Correct or recommended	See
6	Table 3 printed as one table	the two soundings told apart by their first lines; printed separately in Table A.1	§ 2.2
4	percentages of the sea-level *atmosphere* (21, 7, 4.7, 15.5, 1.6 %)	also give oxygen compared with sea-level oxygen (100, 33, 23, 74, 7 %); jet p_{O_2} = 47.6 hPa (4.8 %)	§ 4.5
7	no answer	Hillsborough inversion 1360–1500 m; St John's nearly isothermal to 3 km	§ 6.4
9	no answer	62.10/9.345 = 6.65 °C km^{-1}	§ 7.2
10	Florida tropopause −75.4 °C	−75.6 °C (Table 4)	§ 7.3

Q	The key or sheet says	Correct or recommended	See
13	cold air “floats” because temperature falls at a constant rate	lower density at low pressure, and warming of sinking air by compression: potential temperature increases with height	§ 8.4

A.5 Symbols

Table A.5. The symbols used in Part I. Each equation’s own *where* list repeats the symbols it uses.

Symbol	Meaning	Unit	First used
c_p	specific heat of dry air at constant pressure	$\mathrm{J\,kg^{-1}\,K^{-1}}$	Eq. (8.2)
f	fraction of the atmosphere above a height	–	Eq. (3.4)
g	acceleration due to gravity	$\mathrm{m\,s^{-2}}$	Eq. (3.2)
Γ	lapse rate	$\mathrm{°C\,km^{-1}}$	Eq. (5.1)
Γ_d	dry adiabatic lapse rate	$\mathrm{°C\,km^{-1}}$	Eq. (8.2)
H	scale height	km	Eq. (3.3)
p	pressure	hPa or Pa	Eq. (3.1)
p_0	pressure at sea level	hPa	Eq. (3.3)
$p_{\mathrm{O_2}}$	partial pressure of oxygen	hPa	Eq. (4.1)
R_d	gas constant for dry air	$\mathrm{J\,kg^{-1}\,K^{-1}}$	Eq. (3.1)
ρ	density of the air	$\mathrm{kg\,m^{-3}}$	Eq. (3.1)
T	temperature	K or °C	Eq. (3.1)
θ	potential temperature	K	Eq. (8.3)
$x_{\mathrm{O_2}}$	fraction of the molecules that are oxygen	–	Eq. (4.1)
z	height above sea level	m or km	Eq. (3.2)

A.6 About the figures and the data

Every figure in Part I was drawn for it in the document itself with TikZ and pgfplots, from the data of the sheet and the equations of the text.

- The **halving-rule curves** (Figures 3.2 and 4.1) are Eq. (3.4); the dashed comparison curve and the layered profile of Figure 5.1 are the U.S. Standard Atmosphere (NOAA et al., 1976).
- The **soundings** (Figures 6.1, 7.1 and 8.1) plot the values of Tables 3 and 4 as printed on the sheet, joined by straight lines.
- The **potential temperatures** (Figure 8.2) are computed from the soundings with Eq. (8.3), and the **oxygen partial pressures** (Figure 4.2) with Eq. (4.1).
- The **hydrostatic slab** (Figure 3.1) is a diagram of the forces in Eq. (3.2).
- The **tropopause band** of Figure 7.2 and the **ocean-current sketch** of Figure 9.1 are schematic: they show the pattern, not measurements.

The soundings are single ascents on a single day; they illustrate the structure of the atmosphere and should not be read as climatological averages (§ 9.4).

Table A.1. Table 3 of the sheet: the radiosonde soundings of 21 March 1990, printed separately. Heights in metres, temperatures in °C.

3a. Hillsborough			**3b. St John's**		
p (hPa)	z (m)	T (°C)	p (hPa)	z (m)	T (°C)
1001	38	11.05	1005	140	−1.65
853	1330	−1.25	938	687	−2.05
850	1358	−1.25	700	3006	−2.85
835	1503	−1.05	612	4062	−6.65
735	2504	−7.45	500	5604	−15.85
573	4405	−15.25	458	6263	−19.85
549	4725	−18.05	400	7248	−27.05
500	5414	−22.45	307	9103	−41.05
460	6025	−25.45	250	10 446	−53.45
400	7018	−32.85	212	11 482	−62.05
282	9383	−51.05	200	11 843	−58.05
250	10 164	−49.05	191	12 141	−54.65
216	11 116	−44.65	100	16 254	−56.25
150	13 524	−49.65	88.9	17 004	−55.85
120	14 969	−53.85	73.6	18 190	−56.65
100	16 126	−56.45	50.5	20 566	−54.85
70	18 379	−58.65	30	23 834	−56.25
50	20 482	−58.45	20	26 404	−54.05
30	23 691	−60.05	18.1	27 047	−53.65
20	26 218	−56.25	10	30 828	−54.05

Table A.2. Table 4 of the sheet: the Florida temperature profile.

Height (m)	0	2000	4000	6000	8000	10 000	12 000	14 000	16 000	18 000
T (°C)	27.2	15.2	6.1	−6.7	−17.9	−32.6	−50.5	−64.5	−75.6	−73.3

Appendix B

Practical 2: formulas, derivations, data and corrections

B.1 Formula sheet

Every formula used in this part, with the equation number where it is introduced and explained. Temperatures are in kelvin; angles in degrees.

Quantity	Formula	Where, and what for
Speed of a wave	$c = f\lambda$	Eq. (12.1); frequency from wavelength
Energy of a photon	$\varepsilon = hc/\lambda$	Eq. (12.2); why short waves carry more energy
Planck curve	$B_\lambda = \dfrac{2hc^2}{\lambda^5}\dfrac{1}{e^{hc/\lambda k_B T} - 1}$	Eq. (12.3); the shape of every spectrum
Wien's law	$\lambda_{\max} = C/T$	Eq. (13.1); peak from temperature (Q1, Q3)
Wien's law, rearranged	$T = C/\lambda_{\max}$	Eq. (13.2); temperature from peak (Q1, Q3b)
Stefan–Boltzmann law	$E = \sigma T^4$	Eq. (14.1); total emission (Q2)
Ratio of emissions	$E_1/E_2 = (T_1/T_2)^4$	Eq. (14.2); Q2e
Stefan–Boltzmann, rearranged	$T = (E/\sigma)^{1/4}$	Eq. (17.3); temperature from emission (Q4d)
Inverse-square law	$S = S_E/d^2$	Eq. (16.1); sunlight at each planet (Q3c)
Planetary temperature	$T = \left[\dfrac{S(1-a)}{4\sigma}\right]^{1/4}$	Eq. (16.2); black-body temperature (Q3c)
Absorbed fraction	$1 - a$	Eq. (17.1); Q4a–c
Global average sunlight	$\bar{S} = S/4$	Eq. (17.2); Q4b′
Sun angle	$\beta = 90° - \theta_z$	Eq. (18.1); Q5a
Noon zenith angle	$\theta_z = \lvert\varphi - \delta\rvert$, $\delta = \pm 23.5°$ at the solstices	Eq. (18.4); Q5a
Beam spreading	$A = 1/\sin\beta$	Eq. (18.2); Q5a
Solar intensity	$I = 100\,\%/A = 100\,\% \times \sin\beta$	Eq. (18.3); Q5c
Kelvin to Celsius	$T_{°\mathrm{C}} = T_{\mathrm{K}} - 273.16$ (sheet)	§ 11.2

B.2 Derivations for the curious

These derivations show where the formulas of the practical come from. None is needed to answer the sheet; each is referred to from the chapter where its result is used.

B.2.1 Wien's law from the Planck curve

The peak of the Planck curve, Eq. (12.3), is where its slope is zero. Write $x = hc/(\lambda k_B T)$, so that a short wavelength is a large x. Up to constants that do not depend on λ, Eq. (12.3) becomes

$$B_\lambda \propto \frac{x^5}{e^x - 1}.$$

Setting the derivative with respect to x to zero:

$$\frac{d}{dx}\left(\frac{x^5}{e^x-1}\right) = \frac{5x^4(e^x-1) - x^5 e^x}{(e^x-1)^2} = 0 \quad\Longrightarrow\quad 5(e^x-1) = xe^x \quad\Longrightarrow\quad x = 5\,(1-e^{-x}).$$

This equation has no algebraic solution, but it is easy to solve by repeated substitution: start with $x = 5$, then $x = 5(1-e^{-5}) = 4.966$, then $5(1-e^{-4.966}) = 4.965$, and it settles at $x = 4.965$. Turning back to the wavelength,

$$\lambda_{\max} = \frac{hc}{4.965\,k_B\,T} = \frac{C}{T}, \quad \text{with}$$
$$C = \frac{(6.626\times10^{-34})(2.998\times10^{8})}{4.965\times1.381\times10^{-23}} = 2.8978\times10^{-3}\ \mathrm{m\,K} = 2897.8\ \mu\mathrm{m\,K},$$

which is Eq. (13.1) with the sheet's $C = 2898\,\mu\mathrm{m\,K}$. (Since x falls steadily as λ rises, the peak found in terms of x is the same point as the peak in terms of λ; and λ^{-5} in Eq. (12.3) is a constant times x^5, which is where the x^5 comes from.)

B.2.2 The Stefan–Boltzmann law from the Planck curve

The total power leaving one square metre of a black body is π times the spectral radiance, Eq. (12.3), added up over all wavelengths (the factor π collects the radiation leaving in every direction above the surface):

$$E = \pi \int_0^\infty B_\lambda(T)\,d\lambda.$$

With the same substitution $x = hc/(\lambda k_B T)$ the integral becomes

$$E = \frac{2\pi k_B^4 T^4}{h^3c^2}\int_0^\infty \frac{x^3}{e^x-1}\,dx = \frac{2\pi k_B^4 T^4}{h^3c^2}\times\frac{\pi^4}{15}.$$

All the temperature dependence has come out as T^4, and the rest is the Stefan–Boltzmann constant:

$$\sigma = \frac{2\pi^5 k_B^4}{15\,h^3c^2} = 5.670\times10^{-8}\ \mathrm{W\,m^{-2}\,K^{-4}},$$

which is Eq. (14.1). The value of the integral, $\pi^4/15 = 6.494$, is a standard result.

B.2.3 The black-body temperature of a planet

This is worked through in full in § 16.4: absorbed sunlight on the disc, $S(1-a)\pi R^2$, equals emission from the sphere, $\sigma T^4 4\pi R^2$, which gives Eq. (16.2). Dividing Eq. (16.2) for two planets shows how temperature scales:

$$\frac{T_1}{T_2} = \left(\frac{S_1(1-a_1)}{S_2(1-a_2)}\right)^{1/4},$$

so with equal albedos, a planet at a quarter of the sunlight (twice the distance) is $4^{1/4} = \sqrt{2} = 1.41$ times colder.

B.2.4 Beam spreading

In Figure 18.1, draw a line across the beam at right angles to the rays: its length is the beam width, 1 unit. This line, the strip of ground covered by the beam (length A), and the edge ray make a right-angled triangle. The ground strip is the hypotenuse, and the beam width is the side opposite the angle β at which the rays meet the ground. By the definition of the sine,

$$\sin\beta = \frac{\text{opposite}}{\text{hypotenuse}} = \frac{1}{A} \qquad \Longrightarrow \qquad A = \frac{1}{\sin\beta},$$

which is Eq. (18.2). The intensity per unit of ground is the beam's energy divided by A, which is Eq. (18.3). In words: the energy per unit area is proportional to the sine of the sun angle, or equivalently to the cosine of the zenith angle, since $\sin\beta = \cos\theta_z$.

B.2.5 The noon zenith angle

The local vertical at latitude φ points along the Earth's radius, at an angle φ from the equatorial plane (Figure 18.2). At noon the Sun lies in the same north–south plane, in the direction at angle δ from the equatorial plane, where δ is the latitude at which it is overhead. The Sun's rays are parallel across the Earth, because the Sun is so far away, so the angle between the vertical and the direction of the Sun is the difference of the two angles:

$$\theta_z = |\varphi - \delta|,$$

which is Eq. (18.4). With $\delta = +23.5°$ (June) and $-23.5°$ (December), and a site north of the Tropic of Cancer, this gives the sheet's $\theta_z = \varphi - 23.5°$ and $\theta_z = \varphi + 23.5°$. The value 23.5° is the tilt of the Earth's axis to the plane of its orbit (more precisely 23.44°).

B.3 The spectra of Question 3, recomputed

Each column of the sheet's Question 3 data (Table 15.1) was compared with the Planck curve, Eq. (12.3), by finding the temperature and the scale factor that minimise the relative differences between the two. All three columns fit one scale factor, about 316 times the spectral radiance in $\mathrm{W\,m^{-2}\,sr^{-1}\,\mu m^{-1}}$, and the best-fitting temperatures are

$$T_A = 254.1\ \mathrm{K}, \qquad T_B = 210.0\ \mathrm{K}, \qquad T_C = 184.1\ \mathrm{K},$$

the black-body temperatures of the Earth, Mars and Venus calculated in Question 3c (Table 16.1). Table B.1 sets the sheet's values beside the fitted curves: they agree to within about 0.3 % everywhere, which confirms that the data are Planck curves sampled every 2 μm.

Because the true peaks fall between the 2 μm sampling points, the highest data point in each column lies 0.3–0.6 μm to the right of the true peak. That is why Question 3b, which reads the peak from the data, gives temperatures a few kelvin below the true ones (§ 15.4).

B.4 Corrections to the printed key, at a glance

Each item was found by recomputing the answer sheet (EGM305 module team, 2025a) independently. The section column links to the full explanation. None of the corrections changes the physics the practical teaches.

Table B.1. The data of Question 3 ("sheet") beside the Planck curves for 254.1, 210.0 and 184.1 K on the same scale ("Planck"). Wavelengths in μm.

	Data A		**Data B**		**Data C**	
λ	sheet	Planck, 254.1 K	sheet	Planck, 210.0 K	sheet	Planck, 184.1 K
4	26.199	26.16	1.3338	1.338	0.11992	0.1202
6	386.28	385.9	53.056	53.18	10.648	10.67
8	970.65	969.8	218.86	219.2	65.619	65.7
10	1313.3	1312	398.14	398.6	151.8	151.9
12	1363.5	1363	502.49	503	224.7	224.9
14	1248.6	1248	527.86	528.3	264.23	264.4
16	1074.3	1074	502.41	502.8	273.39	273.5
18	896.25	895.8	452.58	452.9	262.51	262.6
20	737.1	736.7	395.22	395.4	241.03	241.1
22	603.1	602.8	339.25	339.4	215.38	215.5
24	493.45	493.2	288.62	288.8	189.36	189.4
26	404.91	404.7	244.61	244.7	164.92	165
28	333.76	333.6	207.16	207.2	142.9	142.9
30	276.58	276.5	175.67	175.7	123.56	123.6
True peaks, Eq. (13.1) with $C = 2897.8$:			A 11.40 μm,		B 13.80 μm,	C 15.74 μm.

Table B.2. Corrections and clarifications to the printed sheet and answer key.

Q	**The key or sheet says**	**Correct or recommended**	**See**
1	°C = K − 273.16	exact offset 273.15; no effect at this precision	§ 13.3
2b	$(5.67 \times 10^{-8} \times 287.16)^4$; unit "Wm2"	$5.67 \times 10^{-8} \times 287.16^4 = 385.5\,\mathrm{W\,m^{-2}}$ (the number is right)	§ 14.4
2c	5522.8/287.16 = 20 (Celsius over kelvin)	5796/287.16 = 20.2, both in kelvin	§ 14.5
2e	"E changes exponentially"	a power law: $E \propto T^4$	§ 14.7
3	data headed "Irradiance, $\mathrm{W\,m^{-2}}$"	spectral emission per μm, on an arbitrary scale	§ 15.1
3b	A: 241.5 K	right for the 2 μm data; the true curve is 254.1 K	§ 15.4
3d	two parts labelled "3d"	second one called 3e here	§ 11.2
3e(ii)	Venus albedo 0.9	measured Bond albedo 0.77 (227 K); conclusion unchanged	§ 16.7
3e(iii)	"runaway heating"	an intense greenhouse effect, in balance today; "runaway greenhouse" describes its past	§ 16.7
3e(iv)	Mercury the only match	Mars matches too (thin atmosphere)	§ 16.7
4b	reflected $0.3 \times 1368 = 41.04$	$410.4\,\mathrm{W\,m^{-2}}$	§ 17.3
4b	$1368\,\mathrm{W\,m^{-2}}$ "at the equator"	at the top of the atmosphere, on a surface facing the Sun	§ 17.3
4b′	average $341\,\mathrm{W\,m^{-2}}$	1368/4 = 342; 341 is the measured value (solar constant 1365)	§ 17.4

Q	The key or sheet says	Correct or recommended	See
4e	greenhouse gases “prevent all solar energy escaping to space”	they absorb outgoing infrared and re-emit part downward; the Earth still emits what it absorbs	§ 17.7
4g	“Earth does not radiate its own heat”	no internal heat *source*; also a black body at one temperature, in equilibrium	§ 17.9
5a	spreading at 60° N in June “1.2 units”; “90 − 83.5.5”	1.244 units; $90 - 83.5 = 6.5°$	§ 18.6
5c(i)	no final answer	59.5 against 11.3 % per unit: 48.2 points, 5.3 times	§ 18.6
5c(ii)	80.6 and 11.4 % per unit; “69.2 % more”	80.4 and 11.3 % per unit; 69.1 percentage points, 7.1 times (610 % more)	§ 18.6

B.5 Symbols and constants

Table B.3. Every symbol used in this part. Each equation’s own *where* list repeats the symbols it uses.

Symbol	Meaning	Unit	First used
a	albedo: fraction of sunlight reflected	–	Eq. (16.2)
A	beam spreading (the sheet’s “surface area”)	units	Eq. (18.2)
B_λ	spectral radiance	$\mathrm{W\,m^{-2}\,sr^{-1}\,m^{-1}}$	Eq. (12.3)
β	sun angle (solar elevation)	degrees	Eq. (18.1)
c	speed of light	$\mathrm{m\,s^{-1}}$	Eq. (12.1)
C	Wien’s constant	$\mu\mathrm{m\,K}$	Eq. (13.1)
d	distance from the Sun	AU	Eq. (16.1)
δ	solar declination	degrees	Eq. (18.4)
E	emission per square metre (“intensity of radiation”)	$\mathrm{W\,m^{-2}}$	Eq. (14.1)
ϵ	emissivity	–	§ 12.6
ε	energy of one photon	J	Eq. (12.2)
f	frequency	Hz	Eq. (12.1)
h	Planck’s constant	J s	Eq. (12.2)
I	solar intensity per unit of ground	% per unit	Eq. (18.3)
k_B	Boltzmann’s constant	$\mathrm{J\,K^{-1}}$	Eq. (12.3)
λ	wavelength	m or μm	Eq. (12.1)
$\lambda_{\max}$	wavelength of peak emission	μm	Eq. (13.1)
φ	latitude	degrees	Eq. (18.4)
R	radius of a planet	m	§ 16.4
S	solar constant at a planet	$\mathrm{W\,m^{-2}}$	Eq. (16.1)
$\bar{S}$	sunlight averaged over the globe, day and night	$\mathrm{W\,m^{-2}}$	Eq. (17.2)
σ	Stefan–Boltzmann constant	$\mathrm{W\,m^{-2}\ K^{-4}}$	Eq. (14.1)
T	absolute temperature	K	Eq. (13.1)

Symbol	Meaning	Unit	First used
θ_z	zenith angle	degrees	Eq. (18.1)

Table B.4. Constants. Answers in this part use the values on the sheet, so that they match the answer key; the precise values are given for reference.

Quantity	Value on the sheet	Precise value	Source
Wien's constant C	$2898\,\mu\mathrm{m\,K}$	$2897.77\,\mu\mathrm{m\,K}$	Tiesinga et al. (2021)
Stefan–Boltzmann σ	$5.67 \times 10^{-8}\,\mathrm{W\,m^{-2}\,K^{-4}}$	5.670374×10^{-8}	Tiesinga et al. (2021)
Planck's constant h	–	$6.62607015 \times 10^{-34}\,\mathrm{J\,s}$ (exact)	Tiesinga et al. (2021)
Speed of light c	–	$2.99792458 \times 10^{8}\,\mathrm{m\,s^{-1}}$ (exact)	Tiesinga et al. (2021)
Boltzmann's constant k_B	–	$1.380649 \times 10^{-23}\,\mathrm{J\,K^{-1}}$ (exact)	Tiesinga et al. (2021)
Solar constant S	$1368\,\mathrm{W\,m^{-2}}$	$1361\,\mathrm{W\,m^{-2}}$	Kopp and Lean (2011); Prša et al. (2016)
Earth's albedo	0.30	about 0.29–0.30	Forster et al. (2021); Williams (2024)
Kelvin–Celsius offset	273.16	273.15 (exact)	Tiesinga et al. (2021)
Earth's axial tilt	23.5°	23.44°	Stull (2017)
Sun's effective temperature	–	5772 K	Prša et al. (2016)

B.6 About the figures and the data

B.6.1 How the figures were made

Every figure in this part was drawn for it as vector graphics, in the document itself with TikZ and pgfplots, so it stays sharp at any size and prints clearly in black and white (each curve has its own line style or marker as well as its own colour).

- The **Planck curves** (Figures 12.3, 13.1 and 15.1) are computed from Eq. (12.3) with the constants of Table B.4.
- The **Question 3 graph** (Figure 15.1) plots the values printed on the sheet; the curves through them are the fits described in Appendix B.3.
- The **planetary temperatures** (Figure 16.2) are the values of Table 16.1: calculated with Eq. (16.2), and the actual averages as printed on the sheet.
- The **sun-angle sketches** (Figure 18.3) are drawn to the exact angles of Table 18.1 and to one scale, each beam one unit (5 mm) wide, so the lengths of the red strips can be compared directly.
- The **energy flows** (Figure 17.1) are the values of Questions 4b′–4d, with arrow widths proportional to the flows.

B.6.2 Limits to bear in mind

- The black-body model treats each planet as having one temperature, one albedo and no internal heat. It explains the main differences between the planets, not the details of any

one of them.

- The sun-angle calculation gives the intensity at *noon* at the top of the atmosphere. Daily totals also depend on day length, and the energy reaching the ground on cloud and on the length of the path through the air (§ 18.5).
- Values for the planets (albedos, average temperatures) are rounded on the sheet; current measured values are in Williams (2024).

Appendix C

Practical 3: formulas, data, derivations and corrections

C.1 Formula sheet

Table C.1. The relationships used in this part.

Quantity	Formula	Where, and what for
Saturation vapour pressure	$e_s(T) = 6.112 \exp\left[\dfrac{17.67\,T}{T + 243.5}\right]$ hPa (T in °C)	Table 1, dew point
Saturation mixing ratio	$w_s = \dfrac{622\,e_s}{p - e_s}$ g kg^{-1}, with $p = 995.5$ hPa	Table 1
Mixing ratio from RH	$w = w_s \times \mathrm{RH}/100$	Eq. (21.4); Moisture Q3, Q6
Relative humidity	$\mathrm{RH} = 100\,w/w_s = 100\,e/e_s$	Eq. (21.3); Moisture Q1
Vapour pressure from w	$e = \dfrac{w\,p}{622 + w}$	Eq. (21.2); dew point (formula)
Dew point	$T_d = \dfrac{243.5\,\ln(e/6.112)}{17.67 - \ln(e/6.112)}$	Moisture Q5, Q6
Rule of thumb	$\mathrm{RH} \approx 100 - 5\,(T - T_d)$	Eq. (22.1); checks
Cloud base	$z \approx 125\,(T - T_d)$ m	Eq. (22.3); cloud height
Condensed water	$w - w_s(T)$	Eq. (22.2); below dew point
Hypsometric equation	$\Delta z = \dfrac{R_d\,\overline{T}}{g} \ln \dfrac{p_1}{p_2}$	Eq. (24.1); Motion Q1–Q6
Pressure gradient force	$-\dfrac{1}{\rho}\dfrac{\Delta p}{\Delta n} = -g\dfrac{\Delta z}{\Delta n}$ (on a pressure surface)	Eq. (23.1); Chapters 23–26
Coriolis parameter	$f = 2\Omega \sin\varphi$, $\Omega = 7.292 \times 10^{-5}$ s^{-1}	Motion Q7
Coriolis force	fV per unit mass	Motion Q8
Geostrophic wind	$V_g = \dfrac{1}{\rho f}\dfrac{\Delta p}{\Delta n} = \dfrac{g}{f}\dfrac{\Delta z}{\Delta n}$	Eq. (25.1); Motion Q6, Q7
Wind with friction	$\alpha = \arctan(k/f)$, $V = V_g \cos\alpha$	Eq. (26.1); Motion Q9–Q11
Temperature	$T(°\mathrm{C}) = \frac{5}{9}[T(°\mathrm{F}) - 32]$	Eq. (27.1); Map Q12
Pressure code	3 digits: decimal before last; prefix 10 if < 500, 9 if ≥ 500	Map Q12
Knots	$1\,\mathrm{kt} = 0.514\,\mathrm{m\,s^{-1}} = 1.85\,\mathrm{km\,h^{-1}}$	Map Q12
Interpolation	$\text{fraction} = \dfrac{v_{\text{wanted}} - v_1}{v_2 - v_1}$	Map Q14

Constants. $g = 9.81\,\mathrm{m\,s^{-2}}$; $R_d = 287\,\mathrm{J\,kg^{-1}\,K^{-1}}$; $\rho \approx 1.2\,\mathrm{kg\,m^{-3}}$ near the ground; $\Omega =$

$7.292 \times 10^{-5}\,\mathrm{s}^{-1}$; Earth's radius 6371 km.

C.2 Table 1, with the two corrections

Table 1 of the sheet gives the saturation vapour pressure (mb) and the saturation mixing ratio (g/kg) at sea level. The values below are those printed on the sheet. The two shaded rows contain a mis-printed mixing ratio; the fourth column gives the value that fits the formula $w_s = 622\, e_s/(p - e_s)$ with $p = 995.5$ hPa (see Section 21.11).

Table C.2. Saturation vapour pressure e_s and saturation mixing ratio w_s against temperature, as printed on the sheet.

Temp. (°C)	e_s **(mb)**	w_s **printed** ($\mathrm{g\,kg^{-1}}$)	**corrected** ($\mathrm{g\,kg^{-1}}$)
-40	0.189	0.118	
-35	0.314	0.195	
-30	0.519	0.318	
-25	0.807	0.510	
-20	1.254	0.784	
-18	1.488	0.931	
-16	1.760	1.102	
-14	2.076	1.300	
-12	2.441	1.529	
-10	2.863	1.794	
-8	3.348	2.009	2.099
-6	3.906	2.450	
-4	4.545	2.852	
-2	5.275	3.313	
0	6.108	3.189	3.840
2	7.055	4.439	
4	8.129	5.120	
6	9.347	5.894	
8	10.722	6.771	
10	12.272	7.762	
12	14.017	8.882	
14	15.977	10.140	
16	18.173	11.560	
18	20.630	13.162	
20	23.373	14.956	
22	26.430	16.963	
24	29.831	19.210	
26	33.608	21.734	
28	37.796	24.557	
30	42.430	27.694	
32	47.551	31.213	
34	53.200	35.134	

Temp. (°C)	e_s (mb)	w_s printed ($\mathrm{g\,kg^{-1}}$)	corrected ($\mathrm{g\,kg^{-1}}$)
36	59.422	39.502	
38	66.264	44.381	
40	73.777	49.815	

Using the table. (1) To look up, read across from the temperature. (2) To interpolate, take the proportion of the way between two rows. (3) To look up backwards (for a dew point), find the two rows whose w_s bracket the value and interpolate for the temperature. Linear interpolation over a 2-degree step is accurate to about 0.1 %.

C.3 The 34 stations of Figures 8 and 10

The values below were read from the printed Figure 10. Temperatures and dew points are in °F; the pressure is the three-digit code and the decoded value in hPa; cloud cover is in percent (the filled fraction of the circle); the wind is the direction it comes **from** and its speed in knots (counted from the barbs, so approximate). Positions are the latitude and longitude of each station found by fitting a map projection to the printed figure, so they are good to about 10 km. A dash means that the figure does not print the value (station 9 has no dew point).

#	Position	Temp.	Dew	Code	hPa	Sky	Wind	Weather
1	44.6°N, 98.4°W	34	22	110	1011.0	25	NNW 10	
2	44.3°N, 92.9°W	45	43	096	1009.6	100	E 10	rain (continuous)
3	44.3°N, 85.6°W	39	39	154	1015.4	75	ENE 15	
4	42.7°N, 98.7°W	36	28	124	1012.4	50	NW 10	
5	42.5°N, 95.9°W	40	38	088	1008.8	100	NW 20	rain (intermittent)
6	43.4°N, 88.8°W	39	39	084	1008.4	100	SE 20	rain (continuous)
7	42.9°N, 85.8°W	43	41	132	1013.2	60	ESE 20	
8	42.0°N, 91.3°W	48	45	024	1002.4	100	NE 10	rain (continuous)
9	41.2°N, 93.9°W	48	–	026	1002.6	100	WNW 10	rain (continuous)
10	41.6°N, 86.0°W	46	43	124	1012.4	100	E 5	rain (continuous)
11	40.6°N, 83.1°W	45	43	163	1016.3	100	ESE 2	rain (intermittent)
12	40.8°N, 100.3°W	39	31	124	1012.4	0	NW 10	
13	40.8°N, 89.9°W	57	54	029	1002.9	75	SSW 10	
14	40.1°N, 86.4°W	54	45	097	1009.7	100	E 10	
15	39.6°N, 98.5°W	41	30	103	1010.3	10	NW 10	
16	39.7°N, 95.0°W	45	40	069	1006.9	50	NW 25	
17	39.8°N, 92.7°W	60	55	040	1004.0	100	SSW 25	thunder, rain
18	39.1°N, 90.4°W	61	57	044	1004.4	50	SSW 10	rain (continuous)
19	38.7°N, 87.6°W	66	53	099	1009.9	75	S 10	
20	38.0°N, 86.3°W	66	55	111	1011.1	25	S 10	
21	37.7°N, 100.5°W	41	33	126	1012.6	0	NW 10	
22	38.0°N, 97.5°W	42	36	109	1010.9	0	NW 10	
23	37.2°N, 93.7°W	61	57	075	1007.5	100	SSW 10	rain (continuous)
24	36.3°N, 86.5°W	71	60	149	1014.9	50	S 10	

#	Position	Temp.	Dew	Code	hPa	Sky	Wind	Weather
25	36.0°N, 97.3°W	48	38	126	1012.6	0	WNW 10	
26	35.7°N, 89.8°W	70	59	140	1014.0	50	S 10	
27	34.9°N, 100.8°W	49	38	128	1012.8	0	WNW 10	
28	35.0°N, 93.9°W	66	59	114	1011.4	75	SW 10	
29	34.6°N, 92.5°W	66	60	124	1012.4	0	SSW 10	
30	33.3°N, 98.7°W	50	38	120	1012.0	0	NW 10	
31	31.9°N, 97.0°W	52	50	128	1012.8	100	WSW 10	
32	32.5°N, 93.7°W	63	61	153	1015.3	100	S 10	
33	32.6°N, 90.2°W	69	61	171	1017.1	25	SSW 10	
34	33.1°N, 87.0°W	72	64	194	1019.4	0	SSW 10	

If a value differs from your copy of the sheet, trust the sheet. The map figures in this part are redrawn from this table, so correcting the table and regenerating the figures would update them all.

C.4 Derivations for the curious

Nothing in this appendix is needed to answer the sheet, but each result is used in the chapters. They are short.

C.4.1 Mixing ratio and vapour pressure

Treat the dry air and the water vapour as ideal gases at the same temperature T and volume. Their densities are $\rho_d = (p - e)/(R_d T)$ and $\rho_v = e/(R_v T)$, where p is the total pressure, e the vapour pressure and R_d, R_v the gas constants of dry air and of water vapour. The mixing ratio is

$$w = \frac{\rho_v}{\rho_d} = \frac{R_d}{R_v}\,\frac{e}{p - e} = \varepsilon\,\frac{e}{p - e}, \qquad \varepsilon = \frac{R_d}{R_v} = \frac{18.02}{28.96} = 0.622.$$

Multiplied by 1000 this is the $\mathrm{g\,kg^{-1}}$ of Table 1: $w = 622\,e/(p - e)$.

C.4.2 Saturation vapour pressure

The Clausius–Clapeyron equation relates the change of the saturation vapour pressure of water to temperature:

$$\frac{de_s}{dT} = \frac{L_v\, e_s}{R_v\, T^2},$$

where $L_v = 2.5 \times 10^6\ \mathrm{J\,kg^{-1}}$ is the latent heat of vaporisation. At 15 °C this gives $\frac{1}{e_s}\frac{de_s}{dT} = 6.9\,\%\ \mathrm{K^{-1}}$, which is why the curve of Figure 21.1 is steep and bends upward. Integrating with a constant L_v gives an exponential in $-1/T$; fitting the result over the range of the atmosphere gives the convenient empirical form

$$e_s(T) = 6.112 \exp\!\left[\frac{17.67\,T}{T + 243.5}\right]\ \mathrm{hPa} \qquad (T \text{ in } ^\circ\mathrm{C}),$$

which reproduces the values of Table 1 to better than 0.2 % between −20 and 40 °C.

C.4.3 The dew point

The dew point is the temperature at which the saturation vapour pressure equals the actual vapour pressure e. Setting $e_s(T_d) = e$ in the empirical form and solving for T_d gives, with $x = \ln(e/6.112)$,

$$T_d = \frac{243.5\,x}{17.67 - x}.$$

For the sample of Question 6, $w = 21.78\,\mathrm{g\,kg^{-1}}$, $p = 995.5\,\mathrm{hPa}$: $e = 21.78 \times 995.5/(622 + 21.78) = 33.68\,\mathrm{hPa}$, $x = 1.7066$, $T_d = 26.03\,^\circ\mathrm{C}$.

C.4.4 Cloud base from the dew point

A rising unsaturated parcel cools at the dry adiabatic rate $\Gamma_d = g/c_p = 9.8\,\mathrm{K\,km^{-1}}$. Its dew point falls more slowly, because the vapour pressure falls with the pressure: about $1.8\,\mathrm{K\,km^{-1}}$. The gap $T - T_d$ therefore closes at $9.8 - 1.8 = 8\,\mathrm{K\,km^{-1}}$, so the parcel saturates after rising $(T - T_d)/8$ km $\approx 125\,(T - T_d)$ m.

C.4.5 The hypsometric equation

The vertical balance (hydrostatic balance) says that pressure falls with height by the weight of the air: $dp = -\rho g\,dz$. Using the ideal gas law $\rho = p/(R_d T)$,

$$dz = -\frac{R_d T}{g}\frac{dp}{p}.$$

Integrating from p_1 (lower) to p_2 (upper) with the mean temperature $\overline{T}$ of the layer,

$$z_2 - z_1 = \frac{R_d \overline{T}}{g}\ln\frac{p_1}{p_2}.$$

Thickness is proportional to the mean temperature: that is why pressure surfaces rise over warm air.

C.4.6 The pressure gradient force on a pressure surface

At a fixed height the horizontal pressure gradient force per unit mass is $-\frac{1}{\rho}\frac{\partial p}{\partial n}$. On a surface of constant pressure, the height varies with position, and the hydrostatic relation $\partial p/\partial z = -\rho g$ converts the horizontal pressure gradient at fixed height into a height gradient along the surface:

$$-\frac{1}{\rho}\left(\frac{\partial p}{\partial n}\right)_z = -g\left(\frac{\partial z}{\partial n}\right)_p.$$

The density has disappeared: the force depends only on the slope of the pressure surface. This is why the sheet's questions about the slope of the surfaces are questions about the force.

C.4.7 The geostrophic wind

For a wind V blowing along the isobars, the pressure gradient force $g\,\Delta z/\Delta n$ and the Coriolis force fV act at right angles to the wind and to each other's line, in opposite directions. Setting them equal,

$$fV_g = g\frac{\Delta z}{\Delta n} \quad\Longrightarrow\quad V_g = \frac{g}{f}\frac{\Delta z}{\Delta n}.$$

In terms of pressure: $V_g = \Delta p/(\rho f\,\Delta n)$.

C.4.8 The steady wind with friction

Take x east, y north, a pressure gradient force (P_x, P_y), Coriolis acceleration $(fv, -fu)$ and a linear drag $(-ku, -kv)$. In a steady state the three add to zero:

$$P_x + fv - ku = 0, \qquad P_y - fu - kv = 0.$$

Solving the pair for u and v,

$$u = \frac{kP_x + fP_y}{k^2 + f^2}, \qquad v = \frac{kP_y - fP_x}{k^2 + f^2}.$$

For the low on the north side and isobars running east–west ($P_x = 0$, $P_y = P > 0$) this gives $u = fP/(k^2 + f^2)$ and $v = kP/(k^2 + f^2)$: a wind with a component towards the low. The angle to the isobars satisfies $\tan\alpha = v/u = k/f$ and the speed is $\sqrt{u^2 + v^2} = P/\sqrt{k^2 + f^2} = V_g \cos\alpha$, with $V_g = P/f$. For $k = 0$ this is the geostrophic wind.

C.4.9 Pressure at a height between two reported levels

In Question 4 the pressure at 9000 m was estimated from the reported levels. Between two levels with pressures p_a and p_b at heights z_a and z_b, the pressure was interpolated linearly in $\ln p$:

$$\ln p = \ln p_a + (\ln p_b - \ln p_a)\,\frac{z - z_a}{z_b - z_a},$$

which is the exact result for an isothermal layer. Above the highest level an isothermal extension $p = p_{300}\exp[-g(z - z_{300})/(R_d T)]$ was used.

C.5 Weather symbols: quick reference

The symbols of the sheet's Appendix, redrawn. Cover the right-hand descriptions and test yourself.




Figure C.1. Sky cover: no cloud, ≤ 1/10, 2–3/10, 4/10, 5/10, 6/10, 7–8/10, 9/10, 10/10 (overcast), obscured.

C.6 Corrections to the printed key, at a glance

Each item was found by checking the answer key (EGM305 module team, n.d.) against Table 1 and the figures of the sheet. The last column links to the full explanation.




Figure C.2. Wind barbs (staff pointing north, from the north): calm, 1–2, 5, 10, 15, 20, 25, 30, 40, 50, 60, 65, 75 and 105 knots.




Figure C.3. Present weather: intermittent rain, continuous rain, continuous snow, continuous drizzle, fog, thunder heard, thunder with intermittent rain.

Table C.4. Corrections to the Practical 3 answer key and sheet.

Q	The key or sheet says	Correct	See
Moisture Q6	dew point 25 °C	about **26** °C: the sample holds 21.78 g kg^{-1}, more than 25 °C air can hold (20.4 g kg^{-1})	§ 22.4
Map Q12	station model 1: dew point 28 °F	**21** °F, as printed at the bottom left of the model	§ 27.8
Table 1	3.189 g kg^{-1} at 0 °C	**3.840** g kg^{-1} (typing slip; no question uses this row)	§ C.2
Table 1	2.009 g kg^{-1} at −8 °C	**2.099** g kg^{-1} (typing slip; no question uses this row)	§ C.2




Figure C.4. Barometric tendency over the past three hours.

C.7 About the figures, the data and further reading

C.7.1 How the figures were made

Every figure in this part was drawn for it and is vector graphics: the curves and glasses, the force diagrams and the weather maps are all drawn in the document itself with TikZ and pgfplots, so they stay sharp at any size.

- The **saturation curves** are computed from the formulas of Appendix C.1; the dots are the values printed in Table 1.
- The **geostrophic adjustment** (Figures in Chapter 25) is the numerical solution of the equations of motion of a parcel released at rest between two height lines, with a gentle damping that relaxes the velocity towards the geostrophic wind.
- The **four circulation systems** (Chapter 26) are computed from the steady force balance of Appendix C.4, with friction equal to 0.6 times the Coriolis force.
- The **map**: state outlines are boundaries from Natural Earth (public domain), drawn in the Lambert conformal conic projection, fitted to the printed Figure 10 so that the stations fall on the right states. The 34 stations were read from the figure, as described in Chapter 20 and Appendix C.3.
- The **isobars** are contours of a smooth thin-plate-spline surface through the 34 station pressures; the **cloud and rain areas** are contours of distance-weighted averages of the station cover and rain reports; the **streamlines** follow a smoothed wind field made from the 34 barbs. The **fronts** were placed by hand, from the temperature, dew-point and wind contrasts, as an analyst would.

C.7.2 Limits to bear in mind

- Station values are read from a printed figure: wind speeds are estimated by counting barbs, and a few cloud-cover symbols are open to a one-tenth difference of interpretation. Check any value against your sheet.
- The geostrophic-wind estimates between Saskatoon and Monterrey use only two places and rounded heights. They show the right behaviour (increase of wind with height) but not exact winds.
- The fronts and the isobars are *analyses*: reasonable analysts can draw them slightly differently. What matters is that every rule is respected (isobars never cross, fronts follow the contrasts of temperature, moisture and wind).

C.7.3 Further reading

- Ahrens and Henson (2022): the standard introduction; the station model, humidity and the forces on the wind are all treated at the level of the sheet.
- Stull (2017): a free textbook that includes the hypsometric equation, the geostrophic wind and station models with worked examples.
- Wallace and Hobbs (2006): thermodynamics of moist air, the balance of forces, and fronts and cyclones.
- Holton and Hakim (2013): the geostrophic wind, thermal wind and friction at a more advanced level.
- Bolton (1980): the saturation vapour pressure formula used here.
- World Meteorological Organization (2019): the definition of the station model and its codes.

Appendix D

Constants, units and conversions

D.1 Constants

Each part uses the values printed on its practical sheet, so that answers match the key. The precise values are given for reference.

Table D.1. Constants used in the companion.

Quantity	Value used	Precise value	Source	Part
Acceleration due to gravity g	$9.81\,\mathrm{m\,s^{-2}}$	$9.80665\,\mathrm{m\,s^{-2}}$ (standard)	NOAA et al. (1976)	I, III
Gas constant for dry air R_d	$287\,\mathrm{J\,kg^{-1}\,K^{-1}}$	$287.05\,\mathrm{J\,kg^{-1}\,K^{-1}}$	Wallace and Hobbs (2006)	I, III
Specific heat of dry air c_p	$1004\,\mathrm{J\,kg^{-1}\,K^{-1}}$	$1004\,\mathrm{J\,kg^{-1}\,K^{-1}}$	Wallace and Hobbs (2006)	I
Standard sea-level pressure	1000 hPa (Part I sheet)	1013.25 hPa	NOAA et al. (1976)	I
Standard sea-level temperature	15 °C	288.15 K	NOAA et al. (1976)	I
Stefan–Boltzmann constant σ	$5.67 \times 10^{-8}\,\mathrm{W\,m^{-2}\,K^{-4}}$	5.670374×10^{-8}	Tiesinga et al. (2021)	II
Wien's constant C	$2898\,\mu\mathrm{m\,K}$	$2897.77\,\mu\mathrm{m\,K}$	Tiesinga et al. (2021)	II
Planck's constant h	–	$6.62607015 \times 10^{-34}\,\mathrm{J\,s}$ (exact)	Tiesinga et al. (2021)	II
Speed of light c	–	$2.99792458 \times 10^{8}\,\mathrm{m\,s^{-1}}$ (exact)	Tiesinga et al. (2021)	II
Boltzmann's constant k_B	–	$1.380649 \times 10^{-23}\,\mathrm{J\,K^{-1}}$ (exact)	Tiesinga et al. (2021)	II
Solar constant S	$1368\,\mathrm{W\,m^{-2}}$	$1361\,\mathrm{W\,m^{-2}}$	Kopp and Lean (2011)	II
Earth's axial tilt	23.5°	23.44°	Stull (2017)	II
Earth's rotation rate Ω	$7.292 \times 10^{-5}\,\mathrm{s^{-1}}$	$7.292 \times 10^{-5}\,\mathrm{s^{-1}}$	Holton and Hakim (2013)	III
Kelvin–Celsius offset	273.15 (273.16 on the Part II sheet)	273.15 (exact)	Tiesinga et al. (2021)	all

D.2 Units and conversions

Quantity	**Conversion**
Pressure	$1\ \text{hPa} = 1\ \text{mb} = 100\ \text{Pa}$; $1\ \text{mmHg (torr)} = 1.333\ \text{hPa}$
Temperature	$T_\text{K} = T_{\circ\text{C}} + 273.15$; $T_{\circ\text{C}} = \frac{5}{9}(T_{\circ\text{F}} - 32)$
Temperature difference	$1\ \text{K} = 1\ ^\circ\text{C} = 1.8\ ^\circ\text{F}$
Wind speed	$1\ \text{kt} = 0.514\ \text{m s}^{-1} = 1.852\ \text{km h}^{-1}$
Length	$1\ \text{km} = 1000\ \text{m}$; $1\ \mu\text{m} = 10^{-6}\ \text{m}$
Power per area	W m^{-2}: watts per square metre

References

C. Donald Ahrens and Robert Henson. *Meteorology Today: An Introduction to Weather, Climate, and the Environment*. Cengage Learning, Boston, MA, 13th edition, 2022.

David Bolton. The computation of equivalent potential temperature. *Monthly Weather Review*, 108(7):1046–1053, 1980. doi: 10.1175/1520-0493(1980)108<1046:TCOEPT>2.0.CO;2.

J. H. Davies and D. R. Davies. Earth's surface heat flux. *Solid Earth*, 1:5–24, 2010. doi: 10.5194/se-1-5-2010.

EGM305 module team. EGM305 The Atmosphere, lecture 2: Temperature. Lecture slides. Coleraine: School of Geography and Environmental Sciences, Ulster University, 2020.

EGM305 module team. EGM305 practical 2: Solar radiation (radiation and heating). Practical sheet and answer sheet. Coleraine: School of Geography and Environmental Sciences, Ulster University, 2025a.

EGM305 module team. EGM305 practical 1: Vertical structure of the atmosphere. Practical sheet and answer sheet. Coleraine: School of Geography and Environmental Sciences, Ulster University, 2025b.

EGM305 module team. EGM305 The Atmosphere, lecture 1: Introduction to the atmosphere. Lecture slides. Coleraine: School of Geography and Environmental Sciences, Ulster University, 2026.

EGM305 module team. EGM305 practical 3: Atmospheric moisture, atmospheric motion and weather map analysis. Practical sheet and answer key. Coleraine: School of Geography and Environmental Sciences, Ulster University, n.d.

Piers Forster, Trude Storelvmo, Kyle Armour, et al. The Earth's energy budget, climate feedbacks, and climate sensitivity. In V. Masson-Delmotte et al., editors, *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, pages 923–1054. Cambridge University Press, Cambridge, 2021. doi: 10.1017/9781009157896.009.

R. A. Hanel, B. J. Conrath, L. W. Herath, V. G. Kunde, and J. A. Pirraglia. Albedo, internal heat, and energy balance of Jupiter: preliminary results of the Voyager infrared investigation. *Journal of Geophysical Research: Space Physics*, 86(A10):8705–8712, 1981. doi: 10.1029/JA086iA10p08705.

Klaus P. Hoinka. Statistics of the global tropopause pressure. *Monthly Weather Review*, 126(12): 3303–3325, 1998. doi: 10.1175/1520-0493(1998)126<3303:SOTGTP>2.0.CO;2.

James R. Holton and Gregory J. Hakim. *An Introduction to Dynamic Meteorology*. Academic Press, Amsterdam, 5th edition, 2013.

J. F. Kasting. Runaway and moist greenhouse atmospheres and the evolution of Earth and Venus. *Icarus*, 74(3):472–494, 1988. doi: 10.1016/0019-1035(88)90116-9.

Greg Kopp and Judith L. Lean. A new, lower value of total solar irradiance: evidence and climate significance. *Geophysical Research Letters*, 38:L01706, 2011. doi: 10.1029/2010GL045777.

Frederick K. Lutgens, Edward J. Tarbuck, and Redina L. Herman. *The Atmosphere: An Introduction to Meteorology*. Pearson, New York, 14th edition, 2019.

NOAA, NASA, and US Air Force. *U.S. Standard Atmosphere, 1976*. U.S. Government Printing Office, Washington, DC, 1976. NOAA-S/T 76-1562.

Raymond T. Pierrehumbert. Infrared radiation and planetary temperature. *Physics Today*, 64(1): 33–38, 2011. doi: 10.1063/1.3541943.

Max Planck. Ueber das Gesetz der Energieverteilung im Normalspectrum. *Annalen der Physik*,

309(3):553–563, 1901. doi: 10.1002/andp.19013090310.

Andrej Prša, Petr Harmanec, Guillermo Torres, Eric Mamajek, et al. Nominal values for selected solar and planetary quantities: IAU 2015 Resolution B3. *The Astronomical Journal*, 152(2):41, 2016. doi: 10.3847/0004-6256/152/2/41.

Thomas Reichler, Martin Dameris, and Robert Sausen. Determining the tropopause height from gridded data. *Geophysical Research Letters*, 30(20), 2003. doi: 10.1029/2003GL018240.

Roland Stull. *Practical Meteorology: An Algebra-based Survey of Atmospheric Science*. University of British Columbia, Vancouver, 2017. URL `https://www.eoas.ubc.ca/books/Practical_Meteorology/`.

Eite Tiesinga, Peter J. Mohr, David B. Newell, and Barry N. Taylor. CODATA recommended values of the fundamental physical constants: 2018. *Reviews of Modern Physics*, 93(2):025010, 2021. doi: 10.1103/RevModPhys.93.025010.

Kevin E. Trenberth, John T. Fasullo, and Jeffrey Kiehl. Earth's global energy budget. *Bulletin of the American Meteorological Society*, 90(3):311–324, 2009. doi: 10.1175/2008BAMS2634.1.

John M. Wallace and Peter V. Hobbs. *Atmospheric Science: An Introductory Survey*. Elsevier Academic Press, Amsterdam, 2nd edition, 2006.

J. B. West, S. Lahiri, K. H. Maret, R. M. Peters, and C. J. Pizzo. Barometric pressures at extreme altitudes on Mt. Everest: physiological significance. *Journal of Applied Physiology*, 54 (5):1188–1194, 1983. doi: 10.1152/jappl.1983.54.5.1188.

John B. West. Prediction of barometric pressures at high altitudes with the use of model atmospheres. *Journal of Applied Physiology*, 81(4):1850–1854, 1996. doi: 10.1152/jappl.1996.81.4.1850.

David R. Williams. Planetary fact sheets: Mercury, Venus, Earth, Mars, Jupiter. Greenbelt, MD: NASA Goddard Space Flight Center, NSSDCA, 2024. URL `https://nssdc.gsfc.nasa.gov/planetary/factsheet/`. Accessed 21 October 2026.

J.-P. Williams, D. A. Paige, B. T. Greenhagen, and E. Sefton-Nash. The global surface temperatures of the Moon as measured by the Diviner Lunar Radiometer Experiment. *Icarus*, 283:300–325, 2017. doi: 10.1016/j.icarus.2016.08.012.

World Meteorological Organization. *Manual on Codes, Volume I.1: International Codes (WMO-No. 306)*. WMO, Geneva, 2019.

World Meteorological Organization. *Guide to Instruments and Methods of Observation, Volume I: Measurement of Meteorological Variables (WMO-No. 8)*. WMO, Geneva, 2023.