

Fonction	Dérivée	Intervalle
$f(x) = c$	$f'(x) = 0$	$] -\infty, +\infty[$
$f(x) = x$	$f'(x) = 1$	$] -\infty, +\infty[$
$f(x) = x^n$	$f'(x) = nx^{n-1}$	$] -\infty, +\infty[$ si $n \in \mathbb{N}$
$f(x) = \frac{1}{x}$	$f'(x) = -\frac{1}{x^2}$	$] -\infty, 0[$ ou $] 0, +\infty[$
$f(x) = \sqrt{x}$	$f'(x) = \frac{1}{2\sqrt{x}}$	$] 0, +\infty[$
$f(x) = x^\alpha$	$f'(x) = \alpha x^{\alpha-1}$	$] 0, +\infty[$ si $\alpha \in \mathbb{R}$
$f(x) = \ln x$	$f'(x) = \frac{1}{x}$	$] 0, +\infty[$
$f(x) = e^x$	$f'(x) = e^x$	$] -\infty, +\infty[$
$f(x) = \sin x$	$f'(x) = \cos x$	$] -\infty, +\infty[$
$f(x) = \cos x$	$f'(x) = -\sin x$	$] -\infty, +\infty[$
$f(x) = \tan x$	$f'(x) = \frac{1}{\cos^2 x} = 1 + \tan^2 x$	$\left] -\frac{\pi}{2} + k\pi, \frac{\pi}{2} + k\pi \right[$ où $k \in \mathbb{Z}$
$f(x) = \cotan x$	$f'(x) = -\frac{1}{\sin^2 x} = -1 - \cotan^2 x$	$] k\pi, (k+1)\pi[$ où $k \in \mathbb{Z}$

$f(x)$	Primitive
$x^n, n \neq -1$	$\frac{x^{n+1}}{n+1}$
$\frac{1}{x}$	$\ln  x $
$e^x$	$e^x$
$\ln(x)$	$x \ln(x) - x$
$\cos(x)$	$\sin(x)$
$\sin(x)$	$-\cos(x)$
$\frac{1}{1+x^2}$	$\arctan(x)$
$a^x, a > 0, a \neq 1$	$\frac{a^x}{\ln(a)}$

$f(x)$	Sens de variation
$x^n, n > 0$ impair	croissante
$x^n, n > 0$ pair	décroissante, si $x \leq 0$ croissante, si $x \geq 0$
$\frac{1}{x}$	décroissante
$e^x$	croissante
$\ln(x)$	croissante