

Chapter 4 Solutions, Susanna Epp Discrete Math 5th Edition

<https://github.com/spamegg1>

May 17, 2023

Contents

1	Exercise Set 4.1	13
1.1	Exercise 1	13
1.1.1	(a)	13
1.1.2	(b)	13
1.1.3	(c)	13
1.2	Exercise 2	13
1.2.1	(a)	13
1.2.2	(b)	13
1.2.3	(c)	13
1.3	Exercise 3	14
1.3.1	(a)	14
1.3.2	(b)	14
1.3.3	(c)	14
1.4	Exercise 4	14
1.4.1	(a)	14
1.4.2	(b)	14
1.4.3	(c)	15
1.5	Exercise 5	15
1.6	Exercise 6	15
1.7	Exercise 7	15
1.8	Exercise 8	15
1.9	Exercise 9	15
1.10	Exercise 10	16
1.11	Exercise 11	16
1.12	Exercise 12	16
1.13	Exercise 13	16
1.14	Exercise 14	17
1.15	Exercise 15	17
1.16	Exercise 16	17
1.17	Exercise 17	17

1.18	Exercise 18	17
1.19	Exercise 19	18
1.20	Exercise 20	18
1.21	Exercise 21	18
1.22	Exercise 22	18
1.23	Exercise 23	19
1.24	Exercise 24	19
1.25	Exercise 25	19
1.26	Exercise 26	20
1.27	Exercise 27	20
1.28	Exercise 28	20
1.29	Exercise 29	21
1.30	Exercise 30	21
1.31	Exercise 31	22
2	Exercise Set 4.2	22
2.1	Exercise 1	22
2.2	Exercise 2	23
2.3	Exercise 3	24
2.4	Exercise 4	24
2.5	Exercise 5	24
2.6	Exercise 6	25
2.7	Exercise 7	25
2.8	Exercise 8	26
2.9	Exercise 9	26
2.10	Exercise 10	27
2.11	Exercise 11	27
2.12	Exercise 12	27
2.13	Exercise 13	28
2.14	Exercise 14	28
2.15	Exercise 15	28
2.16	Exercise 16	28
2.17	Exercise 17	29
2.18	Exercise 18	29
2.19	Exercise 19	30
2.20	Exercise 20	31
2.21	Exercise 21	31
2.22	Exercise 22	31
2.23	Exercise 23	32
2.24	Exercise 24	32
2.25	Exercise 25	32
2.26	Exercise 26	32
2.27	Exercise 27	33
2.28	Exercise 28	33
2.29	Exercise 29	33
2.30	Exercise 30	33

2.31	Exercise 31	33
2.32	Exercise 32	34
2.33	Exercise 33	34
2.34	Exercise 34	34
2.35	Exercise 35	34
2.36	Exercise 36	35
2.37	Exercise 37	35
2.38	Exercise 38	35
2.39	Exercise 39	36
2.40	Exercise 40	36
2.41	Exercise 41	36
3	Exercise Set 4.3	36
3.1	Exercise 1	36
3.2	Exercise 2	36
3.3	Exercise 3	36
3.4	Exercise 4	36
3.5	Exercise 5	37
3.6	Exercise 6	37
3.7	Exercise 7	37
3.8	Exercise 8	37
3.8.1	(a)	37
3.8.2	(b)	37
3.8.3	(c)	37
3.9	Exercise 9	37
3.10	Exercise 10	37
3.11	Exercise 11	37
3.12	Exercise 12	37
3.13	Exercise 13	37
3.13.1	(a)	37
3.13.2	(b)	38
3.14	Exercise 14	38
3.14.1	(a)	38
3.14.2	(b)	38
3.15	Exercise 15	38
3.16	Exercise 16	38
3.17	Exercise 17	38
3.18	Exercise 18	38
3.19	Exercise 19	38
3.20	Exercise 20	38
3.21	Exercise 21	38
3.22	Exercise 22	38
3.23	Exercise 23	39
3.24	Exercise 24	39
3.25	Exercise 25	39
3.26	Exercise 26	39

3.27	Exercise 27	39
3.28	Exercise 28	39
3.29	Exercise 29	39
3.30	Exercise 30	39
3.31	Exercise 31	39
3.32	Exercise 32	39
3.33	Exercise 33	39
3.33.1	(a)	39
3.33.2	(b)	40
3.34	Exercise 34	40
3.34.1	(a)	40
3.34.2	(b)	40
3.35	Exercise 35	40
3.36	Exercise 36	40
3.37	Exercise 37	40
3.38	Exercise 38	40
3.39	Exercise 39	40
4	Exercise Set 4.4	40
4.1	Exercise 1	40
4.2	Exercise 2	40
4.3	Exercise 3	40
4.4	Exercise 4	41
4.5	Exercise 5	41
4.6	Exercise 6	41
4.7	Exercise 7	41
4.8	Exercise 8	41
4.9	Exercise 9	41
4.10	Exercise 10	41
4.11	Exercise 11	41
4.12	Exercise 12	41
4.13	Exercise 13	41
4.14	Exercise 14	41
4.15	Exercise 15	41
4.16	Exercise 16	42
4.17	Exercise 17	42
4.18	Exercise 18	42
4.18.1	(a)	42
4.18.2	(b)	42
4.19	Exercise 19	42
4.20	Exercise 20	42
4.21	Exercise 21	42
4.22	Exercise 22	42
4.23	Exercise 23	42
4.24	Exercise 24	42
4.25	Exercise 25	42

4.26	Exercise 26	43
4.27	Exercise 27	43
4.28	Exercise 28	43
4.29	Exercise 29	43
4.30	Exercise 30	43
4.31	Exercise 31	43
4.32	Exercise 32	43
4.33	Exercise 33	43
4.34	Exercise 34	43
4.35	Exercise 35	43
4.36	Exercise 36	43
4.36.1	(a)	43
4.36.2	(b)	44
4.36.3	(c)	44
4.36.4	(d)	44
4.37	Exercise 37	44
4.37.1	(a)	44
4.37.2	(b)	44
4.37.3	(c)	44
4.38	Exercise 38	44
4.38.1	(a)	44
4.38.2	(b)	44
4.38.3	(c)	44
4.38.4	(d)	44
4.39	Exercise 39	44
4.39.1	(a)	44
4.39.2	(b)	45
4.40	Exercise 40	45
4.40.1	(a)	45
4.40.2	(b)	45
4.41	Exercise 41	45
4.42	Exercise 42	45
4.42.1	(a)	45
4.42.2	(b)	45
4.42.3	(c)	45
4.43	Exercise 43	45
4.44	Exercise 44	45
4.45	Exercise 45	45
4.46	Exercise 46	45
4.47	Exercise 47	46
4.48	Exercise 48	46
4.49	Exercise 49	46
4.50	Exercise 50	46
5	Exercise Set 4.5	46
5.1	Exercise 1	46

5.2	Exercise 2	46
5.3	Exercise 3	46
5.4	Exercise 4	46
5.5	Exercise 5	46
5.6	Exercise 6	46
5.7	Exercise 7	46
	5.7.1 (a)	46
	5.7.2 (b)	47
5.8	Exercise 8	47
	5.8.1 (a)	47
	5.8.2 (b)	47
5.9	Exercise 9	47
	5.9.1 (a)	47
	5.9.2 (b)	47
5.10	Exercise 10	47
	5.10.1 (a)	47
	5.10.2 (b)	47
5.11	Exercise 11	47
	5.11.1 (a)	47
	5.11.2 (b)	47
	5.11.3 (c)	47
5.12	Exercise 12	48
5.13	Exercise 13	48
5.14	Exercise 14	48
5.15	Exercise 15	48
5.16	Exercise 16	48
5.17	Exercise 17	48
5.18	Exercise 18	48
	5.18.1 (a)	48
	5.18.2 (b)	48
5.19	Exercise 19	48
5.20	Exercise 20	48
5.21	Exercise 21	48
5.22	Exercise 22	49
5.23	Exercise 23	49
5.24	Exercise 24	49
5.25	Exercise 25	49
5.26	Exercise 26	49
5.27	Exercise 27	49
5.28	Exercise 28	49
	5.28.1 (a)	49
	5.28.2 (b)	49
5.29	Exercise 29	49
	5.29.1 (a)	49
	5.29.2 (b)	49
5.30	Exercise 30	49

5.30.1	(a)	49
5.30.2	(b)	50
5.31	Exercise 31	50
5.31.1	(a)	50
5.31.2	(b)	50
5.31.3	(c)	50
5.32	Exercise 32	50
5.33	Exercise 33	50
5.34	Exercise 34	50
5.35	Exercise 35	50
5.36	Exercise 36	50
5.37	Exercise 37	50
5.38	Exercise 38	50
5.39	Exercise 39	51
5.40	Exercise 40	51
5.41	Exercise 41	51
5.42	Exercise 42	51
5.43	Exercise 43	51
5.44	Exercise 44	51
5.44.1	(a)	51
5.44.2	(b)	51
5.44.3	(c)	51
5.45	Exercise 45	51
5.46	Exercise 46	51
5.47	Exercise 47	51
5.48	Exercise 48	52
5.49	Exercise 49	52
5.50	Exercise 50	52
6	Exercise Set 4.6	52
6.1	Exercise 1	52
6.2	Exercise 2	52
6.3	Exercise 3	52
6.4	Exercise 4	52
6.5	Exercise 5	52
6.6	Exercise 6	52
6.7	Exercise 7	52
6.8	Exercise 8	52
6.9	Exercise 9	53
6.10	Exercise 10	53
6.10.1	(a)	53
6.10.2	(b)	53
6.11	Exercise 11	53
6.12	Exercise 12	53
6.13	Exercise 13	53
6.14	Exercise 14	53

6.15	Exercise 15	53
6.16	Exercise 16	53
6.17	Exercise 17	53
6.18	Exercise 18	53
6.19	Exercise 19	54
6.20	Exercise 20	54
6.21	Exercise 21	54
6.22	Exercise 22	54
6.23	Exercise 23	54
6.24	Exercise 24	54
6.25	Exercise 25	54
6.26	Exercise 26	54
6.27	Exercise 27	54
6.28	Exercise 28	54
6.29	Exercise 29	54
6.30	Exercise 30	54
6.31	Exercise 31	55
6.32	Exercise 32	55
6.33	Exercise 33	55
7	Exercise Set 4.7	55
7.1	Exercise 1	55
7.2	Exercise 2	55
7.3	Exercise 3	55
7.4	Exercise 4	55
7.5	Exercise 5	55
7.6	Exercise 6	55
7.7	Exercise 7	55
7.8	Exercise 8	55
7.9	Exercise 9	56
	7.9.1 (a)	56
	7.9.2 (b)	56
7.10	Exercise 10	56
7.11	Exercise 11	56
7.12	Exercise 12	56
	7.12.1 (a)	56
	7.12.2 (b)	56
7.13	Exercise 13	56
	7.13.1 (a)	56
	7.13.2 (b)	56
7.14	Exercise 14	56
	7.14.1 (a)	56
	7.14.2 (b)	56
7.15	Exercise 15	57
7.16	Exercise 16	57
7.17	Exercise 17	57

7.18	Exercise 18	57
7.19	Exercise 19	57
7.20	Exercise 20	57
7.21	Exercise 21	57
7.21.1	(a)	57
7.21.2	(b)	57
7.22	Exercise 22	57
7.22.1	(a)	57
7.22.2	(b)	57
7.23	Exercise 23	57
7.24	Exercise 24	58
7.25	Exercise 25	58
7.26	Exercise 26	58
7.27	Exercise 27	58
7.28	Exercise 28	58
7.29	Exercise 29	58
7.30	Exercise 30	58
7.30.1	(a)	58
7.30.2	(b)	58
7.31	Exercise 31	58
7.31.1	(a)	58
7.31.2	(b)	58
7.31.3	(c)	58
7.32	Exercise 32	59
7.33	Exercise 33	59
7.34	Exercise 34	59
7.34.1	(a)	59
7.34.2	(b)	59
7.34.3	(c)	59
7.34.4	(d)	59
7.35	Exercise 35	59
7.36	Exercise 36	59
8	Exercise Set 4.8	59
8.1	Exercise 1	59
8.2	Exercise 2	59
8.3	Exercise 3	59
8.4	Exercise 4	60
8.5	Exercise 5	60
8.6	Exercise 6	60
8.7	Exercise 7	60
8.8	Exercise 8	60
8.9	Exercise 9	60
8.10	Exercise 10	60
8.11	Exercise 11	60
8.12	Exercise 12	60

8.13	Exercise 13	60
8.14	Exercise 14	60
8.15	Exercise 15	60
8.16	Exercise 16	61
8.17	Exercise 17	61
8.18	Exercise 18	61
8.18.1	(a)	61
8.18.2	(b)	61
8.19	Exercise 19	61
8.19.1	(a)	61
8.19.2	(b)	61
8.19.3	(c)	61
8.20	Exercise 20	61
8.21	Exercise 21	61
8.22	Exercise 22	61
8.23	Exercise 23	61
8.24	Exercise 24	62
8.25	Exercise 25	62
8.26	Exercise 26	62
8.27	Exercise 27	62
8.28	Exercise 28	62
8.29	Exercise 29	62
8.30	Exercise 30	62
8.30.1	(a)	62
8.30.2	(b)	62
8.31	Exercise 31	62
8.32	Exercise 32	62
8.33	Exercise 33	62
8.34	Exercise 34	63
8.34.1	(a)	63
8.34.2	(b)	63
8.35	Exercise 35	63
8.36	Exercise 36	63
8.37	Exercise 37	63
8.38	Exercise 38	63
9	Exercise Set 4.9	63
9.1	Exercise 1	63
9.2	Exercise 2	63
9.3	Exercise 3	63
9.4	Exercise 4	63
9.5	Exercise 5	63
9.6	Exercise 6	64
9.7	Exercise 7	64
9.8	Exercise 8	64
9.9	Exercise 9	64

9.10	Exercise 10	64
9.11	Exercise 11	64
9.12	Exercise 12	64
9.13	Exercise 13	64
9.14	Exercise 14	64
9.14.1	(a)	64
9.14.2	(b)	64
9.15	Exercise 15	64
9.15.1	(a)	64
9.15.2	(b)	65
9.16	Exercise 16	65
9.16.1	(a)	65
9.16.2	(b)	65
9.17	Exercise 17	65
9.18	Exercise 18	65
9.19	Exercise 19	65
9.20	Exercise 20	65
9.20.1	(a)	65
9.20.2	(b)	65
9.21	Exercise 21	65
9.21.1	(a)	65
9.21.2	(b)	65
9.21.3	(c)	65
9.22	Exercise 22	66
9.23	Exercise 23	66
9.23.1	(a)	66
9.23.2	(b)	66
9.23.3	(c)	66
9.23.4	(d)	66
9.23.5	(e)	66
9.23.6	(f)	66
9.24	Exercise 24	66
9.24.1	(a)	66
9.24.2	(b)	66
9.24.3	(c)	66
9.24.4	(d)	66
9.24.5	(e)	67
9.24.6	(f)	67
9.25	Exercise 25	67
10	Exercise Set 4.10	67
10.1	Exercise 1	67
10.2	Exercise 2	67
10.3	Exercise 3	67
10.3.1	(a)	67
10.3.2	(b)	67

10.4 Exercise 4	67
10.5 Exercise 5	67
10.6 Exercise 6	67
10.7 Exercise 7	67
10.8 Exercise 8	68
10.8.1 (a)	68
10.8.2 (b)	68
10.9 Exercise 9	68
10.10 Exercise 10	68
10.11 Exercise 11	68
10.12 Exercise 12	68
10.13 Exercise 13	68
10.14 Exercise 14	68
10.15 Exercise 15	68
10.16 Exercise 16	68
10.17 Exercise 17	68
10.18 Exercise 18	69
10.19 Exercise 19	69
10.20 Exercise 20	69
10.21 Exercise 21	69
10.22 Exercise 22	69
10.23 Exercise 23	69
10.23.1 (a)	69
10.23.2 (b)	69
10.24 Exercise 24	69
10.25 Exercise 25	69
10.25.1 (a)	69
10.25.2 (b)	69
10.26 Exercise 26	69
10.26.1 (a)	69
10.26.2 (b)	70
10.27 Exercise 27	70
10.27.1 (a)	70
10.27.2 (b)	70
10.27.3 (c)	70
10.28 Exercise 28	70
10.28.1 (a)	70
10.28.2 (b)	70
10.28.3 (c)	70
10.29 Exercise 29	70
10.30 Exercise 30	70
10.31 Exercise 31	70
10.32 Exercise 32	70

1 Exercise Set 4.1

1.1 Exercise 1

Assume that k is a particular integer.

1.1.1 (a)

Is -17 an odd integer?

Proof. Yes: $-17 = 2(-9) + 1$.

□

1.1.2 (b)

Is 0 neither even nor odd?

Proof. No. 0 is even because $0 = 0 \cdot 2$.

□

1.1.3 (c)

Is $2k - 1$ odd?

Proof. Yes: $2k - 1 = 2(k - 1) + 1$ and $k - 1$ is an integer because it is a difference of integers.

□

1.2 Exercise 2

Assume that c is a particular integer.

1.2.1 (a)

Is $-6c$ an even integer?

Proof. Yes, because $-6c = 2 \cdot (-3c) = 2k$ where $k = -3c$ is an integer.

□

1.2.2 (b)

Is $8c + 5$ an odd integer?

Proof. Yes, because $8c + 5 = 2(4c + 2) + 1$ and $k = 4c + 2$ is an integer.

□

1.2.3 (c)

Is $(c^2 + 1) - (c^2 - 1) - 2$ an even integer?

Proof. Yes, because it equals 0 : $(c^2 + 1) - (c^2 - 1) - 2 = c^2 + 1 - c^2 + 1 - 2 = 2 - 2 = 0$.

□

1.3 Exercise 3

Assume that m and n are particular integers.

1.3.1 (a)

Is $6m + 8n$ even?

Proof. Yes: $6m + 8n = 2(3m + 4n)$ and $(3m + 4n)$ is an integer because $3, 4, m$, and n are integers, and products and sums of integers are integers. \square

1.3.2 (b)

Is $10mn + 7$ odd?

Proof. Yes: $10mn + 7 = 2(5mn + 3) + 1$ and $5mn + 3$ is an integer because $3, 5, m$, and n are integers, and products and sums of integers are integers. \square

1.3.3 (c)

If $m > n > 0$, is $m^2 - n^2$ composite?

Proof. Not necessarily. For instance, if $m = 3$ and $n = 2$, then $m^2 - n^2 = 9 - 4 = 5$, which is prime. (However, $m^2 - n^2$ is composite for many values of m and n because of the identity $m^2 - n^2 = (m - n)(m + n)$.) \square

1.4 Exercise 4

Assume that r and s are particular integers.

1.4.1 (a)

Is $4rs$ even?

Proof. Yes: $4rs = 2(2rs)$ and $2rs$ is an integer because $2, r, s$ are integers, and products of integers are integers. \square

1.4.2 (b)

Is $6r + 4s^2 + 3$ odd?

Proof. Yes: $6r + 4s^2 + 3 = 2(3r + 2s^2 + 1) + 1$ and $3r + 2s^2 + 1$ is an integer because $3, r, 2, s, 1$ are integers and products and sums of integers are integers. \square

1.4.3 (c)

If r and s are both positive, is $r^2 + 2rs + s^2$ composite?

Proof. Yes: $r^2 + 2rs + s^2 = (r + s)(r + s)$ and $r + s \geq 2$, therefore $r^2 + 2rs + s^2$ is a product of two integers both of which are greater than 1. \square

Prove the statements in 5–11.

1.5 Exercise 5

There are integers m and n such that $m > 1$ and $n > 1$ and $\frac{1}{m} + \frac{1}{n}$ is an integer.

Proof. For example, let $m = n = 2$. Then m and n are integers such that $m > 1$ and $n > 1$ and $\frac{1}{m} + \frac{1}{n} = \frac{1}{2} + \frac{1}{2} = 1$ which is an integer. \square

1.6 Exercise 6

There are distinct integers m and n such that $\frac{1}{m} + \frac{1}{n}$ is an integer.

Proof. For example, let $m = 1, n = -1$. Then m and n are integers such that $\frac{1}{m} + \frac{1}{n} = \frac{1}{1} - \frac{1}{1} = 0$ which is an integer. \square

1.7 Exercise 7

There are real numbers a and b such that $\sqrt{a+b} = \sqrt{a}\sqrt{b}$.

Proof. For example, let $a = 0, b = 0$. Then a and b are real numbers such that $\sqrt{a+b} = \sqrt{0+0} = 0 = \sqrt{0} + \sqrt{0} = \sqrt{a} + \sqrt{b}$. \square

1.8 Exercise 8

There is an integer $n > 5$ such that $2^n - 1$ is prime.

Proof. For example, let $n = 7$. Then n is an integer such that $n > 5$ and $2^n - 1 = 127$, which is prime. \square

1.9 Exercise 9

There is a real number x such that $x > 1$ and $2^x > x^{10}$.

Proof. For example, take $x = 80$. Then

$$x^{10} = 80^{10} = 8^{10} \cdot 10^{10} = (2^3)^{10} \cdot 10^{10} = 2^{30} \cdot 10^{10}.$$

We have $2^{50} \approx 1,125899907 \cdot 10^{15} > 10^{10}$. So $2^{80} = 2^{30} \cdot 2^{50} > 2^{30} \cdot 10^{10} = 80^{10}$.

Therefore $x = 80$ is a real number such that $x > 1$ and $2^x > x^{10}$. \square

Definition: An integer n is called a **perfect square** if, and only if, $n = k^2$ for some integer k .

1.10 Exercise 10

There is a perfect square that can be written as the sum of two other perfect squares.

Proof. For example, 25, 9, and 16 are all perfect squares, because $25 = 5^2$, $9 = 3^2$, and $16 = 4^2$, and $25 = 9 + 16$. Thus 25 is a perfect square that can be written as a sum of two other perfect squares. \square

1.11 Exercise 11

There is an integer n such that $2n^2 - 5n + 2$ is prime.

Proof. For example, take $n = 3$. Then $2n^2 - 5n + 2 = 18 - 15 + 2 = 5$ is prime. (You can find this value of n by either starting at $n = 1$ and using trial and error, or noticing that $2n^2 - 5n + 2 = (2n - 1)(n - 2)$, so, for this to be prime, one of the factors has to be 1.) \square

In 12 – 13, (a) write a negation for the given statement, and (b) use a counterexample to disprove the given statement. explain how the counterexample actually shows that the given statement is false.

1.12 Exercise 12

For all real numbers a and b , if $a < b$ then $a^2 < b^2$.

Proof. a. *Negation for the statement:* There exist real numbers a and b such that $a < b$ and $a^2 \not< b^2$.

b. *Counterexample for the statement:* Let $a = -2$ and $b = -1$. Then $a < b$ because $-2 < -1$, but $a^2 \not< b^2$ because $(-2)^2 = 4$ and $(-1)^2 = 1$ and $4 \not< 1$. [So the hypothesis of the statement is true and its conclusion is false.] \square

1.13 Exercise 13

For every integer n , if n is odd then $\frac{n-1}{2}$ is odd.

Proof. a. *Negation for the statement:* There exists an integer n such that n is odd and $\frac{n-1}{2}$ is not odd.

b. *Counterexample for the statement:* Let $n = 5$. Then n is odd because $5 = 2 \cdot 2 + 1$, but $\frac{n-1}{2}$ is not odd because $\frac{5-1}{2} = 2 = 2 \cdot 1$ is even. [So the hypothesis of the statement is true and its conclusion is false.] \square

Disprove each of the statements in 14 – 16 by giving a counterexample. In each case explain how the counterexample actually disproves the statement.

1.14 Exercise 14

For all integers m and n , if $2m + n$ is odd then m and n are both odd.

Proof. Counterexample: Let $m = 2$ and $n = 1$. Then $2m + n = 2 \cdot 2 + 1 = 5$, which is odd. But m is not odd, and so it is false that both m and n are odd. [This is one counterexample among many.] \square

1.15 Exercise 15

For every integer p , if p is prime then $p^2 - 1$ is even.

Proof. Counterexample: Let $p = 2$ which is prime, but $p^2 - 1 = 2^2 - 1 = 4 - 1 = 3$ is not even. [This is the only counterexample! For every other prime p , $p^2 - 1$ is even.] \square

1.16 Exercise 16

For every integer n , if n is even then $n^2 + 1$ is prime.

Proof. Counterexample: Let $n = 8$. Then n is even. But $n^2 + 1 = 65 = 13 \cdot 5$ is not prime. [This is one counterexample among many.] \square

In 17 – 20, determine whether the property is true for all integers, true for no integers, or true for some integers and false for other integers. Justify your answers.

1.17 Exercise 17

$$(a + b)^2 = a^2 + b^2$$

Proof. This property is true for some integers and false for other integers. For instance, if $a = 0$ and $b = 1$, the property is true because $(0 + 1)^2 = 0^2 + 1^2$, but if $a = 1$ and $b = 1$, the property is false because $(1 + 1)^2 = 4$ and $1^2 + 1^2 = 2$ and $4 \neq 2$. \square

1.18 Exercise 18

$$\frac{a}{b} + \frac{c}{d} = \frac{a + c}{b + d}$$

Proof. True for some integers, false for others. For example, if $a = c = 0$ and $b = d = 1$ then $\frac{0}{1} + \frac{0}{1} = 0 = \frac{0 + 0}{1 + 1}$ is true. But if $a = 1, b = 2, c = 3$ and $d = 4$ then

$$\frac{a}{b} + \frac{c}{d} = \frac{1}{2} + \frac{3}{4} = \frac{5}{4} \neq \frac{4}{6} = \frac{1 + 3}{2 + 4} = \frac{a + c}{b + d}.$$

\square

1.19 Exercise 19

$$-a^n = (-a)^n$$

Hint: This property is true for some integers and false for other integers. To justify this answer you need to find examples of both.

Proof. True for some integers: let $a = 0, n = 1$. Then $-0^1 = -0 = 0$ and $(-0)^1 = 0^1 = 0$ so the equality holds. When $a = n = 2$ it is false: $-2^2 = -4$ but $(-2)^2 = 4$ and $-4 \neq 4$. \square

1.20 Exercise 20

The average of any two odd integers is odd.

Proof. True for some, false for others. For example, 3 and 7 are both odd, and their average $(3+7)/2 = 5$ is odd. But 3 and 5 are both odd, and their average is $(3+5)/2 = 4$ is even. \square

Prove the statement in 21 and 22 by the method of exhaustion.

1.21 Exercise 21

Every positive even integer less than 26 can be expressed as a sum of three or fewer perfect squares. (For instance, $10 = 1^2 + 3^2$ and $16 = 4^2$.)

Proof. $2 = 1^2 + 1^2$

$$4 = 2^2$$

$$6 = 2^2 + 1^2 + 1^2$$

$$8 = 2^2 + 2^2$$

$$10 = 1^2 + 3^2$$

$$12 = 2^2 + 2^2 + 2^2$$

$$16 = 4^2$$

$$18 = 4^2 + 1^2 + 1^2$$

$$20 = 4^2 + 2^2$$

$$22 = 2^2 + 2^2 + 3^2$$

$$24 = 4^2 + 2^2 + 2^2$$

\square

1.22 Exercise 22

For each integer n with $1 \leq n \leq 10$, $n^2 - n + 11$ is a prime number.

Proof. $1^2 - 1 + 11 = 11$ is prime.

$2^2 - 2 + 11 = 13$ is prime.

$3^2 - 3 + 11 = 17$ is prime.

$4^2 - 4 + 11 = 23$ is prime.

$5^2 - 5 + 11 = 31$ is prime.

$6^2 - 6 + 11 = 41$ is prime.

$7^2 - 7 + 11 = 53$ is prime.

$8^2 - 8 + 11 = 67$ is prime.

$9^2 - 9 + 11 = 83$ is prime.

$10^2 - 10 + 11 = 101$ is prime. □

Each of the statements in 23 – 26 is true. For each, (a) rewrite the statement with the quantification implicit as If _____, then _____, and (b) write the first sentence of a proof (the “starting point”) and the last sentence of a proof (the “conclusion to be shown”). (Note that you do not need to understand the statements in order to be able to do these exercises.)

1.23 Exercise 23

For every integer m , if $m > 1$ then $0 < \frac{1}{m} < 1$.

Proof. a. If an integer is greater than 1, then its reciprocal is between 0 and 1.

b. *Start of proof:* Suppose m is any integer such that $m > 1$. *Conclusion to be shown:* $0 < 1/m < 1$. □

1.24 Exercise 24

For every real number x , if $x > 1$ then $x^2 > x$.

Proof. a. If a real number is greater than 1, then its square is greater than itself.

b. *Start of proof:* Suppose x is any real number such that $x > 1$. *Conclusion to be shown:* $x^2 > x$. □

1.25 Exercise 25

For all integers m and n , if $mn = 1$ then $m = n = 1$ or $m = n = -1$.

Proof. a. If the product of two integers is 1, then either both are 1 or both are -1 .

b. *Start of proof:* Suppose m and n are any integers with $mn = 1$. *Conclusion to be shown:* $m = n = 1$ or $m = n = -1$. □

1.26 Exercise 26

For every real number x , if $0 < x < 1$ then $x^2 < x$.

Proof. a. If a real number is strictly between 0 and 1, then its square is less than itself.

b. *Start of proof:* Suppose x is any real number such that $0 < x < 1$. *Conclusion to be shown:* $x^2 < x$. \square

1.27 Exercise 27

Fill in the blanks in the following proof.

Theorem: For every odd integer n , n^2 is odd.

Proof: Suppose n is any (a) _____. By definition of odd, $n = 2k + 1$ for some integer k . Then

$$\begin{aligned} n^2 &= \text{(b) } (\text{_____})^2 && \text{by substitution} \\ &= 4k^2 + 4k + 1 && \text{by multiplying} \\ &= 2(2k^2 + 2k) + 1 && \text{by factoring out a 2} \end{aligned}$$

Now $2k^2 + 2k$ is an integer because it is a sum of products of integers. Therefore, n^2 equals $2 \cdot (\text{an integer}) + 1$, and so (c) _____ is odd by definition of odd.

Because we have not assumed anything about n except that it is an odd integer, it follows from the principle of (d) _____ that for every odd integer n , n^2 is odd.

Proof. (a) particular but arbitrarily chosen odd integer (b) $2k + 1$ (c) n^2 (d) universal generalization \square

In each of 28 – 31:

a. Rewrite the theorem in three different ways: as \forall _____, if _____, then _____; as \forall _____, _____ (without using the words if or then), and as If _____, then _____ (without using an explicit universal quantifier).

b. Fill in the blanks in the proof of the theorem.

1.28 Exercise 28

Theorem: The sum of any two odd integers is even.

Proof: Suppose m and n are any [particular but arbitrarily chosen] odd integers.

[We must show that $m + n$ is even.]

By (a) _____, $m = 2r + 1$ and $n = 2s + 1$ for some integers r and s . Then

$$\begin{aligned} m + n &= (2r + 1) + (2s + 1) && \text{by (b) _____} \\ &= 2r + 2s + 2 \\ &= 2(r + s + 1) && \text{by algebra} \end{aligned}$$

Let $u = r + s + 1$. Then u is an integer because r, s and 1 are integers and because (c) ____ .

Hence $m + n = 2u$, where u is an integer, and so, by (d) ____ , $m + n$ is even [as was to be shown].

Proof. a. \forall integers m and n , if m and n are odd then $m + n$ is even.

\forall odd integers m and n , $m + n$ is even.

If m and n are any odd integers, then $m + n$ is even.

b. (a) definition of odd, (b) substitution, (c) any sum of integers is an integer, (d) definition of even \square

1.29 Exercise 29

Theorem: The negative of any even integer is even.

Proof: Suppose n is any [particular but arbitrarily chosen] even integer.

[We must show that $-n$ is even.]

By (a) ____ , $n = 2k$ for some integer k .

Then

$$\begin{aligned} -n &= -(2k) \text{ by (b) } ______ \\ &= 2(-k) \text{ by algebra} \end{aligned}$$

Let $r = -k$. Then r is an integer because -1 and k are integers and (c) ____ .

Hence $-n = 2r$, where r is an integer, and so $-n$ is even by (d) ____ [as was to be shown].

Proof. a. \forall integer n , if n is even then $-n$ is even.

\forall even integer n , $-n$ is even.

If n is any even integer, then $-n$ is even.

b. (a) definition of even, (b) substitution, (c) any product of integers is an integer, (d) definition of even \square

1.30 Exercise 30

Theorem: The sum of any even integer and any odd integer is odd.

Proof: Suppose m is any even integer and n is any (a) ____ . By definition of even, $m = 2r$ for some (b) ____ , and by definition of odd, $n = 2s + 1$ for some integer s . By substitution and algebra,

$$m + n = \text{(c) } ______ = 2(r + s) + 1$$

Since r and s are integers, so is their sum $r + s$. Hence $m + n$ has the form twice some integer plus one, and so, by (d) _____ by definition of odd.

Proof. a. \forall integers m and n , if m is even and n is odd, then $m + n$ is odd.

\forall even integers m and odd integers n , $m + n$ is odd.

If m is any even integer and n is any odd integer, then $m + n$ is odd.

b. (a) any odd integer (b) integer r (c) $2r + (2s + 1)$ (d) $m + n$ is odd □

1.31 Exercise 31

Theorem: Whenever n is an odd integer, $5n^2 + 7$ is even.

Proof: Suppose n is any *[particular but arbitrarily chosen]* odd integer.

[We must show that $5n^2 + 7$ is even.]

By definition of odd, $n =$ (a) _____, for some integer k .

Then

$$\begin{aligned} 5n^2 + 7 &= \text{(b) } \underline{\hspace{2cm}} && \text{substitution} \\ &= 5(4k^2 + 4k + 1) + 7 \\ &= 20k^2 + 20k + 12 \\ &= 2(10k^2 + 10k + 6) && \text{by algebra} \end{aligned}$$

Let $t =$ (c) _____. Then t is an integer because products and sums of integers are integers.

Hence $5n^2 + 7 = 2t$, where t is an integer, and thus (d) _____ by definition of even *[as was to be shown]*.

Proof. a. \forall integer n , if n is odd, then $5n^2 + 7$ is even.

\forall odd integer n , $5n^2 + 7$ is even.

If n is any odd integer, then $5n^2 + 7$ is even.

b. (a) $2k + 1$ (b) $5(2k + 1)^2 + 7$ (c) $10k^2 + 10k + 6$ (d) $5n^2 + 7$ is even □

2 Exercise Set 4.2

Prove the statements in 1 – 11. In each case use only the definitions of the terms and the assumptions listed on page 161, not any previously established properties of odd and even integers. Follow the directions given in this section for writing proofs of universal statements.

2.1 Exercise 1

For every integer n , if n is odd then $3n + 5$ is even.

Proof. Suppose n is any [particular but arbitrarily chosen] odd integer.

[We must show that $3n + 5$ is even. By definition of even, this means we must show that $3n + 5 = 2 \cdot (\text{some integer})$.]

By definition of odd, $n = 2r + 1$, for some integer r .

Then

$$\begin{aligned} 3n + 5 &= 3(2r + 1) + 5 && \text{by substitution} \\ &= 6r + 3 + 5 \\ &= 6r + 8 \\ &= 2(3r + 4) && \text{by algebra} \end{aligned}$$

[Idea for the rest of the proof: We want to show that $3n + 5 = 2 \cdot (\text{some integer})$. At this point we know that $3n + 5 = 2(3r + 4)$. So is $3r + 4$ an integer? Yes, because products and sums of integers are integers.]

Let $k = 3r + 4$.

Then k is an integer because products and sums of integers are integers.

Hence $3n + 5 = 2(3r + 4) = 2k$ where k is an integer. Hence by definition of even $3n + 5$ is even [as was to be shown]. \square

2.2 Exercise 2

For every integer m , if m is even then $3m + 5$ is odd.

Proof. Suppose m is any [particular but arbitrarily chosen] even integer.

[We must show that $3m + 5$ is odd. By definition of odd, this means we must show that $3m + 5 = 2 \cdot (\text{some integer}) + 1$.]

By definition of even, $m = 2r$, for some integer r .

Then

$$\begin{aligned} 3m + 5 &= 3(2r) + 5 && \text{by substitution} \\ &= 6r + 5 \\ &= 6r + 4 + 1 \\ &= 2(3r + 2) + 1 && \text{by algebra} \end{aligned}$$

[Idea for the rest of the proof: We want to show that $3m + 5 = 2 \cdot (\text{some integer}) + 1$. At this point we know that $3m + 5 = 2(3r + 2) + 1$. So is $3r + 2$ an integer? Yes, because products and sums of integers are integers.]

Let $k = 3r + 2$.

Then k is an integer because products and sums of integers are integers.

Hence $3m + 5 = 2(3r + 2) + 1 = 2k + 1$ where k is an integer. Hence by definition of odd $3n + 5$ is odd [as was to be shown]. \square

2.3 Exercise 3

For every integer n , $2n - 1$ is odd.

Proof. Suppose n is any [particular but arbitrarily chosen] integer.

[We must show that $2n - 1$ is odd. By definition of odd, this means we must show that $2n - 1 = 2 \cdot (\text{some integer}) + 1$.]

Then

$$\begin{aligned} 2n - 1 &= 2n - 2 + 2 - 1 && \text{because } -2 + 2 = 0 \\ &= 2(n - 1) + 2 - 1 \\ &= 2(n - 1) + 1 && \text{by algebra} \end{aligned}$$

Let $k = n - 1$.

Then k is an integer because the difference of two integers (n and 1) is an integer.

Hence $2n - 1 = 2(n - 1) + 1 = 2k + 1$ where k is an integer, and thus by definition of odd $2n - 1$ is odd [as was to be shown]. \square

2.4 Exercise 4

The difference of any even integer minus any odd integer is odd.

Proof. Suppose a is any even integer and b is any odd integer. [We must show that $a - b$ is odd.] By definition of even and odd, $a = 2r$ and $b = 2s + 1$, for some integers r, s . By substitution and algebra,

$$a - b = 2r - (2s + 1) = 2r - 2s - 1 = 2r - 2s - 2 + 2 - 1 = 2(r - s - 1) + 1$$

Let $t = r - s - 1$. Then t is an integer because differences of integers are integers.

Thus $a - b = 2t + 1$ where t is an integer, and so by definition of odd $a - b$ is odd [as was to be shown]. \square

2.5 Exercise 5

If a and b are any odd integers, then $a^2 + b^2$ is even.

Proof. Suppose a, b are any [particular but arbitrarily chosen] odd integers.

[We must show that $a^2 + b^2$ is even.]

By definition of odd, $a = 2r + 1$ and $b = 2s + 1$, for some integers r, s .

Then

$$\begin{aligned}
a^2 + b^2 &= (2r + 1)^2 + (2s + 1)^2 && \text{by substitution} \\
&= (4r^2 + 4r + 1) + (4s^2 + 4s + 1) && \text{by multiplying} \\
&= 4r^2 + 4r + 4s^2 + 4s + 2 && \text{by adding} \\
&= 2(2r^2 + 2r + 2s^2 + 2s + 1) && \text{by factoring out}
\end{aligned}$$

Let $k = 2r^2 + 2r + 2s^2 + 2s + 1$.

Then k is an integer because squares, products and sums of integers are integers.

Hence $a^2 + b^2 = 2k$ where k is an integer, and thus by definition of even $a^2 + b^2$ is even [as was to be shown]. \square

2.6 Exercise 6

If k is any odd integer and m is any even integer, then $k^2 + m^2$ is odd.

Proof. Suppose k is any odd integer and m is any even integer.

[We must show that $k^2 + m^2$ is odd.]

By definition of odd and even, $k = 2a + 1$ and $m = 2b$, for some integers a, b . Then

$$\begin{aligned}
k^2 + m^2 &= (2a + 1)^2 + (2b)^2 && \text{by substitution} \\
&= 4a^2 + 4a + 1 + 4b^2 \\
&= 4(a^2 + a + b^2) + 1 \\
&= 2(2a^2 + 2a + 2b^2) + 1 && \text{by algebra}
\end{aligned}$$

But $2a^2 + 2a + 2b^2$ is an integer because it is a sum of products of integers. Thus $k^2 + m^2$ is twice an integer plus 1, and so $k^2 + m^2$ is odd [as was to be shown]. \square

2.7 Exercise 7

The difference between the squares of any two consecutive integers is odd.

Proof. Suppose m and n are any [particular but arbitrarily chosen] two consecutive integers.

[We must show that $m^2 - n^2$ (or $n^2 - m^2$) is odd.]

By definition of consecutive, $m = k$ and $n = k + 1$, for some integer k .

Then

$$\begin{aligned}
m^2 - n^2 &= k^2 - (k + 1)^2 && \text{by substitution} \\
&= k^2 - (k^2 + 2k + 1) \\
&= k^2 - k^2 - 2k - 1 \\
&= -2k - 1 \\
&= -2k - 2 + 2 - 1 \\
&= 2(-k - 1) + 1 && \text{by algebra}
\end{aligned}$$

Let $r = -k - 1$. Then r is an integer because it is a difference of integers.

Hence $m^2 - n^2 = 2r + 1$ where r is an integer, and thus by definition of odd $m^2 - n^2$ is odd [as was to be shown]. \square

2.8 Exercise 8

For any integers m and n , if m is even and n is odd then $5m + 3n$ is odd.

Proof. Suppose m is any even integer and n is any odd integer.

[We must show that $5m + 3n$ is odd.]

By definition of even and odd, $m = 2r$ and $n = 2s + 1$, for some integers r, s .

Then

$$\begin{aligned} 5m + 3n &= 5(2r) + 3(2s + 1) && \text{by substitution} \\ &= 10r + 6s + 3 \\ &= 10r + 6s + 2 + 1 \\ &= 2(5r + 3s + 1) + 1 && \text{by algebra} \end{aligned}$$

Let $k = 5r + 3s + 1$.

Then k is an integer because it is a sum of products of integers.

Hence $5m + 3n = 2k + 1$ where k is an integer, and thus by definition of odd $5m + 3n$ is odd [as was to be shown]. \square

2.9 Exercise 9

If an integer greater than 4 is a perfect square, then the immediately preceding integer is not prime.

Proof. Suppose n is any integer greater than 4 that is a perfect square.

[We must show that $n - 1$ is not prime, in other words, $n - 1$ is composite.]

By definition of perfect square, $n = k^2$, for some integer k .

Without loss of generality, we may assume $k > 0$, because $n = k^2 = (-k)^2$, and if k is negative, we can replace it with $-k$ which is positive.

Since $n > 4$ we have $n - 4 > 0$. So $k^2 - 4 > 0$. So $(k - 2)(k + 2) > 0$. So either $k - 2$ and $k + 2$ are both negative, or they are both positive. Since $k > 0$, $k + 2 > 2 > 0$, so they have to be both positive. Therefore, $k - 2 > 0$ so $k > 2$.

Then

$$\begin{aligned} n - 1 &= k^2 - 1 && \text{by substitution} \\ &= (k - 1)(k + 1) && \text{by algebra} \end{aligned}$$

Since $k > 2$ we have both $k - 1 > 1$ and $k + 1 > 3 > 1$.

Hence $n - 1$ is a product of two positive integers both greater than 1, and thus by definition of composite $n - 1$ is composite [as was to be shown]. \square

2.10 Exercise 10

If n is any even integer, then $(-1)^n = 1$.

Proof. Suppose n is any even integer. [We must show that $(-1)^n = 1$.]

By definition of even, $n = 2k$, for some integer k .

Then by the laws of exponents from algebra $(-1)^n = (-1)^{2k} = ((-1)^2)^k = 1^k = 1$, [as was to be shown]. \square

2.11 Exercise 11

If n is any odd integer, then $(-1)^n = -1$.

Proof. Suppose n is any odd integer. [We must show that $(-1)^n = -1$.]

By definition of odd, $n = 2k + 1$, for some integer k .

Then by the laws of exponents from algebra

$$(-1)^n = (-1)^{2k+1} = (-1)^{2k} \cdot (-1) = ((-1)^2)^k \cdot (-1) = 1^k \cdot (-1) = 1 \cdot (-1) = -1$$

[as was to be shown]. \square

Prove that the statements in 12 – 14 are false.

2.12 Exercise 12

There exists an integer $m \geq 3$ such that $m^2 - 1$ is prime.

Proof. To prove the given statement is false, we prove that its negation is true.

The negation of the statement is “For every integer $m \geq 3$, $m^2 - 1$ is not prime.”

Proof of the negation: Suppose m is any integer with $m \geq 3$.

By basic algebra, $m^2 - 1 = (m - 1)(m + 1)$.

Because $m \geq 3$, both $m - 1$ and $m + 1$ are positive integers greater than 1, and each is smaller than $m^2 - 1$.

So $m^2 - 1$ is a product of two smaller positive integers, each greater than 1, and hence $m^2 - 1$ is not prime. \square

2.13 Exercise 13

There exists an integer n such that $6n + 27$ is prime.

Proof. To prove the given statement is false, we prove that its negation is true.

The negation of the statement is “For every integer n , $6n + 27$ is not prime. In other words, $6n + 27$ is composite.”

Proof of the negation: Suppose n is any integer. By basic algebra, $6n + 27 = 3(2n + 9)$. Hence $6n + 27$ is the product of two integers greater than 1. Therefore by definition of composite, $6n + 27$ is composite. \square

2.14 Exercise 14

There exists an integer $k \geq 4$ such that $2k^2 - 5k + 2$ is prime.

Proof. To prove the given statement is false, we prove that its negation is true.

The negation of the statement is “For every integer $k \geq 4$, $2k^2 - 5k + 2$ is composite.”

Proof of the negation: Suppose k is any integer.

By basic algebra, $2k^2 - 5k + 2 = (2k - 1)(k - 2)$.

Because $k \geq 4$, $2k - 1 \geq 7$ and $k - 2 \geq 2$. So both $2k - 1$ and $k - 2$ are integers greater than 1.

Hence $2k^2 - 5k + 2$ is the product of two integers greater than 1. Therefore by definition of composite, $2k^2 - 5k + 2$ is composite. \square

Find the mistakes in the “proofs” shown in 15 – 19.

2.15 Exercise 15

Theorem: For every integer k , if $k > 0$ then $k^2 + 2k + 1$ is composite.

“Proof: For $k = 2$, $k > 0$ and $k^2 + 2k + 1 = 2^2 + 2 \cdot 2 + 1 = 9$. And since $9 = 3 \cdot 3$, then 9 is composite. Hence the theorem is true.”

Proof. The incorrect proof just shows the theorem to be true in the one case where $k = 2$. A real proof must show that it is true for *every* integer $k > 0$. \square

2.16 Exercise 16

Theorem: The difference between any odd integer and any even integer is odd.

“Proof: Suppose n is any odd integer, and m is any even integer. By definition of odd, $n = 2k + 1$ where k is an integer, and by definition of even, $m = 2k$ where k is an integer. Then $n - m = (2k + 1) - 2k = 1$, and 1 is odd. Therefore, the difference between any odd integer and any even integer is odd.”

Proof. The mistake in the “proof” is that the same symbol, k , is used to represent two different quantities. By setting $m = 2k$ and $n = 2k + 1$, the proof implies that $n = m + 1$, and thus it deduces the conclusion only for this one situation. When $m = 4$ and $n = 17$, for instance, the computations in the proof indicate that $n - m = 1$, but actually $n - m = 13$. In other words, the proof does not deduce the conclusion for an arbitrarily chosen even integer m and odd integer n , and hence it is invalid. \square

2.17 Exercise 17

Theorem: For every integer k , if $k > 0$ then $k^2 + 2k + 1$ is composite.

Proof: Suppose k is any integer such that $k > 0$. If $k^2 + 2k + 1$ is composite, then $k^2 + 2k + 1 = rs$ for some integers r and s such that

$$1 < r < k^2 + 2k + 1 \text{ and } 1 < s < k^2 + 2k + 1.$$

Since $k^2 + 2k + 1 = rs$ and both r and s are strictly between 1 and $k^2 + 2k + 1$, then $k^2 + 2k + 1$ is not prime. So $k^2 + 2k + 1$ is composite as was to be shown.

Proof. This incorrect proof assumes what is to be proved. The word since in the third sentence is completely unjustified. The second sentence tells only what happens if $k^2 + 2k + 1$ is composite. But at that point in the proof, it has not been established that $k^2 + 2k + 1$ is composite. In fact, that is exactly what is to be proved. \square

2.18 Exercise 18

Theorem: The product of any even integer and any odd integer is even.

Proof: Suppose m is any even integer and n is any odd integer. If $m \cdot n$ is even, then by definition of even there exists an integer r such that $m \cdot n = 2r$.

Also since m is even, there exists an integer p such that $m = 2p$, and since n is odd there exists an integer q such that $n = 2q + 1$.

Thus $mn = (2p)(2q + 1) = 2r$, where r is an integer. By definition of even, then, $m \cdot n$ is even, as was to be shown.”

Proof. The issue is just like in Exercise 17. The proof uses the r value without establishing the existence of r first.

“If $m \cdot n$ is even...” has an unjustified assumption because we haven’t proved that $m \cdot n$ is even yet (that’s what we are *trying to prove*), so its conclusion “... $m \cdot n = 2r$ ” has not been proven.

Therefore the part “ $mn = (2p)(2q + 1) = 2r$, where r is an integer” is unjustified as well.

Discussion:

This is a fairly common form of circular reasoning: assuming what we have to prove. It happens because, at the beginning of the proof, we want to mention to the reader what we want to prove.

What we want to prove has a short, condensed definition (in this case “being even”), so we write out the full definition of what it is that we are *trying to prove* (in this case “the existence of an integer r such that $\dots = 2r$ ”). Again, the purpose of this is to articulate to the reader what we are *trying to prove*.

But then we forget that and continue as if that was already an established fact. The act of writing out the full definition of what we are trying to prove is not the same as actually having proved it.

Using “if $m \cdot n$ is even...” in this case is the problem; it has the *feeling* of using modus ponens on an already established implication with an established premise. But we are just writing out the full definition, instead of using modus ponens.

So it would be better to write: “*We want to prove that $m \cdot n$ is even. In other words, we want to prove that there is an integer r such that $m \cdot n = 2r$.*” Using the words “We want to prove that...” instead of “If...” goes a long way to avoid this common mistake. This way we can “unpack” the definition of what we are trying to prove without assuming it.

Another related problem is to first unpack the definition of what we are trying to prove, then try to “prove backwards”. Say we want to prove A , and we unpack the definition to B . So we have to prove B . But instead, we start by assuming B is true, and apply some algebra or logic to it, to arrive at something else, say E , that is true:

$$A \rightarrow \text{unpack definition} \rightarrow B \rightarrow \text{middle steps} \rightarrow C \rightarrow D \rightarrow E = \text{something true!}$$

But this would only prove that B implies E . In order to establish the truth of B (and hence of A), we would actually have to prove that E implies B ! So all the “steps” from B to E would have to be “reversible”, in other words, logical equivalences (biconditionals):

$$A \leftrightarrow \text{unpack definition} \leftrightarrow B \leftrightarrow \text{middle steps} \leftrightarrow C \leftrightarrow D \leftrightarrow E = \text{something true!}$$

But that is rarely the case! □

2.19 Exercise 19

Theorem: The sum of any two even integers equals $4k$ for some integer k .

“Proof: Suppose m and n are any two even integers. By definition of even, $m = 2k$ for some integer k and $n = 2k$ for some integer k . By substitution,

$$m + n = 2k + 2k = 4k.$$

This is what was to be shown.”

Proof. The problem here is the same as in Exercise 16. The mistake in the “proof” is that the same symbol, k , is used to represent two different quantities. By setting $m = 2k$

and $n = 2k$, the proof implies that $n = m$, and thus it deduces the conclusion only for this one situation.

When $m = 4$ and $n = 20$, for instance, the proof indicates that $n = m = 4$, but actually $n = 20$. In other words, the proof does not deduce the conclusion for an arbitrarily chosen even integer m and an arbitrarily chosen even integer n , and hence it is invalid. \square

In 20 – 38 determine whether the statement is true or false. Justify your answer with a proof or a counterexample, as appropriate. In each case use only the definitions of the terms and the assumptions listed on page 161, not any previously established properties.

2.20 Exercise 20

The product of any two odd integers is odd.

Proof. True. Suppose m and n are any odd integers. [We must show that mn is odd.] By definition of odd, $n = 2r + 1$ and $m = 2s + 1$ for some integers r and s .

Then

$$\begin{aligned} mn &= (2r + 1)(2s + 1) && \text{by substitution} \\ &= 4rs + 2r + 2s + 1 \\ &= 2(2rs + r + s) + 1 && \text{by algebra} \end{aligned}$$

Now $2rs + r + s$ is an integer because products and sums of integers are integers and 2 , r , and s are all integers. Hence $mn = 2 \cdot (\text{some integer}) + 1$, and so, by definition of odd, mn is odd. \square

2.21 Exercise 21

The negative of any odd integer is odd.

Proof. True. Assume n is any odd integer. We want to prove $-n$ is odd.

By definition of odd, $n = 2r + 1$ for some integer r .

Then $-n = -(2r + 1) = -2r - 1 = -2r - 2 + 2 - 1 = 2(-r - 1) + 1$.

Let $k = -r - 1$. Then k is an integer because it is the difference of two integers.

Therefore $-n = 2k + 1$ where k is an integer, hence by definition of odd, $-n$ is odd. \square

2.22 Exercise 22

For all integers a and b , $4a + 5b + 3$ is even.

Proof. False. Counterexample: Let $a = 1$ and $b = 0$.

Then $4a + 5b + 3 = 4 \cdot 1 + 5 \cdot 0 + 3 = 7$, which is odd.

[This is one counterexample among many. Can you find a way to characterize all counterexamples?] \square

2.23 Exercise 23

The product of any even integer and any integer is even.

Proof. True. Suppose m is any even integer and n is any integer. *[We want to prove $m \cdot n$ is even.]*

By definition of even, $m = 2k$ for some integer k .

Then, $m \cdot n = (2k) \cdot n = 2kn = 2(kn)$.

Let $r = kn$. Then r is an integer because it is the product of two integers.

Therefore $m \cdot n = 2r$ where r is an integer. So by definition of even, $m \cdot n$ is even. \square

2.24 Exercise 24

If a sum of two integers is even, then one of the summands is even. (In the expression $a + b$, a and b are called **summands**.)

Proof. False. Counterexample: Let $m = 1$ and $n = 3$.

Then $m + n = 4$ is even, but neither summand m nor summand n is even. \square

2.25 Exercise 25

The difference of any two even integers is even.

Proof. True. Assume m and n are any two even integers. *[We want to prove $m - n$ is even.]*

By definition of even, $m = 2r$ and $n = 2s$ for some integers r, s .

Then $m - n = 2r - 2s = 2(r - s)$. Let $k = r - s$. Then k is an integer because it is the difference of two integers.

Therefore $m - n = 2k$ where k is an integer. So $m - n$ is even by definition of even. \square

2.26 Exercise 26

For all integers a, b , and c , if a, b , and c are consecutive, then $a + b + c$ is even.

Proof. False. Counterexample: Let $a = 2, b = 3, c = 4$. They are consecutive integers but $a + b + c = 9$ which is not even. \square

2.27 Exercise 27

The difference of any two odd integers is even.

Proof. True. Assume m, n are any two odd integers. [We want to prove $m - n$ is even.]

By definition of odd, $m = 2r + 1, n = 2s + 1$ for some integers r, s .

Then $m - n = 2r + 1 - (2s + 1) = 2r + 1 - 2s - 1 = 2r - 2s = 2(r - s)$.

Let $k = r - s$. Then k is an integer because it is the difference of two integers.

So $m - n = 2k$ where k is an integer. Hence $m - n$ is even by definition of even. \square

2.28 Exercise 28

For all integers n and m , if $n - m$ is even then $n^3 - m^3$ is even.

Proof. True. Assume n, m are any integers such that $n - m$ is even. [Want to prove that $n^3 - m^3$ is even.]

By definition of even, $n - m = 2r$ for some integer r . By algebra,

$$n^3 - m^3 = (n - m)(n^2 + nm + m^2) = 2r(n^2 + nm + m^2) = 2(r(n^2 + nm + m^2)).$$

Let $k = r(n^2 + nm + m^2)$. Then k is an integer because it is a sum and product of integers.

So $n^3 - m^3 = 2k$ where k is an integer. So by definition of even, $n^3 - m^3$ is even. \square

2.29 Exercise 29

For every integer n , if n is prime then $(-1)^n = -1$.

Proof. False. Counterexample: Let $n = 2$.

Then n is prime, but $(-1)^n = (-1)^2 = 1 \neq -1$. \square

2.30 Exercise 30

For every integer m , if $m > 2$ then $m^2 - 4$ is composite.

Proof. False. Counterexample: Let $m = 3$. Then $m^2 - 4 = 3^2 - 4 = 9 - 4 = 5$ is prime. not composite. \square

2.31 Exercise 31

For every integer n , $n^2 - n + 11$ is a prime number.

Proof. False. Let $n = 11$. Then $n^2 - n + 11 = 11^2 - 11 + 11 = 11^2$ is not prime. \square

2.32 Exercise 32

For every integer n , $4(n^2 + n + 1) - 3n^2$ is a perfect square.

Proof. True. Suppose n is any integer. Then by algebra

$$4(n^2 + n + 1) - 3n^2 = 4n^2 + 4n + 4 - 3n^2 = n^2 + 4n + 4 = (n + 2)^2$$

Now $(n + 2)^2$ is a perfect square because $n + 2$ is an integer (being a sum of n and 2). Hence $4(n^2 + n + 1) - 3n^2$ is a perfect square, as was to be shown. \square

2.33 Exercise 33

Every positive integer can be expressed as a sum of three or fewer perfect squares.

Proof. False. Counterexample: 7 cannot be written as a sum of three or fewer perfect squares: $7 = 2^2 + 1^2 + 1^2 + 1^2$. \square

2.34 Exercise 34

(Two integers are **consecutive** if, and only if, one is one more than the other.) Any product of four consecutive integers is one less than a perfect square.

Proof. True. Suppose a, b, c, d are any four consecutive integers. [*Want to prove: there is an integer k such that $abcd = k^2 - 1$.*]

By definition of consecutive, there is an integer n such that $a = n, b = n + 1, c = n + 2, d = n + 3$. Then

$$abcd = n(n + 1)(n + 2)(n + 3) = n(n + 3)(n + 1)(n + 2) = (n^2 + 3n)(n^2 + 3n + 2)$$

[*Here we notice a pattern. The two factors differ by 2. So it is reminiscent of $(x - 1)(x + 1) = x^2 - 1^2$ isn't it?*]

By some more algebra,

$$(n^2 + 3n)(n^2 + 3n + 2) = (n^2 + 3n + 1 - 1)(n^2 + 3n + 1 + 1) = (n^2 + 3n + 1)^2 - 1^2$$

Let $k = n^2 + 3n + 1$. Then k is an integer because it is a sum and product of integers. Therefore $abcd = k^2 - 1$ where k is an integer, [*as was to be shown*]. \square

2.35 Exercise 35

If m and n are any positive integers and mn is a perfect square, then m and n are perfect squares.

Proof. False. Counterexample: let $m = n = 2$. Then $mn = 2^2$ is a perfect square. But neither m nor n is a perfect square. \square

2.36 Exercise 36

The difference of the squares of any two consecutive integers is odd.

Proof. True. Assume a, b are any two consecutive integers.

By definition of consecutive, $a = n$ and $b = n + 1$ for some integer n .

Then $b^2 - a^2 = (n + 1)^2 - n^2 = n^2 + 2n + 1 - n^2 = 2n + 1$.

So $b^2 - a^2 = 2k + 1$ where n is an integer. Therefore by definition of odd, $b^2 - a^2$ is odd. \square

2.37 Exercise 37

For all nonnegative real numbers a and b , $\sqrt{ab} = \sqrt{a}\sqrt{b}$. (Note that if x is a nonnegative real number, then there is a unique nonnegative real number y , denoted \sqrt{x} , such that $y^2 = x$.)

Proof. True. Assume a and b are any two nonnegative real numbers. By the information given to us in the parentheses:

1. There is a unique nonnegative real number denoted \sqrt{ab} such that $(\sqrt{ab})^2 = ab$.
2. There is a unique nonnegative real number denoted \sqrt{a} such that $(\sqrt{a})^2 = a$.
3. There is a unique nonnegative real number denoted \sqrt{b} such that $(\sqrt{b})^2 = b$.

Since $ab = a \cdot b$, we have by substitution: $(\sqrt{ab})^2 = (\sqrt{a})^2 \cdot (\sqrt{b})^2$.

By algebra, $(\sqrt{ab})^2 = [(\sqrt{a}) \cdot (\sqrt{b})]^2 = (\sqrt{a}\sqrt{b})^2$. Therefore $(\sqrt{ab})^2 - (\sqrt{a}\sqrt{b})^2 = 0$.

By factoring we get $(\sqrt{ab} - \sqrt{a}\sqrt{b})(\sqrt{ab} + \sqrt{a}\sqrt{b}) = 0$.

So: either $\sqrt{ab} - \sqrt{a}\sqrt{b} = 0$, or $\sqrt{ab} + \sqrt{a}\sqrt{b} = 0$.

If $\sqrt{ab} - \sqrt{a}\sqrt{b} = 0$, then $\sqrt{ab} = \sqrt{a}\sqrt{b}$ [as was to be shown.]

If $\sqrt{ab} + \sqrt{a}\sqrt{b} = 0$, then since both \sqrt{ab} and $\sqrt{a}\sqrt{b}$ are nonnegative, they must be both 0, hence $\sqrt{ab} = \sqrt{a}\sqrt{b}$ again [as was to be shown.] \square

2.38 Exercise 38

For all nonnegative real numbers a and b , $\sqrt{a+b} = \sqrt{a} + \sqrt{b}$.

Proof. False. Counterexample: Let $a = b = 1$. Then

$$\sqrt{a+b} = \sqrt{1+1} = \sqrt{2} \neq 2 = 1+1 = \sqrt{1} + \sqrt{1} = \sqrt{a} + \sqrt{b}$$

\square

2.39 Exercise 39

Suppose that integers m and n are perfect squares. Then $m + n + 2\sqrt{mn}$ is also a perfect square. Why?

Proof. Assume m and n are perfect squares (so they are nonnegative real numbers). By definition of perfect square, $m = r^2$ and $n = s^2$ for some integers r, s . Using Exercise 37 $\sqrt{mn} = \sqrt{m}\sqrt{n}$, we get:

$$m + n + 2\sqrt{mn} = r^2 + s^2 + 2\sqrt{m}\sqrt{n} = r^2 + s^2 + 2rs = (r + s)^2.$$

Let $k = r + s$. k is an integer because it is a sum of integers. So $m + n + 2\sqrt{mn} = k^2$ where k is an integer, therefore $m + n + 2\sqrt{mn}$ is a perfect square by definition. □

2.40 Exercise 40

If p is a prime number, must $2^p - 1$ also be prime? Prove or give a counterexample.

Proof. No. Counterexample: $p = 11$ is prime, but $2^p - 1 = 2^{11} - 1 = 2047 = 13 \cdot 89$ is not prime. □

2.41 Exercise 41

If n is a nonnegative integer, must $2^{2n} + 1$ be prime? Prove or give a counterexample.

Proof. No. Counterexample: Let $n = 3$. Then $2^{2n} + 1 = 2^6 + 1 = 65 = 13 \cdot 5$ is not prime. □

3 Exercise Set 4.3

3.1 Exercise 1

Proof. □

3.2 Exercise 2

Proof. □

3.3 Exercise 3

Proof. □

3.4 Exercise 4

Proof. □

3.5 Exercise 5

Proof.



3.6 Exercise 6

Proof.



3.7 Exercise 7

Proof.



3.8 Exercise 8

3.8.1 (a)

Proof.



3.8.2 (b)

Proof.



3.8.3 (c)

Proof.



3.9 Exercise 9

Proof.



3.10 Exercise 10

Proof.



3.11 Exercise 11

Proof.



3.12 Exercise 12

Proof.



3.13 Exercise 13

3.13.1 (a)

Proof.



3.13.2 (b)

Proof.



3.14 Exercise 14

3.14.1 (a)

Proof.



3.14.2 (b)

Proof.



3.15 Exercise 15

Proof.



3.16 Exercise 16

Proof.



3.17 Exercise 17

Proof.



3.18 Exercise 18

Proof.



3.19 Exercise 19

Proof.



3.20 Exercise 20

Proof.



3.21 Exercise 21

Proof.



3.22 Exercise 22

Proof.



3.23 Exercise 23

Proof.



3.24 Exercise 24

Proof.



3.25 Exercise 25

Proof.



3.26 Exercise 26

Proof.



3.27 Exercise 27

Proof.



3.28 Exercise 28

Proof.



3.29 Exercise 29

Proof.



3.30 Exercise 30

Proof.



3.31 Exercise 31

Proof.



3.32 Exercise 32

Proof.



3.33 Exercise 33

3.33.1 (a)

Proof.



3.33.2 (b)

Proof.



3.34 Exercise 34

3.34.1 (a)

Proof.



3.34.2 (b)

Proof.



3.35 Exercise 35

Proof.



3.36 Exercise 36

Proof.



3.37 Exercise 37

Proof.



3.38 Exercise 38

Proof.



3.39 Exercise 39

Proof.



4 Exercise Set 4.4

4.1 Exercise 1

Proof.



4.2 Exercise 2

Proof.



4.3 Exercise 3

Proof.



4.4 Exercise 4

Proof.



4.5 Exercise 5

Proof.



4.6 Exercise 6

Proof.



4.7 Exercise 7

Proof.



4.8 Exercise 8

Proof.



4.9 Exercise 9

Proof.



4.10 Exercise 10

Proof.



4.11 Exercise 11

Proof.



4.12 Exercise 12

Proof.



4.13 Exercise 13

Proof.



4.14 Exercise 14

Proof.



4.15 Exercise 15

Proof.



4.16 Exercise 16

Proof.



4.17 Exercise 17

Proof.



4.18 Exercise 18

4.18.1 (a)

Proof.



4.18.2 (b)

Proof.



4.19 Exercise 19

Proof.



4.20 Exercise 20

Proof.



4.21 Exercise 21

Proof.



4.22 Exercise 22

Proof.



4.23 Exercise 23

Proof.



4.24 Exercise 24

Proof.



4.25 Exercise 25

Proof.



4.26 Exercise 26

Proof.



4.27 Exercise 27

Proof.



4.28 Exercise 28

Proof.



4.29 Exercise 29

Proof.



4.30 Exercise 30

Proof.



4.31 Exercise 31

Proof.



4.32 Exercise 32

Proof.



4.33 Exercise 33

Proof.



4.34 Exercise 34

Proof.



4.35 Exercise 35

Proof.



4.36 Exercise 36

4.36.1 (a)

Proof.



4.36.2 (b)

Proof.



4.36.3 (c)

Proof.



4.36.4 (d)

Proof.



4.37 Exercise 37

4.37.1 (a)

Proof.



4.37.2 (b)

Proof.



4.37.3 (c)

Proof.



4.38 Exercise 38

4.38.1 (a)

Proof.



4.38.2 (b)

Proof.



4.38.3 (c)

Proof.



4.38.4 (d)

Proof.



4.39 Exercise 39

4.39.1 (a)

Proof.



4.39.2 (b)

Proof.



4.40 Exercise 40

4.40.1 (a)

Proof.



4.40.2 (b)

Proof.



4.41 Exercise 41

Proof.



4.42 Exercise 42

4.42.1 (a)

Proof.



4.42.2 (b)

Proof.



4.42.3 (c)

Proof.



4.43 Exercise 43

Proof.



4.44 Exercise 44

Proof.



4.45 Exercise 45

Proof.



4.46 Exercise 46

Proof.



4.47 Exercise 47

Proof.



4.48 Exercise 48

Proof.



4.49 Exercise 49

Proof.



4.50 Exercise 50

Proof.



5 Exercise Set 4.5

5.1 Exercise 1

Proof.



5.2 Exercise 2

Proof.



5.3 Exercise 3

Proof.



5.4 Exercise 4

Proof.



5.5 Exercise 5

Proof.



5.6 Exercise 6

Proof.



5.7 Exercise 7

5.7.1 (a)

Proof.



5.7.2 (b)

Proof.



5.8 Exercise 8

5.8.1 (a)

Proof.



5.8.2 (b)

Proof.



5.9 Exercise 9

5.9.1 (a)

Proof.



5.9.2 (b)

Proof.



5.10 Exercise 10

5.10.1 (a)

Proof.



5.10.2 (b)

Proof.



5.11 Exercise 11

5.11.1 (a)

Proof.



5.11.2 (b)

Proof.



5.11.3 (c)

Proof.



5.12 Exercise 12

Proof.



5.13 Exercise 13

Proof.



5.14 Exercise 14

Proof.



5.15 Exercise 15

Proof.



5.16 Exercise 16

Proof.



5.17 Exercise 17

Proof.



5.18 Exercise 18

5.18.1 (a)

Proof.



5.18.2 (b)

Proof.



5.19 Exercise 19

Proof.



5.20 Exercise 20

Proof.



5.21 Exercise 21

Proof.



5.22 Exercise 22

Proof.



5.23 Exercise 23

Proof.



5.24 Exercise 24

Proof.



5.25 Exercise 25

Proof.



5.26 Exercise 26

Proof.



5.27 Exercise 27

Proof.



5.28 Exercise 28

5.28.1 (a)

Proof.



5.28.2 (b)

Proof.



5.29 Exercise 29

5.29.1 (a)

Proof.



5.29.2 (b)

Proof.



5.30 Exercise 30

5.30.1 (a)

Proof.



5.30.2 (b)

Proof.



5.31 Exercise 31

5.31.1 (a)

Proof.



5.31.2 (b)

Proof.



5.31.3 (c)

Proof.



5.32 Exercise 32

Proof.



5.33 Exercise 33

Proof.



5.34 Exercise 34

Proof.



5.35 Exercise 35

Proof.



5.36 Exercise 36

Proof.



5.37 Exercise 37

Proof.



5.38 Exercise 38

Proof.



5.39 Exercise 39

Proof.



5.40 Exercise 40

Proof.



5.41 Exercise 41

Proof.



5.42 Exercise 42

Proof.



5.43 Exercise 43

Proof.



5.44 Exercise 44

5.44.1 (a)

Proof.



5.44.2 (b)

Proof.



5.44.3 (c)

Proof.



5.45 Exercise 45

Proof.



5.46 Exercise 46

Proof.



5.47 Exercise 47

Proof.



5.48 Exercise 48

Proof.



5.49 Exercise 49

Proof.



5.50 Exercise 50

Proof.



6 Exercise Set 4.6

6.1 Exercise 1

Proof.



6.2 Exercise 2

Proof.



6.3 Exercise 3

Proof.



6.4 Exercise 4

Proof.



6.5 Exercise 5

Proof.



6.6 Exercise 6

Proof.



6.7 Exercise 7

Proof.



6.8 Exercise 8

Proof.



6.9 Exercise 9

Proof.



6.10 Exercise 10

6.10.1 (a)

Proof.



6.10.2 (b)

Proof.



6.11 Exercise 11

Proof.



6.12 Exercise 12

Proof.



6.13 Exercise 13

Proof.



6.14 Exercise 14

Proof.



6.15 Exercise 15

Proof.



6.16 Exercise 16

Proof.



6.17 Exercise 17

Proof.



6.18 Exercise 18

Proof.



6.19 Exercise 19

Proof.



6.20 Exercise 20

Proof.



6.21 Exercise 21

Proof.



6.22 Exercise 22

Proof.



6.23 Exercise 23

Proof.



6.24 Exercise 24

Proof.



6.25 Exercise 25

Proof.



6.26 Exercise 26

Proof.



6.27 Exercise 27

Proof.



6.28 Exercise 28

Proof.



6.29 Exercise 29

Proof.



6.30 Exercise 30

Proof.



6.31 Exercise 31

Proof.



6.32 Exercise 32

Proof.



6.33 Exercise 33

Proof.



7 Exercise Set 4.7

7.1 Exercise 1

Proof.



7.2 Exercise 2

Proof.



7.3 Exercise 3

Proof.



7.4 Exercise 4

Proof.



7.5 Exercise 5

Proof.



7.6 Exercise 6

Proof.



7.7 Exercise 7

Proof.



7.8 Exercise 8

Proof.



7.9 Exercise 9

7.9.1 (a)

Proof.



7.9.2 (b)

Proof.



7.10 Exercise 10

Proof.



7.11 Exercise 11

Proof.



7.12 Exercise 12

7.12.1 (a)

Proof.



7.12.2 (b)

Proof.



7.13 Exercise 13

7.13.1 (a)

Proof.



7.13.2 (b)

Proof.



7.14 Exercise 14

7.14.1 (a)

Proof.



7.14.2 (b)

Proof.



7.15 Exercise 15

Proof.



7.16 Exercise 16

Proof.



7.17 Exercise 17

Proof.



7.18 Exercise 18

Proof.



7.19 Exercise 19

Proof.



7.20 Exercise 20

Proof.



7.21 Exercise 21

7.21.1 (a)

Proof.



7.21.2 (b)

Proof.



7.22 Exercise 22

7.22.1 (a)

Proof.



7.22.2 (b)

Proof.



7.23 Exercise 23

Proof.



7.24 Exercise 24

Proof.



7.25 Exercise 25

Proof.



7.26 Exercise 26

Proof.



7.27 Exercise 27

Proof.



7.28 Exercise 28

Proof.



7.29 Exercise 29

Proof.



7.30 Exercise 30

7.30.1 (a)

Proof.



7.30.2 (b)

Proof.



7.31 Exercise 31

7.31.1 (a)

Proof.



7.31.2 (b)

Proof.



7.31.3 (c)

Proof.



7.32 Exercise 32

Proof.



7.33 Exercise 33

Proof.



7.34 Exercise 34

7.34.1 (a)

Proof.



7.34.2 (b)

Proof.



7.34.3 (c)

Proof.



7.34.4 (d)

Proof.



7.35 Exercise 35

Proof.



7.36 Exercise 36

Proof.



8 Exercise Set 4.8

8.1 Exercise 1

Proof.



8.2 Exercise 2

Proof.



8.3 Exercise 3

Proof.



8.4 Exercise 4

Proof.



8.5 Exercise 5

Proof.



8.6 Exercise 6

Proof.



8.7 Exercise 7

Proof.



8.8 Exercise 8

Proof.



8.9 Exercise 9

Proof.



8.10 Exercise 10

Proof.



8.11 Exercise 11

Proof.



8.12 Exercise 12

Proof.



8.13 Exercise 13

Proof.



8.14 Exercise 14

Proof.



8.15 Exercise 15

Proof.



8.16 Exercise 16

Proof.



8.17 Exercise 17

Proof.



8.18 Exercise 18

8.18.1 (a)

Proof.



8.18.2 (b)

Proof.



8.19 Exercise 19

8.19.1 (a)

Proof.



8.19.2 (b)

Proof.



8.19.3 (c)

Proof.



8.20 Exercise 20

Proof.



8.21 Exercise 21

Proof.



8.22 Exercise 22

Proof.



8.23 Exercise 23

Proof.



8.24 Exercise 24

Proof.



8.25 Exercise 25

Proof.



8.26 Exercise 26

Proof.



8.27 Exercise 27

Proof.



8.28 Exercise 28

Proof.



8.29 Exercise 29

Proof.



8.30 Exercise 30

8.30.1 (a)

Proof.



8.30.2 (b)

Proof.



8.31 Exercise 31

Proof.



8.32 Exercise 32

Proof.



8.33 Exercise 33

Proof.



8.34 Exercise 34

8.34.1 (a)

Proof.



8.34.2 (b)

Proof.



8.35 Exercise 35

Proof.



8.36 Exercise 36

Proof.



8.37 Exercise 37

Proof.



8.38 Exercise 38

Proof.



9 Exercise Set 4.9

9.1 Exercise 1

Proof.



9.2 Exercise 2

Proof.



9.3 Exercise 3

Proof.



9.4 Exercise 4

Proof.



9.5 Exercise 5

Proof.



9.6 Exercise 6

Proof.



9.7 Exercise 7

Proof.



9.8 Exercise 8

Proof.



9.9 Exercise 9

Proof.



9.10 Exercise 10

Proof.



9.11 Exercise 11

Proof.



9.12 Exercise 12

Proof.



9.13 Exercise 13

Proof.



9.14 Exercise 14

9.14.1 (a)

Proof.



9.14.2 (b)

Proof.



9.15 Exercise 15

9.15.1 (a)

Proof.



9.15.2 (b)

Proof.



9.16 Exercise 16

9.16.1 (a)

Proof.



9.16.2 (b)

Proof.



9.17 Exercise 17

Proof.



9.18 Exercise 18

Proof.



9.19 Exercise 19

Proof.



9.20 Exercise 20

9.20.1 (a)

Proof.



9.20.2 (b)

Proof.



9.21 Exercise 21

9.21.1 (a)

Proof.



9.21.2 (b)

Proof.



9.21.3 (c)

Proof.



9.22 Exercise 22

Proof.



9.23 Exercise 23

9.23.1 (a)

Proof.



9.23.2 (b)

Proof.



9.23.3 (c)

Proof.



9.23.4 (d)

Proof.



9.23.5 (e)

Proof.



9.23.6 (f)

Proof.



9.24 Exercise 24

9.24.1 (a)

Proof.



9.24.2 (b)

Proof.



9.24.3 (c)

Proof.



9.24.4 (d)

Proof.



9.24.5 (e)

Proof.



9.24.6 (f)

Proof.



9.25 Exercise 25

Proof.



10 Exercise Set 4.10

10.1 Exercise 1

Proof.



10.2 Exercise 2

Proof.



10.3 Exercise 3

10.3.1 (a)

Proof.



10.3.2 (b)

Proof.



10.4 Exercise 4

Proof.



10.5 Exercise 5

Proof.



10.6 Exercise 6

Proof.



10.7 Exercise 7

Proof.



10.8 Exercise 8

10.8.1 (a)

Proof.



10.8.2 (b)

Proof.



10.9 Exercise 9

Proof.



10.10 Exercise 10

Proof.



10.11 Exercise 11

Proof.



10.12 Exercise 12

Proof.



10.13 Exercise 13

Proof.



10.14 Exercise 14

Proof.



10.15 Exercise 15

Proof.



10.16 Exercise 16

Proof.



10.17 Exercise 17

Proof.



10.18 Exercise 18

Proof.



10.19 Exercise 19

Proof.



10.20 Exercise 20

Proof.



10.21 Exercise 21

Proof.



10.22 Exercise 22

Proof.



10.23 Exercise 23

10.23.1 (a)

Proof.



10.23.2 (b)

Proof.



10.24 Exercise 24

Proof.



10.25 Exercise 25

10.25.1 (a)

Proof.



10.25.2 (b)

Proof.



10.26 Exercise 26

10.26.1 (a)

Proof.



10.26.2 (b)

Proof.



10.27 Exercise 27

10.27.1 (a)

Proof.



10.27.2 (b)

Proof.



10.27.3 (c)

Proof.



10.28 Exercise 28

10.28.1 (a)

Proof.



10.28.2 (b)

Proof.



10.28.3 (c)

Proof.



10.29 Exercise 29

Proof.



10.30 Exercise 30

Proof.



10.31 Exercise 31

Proof.



10.32 Exercise 32

Proof.

