

Chapter 4 Solutions, Susanna Epp Discrete Math 5th Edition

<https://github.com/spamegg1>

May 20, 2023

Contents

1	Exercise Set 4.1	13
1.1	Exercise 1	13
1.1.1	(a)	13
1.1.2	(b)	13
1.1.3	(c)	13
1.2	Exercise 2	13
1.2.1	(a)	13
1.2.2	(b)	13
1.2.3	(c)	13
1.3	Exercise 3	14
1.3.1	(a)	14
1.3.2	(b)	14
1.3.3	(c)	14
1.4	Exercise 4	14
1.4.1	(a)	14
1.4.2	(b)	14
1.4.3	(c)	15
1.5	Exercise 5	15
1.6	Exercise 6	15
1.7	Exercise 7	15
1.8	Exercise 8	15
1.9	Exercise 9	15
1.10	Exercise 10	16
1.11	Exercise 11	16
1.12	Exercise 12	16
1.13	Exercise 13	16
1.14	Exercise 14	17
1.15	Exercise 15	17
1.16	Exercise 16	17
1.17	Exercise 17	17

1.18	Exercise 18	17
1.19	Exercise 19	18
1.20	Exercise 20	18
1.21	Exercise 21	18
1.22	Exercise 22	18
1.23	Exercise 23	19
1.24	Exercise 24	19
1.25	Exercise 25	19
1.26	Exercise 26	20
1.27	Exercise 27	20
1.28	Exercise 28	20
1.29	Exercise 29	21
1.30	Exercise 30	21
1.31	Exercise 31	22
2	Exercise Set 4.2	22
2.1	Exercise 1	22
2.2	Exercise 2	23
2.3	Exercise 3	24
2.4	Exercise 4	24
2.5	Exercise 5	24
2.6	Exercise 6	25
2.7	Exercise 7	25
2.8	Exercise 8	26
2.9	Exercise 9	26
2.10	Exercise 10	27
2.11	Exercise 11	27
2.12	Exercise 12	27
2.13	Exercise 13	28
2.14	Exercise 14	28
2.15	Exercise 15	28
2.16	Exercise 16	28
2.17	Exercise 17	29
2.18	Exercise 18	29
2.19	Exercise 19	30
2.20	Exercise 20	31
2.21	Exercise 21	31
2.22	Exercise 22	31
2.23	Exercise 23	32
2.24	Exercise 24	32
2.25	Exercise 25	32
2.26	Exercise 26	32
2.27	Exercise 27	33
2.28	Exercise 28	33
2.29	Exercise 29	33
2.30	Exercise 30	33

2.31	Exercise 31	33
2.32	Exercise 32	34
2.33	Exercise 33	34
2.34	Exercise 34	34
2.35	Exercise 35	34
2.36	Exercise 36	35
2.37	Exercise 37	35
2.38	Exercise 38	35
2.39	Exercise 39	36
2.40	Exercise 40	36
2.41	Exercise 41	36
3	Exercise Set 4.3	36
3.1	Exercise 1	36
3.2	Exercise 2	36
3.3	Exercise 3	37
3.4	Exercise 4	37
3.5	Exercise 5	37
3.6	Exercise 6	37
3.7	Exercise 7	37
3.8	Exercise 8	38
3.8.1	(a)	38
3.8.2	(b)	38
3.8.3	(c)	38
3.9	Exercise 9	38
3.10	Exercise 10	38
3.11	Exercise 11	39
3.12	Exercise 12	39
3.13	Exercise 13	39
3.13.1	(a)	39
3.13.2	(b)	40
3.14	Exercise 14	40
3.14.1	(a)	40
3.14.2	(b)	40
3.15	Exercise 15	41
3.16	Exercise 16	41
3.17	Exercise 17	41
3.18	Exercise 18	42
3.19	Exercise 19	42
3.20	Exercise 20	42
3.21	Exercise 21	43
3.22	Exercise 22	43
3.23	Exercise 23	43
3.24	Exercise 24	43
3.25	Exercise 25	44
3.26	Exercise 26	44

3.27	Exercise 27	44
3.28	Exercise 28	44
3.29	Exercise 29	45
3.30	Exercise 30	46
3.31	Exercise 31	46
3.32	Exercise 32	47
3.33	Exercise 33	47
3.33.1	(a)	48
3.33.2	(b)	48
3.34	Exercise 34	48
3.34.1	(a)	48
3.34.2	(b)	49
3.35	Exercise 35	49
3.36	Exercise 36	49
3.37	Exercise 37	49
3.38	Exercise 38	50
3.39	Exercise 39	50
4	Exercise Set 4.4	50
4.1	Exercise 1	50
4.2	Exercise 2	50
4.3	Exercise 3	51
4.4	Exercise 4	51
4.5	Exercise 5	51
4.6	Exercise 6	51
4.7	Exercise 7	51
4.8	Exercise 8	51
4.9	Exercise 9	51
4.10	Exercise 10	52
4.11	Exercise 11	52
4.12	Exercise 12	52
4.13	Exercise 13	52
4.14	Exercise 14	52
4.15	Exercise 15	53
4.16	Exercise 16	53
4.17	Exercise 17	53
4.18	Exercise 18	53
4.18.1	(a)	53
4.18.2	(b)	54
4.19	Exercise 19	54
4.20	Exercise 20	54
4.21	Exercise 21	54
4.22	Exercise 22	55
4.23	Exercise 23	55
4.24	Exercise 24	55
4.25	Exercise 25	55

4.26	Exercise 26	56
4.27	Exercise 27	56
4.28	Exercise 28	56
4.29	Exercise 29	56
4.30	Exercise 30	56
4.31	Exercise 31	56
4.32	Exercise 32	57
4.33	Exercise 33	57
4.34	Exercise 34	57
4.35	Exercise 35	57
4.36	Exercise 36	58
4.36.1	(a)	58
4.36.2	(b)	58
4.36.3	(c)	58
4.36.4	(d)	58
4.37	Exercise 37	58
4.37.1	(a)	59
4.37.2	(b)	59
4.37.3	(c)	59
4.38	Exercise 38	59
4.38.1	(a)	59
4.38.2	(b)	59
4.38.3	(c)	59
4.38.4	(d)	59
4.39	Exercise 39	60
4.39.1	(a)	60
4.39.2	(b)	60
4.40	Exercise 40	60
4.40.1	(a)	60
4.40.2	(b)	61
4.41	Exercise 41	61
4.42	Exercise 42	61
4.42.1	(a)	61
4.42.2	(b)	61
4.42.3	(c)	61
4.43	Exercise 43	62
4.44	Exercise 44	62
4.45	Exercise 45	63
4.46	Exercise 46	63
4.47	Exercise 47	64
4.48	Exercise 48	65
4.49	Exercise 49	65
4.50	Exercise 50	65
5	Exercise Set 4.5	66
5.1	Exercise 1	66

5.2	Exercise 2	66
5.3	Exercise 3	66
5.4	Exercise 4	66
5.5	Exercise 5	66
5.6	Exercise 6	66
5.7	Exercise 7	66
	5.7.1 (a)	66
	5.7.2 (b)	66
5.8	Exercise 8	66
	5.8.1 (a)	66
	5.8.2 (b)	67
5.9	Exercise 9	67
	5.9.1 (a)	67
	5.9.2 (b)	67
5.10	Exercise 10	67
	5.10.1 (a)	67
	5.10.2 (b)	67
5.11	Exercise 11	67
	5.11.1 (a)	67
	5.11.2 (b)	67
	5.11.3 (c)	67
5.12	Exercise 12	67
5.13	Exercise 13	67
5.14	Exercise 14	67
5.15	Exercise 15	68
5.16	Exercise 16	68
5.17	Exercise 17	68
5.18	Exercise 18	68
	5.18.1 (a)	68
	5.18.2 (b)	68
5.19	Exercise 19	68
5.20	Exercise 20	68
5.21	Exercise 21	68
5.22	Exercise 22	68
5.23	Exercise 23	68
5.24	Exercise 24	68
5.25	Exercise 25	69
5.26	Exercise 26	69
5.27	Exercise 27	69
5.28	Exercise 28	69
	5.28.1 (a)	69
	5.28.2 (b)	69
5.29	Exercise 29	69
	5.29.1 (a)	69
	5.29.2 (b)	69
5.30	Exercise 30	69

5.30.1	(a)	69
5.30.2	(b)	69
5.31	Exercise 31	69
5.31.1	(a)	69
5.31.2	(b)	70
5.31.3	(c)	70
5.32	Exercise 32	70
5.33	Exercise 33	70
5.34	Exercise 34	70
5.35	Exercise 35	70
5.36	Exercise 36	70
5.37	Exercise 37	70
5.38	Exercise 38	70
5.39	Exercise 39	70
5.40	Exercise 40	70
5.41	Exercise 41	70
5.42	Exercise 42	71
5.43	Exercise 43	71
5.44	Exercise 44	71
5.44.1	(a)	71
5.44.2	(b)	71
5.44.3	(c)	71
5.45	Exercise 45	71
5.46	Exercise 46	71
5.47	Exercise 47	71
5.48	Exercise 48	71
5.49	Exercise 49	71
5.50	Exercise 50	71
6	Exercise Set 4.6	72
6.1	Exercise 1	72
6.2	Exercise 2	72
6.3	Exercise 3	72
6.4	Exercise 4	72
6.5	Exercise 5	72
6.6	Exercise 6	72
6.7	Exercise 7	72
6.8	Exercise 8	72
6.9	Exercise 9	72
6.10	Exercise 10	72
6.10.1	(a)	72
6.10.2	(b)	72
6.11	Exercise 11	73
6.12	Exercise 12	73
6.13	Exercise 13	73
6.14	Exercise 14	73

6.15	Exercise 15	73
6.16	Exercise 16	73
6.17	Exercise 17	73
6.18	Exercise 18	73
6.19	Exercise 19	73
6.20	Exercise 20	73
6.21	Exercise 21	73
6.22	Exercise 22	73
6.23	Exercise 23	74
6.24	Exercise 24	74
6.25	Exercise 25	74
6.26	Exercise 26	74
6.27	Exercise 27	74
6.28	Exercise 28	74
6.29	Exercise 29	74
6.30	Exercise 30	74
6.31	Exercise 31	74
6.32	Exercise 32	74
6.33	Exercise 33	74
7	Exercise Set 4.7	75
7.1	Exercise 1	75
7.2	Exercise 2	75
7.3	Exercise 3	75
7.4	Exercise 4	75
7.5	Exercise 5	75
7.6	Exercise 6	75
7.7	Exercise 7	75
7.8	Exercise 8	75
7.9	Exercise 9	75
	7.9.1 (a)	75
	7.9.2 (b)	75
7.10	Exercise 10	75
7.11	Exercise 11	76
7.12	Exercise 12	76
	7.12.1 (a)	76
	7.12.2 (b)	76
7.13	Exercise 13	76
	7.13.1 (a)	76
	7.13.2 (b)	76
7.14	Exercise 14	76
	7.14.1 (a)	76
	7.14.2 (b)	76
7.15	Exercise 15	76
7.16	Exercise 16	76
7.17	Exercise 17	76

7.18	Exercise 18	76
7.19	Exercise 19	77
7.20	Exercise 20	77
7.21	Exercise 21	77
7.21.1	(a)	77
7.21.2	(b)	77
7.22	Exercise 22	77
7.22.1	(a)	77
7.22.2	(b)	77
7.23	Exercise 23	77
7.24	Exercise 24	77
7.25	Exercise 25	77
7.26	Exercise 26	77
7.27	Exercise 27	77
7.28	Exercise 28	78
7.29	Exercise 29	78
7.30	Exercise 30	78
7.30.1	(a)	78
7.30.2	(b)	78
7.31	Exercise 31	78
7.31.1	(a)	78
7.31.2	(b)	78
7.31.3	(c)	78
7.32	Exercise 32	78
7.33	Exercise 33	78
7.34	Exercise 34	78
7.34.1	(a)	78
7.34.2	(b)	78
7.34.3	(c)	79
7.34.4	(d)	79
7.35	Exercise 35	79
7.36	Exercise 36	79
8	Exercise Set 4.8	79
8.1	Exercise 1	79
8.2	Exercise 2	79
8.3	Exercise 3	79
8.4	Exercise 4	79
8.5	Exercise 5	79
8.6	Exercise 6	79
8.7	Exercise 7	79
8.8	Exercise 8	80
8.9	Exercise 9	80
8.10	Exercise 10	80
8.11	Exercise 11	80
8.12	Exercise 12	80

8.13	Exercise 13	80
8.14	Exercise 14	80
8.15	Exercise 15	80
8.16	Exercise 16	80
8.17	Exercise 17	80
8.18	Exercise 18	80
8.18.1	(a)	80
8.18.2	(b)	81
8.19	Exercise 19	81
8.19.1	(a)	81
8.19.2	(b)	81
8.19.3	(c)	81
8.20	Exercise 20	81
8.21	Exercise 21	81
8.22	Exercise 22	81
8.23	Exercise 23	81
8.24	Exercise 24	81
8.25	Exercise 25	81
8.26	Exercise 26	81
8.27	Exercise 27	82
8.28	Exercise 28	82
8.29	Exercise 29	82
8.30	Exercise 30	82
8.30.1	(a)	82
8.30.2	(b)	82
8.31	Exercise 31	82
8.32	Exercise 32	82
8.33	Exercise 33	82
8.34	Exercise 34	82
8.34.1	(a)	82
8.34.2	(b)	82
8.35	Exercise 35	82
8.36	Exercise 36	83
8.37	Exercise 37	83
8.38	Exercise 38	83
9	Exercise Set 4.9	83
9.1	Exercise 1	83
9.2	Exercise 2	83
9.3	Exercise 3	83
9.4	Exercise 4	83
9.5	Exercise 5	83
9.6	Exercise 6	83
9.7	Exercise 7	83
9.8	Exercise 8	83
9.9	Exercise 9	84

9.10	Exercise 10	84
9.11	Exercise 11	84
9.12	Exercise 12	84
9.13	Exercise 13	84
9.14	Exercise 14	84
9.14.1	(a)	84
9.14.2	(b)	84
9.15	Exercise 15	84
9.15.1	(a)	84
9.15.2	(b)	84
9.16	Exercise 16	84
9.16.1	(a)	84
9.16.2	(b)	84
9.17	Exercise 17	85
9.18	Exercise 18	85
9.19	Exercise 19	85
9.20	Exercise 20	85
9.20.1	(a)	85
9.20.2	(b)	85
9.21	Exercise 21	85
9.21.1	(a)	85
9.21.2	(b)	85
9.21.3	(c)	85
9.22	Exercise 22	85
9.23	Exercise 23	85
9.23.1	(a)	85
9.23.2	(b)	85
9.23.3	(c)	86
9.23.4	(d)	86
9.23.5	(e)	86
9.23.6	(f)	86
9.24	Exercise 24	86
9.24.1	(a)	86
9.24.2	(b)	86
9.24.3	(c)	86
9.24.4	(d)	86
9.24.5	(e)	86
9.24.6	(f)	86
9.25	Exercise 25	86
10	Exercise Set 4.10	87
10.1	Exercise 1	87
10.2	Exercise 2	87
10.3	Exercise 3	87
10.3.1	(a)	87
10.3.2	(b)	87

10.4 Exercise 4	87
10.5 Exercise 5	87
10.6 Exercise 6	87
10.7 Exercise 7	87
10.8 Exercise 8	87
10.8.1 (a)	87
10.8.2 (b)	87
10.9 Exercise 9	88
10.10 Exercise 10	88
10.11 Exercise 11	88
10.12 Exercise 12	88
10.13 Exercise 13	88
10.14 Exercise 14	88
10.15 Exercise 15	88
10.16 Exercise 16	88
10.17 Exercise 17	88
10.18 Exercise 18	88
10.19 Exercise 19	88
10.20 Exercise 20	88
10.21 Exercise 21	89
10.22 Exercise 22	89
10.23 Exercise 23	89
10.23.1 (a)	89
10.23.2 (b)	89
10.24 Exercise 24	89
10.25 Exercise 25	89
10.25.1 (a)	89
10.25.2 (b)	89
10.26 Exercise 26	89
10.26.1 (a)	89
10.26.2 (b)	89
10.27 Exercise 27	89
10.27.1 (a)	89
10.27.2 (b)	90
10.27.3 (c)	90
10.28 Exercise 28	90
10.28.1 (a)	90
10.28.2 (b)	90
10.28.3 (c)	90
10.29 Exercise 29	90
10.30 Exercise 30	90
10.31 Exercise 31	90
10.32 Exercise 32	90

1 Exercise Set 4.1

1.1 Exercise 1

Assume that k is a particular integer.

1.1.1 (a)

Is -17 an odd integer?

Proof. Yes: $-17 = 2(-9) + 1$.

□

1.1.2 (b)

Is 0 neither even nor odd?

Proof. No. 0 is even because $0 = 0 \cdot 2$.

□

1.1.3 (c)

Is $2k - 1$ odd?

Proof. Yes: $2k - 1 = 2(k - 1) + 1$ and $k - 1$ is an integer because it is a difference of integers.

□

1.2 Exercise 2

Assume that c is a particular integer.

1.2.1 (a)

Is $-6c$ an even integer?

Proof. Yes, because $-6c = 2 \cdot (-3c) = 2k$ where $k = -3c$ is an integer.

□

1.2.2 (b)

Is $8c + 5$ an odd integer?

Proof. Yes, because $8c + 5 = 2(4c + 2) + 1$ and $k = 4c + 2$ is an integer.

□

1.2.3 (c)

Is $(c^2 + 1) - (c^2 - 1) - 2$ an even integer?

Proof. Yes, because it equals 0 : $(c^2 + 1) - (c^2 - 1) - 2 = c^2 + 1 - c^2 + 1 - 2 = 2 - 2 = 0$.

□

1.3 Exercise 3

Assume that m and n are particular integers.

1.3.1 (a)

Is $6m + 8n$ even?

Proof. Yes: $6m + 8n = 2(3m + 4n)$ and $(3m + 4n)$ is an integer because $3, 4, m$, and n are integers, and products and sums of integers are integers. \square

1.3.2 (b)

Is $10mn + 7$ odd?

Proof. Yes: $10mn + 7 = 2(5mn + 3) + 1$ and $5mn + 3$ is an integer because $3, 5, m$, and n are integers, and products and sums of integers are integers. \square

1.3.3 (c)

If $m > n > 0$, is $m^2 - n^2$ composite?

Proof. Not necessarily. For instance, if $m = 3$ and $n = 2$, then $m^2 - n^2 = 9 - 4 = 5$, which is prime. (However, $m^2 - n^2$ is composite for many values of m and n because of the identity $m^2 - n^2 = (m - n)(m + n)$.) \square

1.4 Exercise 4

Assume that r and s are particular integers.

1.4.1 (a)

Is $4rs$ even?

Proof. Yes: $4rs = 2(2rs)$ and $2rs$ is an integer because $2, r, s$ are integers, and products of integers are integers. \square

1.4.2 (b)

Is $6r + 4s^2 + 3$ odd?

Proof. Yes: $6r + 4s^2 + 3 = 2(3r + 2s^2 + 1) + 1$ and $3r + 2s^2 + 1$ is an integer because $3, r, 2, s, 1$ are integers and products and sums of integers are integers. \square

1.4.3 (c)

If r and s are both positive, is $r^2 + 2rs + s^2$ composite?

Proof. Yes: $r^2 + 2rs + s^2 = (r + s)(r + s)$ and $r + s \geq 2$, therefore $r^2 + 2rs + s^2$ is a product of two integers both of which are greater than 1. \square

Prove the statements in 5–11.

1.5 Exercise 5

There are integers m and n such that $m > 1$ and $n > 1$ and $\frac{1}{m} + \frac{1}{n}$ is an integer.

Proof. For example, let $m = n = 2$. Then m and n are integers such that $m > 1$ and $n > 1$ and $\frac{1}{m} + \frac{1}{n} = \frac{1}{2} + \frac{1}{2} = 1$ which is an integer. \square

1.6 Exercise 6

There are distinct integers m and n such that $\frac{1}{m} + \frac{1}{n}$ is an integer.

Proof. For example, let $m = 1, n = -1$. Then m and n are integers such that $\frac{1}{m} + \frac{1}{n} = \frac{1}{1} - \frac{1}{1} = 0$ which is an integer. \square

1.7 Exercise 7

There are real numbers a and b such that $\sqrt{a+b} = \sqrt{a}\sqrt{b}$.

Proof. For example, let $a = 0, b = 0$. Then a and b are real numbers such that $\sqrt{a+b} = \sqrt{0+0} = 0 = \sqrt{0} + \sqrt{0} = \sqrt{a} + \sqrt{b}$. \square

1.8 Exercise 8

There is an integer $n > 5$ such that $2^n - 1$ is prime.

Proof. For example, let $n = 7$. Then n is an integer such that $n > 5$ and $2^n - 1 = 127$, which is prime. \square

1.9 Exercise 9

There is a real number x such that $x > 1$ and $2^x > x^{10}$.

Proof. For example, take $x = 80$. Then

$$x^{10} = 80^{10} = 8^{10} \cdot 10^{10} = (2^3)^{10} \cdot 10^{10} = 2^{30} \cdot 10^{10}.$$

We have $2^{50} \approx 1,125899907 \cdot 10^{15} > 10^{10}$. So $2^{80} = 2^{30} \cdot 2^{50} > 2^{30} \cdot 10^{10} = 80^{10}$.

Therefore $x = 80$ is a real number such that $x > 1$ and $2^x > x^{10}$. \square

Definition: An integer n is called a **perfect square** if, and only if, $n = k^2$ for some integer k .

1.10 Exercise 10

There is a perfect square that can be written as the sum of two other perfect squares.

Proof. For example, 25, 9, and 16 are all perfect squares, because $25 = 5^2$, $9 = 3^2$, and $16 = 4^2$, and $25 = 9 + 16$. Thus 25 is a perfect square that can be written as a sum of two other perfect squares. \square

1.11 Exercise 11

There is an integer n such that $2n^2 - 5n + 2$ is prime.

Proof. For example, take $n = 3$. Then $2n^2 - 5n + 2 = 18 - 15 + 2 = 5$ is prime. (You can find this value of n by either starting at $n = 1$ and using trial and error, or noticing that $2n^2 - 5n + 2 = (2n - 1)(n - 2)$, so, for this to be prime, one of the factors has to be 1.) \square

In 12 – 13, (a) write a negation for the given statement, and (b) use a counterexample to disprove the given statement. explain how the counterexample actually shows that the given statement is false.

1.12 Exercise 12

For all real numbers a and b , if $a < b$ then $a^2 < b^2$.

Proof. a. *Negation for the statement:* There exist real numbers a and b such that $a < b$ and $a^2 \not< b^2$.

b. *Counterexample for the statement:* Let $a = -2$ and $b = -1$. Then $a < b$ because $-2 < -1$, but $a^2 \not< b^2$ because $(-2)^2 = 4$ and $(-1)^2 = 1$ and $4 \not< 1$. [So the hypothesis of the statement is true and its conclusion is false.] \square

1.13 Exercise 13

For every integer n , if n is odd then $\frac{n-1}{2}$ is odd.

Proof. a. *Negation for the statement:* There exists an integer n such that n is odd and $\frac{n-1}{2}$ is not odd.

b. *Counterexample for the statement:* Let $n = 5$. Then n is odd because $5 = 2 \cdot 2 + 1$, but $\frac{n-1}{2}$ is not odd because $\frac{5-1}{2} = 2 = 2 \cdot 1$ is even. [So the hypothesis of the statement is true and its conclusion is false.] \square

Disprove each of the statements in 14 – 16 by giving a counterexample. In each case explain how the counterexample actually disproves the statement.

1.14 Exercise 14

For all integers m and n , if $2m + n$ is odd then m and n are both odd.

Proof. Counterexample: Let $m = 2$ and $n = 1$. Then $2m + n = 2 \cdot 2 + 1 = 5$, which is odd. But m is not odd, and so it is false that both m and n are odd. [This is one counterexample among many.] \square

1.15 Exercise 15

For every integer p , if p is prime then $p^2 - 1$ is even.

Proof. Counterexample: Let $p = 2$ which is prime, but $p^2 - 1 = 2^2 - 1 = 4 - 1 = 3$ is not even. [This is the only counterexample! For every other prime p , $p^2 - 1$ is even.] \square

1.16 Exercise 16

For every integer n , if n is even then $n^2 + 1$ is prime.

Proof. Counterexample: Let $n = 8$. Then n is even. But $n^2 + 1 = 65 = 13 \cdot 5$ is not prime. [This is one counterexample among many.] \square

In 17 – 20, determine whether the property is true for all integers, true for no integers, or true for some integers and false for other integers. Justify your answers.

1.17 Exercise 17

$$(a + b)^2 = a^2 + b^2$$

Proof. This property is true for some integers and false for other integers. For instance, if $a = 0$ and $b = 1$, the property is true because $(0 + 1)^2 = 0^2 + 1^2$, but if $a = 1$ and $b = 1$, the property is false because $(1 + 1)^2 = 4$ and $1^2 + 1^2 = 2$ and $4 \neq 2$. \square

1.18 Exercise 18

$$\frac{a}{b} + \frac{c}{d} = \frac{a + c}{b + d}$$

Proof. True for some integers, false for others. For example, if $a = c = 0$ and $b = d = 1$ then $\frac{0}{1} + \frac{0}{1} = 0 = \frac{0 + 0}{1 + 1}$ is true. But if $a = 1, b = 2, c = 3$ and $d = 4$ then

$$\frac{a}{b} + \frac{c}{d} = \frac{1}{2} + \frac{3}{4} = \frac{5}{4} \neq \frac{4}{6} = \frac{1 + 3}{2 + 4} = \frac{a + c}{b + d}.$$

\square

1.19 Exercise 19

$$-a^n = (-a)^n$$

Hint: This property is true for some integers and false for other integers. To justify this answer you need to find examples of both.

Proof. True for some integers: let $a = 0, n = 1$. Then $-0^1 = -0 = 0$ and $(-0)^1 = 0^1 = 0$ so the equality holds. When $a = n = 2$ it is false: $-2^2 = -4$ but $(-2)^2 = 4$ and $-4 \neq 4$. \square

1.20 Exercise 20

The average of any two odd integers is odd.

Proof. True for some, false for others. For example, 3 and 7 are both odd, and their average $(3+7)/2 = 5$ is odd. But 3 and 5 are both odd, and their average is $(3+5)/2 = 4$ is even. \square

Prove the statement in 21 and 22 by the method of exhaustion.

1.21 Exercise 21

Every positive even integer less than 26 can be expressed as a sum of three or fewer perfect squares. (For instance, $10 = 1^2 + 3^2$ and $16 = 4^2$.)

Proof. $2 = 1^2 + 1^2$

$$4 = 2^2$$

$$6 = 2^2 + 1^2 + 1^2$$

$$8 = 2^2 + 2^2$$

$$10 = 1^2 + 3^2$$

$$12 = 2^2 + 2^2 + 2^2$$

$$16 = 4^2$$

$$18 = 4^2 + 1^2 + 1^2$$

$$20 = 4^2 + 2^2$$

$$22 = 2^2 + 2^2 + 3^2$$

$$24 = 4^2 + 2^2 + 2^2$$

\square

1.22 Exercise 22

For each integer n with $1 \leq n \leq 10$, $n^2 - n + 11$ is a prime number.

Proof. $1^2 - 1 + 11 = 11$ is prime.

$2^2 - 2 + 11 = 13$ is prime.

$3^2 - 3 + 11 = 17$ is prime.

$4^2 - 4 + 11 = 23$ is prime.

$5^2 - 5 + 11 = 31$ is prime.

$6^2 - 6 + 11 = 41$ is prime.

$7^2 - 7 + 11 = 53$ is prime.

$8^2 - 8 + 11 = 67$ is prime.

$9^2 - 9 + 11 = 83$ is prime.

$10^2 - 10 + 11 = 101$ is prime. □

Each of the statements in 23 – 26 is true. For each, (a) rewrite the statement with the quantification implicit as If _____, then _____, and (b) write the first sentence of a proof (the “starting point”) and the last sentence of a proof (the “conclusion to be shown”). (Note that you do not need to understand the statements in order to be able to do these exercises.)

1.23 Exercise 23

For every integer m , if $m > 1$ then $0 < \frac{1}{m} < 1$.

Proof. a. If an integer is greater than 1, then its reciprocal is between 0 and 1.

b. *Start of proof:* Suppose m is any integer such that $m > 1$. *Conclusion to be shown:* $0 < 1/m < 1$. □

1.24 Exercise 24

For every real number x , if $x > 1$ then $x^2 > x$.

Proof. a. If a real number is greater than 1, then its square is greater than itself.

b. *Start of proof:* Suppose x is any real number such that $x > 1$. *Conclusion to be shown:* $x^2 > x$. □

1.25 Exercise 25

For all integers m and n , if $mn = 1$ then $m = n = 1$ or $m = n = -1$.

Proof. a. If the product of two integers is 1, then either both are 1 or both are -1 .

b. *Start of proof:* Suppose m and n are any integers with $mn = 1$. *Conclusion to be shown:* $m = n = 1$ or $m = n = -1$. □

1.26 Exercise 26

For every real number x , if $0 < x < 1$ then $x^2 < x$.

Proof. a. If a real number is strictly between 0 and 1, then its square is less than itself.

b. *Start of proof:* Suppose x is any real number such that $0 < x < 1$. *Conclusion to be shown:* $x^2 < x$. \square

1.27 Exercise 27

Fill in the blanks in the following proof.

Theorem: For every odd integer n , n^2 is odd.

Proof: Suppose n is any (a) _____. By definition of odd, $n = 2k + 1$ for some integer k . Then

$$\begin{aligned} n^2 &= \text{(b) } (\text{_____})^2 && \text{by substitution} \\ &= 4k^2 + 4k + 1 && \text{by multiplying} \\ &= 2(2k^2 + 2k) + 1 && \text{by factoring out a 2} \end{aligned}$$

Now $2k^2 + 2k$ is an integer because it is a sum of products of integers. Therefore, n^2 equals $2 \cdot (\text{an integer}) + 1$, and so (c) _____ is odd by definition of odd.

Because we have not assumed anything about n except that it is an odd integer, it follows from the principle of (d) _____ that for every odd integer n , n^2 is odd.

Proof. (a) particular but arbitrarily chosen odd integer (b) $2k + 1$ (c) n^2 (d) universal generalization \square

In each of 28 – 31:

a. Rewrite the theorem in three different ways: as \forall _____, if _____, then _____; as \forall _____, _____ (without using the words if or then), and as If _____, then _____ (without using an explicit universal quantifier).

b. Fill in the blanks in the proof of the theorem.

1.28 Exercise 28

Theorem: The sum of any two odd integers is even.

Proof: Suppose m and n are any [particular but arbitrarily chosen] odd integers.

[We must show that $m + n$ is even.]

By (a) _____, $m = 2r + 1$ and $n = 2s + 1$ for some integers r and s . Then

$$\begin{aligned} m + n &= (2r + 1) + (2s + 1) && \text{by (b) _____} \\ &= 2r + 2s + 2 \\ &= 2(r + s + 1) && \text{by algebra} \end{aligned}$$

Let $u = r + s + 1$. Then u is an integer because r, s and 1 are integers and because (c) ____ .

Hence $m + n = 2u$, where u is an integer, and so, by (d) ____ , $m + n$ is even [as was to be shown].

Proof. a. \forall integers m and n , if m and n are odd then $m + n$ is even.

\forall odd integers m and n , $m + n$ is even.

If m and n are any odd integers, then $m + n$ is even.

b. (a) definition of odd, (b) substitution, (c) any sum of integers is an integer, (d) definition of even \square

1.29 Exercise 29

Theorem: The negative of any even integer is even.

Proof: Suppose n is any [particular but arbitrarily chosen] even integer.

[We must show that $-n$ is even.]

By (a) ____ , $n = 2k$ for some integer k .

Then

$$\begin{aligned} -n &= -(2k) \text{ by (b) } ______ \\ &= 2(-k) \text{ by algebra} \end{aligned}$$

Let $r = -k$. Then r is an integer because -1 and k are integers and (c) ____ .

Hence $-n = 2r$, where r is an integer, and so $-n$ is even by (d) ____ [as was to be shown].

Proof. a. \forall integer n , if n is even then $-n$ is even.

\forall even integer n , $-n$ is even.

If n is any even integer, then $-n$ is even.

b. (a) definition of even, (b) substitution, (c) any product of integers is an integer, (d) definition of even \square

1.30 Exercise 30

Theorem: The sum of any even integer and any odd integer is odd.

Proof: Suppose m is any even integer and n is any (a) ____ . By definition of even, $m = 2r$ for some (b) ____ , and by definition of odd, $n = 2s + 1$ for some integer s . By substitution and algebra,

$$m + n = \text{(c) } ______ = 2(r + s) + 1$$

Since r and s are integers, so is their sum $r + s$. Hence $m + n$ has the form twice some integer plus one, and so, by (d) _____ by definition of odd.

Proof. a. \forall integers m and n , if m is even and n is odd, then $m + n$ is odd.

\forall even integers m and odd integers n , $m + n$ is odd.

If m is any even integer and n is any odd integer, then $m + n$ is odd.

b. (a) any odd integer (b) integer r (c) $2r + (2s + 1)$ (d) $m + n$ is odd □

1.31 Exercise 31

Theorem: Whenever n is an odd integer, $5n^2 + 7$ is even.

Proof: Suppose n is any *[particular but arbitrarily chosen]* odd integer.

[We must show that $5n^2 + 7$ is even.]

By definition of odd, $n =$ (a) _____, for some integer k .

Then

$$\begin{aligned} 5n^2 + 7 &= \text{(b) } \underline{\hspace{2cm}} && \text{substitution} \\ &= 5(4k^2 + 4k + 1) + 7 \\ &= 20k^2 + 20k + 12 \\ &= 2(10k^2 + 10k + 6) && \text{by algebra} \end{aligned}$$

Let $t =$ (c) _____. Then t is an integer because products and sums of integers are integers.

Hence $5n^2 + 7 = 2t$, where t is an integer, and thus (d) _____ by definition of even *[as was to be shown]*.

Proof. a. \forall integer n , if n is odd, then $5n^2 + 7$ is even.

\forall odd integer n , $5n^2 + 7$ is even.

If n is any odd integer, then $5n^2 + 7$ is even.

b. (a) $2k + 1$ (b) $5(2k + 1)^2 + 7$ (c) $10k^2 + 10k + 6$ (d) $5n^2 + 7$ is even □

2 Exercise Set 4.2

Prove the statements in 1 – 11. In each case use only the definitions of the terms and the assumptions listed on page 161, not any previously established properties of odd and even integers. Follow the directions given in this section for writing proofs of universal statements.

2.1 Exercise 1

For every integer n , if n is odd then $3n + 5$ is even.

Proof. Suppose n is any [particular but arbitrarily chosen] odd integer.

[We must show that $3n + 5$ is even. By definition of even, this means we must show that $3n + 5 = 2 \cdot (\text{some integer})$.]

By definition of odd, $n = 2r + 1$, for some integer r .

Then

$$\begin{aligned} 3n + 5 &= 3(2r + 1) + 5 && \text{by substitution} \\ &= 6r + 3 + 5 \\ &= 6r + 8 \\ &= 2(3r + 4) && \text{by algebra} \end{aligned}$$

[Idea for the rest of the proof: We want to show that $3n + 5 = 2 \cdot (\text{some integer})$. At this point we know that $3n + 5 = 2(3r + 4)$. So is $3r + 4$ an integer? Yes, because products and sums of integers are integers.]

Let $k = 3r + 4$.

Then k is an integer because products and sums of integers are integers.

Hence $3n + 5 = 2(3r + 4) = 2k$ where k is an integer. Hence by definition of even $3n + 5$ is even [as was to be shown]. \square

2.2 Exercise 2

For every integer m , if m is even then $3m + 5$ is odd.

Proof. Suppose m is any [particular but arbitrarily chosen] even integer.

[We must show that $3m + 5$ is odd. By definition of odd, this means we must show that $3m + 5 = 2 \cdot (\text{some integer}) + 1$.]

By definition of even, $m = 2r$, for some integer r .

Then

$$\begin{aligned} 3m + 5 &= 3(2r) + 5 && \text{by substitution} \\ &= 6r + 5 \\ &= 6r + 4 + 1 \\ &= 2(3r + 2) + 1 && \text{by algebra} \end{aligned}$$

[Idea for the rest of the proof: We want to show that $3m + 5 = 2 \cdot (\text{some integer}) + 1$. At this point we know that $3m + 5 = 2(3r + 2) + 1$. So is $3r + 2$ an integer? Yes, because products and sums of integers are integers.]

Let $k = 3r + 2$.

Then k is an integer because products and sums of integers are integers.

Hence $3m + 5 = 2(3r + 2) + 1 = 2k + 1$ where k is an integer. Hence by definition of odd $3n + 5$ is odd [as was to be shown]. \square

2.3 Exercise 3

For every integer n , $2n - 1$ is odd.

Proof. Suppose n is any [particular but arbitrarily chosen] integer.

[We must show that $2n - 1$ is odd. By definition of odd, this means we must show that $2n - 1 = 2 \cdot (\text{some integer}) + 1$.]

Then

$$\begin{aligned} 2n - 1 &= 2n - 2 + 2 - 1 && \text{because } -2 + 2 = 0 \\ &= 2(n - 1) + 2 - 1 \\ &= 2(n - 1) + 1 && \text{by algebra} \end{aligned}$$

Let $k = n - 1$.

Then k is an integer because the difference of two integers (n and 1) is an integer.

Hence $2n - 1 = 2(n - 1) + 1 = 2k + 1$ where k is an integer, and thus by definition of odd $2n - 1$ is odd [as was to be shown]. \square

2.4 Exercise 4

The difference of any even integer minus any odd integer is odd.

Proof. Suppose a is any even integer and b is any odd integer. [We must show that $a - b$ is odd.] By definition of even and odd, $a = 2r$ and $b = 2s + 1$, for some integers r, s . By substitution and algebra,

$$a - b = 2r - (2s + 1) = 2r - 2s - 1 = 2r - 2s - 2 + 2 - 1 = 2(r - s - 1) + 1$$

Let $t = r - s - 1$. Then t is an integer because differences of integers are integers.

Thus $a - b = 2t + 1$ where t is an integer, and so by definition of odd $a - b$ is odd [as was to be shown]. \square

2.5 Exercise 5

If a and b are any odd integers, then $a^2 + b^2$ is even.

Proof. Suppose a, b are any [particular but arbitrarily chosen] odd integers.

[We must show that $a^2 + b^2$ is even.]

By definition of odd, $a = 2r + 1$ and $b = 2s + 1$, for some integers r, s .

Then

$$\begin{aligned}
a^2 + b^2 &= (2r + 1)^2 + (2s + 1)^2 && \text{by substitution} \\
&= (4r^2 + 4r + 1) + (4s^2 + 4s + 1) && \text{by multiplying} \\
&= 4r^2 + 4r + 4s^2 + 4s + 2 && \text{by adding} \\
&= 2(2r^2 + 2r + 2s^2 + 2s + 1) && \text{by factoring out}
\end{aligned}$$

Let $k = 2r^2 + 2r + 2s^2 + 2s + 1$.

Then k is an integer because squares, products and sums of integers are integers.

Hence $a^2 + b^2 = 2k$ where k is an integer, and thus by definition of even $a^2 + b^2$ is even [as was to be shown]. \square

2.6 Exercise 6

If k is any odd integer and m is any even integer, then $k^2 + m^2$ is odd.

Proof. Suppose k is any odd integer and m is any even integer.

[We must show that $k^2 + m^2$ is odd.]

By definition of odd and even, $k = 2a + 1$ and $m = 2b$, for some integers a, b . Then

$$\begin{aligned}
k^2 + m^2 &= (2a + 1)^2 + (2b)^2 && \text{by substitution} \\
&= 4a^2 + 4a + 1 + 4b^2 \\
&= 4(a^2 + a + b^2) + 1 \\
&= 2(2a^2 + 2a + 2b^2) + 1 && \text{by algebra}
\end{aligned}$$

But $2a^2 + 2a + 2b^2$ is an integer because it is a sum of products of integers. Thus $k^2 + m^2$ is twice an integer plus 1, and so $k^2 + m^2$ is odd [as was to be shown]. \square

2.7 Exercise 7

The difference between the squares of any two consecutive integers is odd.

Proof. Suppose m and n are any [particular but arbitrarily chosen] two consecutive integers.

[We must show that $m^2 - n^2$ (or $n^2 - m^2$) is odd.]

By definition of consecutive, $m = k$ and $n = k + 1$, for some integer k .

Then

$$\begin{aligned}
m^2 - n^2 &= k^2 - (k + 1)^2 && \text{by substitution} \\
&= k^2 - (k^2 + 2k + 1) \\
&= k^2 - k^2 - 2k - 1 \\
&= -2k - 1 \\
&= -2k - 2 + 2 - 1 \\
&= 2(-k - 1) + 1 && \text{by algebra}
\end{aligned}$$

Let $r = -k - 1$. Then r is an integer because it is a difference of integers.

Hence $m^2 - n^2 = 2r + 1$ where r is an integer, and thus by definition of odd $m^2 - n^2$ is odd [as was to be shown]. \square

2.8 Exercise 8

For any integers m and n , if m is even and n is odd then $5m + 3n$ is odd.

Proof. Suppose m is any even integer and n is any odd integer.

[We must show that $5m + 3n$ is odd.]

By definition of even and odd, $m = 2r$ and $n = 2s + 1$, for some integers r, s .

Then

$$\begin{aligned} 5m + 3n &= 5(2r) + 3(2s + 1) && \text{by substitution} \\ &= 10r + 6s + 3 \\ &= 10r + 6s + 2 + 1 \\ &= 2(5r + 3s + 1) + 1 && \text{by algebra} \end{aligned}$$

Let $k = 5r + 3s + 1$.

Then k is an integer because it is a sum of products of integers.

Hence $5m + 3n = 2k + 1$ where k is an integer, and thus by definition of odd $5m + 3n$ is odd [as was to be shown]. \square

2.9 Exercise 9

If an integer greater than 4 is a perfect square, then the immediately preceding integer is not prime.

Proof. Suppose n is any integer greater than 4 that is a perfect square.

[We must show that $n - 1$ is not prime, in other words, $n - 1$ is composite.]

By definition of perfect square, $n = k^2$, for some integer k .

Without loss of generality, we may assume $k > 0$, because $n = k^2 = (-k)^2$, and if k is negative, we can replace it with $-k$ which is positive.

Since $n > 4$ we have $n - 4 > 0$. So $k^2 - 4 > 0$. So $(k - 2)(k + 2) > 0$. So either $k - 2$ and $k + 2$ are both negative, or they are both positive. Since $k > 0$, $k + 2 > 2 > 0$, so they have to be both positive. Therefore, $k - 2 > 0$ so $k > 2$.

Then

$$\begin{aligned} n - 1 &= k^2 - 1 && \text{by substitution} \\ &= (k - 1)(k + 1) && \text{by algebra} \end{aligned}$$

Since $k > 2$ we have both $k - 1 > 1$ and $k + 1 > 3 > 1$.

Hence $n - 1$ is a product of two positive integers both greater than 1, and thus by definition of composite $n - 1$ is composite [as was to be shown]. \square

2.10 Exercise 10

If n is any even integer, then $(-1)^n = 1$.

Proof. Suppose n is any even integer. [We must show that $(-1)^n = 1$.]

By definition of even, $n = 2k$, for some integer k .

Then by the laws of exponents from algebra $(-1)^n = (-1)^{2k} = ((-1)^2)^k = 1^k = 1$, [as was to be shown]. \square

2.11 Exercise 11

If n is any odd integer, then $(-1)^n = -1$.

Proof. Suppose n is any odd integer. [We must show that $(-1)^n = -1$.]

By definition of odd, $n = 2k + 1$, for some integer k .

Then by the laws of exponents from algebra

$$(-1)^n = (-1)^{2k+1} = (-1)^{2k} \cdot (-1) = ((-1)^2)^k \cdot (-1) = 1^k \cdot (-1) = 1 \cdot (-1) = -1$$

[as was to be shown]. \square

Prove that the statements in 12 – 14 are false.

2.12 Exercise 12

There exists an integer $m \geq 3$ such that $m^2 - 1$ is prime.

Proof. To prove the given statement is false, we prove that its negation is true.

The negation of the statement is “For every integer $m \geq 3$, $m^2 - 1$ is not prime.”

Proof of the negation: Suppose m is any integer with $m \geq 3$.

By basic algebra, $m^2 - 1 = (m - 1)(m + 1)$.

Because $m \geq 3$, both $m - 1$ and $m + 1$ are positive integers greater than 1, and each is smaller than $m^2 - 1$.

So $m^2 - 1$ is a product of two smaller positive integers, each greater than 1, and hence $m^2 - 1$ is not prime. \square

2.13 Exercise 13

There exists an integer n such that $6n + 27$ is prime.

Proof. To prove the given statement is false, we prove that its negation is true.

The negation of the statement is “For every integer n , $6n + 27$ is not prime. In other words, $6n + 27$ is composite.”

Proof of the negation: Suppose n is any integer. By basic algebra, $6n + 27 = 3(2n + 9)$. Hence $6n + 27$ is the product of two integers greater than 1. Therefore by definition of composite, $6n + 27$ is composite. \square

2.14 Exercise 14

There exists an integer $k \geq 4$ such that $2k^2 - 5k + 2$ is prime.

Proof. To prove the given statement is false, we prove that its negation is true.

The negation of the statement is “For every integer $k \geq 4$, $2k^2 - 5k + 2$ is composite.”

Proof of the negation: Suppose k is any integer.

By basic algebra, $2k^2 - 5k + 2 = (2k - 1)(k - 2)$.

Because $k \geq 4$, $2k - 1 \geq 7$ and $k - 2 \geq 2$. So both $2k - 1$ and $k - 2$ are integers greater than 1.

Hence $2k^2 - 5k + 2$ is the product of two integers greater than 1. Therefore by definition of composite, $2k^2 - 5k + 2$ is composite. \square

Find the mistakes in the “proofs” shown in 15 – 19.

2.15 Exercise 15

Theorem: For every integer k , if $k > 0$ then $k^2 + 2k + 1$ is composite.

“Proof: For $k = 2$, $k > 0$ and $k^2 + 2k + 1 = 2^2 + 2 \cdot 2 + 1 = 9$. And since $9 = 3 \cdot 3$, then 9 is composite. Hence the theorem is true.”

Proof. The incorrect proof just shows the theorem to be true in the one case where $k = 2$. A real proof must show that it is true for *every* integer $k > 0$. \square

2.16 Exercise 16

Theorem: The difference between any odd integer and any even integer is odd.

“Proof: Suppose n is any odd integer, and m is any even integer. By definition of odd, $n = 2k + 1$ where k is an integer, and by definition of even, $m = 2k$ where k is an integer. Then $n - m = (2k + 1) - 2k = 1$, and 1 is odd. Therefore, the difference between any odd integer and any even integer is odd.”

Proof. The mistake in the “proof” is that the same symbol, k , is used to represent two different quantities. By setting $m = 2k$ and $n = 2k + 1$, the proof implies that $n = m + 1$, and thus it deduces the conclusion only for this one situation. When $m = 4$ and $n = 17$, for instance, the computations in the proof indicate that $n - m = 1$, but actually $n - m = 13$. In other words, the proof does not deduce the conclusion for an arbitrarily chosen even integer m and odd integer n , and hence it is invalid. \square

2.17 Exercise 17

Theorem: For every integer k , if $k > 0$ then $k^2 + 2k + 1$ is composite.

Proof: Suppose k is any integer such that $k > 0$. If $k^2 + 2k + 1$ is composite, then $k^2 + 2k + 1 = rs$ for some integers r and s such that

$$1 < r < k^2 + 2k + 1 \text{ and } 1 < s < k^2 + 2k + 1.$$

Since $k^2 + 2k + 1 = rs$ and both r and s are strictly between 1 and $k^2 + 2k + 1$, then $k^2 + 2k + 1$ is not prime. So $k^2 + 2k + 1$ is composite as was to be shown.

Proof. This incorrect proof assumes what is to be proved. The word since in the third sentence is completely unjustified. The second sentence tells only what happens if $k^2 + 2k + 1$ is composite. But at that point in the proof, it has not been established that $k^2 + 2k + 1$ is composite. In fact, that is exactly what is to be proved. \square

2.18 Exercise 18

Theorem: The product of any even integer and any odd integer is even.

Proof: Suppose m is any even integer and n is any odd integer. If $m \cdot n$ is even, then by definition of even there exists an integer r such that $m \cdot n = 2r$.

Also since m is even, there exists an integer p such that $m = 2p$, and since n is odd there exists an integer q such that $n = 2q + 1$.

Thus $mn = (2p)(2q + 1) = 2r$, where r is an integer. By definition of even, then, $m \cdot n$ is even, as was to be shown.”

Proof. The issue is just like in Exercise 17. The proof uses the r value without establishing the existence of r first.

“If $m \cdot n$ is even...” has an unjustified assumption because we haven’t proved that $m \cdot n$ is even yet (that’s what we are *trying to prove*), so its conclusion “... $m \cdot n = 2r$ ” has not been proven.

Therefore the part “ $mn = (2p)(2q + 1) = 2r$, where r is an integer” is unjustified as well.

Discussion:

This is a fairly common form of circular reasoning: assuming what we have to prove. It happens because, at the beginning of the proof, we want to mention to the reader what we want to prove.

What we want to prove has a short, condensed definition (in this case “being even”), so we write out the full definition of what it is that we are *trying to prove* (in this case “the existence of an integer r such that $\dots = 2r$ ”). Again, the purpose of this is to articulate to the reader what we are *trying to prove*.

But then we forget that and continue as if that was already an established fact. The act of writing out the full definition of what we are trying to prove is not the same as actually having proved it.

Using “if $m \cdot n$ is even...” in this case is the problem; it has the *feeling* of using modus ponens on an already established implication with an established premise. But we are just writing out the full definition, instead of using modus ponens.

So it would be better to write: “*We want to prove that $m \cdot n$ is even. In other words, we want to prove that there is an integer r such that $m \cdot n = 2r$.*” Using the words “We want to prove that...” instead of “If...” goes a long way to avoid this common mistake. This way we can “unpack” the definition of what we are trying to prove without assuming it.

Another related problem is to first unpack the definition of what we are trying to prove, then try to “prove backwards”. Say we want to prove A , and we unpack the definition to B . So we have to prove B . But instead, we start by assuming B is true, and apply some algebra or logic to it, to arrive at something else, say E , that is true:

$$A \rightarrow \text{unpack definition} \rightarrow B \rightarrow \text{middle steps} \rightarrow C \rightarrow D \rightarrow E = \text{something true!}$$

But this would only prove that B implies E . In order to establish the truth of B (and hence of A), we would actually have to prove that E implies B ! So all the “steps” from B to E would have to be “reversible”, in other words, logical equivalences (biconditionals):

$$A \leftrightarrow \text{unpack definition} \leftrightarrow B \leftrightarrow \text{middle steps} \leftrightarrow C \leftrightarrow D \leftrightarrow E = \text{something true!}$$

But that is rarely the case! □

2.19 Exercise 19

Theorem: The sum of any two even integers equals $4k$ for some integer k .

“Proof: Suppose m and n are any two even integers. By definition of even, $m = 2k$ for some integer k and $n = 2k$ for some integer k . By substitution,

$$m + n = 2k + 2k = 4k.$$

This is what was to be shown.”

Proof. The problem here is the same as in Exercise 16. The mistake in the “proof” is that the same symbol, k , is used to represent two different quantities. By setting $m = 2k$

and $n = 2k$, the proof implies that $n = m$, and thus it deduces the conclusion only for this one situation.

When $m = 4$ and $n = 20$, for instance, the proof indicates that $n = m = 4$, but actually $n = 20$. In other words, the proof does not deduce the conclusion for an arbitrarily chosen even integer m and an arbitrarily chosen even integer n , and hence it is invalid. \square

In 20 – 38 determine whether the statement is true or false. Justify your answer with a proof or a counterexample, as appropriate. In each case use only the definitions of the terms and the assumptions listed on page 161, not any previously established properties.

2.20 Exercise 20

The product of any two odd integers is odd.

Proof. True. Suppose m and n are any odd integers. [We must show that mn is odd.] By definition of odd, $n = 2r + 1$ and $m = 2s + 1$ for some integers r and s .

Then

$$\begin{aligned} mn &= (2r + 1)(2s + 1) && \text{by substitution} \\ &= 4rs + 2r + 2s + 1 \\ &= 2(2rs + r + s) + 1 && \text{by algebra} \end{aligned}$$

Now $2rs + r + s$ is an integer because products and sums of integers are integers and 2 , r , and s are all integers. Hence $mn = 2 \cdot (\text{some integer}) + 1$, and so, by definition of odd, mn is odd. \square

2.21 Exercise 21

The negative of any odd integer is odd.

Proof. True. Assume n is any odd integer. We want to prove $-n$ is odd.

By definition of odd, $n = 2r + 1$ for some integer r .

Then $-n = -(2r + 1) = -2r - 1 = -2r - 2 + 2 - 1 = 2(-r - 1) + 1$.

Let $k = -r - 1$. Then k is an integer because it is the difference of two integers.

Therefore $-n = 2k + 1$ where k is an integer, hence by definition of odd, $-n$ is odd. \square

2.22 Exercise 22

For all integers a and b , $4a + 5b + 3$ is even.

Proof. False. Counterexample: Let $a = 1$ and $b = 0$.

Then $4a + 5b + 3 = 4 \cdot 1 + 5 \cdot 0 + 3 = 7$, which is odd.

[This is one counterexample among many. Can you find a way to characterize all counterexamples?] □

2.23 Exercise 23

The product of any even integer and any integer is even.

Proof. True. Suppose m is any even integer and n is any integer. [We want to prove $m \cdot n$ is even.]

By definition of even, $m = 2k$ for some integer k .

Then, $m \cdot n = (2k) \cdot n = 2kn = 2(kn)$.

Let $r = kn$. Then r is an integer because it is the product of two integers.

Therefore $m \cdot n = 2r$ where r is an integer. So by definition of even, $m \cdot n$ is even. □

2.24 Exercise 24

If a sum of two integers is even, then one of the summands is even. (In the expression $a + b$, a and b are called **summands**.)

Proof. False. Counterexample: Let $m = 1$ and $n = 3$.

Then $m + n = 4$ is even, but neither summand m nor summand n is even. □

2.25 Exercise 25

The difference of any two even integers is even.

Proof. True. Assume m and n are any two even integers. [We want to prove $m - n$ is even.]

By definition of even, $m = 2r$ and $n = 2s$ for some integers r, s .

Then $m - n = 2r - 2s = 2(r - s)$. Let $k = r - s$. Then k is an integer because it is the difference of two integers.

Therefore $m - n = 2k$ where k is an integer. So $m - n$ is even by definition of even. □

2.26 Exercise 26

For all integers a, b , and c , if a, b , and c are consecutive, then $a + b + c$ is even.

Proof. False. Counterexample: Let $a = 2, b = 3, c = 4$. They are consecutive integers but $a + b + c = 9$ which is not even. □

2.27 Exercise 27

The difference of any two odd integers is even.

Proof. True. Assume m, n are any two odd integers. [We want to prove $m - n$ is even.]

By definition of odd, $m = 2r + 1, n = 2s + 1$ for some integers r, s .

Then $m - n = 2r + 1 - (2s + 1) = 2r + 1 - 2s - 1 = 2r - 2s = 2(r - s)$.

Let $k = r - s$. Then k is an integer because it is the difference of two integers.

So $m - n = 2k$ where k is an integer. Hence $m - n$ is even by definition of even. \square

2.28 Exercise 28

For all integers n and m , if $n - m$ is even then $n^3 - m^3$ is even.

Proof. True. Assume n, m are any integers such that $n - m$ is even. [Want to prove that $n^3 - m^3$ is even.]

By definition of even, $n - m = 2r$ for some integer r . By algebra,

$$n^3 - m^3 = (n - m)(n^2 + nm + m^2) = 2r(n^2 + nm + m^2) = 2(r(n^2 + nm + m^2)).$$

Let $k = r(n^2 + nm + m^2)$. Then k is an integer because it is a sum and product of integers.

So $n^3 - m^3 = 2k$ where k is an integer. So by definition of even, $n^3 - m^3$ is even. \square

2.29 Exercise 29

For every integer n , if n is prime then $(-1)^n = -1$.

Proof. False. Counterexample: Let $n = 2$.

Then n is prime, but $(-1)^n = (-1)^2 = 1 \neq -1$. \square

2.30 Exercise 30

For every integer m , if $m > 2$ then $m^2 - 4$ is composite.

Proof. False. Counterexample: Let $m = 3$. Then $m^2 - 4 = 3^2 - 4 = 9 - 4 = 5$ is prime. not composite. \square

2.31 Exercise 31

For every integer n , $n^2 - n + 11$ is a prime number.

Proof. False. Let $n = 11$. Then $n^2 - n + 11 = 11^2 - 11 + 11 = 11^2$ is not prime. \square

2.32 Exercise 32

For every integer n , $4(n^2 + n + 1) - 3n^2$ is a perfect square.

Proof. True. Suppose n is any integer. Then by algebra

$$4(n^2 + n + 1) - 3n^2 = 4n^2 + 4n + 4 - 3n^2 = n^2 + 4n + 4 = (n + 2)^2$$

Now $(n + 2)^2$ is a perfect square because $n + 2$ is an integer (being a sum of n and 2). Hence $4(n^2 + n + 1) - 3n^2$ is a perfect square, as was to be shown. \square

2.33 Exercise 33

Every positive integer can be expressed as a sum of three or fewer perfect squares.

Proof. False. Counterexample: 7 cannot be written as a sum of three or fewer perfect squares: $7 = 2^2 + 1^2 + 1^2 + 1^2$. \square

2.34 Exercise 34

(Two integers are **consecutive** if, and only if, one is one more than the other.) Any product of four consecutive integers is one less than a perfect square.

Proof. True. Suppose a, b, c, d are any four consecutive integers. [*Want to prove: there is an integer k such that $abcd = k^2 - 1$.*]

By definition of consecutive, there is an integer n such that $a = n, b = n + 1, c = n + 2, d = n + 3$. Then

$$abcd = n(n + 1)(n + 2)(n + 3) = n(n + 3)(n + 1)(n + 2) = (n^2 + 3n)(n^2 + 3n + 2)$$

[*Here we notice a pattern. The two factors differ by 2. So it is reminiscent of $(x - 1)(x + 1) = x^2 - 1^2$ isn't it?*]

By some more algebra,

$$(n^2 + 3n)(n^2 + 3n + 2) = (n^2 + 3n + 1 - 1)(n^2 + 3n + 1 + 1) = (n^2 + 3n + 1)^2 - 1^2$$

Let $k = n^2 + 3n + 1$. Then k is an integer because it is a sum and product of integers. Therefore $abcd = k^2 - 1$ where k is an integer, [*as was to be shown*]. \square

2.35 Exercise 35

If m and n are any positive integers and mn is a perfect square, then m and n are perfect squares.

Proof. False. Counterexample: let $m = n = 2$. Then $mn = 2^2$ is a perfect square. But neither m nor n is a perfect square. \square

2.36 Exercise 36

The difference of the squares of any two consecutive integers is odd.

Proof. True. Assume a, b are any two consecutive integers.

By definition of consecutive, $a = n$ and $b = n + 1$ for some integer n .

Then $b^2 - a^2 = (n + 1)^2 - n^2 = n^2 + 2n + 1 - n^2 = 2n + 1$.

So $b^2 - a^2 = 2k + 1$ where n is an integer. Therefore by definition of odd, $b^2 - a^2$ is odd. \square

2.37 Exercise 37

For all nonnegative real numbers a and b , $\sqrt{ab} = \sqrt{a}\sqrt{b}$. (Note that if x is a nonnegative real number, then there is a unique nonnegative real number y , denoted \sqrt{x} , such that $y^2 = x$.)

Proof. True. Assume a and b are any two nonnegative real numbers. By the information given to us in the parentheses:

1. There is a unique nonnegative real number denoted \sqrt{ab} such that $(\sqrt{ab})^2 = ab$.
2. There is a unique nonnegative real number denoted \sqrt{a} such that $(\sqrt{a})^2 = a$.
3. There is a unique nonnegative real number denoted \sqrt{b} such that $(\sqrt{b})^2 = b$.

Since $ab = a \cdot b$, we have by substitution: $(\sqrt{ab})^2 = (\sqrt{a})^2 \cdot (\sqrt{b})^2$.

By algebra, $(\sqrt{ab})^2 = [(\sqrt{a}) \cdot (\sqrt{b})]^2 = (\sqrt{a}\sqrt{b})^2$. Therefore $(\sqrt{ab})^2 - (\sqrt{a}\sqrt{b})^2 = 0$.

By factoring we get $(\sqrt{ab} - \sqrt{a}\sqrt{b})(\sqrt{ab} + \sqrt{a}\sqrt{b}) = 0$.

So: either $\sqrt{ab} - \sqrt{a}\sqrt{b} = 0$, or $\sqrt{ab} + \sqrt{a}\sqrt{b} = 0$ (by the Zero Product Property T11).

If $\sqrt{ab} - \sqrt{a}\sqrt{b} = 0$, then $\sqrt{ab} = \sqrt{a}\sqrt{b}$ [as was to be shown.]

If $\sqrt{ab} + \sqrt{a}\sqrt{b} = 0$, then since both \sqrt{ab} and $\sqrt{a}\sqrt{b}$ are nonnegative, they must be both 0, hence $\sqrt{ab} = \sqrt{a}\sqrt{b}$ again [as was to be shown.] \square

2.38 Exercise 38

For all nonnegative real numbers a and b , $\sqrt{a+b} = \sqrt{a} + \sqrt{b}$.

Proof. False. Counterexample: Let $a = b = 1$. Then

$$\sqrt{a+b} = \sqrt{1+1} = \sqrt{2} \neq 2 = 1+1 = \sqrt{1} + \sqrt{1} = \sqrt{a} + \sqrt{b}$$

\square

2.39 Exercise 39

Suppose that integers m and n are perfect squares. Then $m + n + 2\sqrt{mn}$ is also a perfect square. Why?

Proof. Assume m and n are perfect squares (so they are nonnegative real numbers). By definition of perfect square, $m = r^2$ and $n = s^2$ for some integers r, s . Using Exercise 37 $\sqrt{mn} = \sqrt{m}\sqrt{n}$, we get:

$$m + n + 2\sqrt{mn} = r^2 + s^2 + 2\sqrt{m}\sqrt{n} = r^2 + s^2 + 2rs = (r + s)^2.$$

Let $k = r + s$. k is an integer because it is a sum of integers. So $m + n + 2\sqrt{mn} = k^2$ where k is an integer, therefore $m + n + 2\sqrt{mn}$ is a perfect square by definition. □

2.40 Exercise 40

If p is a prime number, must $2^p - 1$ also be prime? Prove or give a counterexample.

Proof. No. Counterexample: $p = 11$ is prime, but $2^p - 1 = 2^{11} - 1 = 2047 = 13 \cdot 89$ is not prime. □

2.41 Exercise 41

If n is a nonnegative integer, must $2^{2n} + 1$ be prime? Prove or give a counterexample.

Proof. No. Counterexample: Let $n = 3$. Then $2^{2n} + 1 = 2^6 + 1 = 65 = 13 \cdot 5$ is not prime. □

3 Exercise Set 4.3

The numbers in 1 – 7 are all rational. Write each number as a ratio of two integers.

3.1 Exercise 1

$$-\frac{35}{6}$$

Proof. $-\frac{35}{6} = \frac{-35}{6}$ □

3.2 Exercise 2

4.6037

Proof. $4.6037 = \frac{46037}{10000}$ □

3.3 Exercise 3

$$\frac{4}{5} + \frac{2}{9}$$

Proof. $\frac{4}{5} + \frac{2}{9} = \frac{4 \cdot 9 + 5 \cdot 2}{5 \cdot 9} = \frac{46}{45}$

□

3.4 Exercise 4

0.37373737...

Proof. Let $x = 0.373737 \dots$

Then $100x = 37.373737 \dots$, so $100x - x = 37.373737 \dots - 0.373737 \dots = 37$.

Thus $99x = 37$, and hence $x = \frac{37}{99}$.

□

3.5 Exercise 5

0.56565656...

Proof. Let $x = 0.565656 \dots$

Then $100x = 56.565656 \dots$ and so $100x - x = 56.565656 \dots - 0.565656 \dots = 56$.

Thus $99x = 56$, and hence $x = \frac{56}{99}$.

□

3.6 Exercise 6

320.5492492492...

Proof. Let $x = 320.5492492492 \dots$

Then $10000x = 3205492.492492 \dots$ and $10x = 3205.492492 \dots$, so

$10000x - 10x = 3205492.492492 \dots - 3205.492492 \dots = 3205492 - 3205 = 3202287$.

Thus $9990x = 3202287$, and hence $x = \frac{3202287}{9990}$.

□

3.7 Exercise 7

52.4672167216721...

Proof. Let $x = 52.467216721 \dots$

Then $100000x = 5246721.67216721 \dots$ and $10x = 524.67216721 \dots$, so

$100000x - 10x = 5246721.67216721 \dots - 524.67216721 \dots = 5246721 - 524 = 5246197$.

Thus $99990x = 5246197$, and hence $x = \frac{5246197}{99990}$.

□

3.8 Exercise 8

The zero product property, says that if a product of two real numbers is 0, then one of the numbers must be 0.

3.8.1 (a)

Write this property formally using quantifiers and variables.

Proof. \forall real numbers x, y , if $xy = 0$ then $x = 0$ or $y = 0$. □

3.8.2 (b)

Write the contrapositive of your answer to part (a).

Proof. \forall real numbers x, y , if $x \neq 0$ and $y \neq 0$ then $xy \neq 0$. □

3.8.3 (c)

Write an informal version (without quantifier symbols or variables) for your answer to part (b).

Proof. The product of two nonzero real numbers is nonzero. □

3.9 Exercise 9

Assume that a and b are both integers and that $a \neq 0$ and $b \neq 0$. Explain why $(b - a)/(ab^2)$ must be a rational number.

Proof. Given that a and b are integers, both $b - a$ and ab^2 are integers (since differences and products of integers are integers). Also, by the zero product property, $ab^2 \neq 0$ because neither a nor b is zero. Hence $(b - a)/(ab^2)$ is a quotient of two integers with a nonzero denominator, and so it is rational. □

3.10 Exercise 10

Assume that m and n are both integers and that $n \neq 0$. Explain why $(5m - 12n)/(4n)$ must be a rational number.

Proof. Given that m and n are integers, both $5m - 12n$ and $4n$ are integers (since differences and products of integers are integers). Also, by the zero product property, $4n \neq 0$ because neither 4 nor n is zero. Hence $(5m - 12n)/(4n)$ is a quotient of two integers with a nonzero denominator, and so it is rational. □

3.11 Exercise 11

Prove that every integer is a rational number.

Proof. Suppose n is any [particular but arbitrarily chosen] integer. Then $n = n \cdot 1$, and so $n = n/1$ by dividing both sides by 1. Now n and 1 are both integers, and $1 \neq 0$. Hence n can be written as a quotient of integers with a nonzero denominator, and so n is rational. \square

3.12 Exercise 12

Let S be the statement “The square of any rational number is rational.” A formal version of S is “For every rational number r , r^2 is rational.” Fill in the blanks in the proof for S .

Proof: Suppose that r is (a) _____. By definition of rational, $r = a/b$ for some (b) _____ with $b \neq 0$. By substitution,

$$r^2 = (c) \text{_____} = a^2/b^2.$$

Since a and b are both integers, so are the products a^2 and (d) _____. Also $b^2 \neq 0$ by the (e) _____. Hence r^2 is a ratio of two integers with a nonzero denominator, and so (f) _____ by definition of rational.

Proof. (a) any [particular but arbitrarily chosen] rational number

(b) integers a and b

(c) $(a/b)^2$

(d) b^2

(e) zero product property

(f) r^2 is rational \square

3.13 Exercise 13

Consider the following statement: The negative of any rational number is rational.

3.13.1 (a)

Write the statement formally using a quantifier and a variable.

Proof. \forall real number r , if r is rational then $-r$ is rational.

Or: $\forall r$, if r is a rational number then $-r$ is rational.

Or: \forall rational number r , $-r$ is rational. \square

3.13.2 (b)

Determine whether the statement is true or false and justify your answer.

Proof. The statement is true. Suppose r is a [particular but arbitrarily chosen] rational number. [We must show that $-r$ is rational.] By definition of rational, $r = a/b$ for some integers a and b with $b \neq 0$. Then by substitution and algebra,

$$-r = -\frac{a}{b} = \frac{-a}{b}$$

Now since a is an integer, so is $-a$ (being the product of -1 and a). Hence $-r$ is a quotient of integers with a nonzero denominator, and so $-r$ is rational [as was to be shown]. \square

3.14 Exercise 14

Consider the statement: The cube of any rational number is a rational number.

3.14.1 (a)

Write the statement formally using a quantifier and a variable.

Proof. \forall rational r , r^3 is rational. \square

3.14.2 (b)

Determine whether the statement is true or false and justify your answer.

Proof. The statement is true. Suppose r is a [particular but arbitrarily chosen] rational number. [We must show that r^3 is rational.] By definition of rational, $r = a/b$ for some integers a and b with $b \neq 0$. Then by substitution and algebra,

$$r^3 = \left(\frac{a}{b}\right)^3 = \frac{a^3}{b^3}$$

Now since a, b are integers, so are a^3 and b^3 (being the products of a and b). Moreover, since $b \neq 0$, by the Zero Product Property, $b^3 \neq 0$.

Hence r^3 is a quotient of integers with a nonzero denominator, and so r^3 is rational [as was to be shown]. \square

Determine which of the statements in 15 – 19 are true and which are false. prove each true statement directly from the definitions, and give a counterexample for each false statement. For a statement that is false, determine whether a small change would make it true. If so, make the change and prove the new statement. Follow the directions for writing proofs on page 173.

3.15 Exercise 15

The product of any two rational numbers is a rational number.

Proof. Suppose r and s are rational numbers. By definition of rational, $r = a/b$ and $s = c/d$ for some integers a, b, c , and d with $b \neq 0$ and $d \neq 0$. Then by substitution and algebra,

$$rs = \frac{a}{b} \cdot \frac{c}{d} = \frac{ac}{bd}$$

Now ac and bd are both integers (being products of integers) and $bd \neq 0$ (by the zero product property). Hence rs is a quotient of integers with a nonzero denominator, and so, by definition of rational, rs is rational. \square

3.16 Exercise 16

The quotient of any two rational numbers is a rational number.

Proof. Counterexample: Let r be any rational number and $s = 0$. Then r and s are both rational, but the quotient of r divided by s is not a real number and therefore is not a rational number.

Revised statement to be proved: For all rational numbers r and s , if $s \neq 0$ then r/s is rational.

Suppose r, s are rational numbers such that $s \neq 0$. [*Want to prove r/s is rational.*]

By definition of rational, $r = a/b$ and $s = c/d$ for some integers a, b, c, d where $b \neq 0, d \neq 0$. Since $s \neq 0$ we also have $c \neq 0$. Then by algebra

$$\frac{r}{s} = \frac{a/b}{c/d} = \frac{ad}{bc}$$

Now ad and bc are integers because they are products of integers. Since $b \neq 0$ and $c \neq 0$, by Zero Product Property $bc \neq 0$.

Let $m = ad$ and $n = bc$. So m and n are integers with $n \neq 0$, and $r/s = m/n$. Therefore by definition of rational, r/s is rational. \square

3.17 Exercise 17

The difference of any two rational numbers is a rational number.

Proof. True. Suppose r, s are rational numbers. [*Want to prove $r - s$ is rational.*]

By definition of rational, $r = a/b$ and $s = c/d$ for some integers a, b, c, d where $b \neq 0, d \neq 0$. Then by algebra

$$r - s = \frac{a}{b} - \frac{c}{d} = \frac{ad - bc}{bd}$$

Now $ad - bc$ and bd are integers because they are products and differences of integers. Since $b \neq 0$ and $d \neq 0$, by Zero Product Property $bd \neq 0$.

Let $m = ad - bc$ and $n = bd$. So m and n are integers with $n \neq 0$, and $r - s = m/n$. Therefore by definition of rational, $r - s$ is rational. \square

3.18 Exercise 18

If r and s are any two rational numbers, then $\frac{r+s}{2}$ is rational.

Proof. True. The proof is very similar to Exercises 16 and 17. The crucial steps are

$$\frac{r+s}{2} = \frac{\frac{a}{b} + \frac{c}{d}}{2} = \frac{(ad+bc)/bd}{2} = \frac{ad+bc}{2bd}$$

and noticing that $ad+bc$ and $2bd$ are integers, and $2bd \neq 0$. \square

3.19 Exercise 19

For all real numbers a and b , if $a < b$ then $a < \frac{a+b}{2} < b$. (You may use the properties of inequalities in T17-T27 of Appendix A.)

Proof. True. Suppose a, b are any two real numbers such that $a < b$. [We need to prove two inequalities: $a < \frac{a+b}{2}$ and $\frac{a+b}{2} < b$.]

Since $a < b$ we have $a + a < a + b$ by T19. So $2a < a + b$. Then $a < \frac{a+b}{2}$ by T20. This proves the first inequality.

Since $a < b$ we have $a + b < b + b$ by T19. So $a + b < 2b$. Then $\frac{a+b}{2} < b$ by T20. This proves the second inequality. \square

3.20 Exercise 20

Use the results of exercises 18 and 19 to prove that given any two rational numbers r and s with $r < s$, there is another rational number between r and s . An important consequence is that there are infinitely many rational numbers in between any two distinct rational numbers. See Section 7.4.

Proof. Assume r and s are any two rational numbers with $r < s$. [Want to prove: $r < t < s$ for some rational number t .]

Let $t = \frac{r+s}{2}$. By Exercise 18, t is a rational number. By Exercise 19, $r < t < s$, [as was to be shown.] \square

Use the properties of even and odd integers that are listed in Example 4.3.3 to do exercises 21 – 23. Indicate which properties you use to justify your reasoning.

3.21 Exercise 21

True or false? If m is any even integer and n is any odd integer, then $m^2 + 3n$ is odd. Explain.

Proof. True.

m is even. An even integer times an even integer is even, therefore m^2 is even.

3 and n are both odd. An odd integer times an odd integer is odd, therefore $3n$ is odd.

m^2 is even. $3n$ is odd. An even integer plus an odd integer is odd, therefore $m^2 + 3n$ is odd. \square

3.22 Exercise 22

True or false? If a is any odd integer, then $a^2 + a$ is even. Explain.

Proof. True. $a^2 + a = a(a + 1)$. Since a is odd, $a + 1$ is even. Odd times even is even, therefore $a(a + 1)$ is even. So $a^2 + a$ is even. \square

3.23 Exercise 23

True or false? If k is any even integer and m is any odd integer, then $(k + 2)^2 - (m - 1)^2$ is even. Explain.

Proof. True.

k is even, so $k + 2$ is even. Even squared is even, so $(k + 2)^2$ is even.

m is odd, so $m - 1$ is even. Even squared is even, so $(m - 1)^2$ is even.

Even minus even is even, so $(k + 2)^2 - (m - 1)^2$ is even.

Another solution. By algebra:

$$(k + 2)^2 - (m - 1)^2 = (k + 2 - (m - 1))(k + 2 + m - 1) = (k - m + 3)(k + m + 1)$$

Now $k - m + 3$ is even $-$ odd $+$ odd $=$ even. Even times anything is even, therefore $(k - m + 3)(k + m + 1)$ is even. So $(k + 2)^2 - (m - 1)^2$ is even. \square

Derive the statements in 24 – 26 as corollaries of theorems 4.3.1, 4.3.2, and the results of exercises 12, 13, 14, 15, and 17.

3.24 Exercise 24

For any rational numbers r and s , $2r + 3s$ is rational.

Proof. Suppose r and s are any rational numbers. By Theorem 4.3.1, both 2 and 3 are rational, and so, by Exercise 15, both $2r$ and $3s$ are rational. Hence, by Theorem 4.3.2, $2r + 3s$ is rational. \square

3.25 Exercise 25

If r is any rational number, then $3r^2 - 2r + 4$ is rational.

Proof. Suppose r is any rational number. By Exercise 12, r^2 is rational. By Theorem 4.3.1, 2, 3, 4 are all rational. By Exercise 15, $3r^2$ and $2r$ are rational. By Exercise 17, $3r^2 - 2r$ is rational. So by Theorem 4.3.2, $3r^2 - 2r + 4$ is rational. \square

3.26 Exercise 26

For any rational number s , $5s^3 + 8s^2 - 7$ is rational.

Proof. Assume s is any rational number. By Theorem 4.3.1, 5, 8, 7 are rational, and by Exercise 13, -7 is rational. By Exercise 14, s^3 is rational. By Exercise 12, s^2 is rational. By Exercise 15, $5s^3$ and $8s^2$ are rational. Therefore by Theorem 4.3.2, $5s^3 + 8s^2 - 7$ is rational. \square

3.27 Exercise 27

It is a fact that if n is any nonnegative integer, then

$$1 + \frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \cdots + \frac{1}{2^n} = \frac{1 - (1/2^{n+1})}{1 - (1/2)}$$

(A more general form of this statement is proved in Section 5.2.) Is the right-hand side of this equation rational? If so, express it as a ratio of two integers.

Proof.

$$x = \frac{1 - \frac{1}{2^{n+1}}}{1 - \frac{1}{2}} = \frac{\frac{2^{n+1} - 1}{2^{n+1}}}{\frac{1}{2}} = \frac{2^{n+1} - 1}{2^{n+1}} \cdot \frac{2}{1} = \frac{2^{n+1} - 1}{2^n}$$

Now $2^{n+1} - 1$ and 2^n are both integers (since n is a nonnegative integer) and $2^n \neq 0$ by the zero product property. Therefore, x is rational. \square

3.28 Exercise 28

Suppose a, b, c , and d are integers and $a \neq c$. Suppose also that x is a real number that satisfies the equation

$$\frac{ax + b}{cx + d} = 1.$$

Must x be rational? If so, express x as a ratio of two integers.

Proof.

$$\frac{ax + b}{cx + d} = 1 \rightarrow ax + b = cx + d \rightarrow ax - cx + b - d = 0 \rightarrow x(a - c) = d - b \rightarrow x = \frac{d - b}{a - c}$$

x is rational because both $d - b$ and $a - c$ are rational, and because $a - c \neq 0$ (since $a \neq c$). \square

3.29 Exercise 29

Suppose a, b , and c are integers and x, y , and z are nonzero real numbers that satisfy the following equations:

$$\frac{xy}{x+y} = a, \frac{xz}{x+z} = b, \frac{yz}{y+z} = c$$

Is x rational? If so, express it as ratio of two integers.

Proof. Taking the reciprocals of both sides of the first equation:

$$\frac{x+y}{xy} = \frac{1}{a}$$

Now we split the first fraction into two, and simplify:

$$\frac{x+y}{xy} = \frac{x}{xy} + \frac{y}{xy} = \frac{1}{y} + \frac{1}{x}$$

Therefore $\frac{1}{y} + \frac{1}{x} = \frac{1}{a}$. By performing the same steps on the other two equations, we see that $\frac{1}{z} + \frac{1}{x} = \frac{1}{b}$ and $\frac{1}{z} + \frac{1}{y} = \frac{1}{c}$.

For the sake of simplicity let's do some renaming: let

$$X = 1/x, Y = 1/y, Z = 1/z, A = 1/a, B = 1/b, C = 1/c$$

So the three new equations we derived above become:

$$Y + X = A \quad (1)$$

$$Z + X = B \quad (2)$$

$$Z + Y = C \quad (3)$$

From (1) we get (4): $Y = A - X$ and from (2) we get (5): $Z = B - X$.

Substituting (4) and (5) back into (3) we get: $(B - X) + (A - X) = C$.

So $B + A - 2X = C$, then $B + A - C = 2X$ and $\frac{1}{2}(B + A - C) = X$. Using our old variable names, we get:

$$\frac{1}{2} \left(\frac{1}{b} + \frac{1}{a} - \frac{1}{c} \right) = \frac{1}{x}$$

Rewriting:

$$\frac{1}{2b} + \frac{1}{2a} - \frac{1}{2c} = \frac{1}{x}$$

Getting a common denominator, then adding:

$$\frac{ac}{2abc} + \frac{bc}{2abc} - \frac{ab}{2abc} = \frac{ac + bc - ab}{2abc} = \frac{1}{x}$$

Finally, taking reciprocals of both sides, we get x as a ratio of two integers:

$$\frac{2abc}{ac + bc - ab} = x$$

□

3.30 Exercise 30

Prove that if one solution for a quadratic equation of the form $x^2 + bx + c = 0$ is rational (where b and c are rational), then the other solution is also rational. (Use the fact that if the solutions of the equation are r and s , then $x^2 + bx + c = (x - r)(x - s)$.)

Proof. Assume $x^2 + bx + c = 0$ has two solutions r, s where one of them, r , is rational. [Want to prove: s is also rational.]

We are given the fact that $x^2 + bx + c = (x - r)(x - s)$. This holds true for all real numbers x . Solving for s , we get

$$\frac{x^2 + bx + c}{x - r} = x - s \implies \frac{x^2 + bx + c}{x - r} - x = -s \implies -\frac{x^2 + bx + c}{x - r} + x = s$$

This equality is true for all real x except $x = r$ (because then division by $x - r$ would be illegal). So, let's substitute an x value that is rational and different than r , say $r + 1$. Then we get:

$$-\frac{(r + 1)^2 + b(r + 1) + c}{r + 1 - r} + r + 1 = s \implies -(r + 1)^2 + b(r + 1) + c + r + 1 = s$$

Now we use the facts established in the Exercises that sums, negatives, products and squares of rational numbers are rational. Since $r, b, c, 1$ are all rational, this implies that s is rational. \square

3.31 Exercise 31

Prove that if a real number c satisfies a polynomial equation of the form

$$r_3x^3 + r_2x^2 + r_1x + r_0 = 0$$

where r_0, r_1, r_2, r_3 are rational numbers, then c satisfies an equation of the form

$$n_3x^3 + n_2x^2 + n_1x + n_0 = 0$$

where n_0, n_1, n_2, n_3 are integers.

Proof. Suppose c is a real number such that $r_3c^3 + r_2c^2 + r_1c + r_0 = 0$, where r_0, r_1, r_2, r_3 are rational numbers.

By definition of rational, $r_0 = a_0/b_0, r_1 = a_1/b_1, r_2 = a_2/b_2, r_3 = a_3/b_3$ for some integers a_0, a_1, a_2, a_3 and some nonzero integers b_0, b_1, b_2, b_3 . By substitution,

$$\begin{aligned} r_3c^3 + r_2c^2 + r_1c + r_0 &= \frac{a_3}{b_3}c^3 + \frac{a_2}{b_2}c^2 + \frac{a_1}{b_1}c + \frac{a_0}{b_0} \\ &= \frac{b_0b_1b_2a_3}{b_0b_1b_2b_3}c^3 + \frac{b_0b_1b_3a_2}{b_0b_1b_2b_3}c^2 + \frac{b_0b_2b_3a_1}{b_0b_1b_2b_3}c + \frac{b_1b_2b_3a_0}{b_0b_1b_2b_3} \\ &= 0. \end{aligned}$$

Multiplying both sides by $b_0b_1b_2b_3$ gives

$$b_0b_1b_2a_3 \cdot c^3 + b_0b_1b_3a_2 \cdot c^2 + b_0b_2b_3a_1 \cdot c + b_1b_2b_3a_0 = 0$$

Let $n_3 = b_0b_1b_2a_3$, $n_2 = b_0b_1b_3a_2$, $n_1 = b_0b_2b_3a_1$, $n_0 = b_1b_2b_3a_0$. Then n_3, n_2, n_1, n_0 are all integers (being products of integers). Hence c satisfies the equation

$$n_3 \cdot c^3 + n_2 \cdot c^2 + n_1 \cdot c + n_0 = 0$$

where n_3, n_2, n_1, n_0 are all integers, [as was to be shown.] □

Definition: A number c is called a **root** of a polynomial $p(x)$ if, and only if, $p(c) = 0$.

3.32 Exercise 32

Prove that for every real number c , if c is a root of a polynomial with rational coefficients, then c is a root of a polynomial with integer coefficients.

Proof. The proof is extremely similar to Exercise 31. Assume $p(c) = 0$ where:

c is any real number, and $p(x) = \frac{a_n}{b_n}x^n + \cdots + \frac{a_1}{b_1}x + \frac{a_0}{b_0}$ is a polynomial with rational coefficients (so a_0, \dots, a_n are all integers and b_0, \dots, b_n are all nonzero integers).

So c satisfies the equation

$$\frac{a_n}{b_n}c^n + \cdots + \frac{a_1}{b_1}c + \frac{a_0}{b_0} = 0$$

Multiply both sides by $L = b_0b_1 \cdots b_{n-1}b_n$:

$$\frac{a_nL}{b_n}c^n + \cdots + \frac{a_1L}{b_1}c + \frac{a_0L}{b_0} = 0$$

Notice that $\frac{a_nL}{b_n}, \dots, \frac{a_0L}{b_0}$ are all integers (because all the denominators can be cancelled out with one of the factors of L). So c is the root of a polynomial

$$q(x) = \frac{a_nL}{b_n}x^n + \cdots + \frac{a_1L}{b_1}x + \frac{a_0L}{b_0}$$

where $q(x)$ has all integer coefficients, [as was to be shown.] □

Use the properties of even and odd integers that are listed in example 4.3.3 to do exercises 33 and 34.

3.33 Exercise 33

When expressions of the form $(x - r)(x - s)$ are multiplied out, a quadratic polynomial is obtained. For instance, $(x - 2)(x - (-7)) = (x - 2)(x + 7) = x^2 + 5x - 14$.

3.33.1 (a)

What can be said about the coefficients of the polynomial obtained by multiplying out $(x-r)(x-s)$ when both r and s are odd integers? When both r and s are even integers? When one of r and s is even and the other is odd?

Proof. Note that $(x-r)(x-s) = x^2 - (r+s)x + rs$.

If both r and s are odd, then $r+s$ is even and rs is odd. So the coefficient of x^2 is 1 (odd), the coefficient of x is even, and the constant coefficient, rs , is odd.

If both r and s are even, then $r+s$ is even and rs is even. So the coefficient of x^2 is 1 (odd), the coefficient of x is even, and the constant coefficient, rs , is even.

If one of r and s is even and the other is odd, then $r+s$ is odd and rs is even. So the coefficient of x^2 is 1 (odd), the coefficient of x is odd, and the constant coefficient, rs , is even. \square

3.33.2 (b)

It follows from part (a) that $x^2 - 1253x + 255$ cannot be written as a product of two polynomials with integer coefficients. Explain why this is so.

Proof. Assume $x^2 - 1253x + 255 = (x-r)(x-s)$ where r, s are real numbers. So $r+s = 1253$ and $rs = 255$. If r, s are both integers, then since $rs = 255$, r and s must be both odd. But this is impossible, because then $r+s$ would be even, but 1253 is not even! Therefore r, s cannot be both integers. \square

3.34 Exercise 34

Observe that

$$(x-r)(x-s)(x-t) = x^3 - (r+s+t)x^2 + (rs+rt+st)x - rst$$

3.34.1 (a)

Derive a result for cubic polynomials similar to the result in part (a) of exercise 33 for quadratic polynomials.

Proof. If r, s, t are all odd, then the constant coefficient is odd, the coefficient of x is odd, the coefficient of x^2 is odd, and the coefficient of x^3 is odd (it's 1).

If exactly one of r, s, t is even, then the constant coefficient is even, the coefficient of x is odd, the coefficient of x^2 is even, and the coefficient of x^3 is odd (it's 1).

If exactly two of r, s, t are even, then the constant coefficient is even, the coefficient of x is even, the coefficient of x^2 is odd, and the coefficient of x^3 is odd (it's 1).

If r, s, t are all even, then the constant coefficient is even, the coefficient of x is even, the coefficient of x^2 is even, and the coefficient of x^3 is odd (it's 1). \square

3.34.2 (b)

Can $x^3 + 7x^2 - 8x - 27$ be written as a product of three polynomials with integer coefficients? Explain.

Proof. Assume $x^3 + 7x^2 - 8x - 27 = (x - r)(x - s)(x - t)$ where r, s, t are integers.

The coefficient of x^2 is 7, which is odd. So by part (a), either r, s, t are all odd, or exactly two of them are even.

They can't be all odd, because then the coefficient of x would be odd, but it's -8 which is even.

If exactly two of them are even, then the constant coefficient would have to be even, but it's -27 which is odd.

So it's impossible for r, s, t to be all integers. □

In 35 – 39 find the mistakes in the “proofs” that the sum of any two rational numbers is a rational number.

3.35 Exercise 35

“Proof: Any two rational numbers produce a rational number when added together. So if r and s are particular but arbitrarily chosen rational numbers, then $r + s$ is rational.”

Proof. This “proof” assumes what is to be proved. □

3.36 Exercise 36

“Proof: Let rational numbers $r = \frac{1}{4}$ and $s = \frac{1}{2}$ be given. Then $r + s = \frac{1}{4} + \frac{1}{2} = \frac{3}{4}$, which is a rational number. This is what was to be shown.”

Proof. This “proof” argues from a single example. It does not establish the result for *[arbitrarily chosen]* rational numbers. □

3.37 Exercise 37

“Proof: Suppose r and s are rational numbers. By definition of rational, $r = a/b$ for some integers a and b with $b \neq 0$, and $s = a/b$ for some integers a and b with $b \neq 0$. Then $r + s = \frac{a}{b} + \frac{a}{b} = \frac{2a}{b}$.

Let $p = 2a$. Then p is an integer since it is a product of integers. Hence $r + s = p/b$, where p and b are integers and $b \neq 0$. Thus $r + s$ is a rational number by definition of rational. This is what was to be shown.”

Proof. By setting both r and s equal to a/b , this incorrect proof violates the requirement that r and s be arbitrarily chosen rational numbers. If both r and s equal a/b , then $r = s$. □

3.38 Exercise 38

“Proof: Suppose r and s are rational numbers. Then $r = a/b$ and $s = c/d$ for some integers a, b, c , and d with $b \neq 0$ and $d \neq 0$ (by definition of rational). Then

$$r = \frac{a}{b} + \frac{c}{d}$$

But this is a sum of two fractions, which is a fraction. So $r + s$ is a rational number since a rational number is a fraction.”

Proof. This “proof” does not establish that “the sum of two fractions is a fraction”. Also, “a rational number is a fraction” is ambiguous. A rational number is a quotient of two integers (with nonzero denominator). But “a fraction” could be a fraction of non-integer numbers too. \square

3.39 Exercise 39

“Proof: Suppose r and s are rational numbers. If $r + s$ is rational, then by definition of rational $r + s = a/b$ for some integers a and b with $b \neq 0$.

Also since r and s are rational, $r = i/j$ and $s = m/n$ for some integers i, j, m , and n with $j \neq 0$ and $n \neq 0$. It follows that

$$r + s = \frac{i}{j} + \frac{m}{n} = \frac{a}{b}$$

which is a quotient of two integers with a nonzero denominator. Hence it is a rational number. This is what was to be shown.”

Proof. This “proof” assumes what is to be proved. \square

4 Exercise Set 4.4

Give a reason for your answer in each of 1 – 13. assume that all variables represent integers.

4.1 Exercise 1

Is 52 divisible by 13?

Proof. Yes, $52 = 13 \cdot 4$ \square

4.2 Exercise 2

Does $7 \mid 56$?

Proof. Yes, $56 = 7 \cdot 8$ \square

4.3 Exercise 3

Does $5 \mid 0$?

Proof. Yes, $0 = 5 \cdot 0$ □

4.4 Exercise 4

Does 3 divide $(3k + 1)(3k + 2)(3k + 3)$?

Proof. Yes, $(3k + 1)(3k + 2)(3k + 3) = 3[(3k + 1)(3k + 2)(k + 1)]$, and $(3k + 1)(3k + 2)(k + 1)$ is an integer because k is an integer and sums and products of integers are integers. □

4.5 Exercise 5

Is $6m(2m + 10)$ divisible by 4?

Proof. Yes: $6m(2m + 10) = (2(3m))(2(m + 5)) = 4(3m)(m + 5)$, and $(3m)(m + 5)$ is an integer because m is an integer and sums and products of integers are integers. □

4.6 Exercise 6

Is 29 a multiple of 3?

Proof. No, $29/3 \approx 9.67$, which is not an integer. □

4.7 Exercise 7

Is -3 a factor of 66?

Proof. Yes, $66 = (-3)(-22)$. □

4.8 Exercise 8

Is $6a(a + b)$ a multiple of $3a$?

Proof. Yes, $6a(a + b) = 3a[2(a + b)]$, and $2(a + b)$ is an integer because a and b are integers and sums and products of integers are integers. □

4.9 Exercise 9

Is 4 a factor of $2a \cdot 34b$?

Proof. Yes: $2a \cdot 34b = 2a \cdot (2 \cdot (17b)) = 4(a \cdot (17b))$ □

4.10 Exercise 10

Does $7 \mid 34$?

Proof. No, $34/7 \approx 4.86$, which is not an integer. □

4.11 Exercise 11

Does $13 \mid 73$?

Proof. No, $73/13 \approx 5.61$, which is not an integer. □

4.12 Exercise 12

If $n = 4k + 1$, does 8 divide $n^2 - 1$?

Proof. Yes, $n^2 - 1 = (4k + 1)^2 - 1 = (16k^2 + 8k + 1) - 1 = 16k^2 + 8k = 8(2k^2 + k)$, and $2k^2 + k$ is an integer because k is an integer and sums and products of integers are integers. □

4.13 Exercise 13

If $n = 4k + 3$, does 8 divide $n^2 - 1$?

Proof. Yes, $n^2 - 1 = (4k + 3)^2 - 1 = (16k^2 + 24k + 9) - 1 = 16k^2 + 24k + 8 = 8(2k^2 + 3k + 1)$, and $2k^2 + 3k + 1$ is an integer because k is an integer and sums and products of integers are integers. □

4.14 Exercise 14

Fill in the blanks in the following proof that for all integers a and b , if $a \mid b$ then $a \mid (-b)$.

Proof: Suppose a and b are any integers such that (a) _____. By definition of divisibility, there exists an integer r such that (b) _____. By substitution,

$$-b = -(ar) = a(-r).$$

Let $t =$ (c) _____. Then t is an integer because $t = (-1) \cdot r$, and both -1 and r are integers. Thus, by substitution, $-b = at$, where t is an integer, and so by definition of divisibility, (d) _____, as was to be shown.

Proof. (a) $a \mid b$ (b) $b = a \cdot r$ (c) $-r$ (d) $a \mid (-b)$ □

Prove statements 15 – 17 directly from the definition of divisibility.

4.15 Exercise 15

For all integers a, b , and c , if $a \mid b$ and $a \mid c$ then $a \mid (b + c)$.

Proof. Suppose a, b , and c are any integers such that $a \mid b$ and $a \mid c$. [We must show that $a \mid (b + c)$.]

By definition of divides, $b = ar$ and $c = as$ for some integers r and s . Then $b + c = ar + as = a(r + s)$ by algebra.

Let $t = r + s$. Then t is an integer (being a sum of integers), and thus $b + c = at$ where t is an integer. By definition of divides, then, $a \mid (b + c)$ [as was to be shown]. \square

4.16 Exercise 16

For all integers a, b , and c , if $a \mid b$ and $a \mid c$ then $a \mid (b - c)$.

Proof. Suppose a, b , and c are any integers such that $a \mid b$ and $a \mid c$. [We must show that $a \mid (b - c)$.]

By definition of divides, $b = ar$ and $c = as$ for some integers r and s . Then $b - c = ar - as = a(r - s)$ by algebra.

Let $t = r - s$. Then t is an integer (being a sum of integers), and thus $b - c = at$ where t is an integer. By definition of divides, then, $a \mid (b - c)$ [as was to be shown]. \square

4.17 Exercise 17

For all integers a, b, c , and d , if $a \mid c$ and $b \mid d$ then $ab \mid cd$.

Proof. Suppose a, b, c , and d are any integers such that $a \mid c$ and $b \mid d$. [We must show that $ab \mid cd$.]

By definition of divides, $c = ar$ and $d = bs$ for some integers r and s . Then $cd = (ar)(bs) = ab(rs)$ by algebra.

Let $t = rs$. Then t is an integer (being a product of integers), and thus $cd = abt$ where t is an integer. By definition of divides, then, $ab \mid cd$ [as was to be shown]. \square

4.18 Exercise 18

Consider the following statement: The negative of any multiple of 3 is a multiple of 3.

4.18.1 (a)

Write the statement formally using a quantifier and a variable.

Proof. \forall integers n if n is a multiple of 3 then $-n$ is a multiple of 3. \square

4.18.2 (b)

Determine whether the statement is true or false and justify your answer.

Proof. The statement is true. Suppose n is any integer that is a multiple of 3. [We must show that $-n$ is a multiple of 3.]

By definition of multiple, $n = 3k$ for some integer k . Then $-n = -(3k) = 3(-k)$ by substitution and by algebra.

Now $-k$ is an integer because k is. Hence, by definition of multiple, $-n$ is a multiple of 3 [as was to be shown]. \square

4.19 Exercise 19

Show that the following statement is false: For all integers a and b , if $3 \mid (a + b)$ then $3 \mid (a - b)$.

Proof. Counterexample: Let $a = 2$ and $b = 1$. Then $a + b = 2 + 1 = 3$, and so $3 \mid (a + b)$ because $3 = 3 \cdot 1$.

On the other hand, $a - b = 2 - 1 = 1$, and $3 \nmid 1$ because $1/3$ is not an integer. Thus $3 \nmid (a - b)$. [So the hypothesis of the statement is true and its conclusion is false.] \square

For each statement in 20 – 31, determine whether the statement is true or false. prove the statement directly from the definitions if it is true, and give a counterexample if it is false.

4.20 Exercise 20

The sum of any three consecutive integers is divisible by 3.

Proof. True. Assume a, b, c are any three consecutive integers. By definition of consecutive, $a = n, b = n + 1, c = n + 2$ for some integer n . Then $a + b + c = n + n + 1 + n + 2 = 3n + 6 = 3(n + 2)$. Let $t = n + 2$. Then t is an integer (being a sum of integers). So $a + b + c = 3t$ where t is an integer. So by definition of divisibility, $a + b + c$ is divisible by 3. \square

4.21 Exercise 21

The product of any two even integers is a multiple of 4.

Proof. True. Assume a, b are any two integers. By definition of even, $a = 2r$ and $b = 2s$ for some integers r, s . Then $ab = (2r)(2s) = 4rs$. Let $t = rs$. Then t is an integer (being a product of integers). So $ab = 4t$ where t is an integer. So ab is a multiple of 4. \square

4.22 Exercise 22

A necessary condition for an integer to be divisible by 6 is that it be divisible by 2.

Proof. Rewriting the statement, we get:

\forall integers n , if n is divisible by 6 then n is divisible by 2.

True. Assume n is any integer divisible by 6. By definition of divisibility, $n = 6m$ for some integer m . Then $n = 6m = 2(3m)$. Let $t = 3m$. Then t is an integer because it is a product of integers. So $n = 2t$ where t is an integer, therefore n is divisible by 2 by the definition of divisibility. \square

4.23 Exercise 23

A sufficient condition for an integer to be divisible by 8 is that it be divisible by 16.

Proof. Rewriting the statement, we get:

\forall integers n , if n is divisible by 16 then n is divisible by 8.

True. Assume n is any integer divisible by 16. By definition of divisibility, $n = 16m$ for some integer m . Then $n = 16m = 8(2m)$. Let $t = 2m$. Then t is an integer because it is a product of integers. So $n = 8t$ where t is an integer, therefore n is divisible by 8 by the definition of divisibility. \square

4.24 Exercise 24

For all integers a, b , and c , if $a \mid b$ and $a \mid c$ then $a \mid (2b - 3c)$.

Proof. The statement is true. Suppose a, b , and c are any integers such that $a \mid b$ and $a \mid c$. [We must show that $a \mid (2b - 3c)$.]

By definition of divisibility, we know that $b = am$ and $c = an$ for some integers m and n .

It follows that $2b - 3c = 2(am) - 3(an)$ (by substitution) $= a(2m - 3n)$ (by basic algebra).

Let $t = 2m - 3n$. Then t is an integer because it is a difference of products of integers. Hence $2b - 3c = at$, where t is an integer, and so $a \mid (2b - 3c)$ by definition of divisibility [as was to be shown]. \square

4.25 Exercise 25

For all integers a, b , and c , if a is a factor of c and b is a factor of c then ab is a factor of c .

Proof. The statement is false. Counterexample: Let $a = 2, b = 8$, and $c = 8$. Then a is a factor of c because $8 = 2 \cdot 4$ and b is a factor of c because $8 = 1 \cdot 8$, but $ab = 16$ and 16 is not a factor of 8 because $8 \neq 16 \cdot k$ for any integer k since $8/16 = 1/2$. \square

4.26 Exercise 26

For all integers a, b , and c , if $ab \mid c$ then $a \mid c$ and $b \mid c$.

Proof. True. Assume a, b, c are any three integers such that $ab \mid c$. By definition of divides, $abm = c$ for some integer m . Let $t = bm$ and $s = am$. Then t and s are integers (being products of integers). So $c = at$ and $c = bs$ where t and s are integers. Therefore by definition of divides, $a \mid c$ and $b \mid c$. \square

4.27 Exercise 27

For all integers a, b , and c , if $a \mid (b + c)$ then $a \mid b$ or $a \mid c$.

Proof. False Counterexample: Let $a = 6, b = 2, c = 4$. Then $b + c = 6$ and so $b + c = 6 = 6 \cdot 1 = a \cdot 1$ therefore $a \mid (b + c)$. However, $a \nmid b$ because $6 \nmid 2$ because $2/6$ is not an integer; similarly $a \nmid c$ because $6 \nmid 4$ because $4/6$ is not an integer. \square

4.28 Exercise 28

For all integers a, b , and c , if $a \mid bc$ then $a \mid b$ or $a \mid c$.

Proof. False Counterexample: Let $a = 6, b = 2, c = 3$. Then $bc = 6$ and so $bc = 6 = 6 \cdot 1 = a \cdot 1$ therefore $a \mid bc$. However, $a \nmid b$ because $6 \nmid 2$ because $2/6$ is not an integer; similarly $a \nmid c$ because $6 \nmid 3$ because $3/6$ is not an integer. \square

4.29 Exercise 29

For all integers a and b , if $a \mid b$ then $a^2 \mid b^2$.

Proof. True. Suppose a, b are any two integers such that $a \mid b$. By definition of divides, $b = ac$ for some integer c . Then $b^2 = (ac)^2 = a^2c^2$. Let $t = c^2$. Then t is an integer (being the square of an integer). So $b^2 = a^2 \cdot t$ where t is an integer. So by definition of divides, $a^2 \mid b^2$. \square

4.30 Exercise 30

For all integers a and n , if $a \mid n^2$ and $a \leq n$ then $a \mid n$.

Proof. False. Counterexample: Let $a = 4, n = 6$. Then $n^2 = 36$, so $a \mid n^2$ because $4 \mid 36$ because $36 = 4 \cdot 9$. But $a \nmid n$ because $4 \nmid 6$ because $6/4$ is not an integer. \square

4.31 Exercise 31

For all integers a and b , if $a \mid 10b$ then $a \mid 10$ or $a \mid b$.

Proof. False. Counterexample: Let $a = 15, b = 3$. Then $10b = 30$, so $a \mid 10b$ because $15 \mid 30$ because $30 = 15 \cdot 2$. But $a \nmid 10$ because $15 \nmid 10$ because $10/15$ is not an integer; similarly $a \nmid b$ because $15 \nmid 3$ because $3/15$ is not an integer. \square

4.32 Exercise 32

A fast-food chain has a contest in which a card with numbers on it is given to each customer who makes a purchase. If some of the numbers on the card add up to 100, then the customer wins \$100. A certain customer receives a card containing the numbers

72, 21, 15, 36, 69, 81, 9, 27, 42, and 63.

Will the customer win \$100? Why or why not?

Proof. Each of these numbers is divisible by 3, and so their sum is also divisible by 3. But 100 is not divisible by 3. Thus the sum cannot equal \$100. \square

4.33 Exercise 33

Is it possible to have a combination of nickels, dimes, and quarters that add up to \$4.72? Explain.

Proof. Impossible. Nickels are 5 cents, dimes are 10 cents, quarters are 25 cents. These are all divisible by 5. Therefore the cents value any combination of nickels, dimes, and quarters is also divisible by 5. But the cents value of \$4.72 is 472, which is not divisible by 5 (it does not end in 0 or 5). \square

4.34 Exercise 34

Consider a string consisting of a 's, b 's, and c 's where the number of b 's is three times the number of a 's and the number of c 's is five times the number of a 's. Prove that the length of the string is divisible by 3.

Proof. Suppose that the number of a 's in the string S is n where n is some integer. Then the number of b 's is $3n$, and the number of c 's is $5n$.

The total length of the string is: number of a 's + number of b 's + number of c 's = $n + 3n + 5n = 9n$. Since $9n = 3(3n)$ and $3n$ is an integer, $9n$ is divisible by 3. \square

4.35 Exercise 35

Two athletes run a circular track at a steady pace so that the first completes one round in 8 minutes and the second in 10 minutes. If they both start from the same spot at 4 p.m., when will be the first time they return to the start together?

Proof. The length of time needed is the least common multiple of 8 and 10, which is 40 minutes. So they will be back at the start together again for the first time at 4:40 p.m. \square

4.36 Exercise 36

It can be shown (see exercises 44 – 48) that an integer is divisible by 3 if, and only if, the sum of its digits is divisible by 3; an integer is divisible by 9 if, and only if, the sum of its digits is divisible by 9; an integer is divisible by 5 if, and only if, its right-most digit is a 5 or a 0; and an integer is divisible by 4 if, and only if, the number formed by its right-most two digits is divisible by 4. Check the following integers for divisibility by 3, 4, 5, and 9.

4.36.1 (a)

637,425,403,705,125

Proof. The sum of the digits is 54, which is divisible by 9. Therefore, 637,425,403,705,125 is divisible by 9 and hence also divisible by 3 (by transitivity of divisibility). Because the rightmost digit is 5, then 637,425,403,705,125 is divisible by 5. And because the two rightmost digits are 25, which is not divisible by 4, then 637,425,403,705,125 is not divisible by 4. \square

4.36.2 (b)

12,858,306,120,312

Proof. $1 + 2 + 8 + 5 + 8 + 3 + 0 + 6 + 1 + 2 + 0 + 3 + 1 + 2 = 42$. It is divisible by 3 (because $42/3 = 14$ is an integer), not divisible by 9 (because $42/9 \approx 4.67$ which is not an integer).

Last 2 digits are 12, so it's divisible by 4 (because $12/4 = 3$ is an integer) and not divisible by 5 (because $2 \neq 0$ or 5). \square

4.36.3 (c)

517,924,440,926,512

Proof. Last two digits are 12, so divisible by 4, not divisible by 5. Sum of digits is 61, so not divisible by 3 or 9. \square

4.36.4 (d)

14,328,083,360,232

Proof. Last two digits are 32, so divisible by 4, not divisible by 5. Sum of digits is 45, so divisible by 3 and 9. \square

4.37 Exercise 37

Use the unique factorization theorem to write the following integers in standard factored form.

4.37.1 (a)

1,176

Proof. $1,176 = 2^3 \cdot 3 \cdot 7^2$

□

4.37.2 (b)

5,733

Proof. $5,733 = 3^2 \cdot 7^2 \cdot 13$

□

4.37.3 (c)

3,675

Proof. $3,675 = 3 \cdot 5^2 \cdot 7^2$

□

4.38 Exercise 38

Let $n = 8,424$.

4.38.1 (a)

Write the prime factorization for n .

Proof. $8,424 = 2^3 \cdot 3^4 \cdot 13$

□

4.38.2 (b)

Write the prime factorization for n^5 .

Proof. $8,424^5 = (2^3 \cdot 3^4 \cdot 13)^5 = 2^{15} \cdot 3^{20} \cdot 13^5$

□

4.38.3 (c)

Is n^5 divisible by 20? Explain.

Proof. The answer is no. $20 = 2^2 \cdot 5$. So if n^5 is divisible by 20, it is divisible by 5. But 5 is not a factor of n^5 , so it is not divisible by 5.

□

4.38.4 (d)

What is the least positive integer m so that $8,424 \cdot m$ is a perfect square?

Proof. The answer is 26.

In order for $8,424 \cdot m$ to be a perfect square, each prime factor must be raised to an even power.

$8,424 = 2^3 \cdot 3^4 \cdot 13$. The prime factor 3 already has an even power of 4. The prime factor 2 has an odd power of 3, the closest even power is 4. The prime factor 13 has an odd power of 1, the closest even power is 2.

So let $m = 2 \cdot 13 = 26$. Then $8,424 \cdot m = 2^3 \cdot 3^4 \cdot 13 \cdot (2 \cdot 13) = 2^4 \cdot 3^4 \cdot 13^2 = (2^2 \cdot 3^2 \cdot 13^1)^2$ is a perfect square.

$m = 26$ is the least positive integer for this; any positive integer smaller than 26 will not result in $8,424 \cdot m$ being a perfect square, since we need to increase the powers of both prime factors 2 and 13 by at least 1. \square

4.39 Exercise 39

Suppose that in standard factored form $a = p_1^{e_1} p_2^{e_2} \dots p_k^{e_k}$, where k is a positive integer; p_1, p_2, \dots, p_k are prime numbers; and e_1, e_2, \dots, e_k are positive integers.

4.39.1 (a)

What is the standard factored form for a^3 ?

Proof. $a^3 = (p_1^{e_1} p_2^{e_2} \dots p_k^{e_k})^3 = p_1^{3e_1} p_2^{3e_2} \dots p_k^{3e_k}$ \square

4.39.2 (b)

Find the least positive integer k such that $2^4 \cdot 3^5 \cdot 7 \cdot 11^2 \cdot k$ is a perfect cube (that is, it equals an integer to the third power). Write the resulting product as a perfect cube.

Proof. Every prime factor must be raised to a power that is a multiple of 3.

So $2^4 \cdot 3^5 \cdot 7 \cdot 11^2$ should be turned into: $2^6 \cdot 3^6 \cdot 7^3 \cdot 11^3$ (because those powers are the closest multiples of 3).

Therefore, let $k = 2^2 \cdot 3^1 \cdot 7^2 \cdot 11^1$. Now

$2^4 \cdot 3^5 \cdot 7 \cdot 11^2 \cdot k = 2^4 \cdot 3^5 \cdot 7 \cdot 11^2 \cdot (2^2 \cdot 3^1 \cdot 7^2 \cdot 11^1) = 2^6 \cdot 3^6 \cdot 7^3 \cdot 11^3 = (2^2 \cdot 3^2 \cdot 7^1 \cdot 11^1)^3$ is a perfect cube. \square

4.40 Exercise 40

4.40.1 (a)

If a and b are integers and $12a = 25b$, does $12 \mid b$? does $25 \mid a$? Explain.

Proof. Because $12a = 25b$, the unique factorization theorem guarantees that the standard factored forms of $12a$ and $25b$ must be the same. Thus $25b$ contains the factors $2^2 \cdot 3 (= 12)$. But since neither 2 nor 3 divides 25, the factors $2^2 \cdot 3$ must all occur in b , and hence $12 \mid b$. Similarly, $12a$ contains the factors $5^2 = 25$, and since 5 is not a factor of 12, the factors 5^2 must occur in a . So $25 \mid a$. \square

4.40.2 (b)

If x and y are integers and $10x = 9y$, does $10 \mid y$? does $9 \mid x$? Explain.

Proof. Because $10x = 9y$, the unique factorization theorem guarantees that the standard factored forms of $10x$ and $9y$ must be the same. Thus $9y$ contains the factors $2 \cdot 5 (= 10)$. But since neither 2 nor 5 divides 9, the factors $2 \cdot 5$ must all occur in y , and hence $10 \mid y$. Similarly, $10x$ contains the factors $3^2 = 9$, and since 3 is not a factor of 10, the factors 3^2 must occur in x . So $9 \mid x$. \square

4.41 Exercise 41

How many zeros are at the end of $45^8 \cdot 88^5$? Explain how you can answer this question without actually computing the number. (Hint: $10 = 2 \cdot 5$.)

Proof. $45^8 \cdot 88^5 = (3^2 \cdot 5)^8 \cdot (2^3 \cdot 11)^5 = 3^{16} \cdot 5^8 \cdot 2^{15} \cdot 11^5$.

Since $10 = 2 \cdot 5$, there are 8 factors of 10 in this number, because 8 is the smaller power between the powers of 2 and 5.

$5^8 \cdot 2^{15}$ can be combined as $(2 \cdot 5)^8 \cdot 2^7 = 10^8 \cdot 2^7$. So 7 powers of 2 are “left over” and they do not contribute to powers of 10 (in other words, zeros at the end of the number). \square

4.42 Exercise 42

If n is an integer and $n > 1$, then $n!$ is the product of n and every other positive integer that is less than n . For example, $5! = 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1$.

4.42.1 (a)

Write $6!$ in standard factored form.

Proof. $6! = 6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1 = 2 \cdot 3 \cdot 5 \cdot 2 \cdot 2 \cdot 3 \cdot 2 = 2^4 \cdot 3^2 \cdot 5$ \square

4.42.2 (b)

Write $20!$ in standard factored form.

Proof. $(20)! = 20 \cdot 19 \cdot 18 \cdot 17 \cdot 16 \cdot 15 \cdot 14 \cdot 13 \cdot 12 \cdot 11 \cdot 10 \cdot 9 \cdot 8 \cdot 7 \cdot 6 \cdot 5 \cdot 4 \cdot 3 \cdot 2$
 $= (2^2 \cdot 5) \cdot 19 \cdot (2 \cdot 3^2) \cdot 17 \cdot (2^4) \cdot (3 \cdot 5) \cdot (2 \cdot 7) \cdot 13 \cdot (2^2 \cdot 3) \cdot 11 \cdot (2 \cdot 5) \cdot (3^2) \cdot (2^3) \cdot 7 \cdot (2 \cdot 3) \cdot 5 \cdot (2^2) \cdot 3 \cdot 2$
 $= 2^{18} \cdot 3^8 \cdot 5^4 \cdot 7^2 \cdot 11 \cdot 13 \cdot 17 \cdot 19$ \square

4.42.3 (c)

Without computing the value of $(20!)^2$ determine how many zeros are at the end of this number when it is written in decimal form. Justify your answer.

Proof. By part (b), $(20!) = 2^{18} \cdot 3^8 \cdot 5^4 \cdot 7^2 \cdot 11 \cdot 13 \cdot 17 \cdot 19$.

So $(20!)^2 = (2^{18} \cdot 3^8 \cdot 5^4 \cdot 7^2 \cdot 11 \cdot 13 \cdot 17 \cdot 19)^2 = 2^{36} \cdot 3^{16} \cdot 5^8 \cdot 7^4 \cdot 11^2 \cdot 13^2 \cdot 17^2 \cdot 19^2$.

To find the number of zeros at the end of this number, we look for the largest power of 10 in it. The largest power of 10 is the smaller of the powers of 2 and 5 (36 and 8), which is 8. So there are 8 zeros at the end. \square

4.43 Exercise 43

At a certain university $2/3$ of the mathematics students and $3/5$ of the computer science students have taken a discrete mathematics course. The number of mathematics students who have taken the course equals the number of computer science students who have taken the course. If there are at least 100 mathematics students at the university, what are the least possible number of mathematics students and the least possible number of computer science students at the university?

Proof. Let m and c denote the number of mathematics and computer science students, respectively.

We are given that $2m/3 = 3c/5$. So $10m = 9c$. By Exercise 40 part (b), we have $10 \mid c$ and $9 \mid m$.

The smallest possible value for m such that $9 \mid m$ and $m \geq 100$ is $m = 108$. (99 is divisible by 9, then 108, then 117...)

Putting this back into the equation we get $10(108) = 9c$ and solving for c we get $c = 10(108)/9 = 120$. \square

Definition: Given any nonnegative integer n , the **decimal representation** of n is an expression of the form

$$d_k d_{k-1} \cdots d_2 d_1 d_0,$$

where k is a nonnegative integer, $d_0, d_1, d_2, \dots, d_k$ (called the **decimal digits** of n) are integers from 0 to 9 inclusive, $d_k \neq 0$ unless $n = 0$ and $k = 0$, and

$$n = d_k \cdot 10^k + d_{k-1} \cdot 10^{k-1} + \cdots + d_2 \cdot 10^2 + d_1 \cdot 10 + d_0.$$

(For example, $2,503 = 2 \cdot 10^3 + 5 \cdot 10^2 + 0 \cdot 10 + 3$.)

4.44 Exercise 44

Prove that if n is any nonnegative integer whose decimal representation ends in 0, then $5 \mid n$. (Hint: If the decimal representation of a nonnegative integer n ends in d_0 , then $n = 10m + d_0$ for some integer m .)

Proof. Suppose n is a nonnegative integer whose decimal representation ends in 0. Then $n = 10m + 0 = 10m$ for some integer m . Factoring out a 5 yields $n = 10m = 5(2m)$,

and $2m$ is an integer since m is an integer. Hence $10m$ is divisible by 5, which is what was to be shown. \square

4.45 Exercise 45

Prove that if n is any nonnegative integer whose decimal representation ends in 5, then $5 \mid n$.

Proof. Assume that if n is any nonnegative integer whose decimal representation ends in 5. Then by definition,

$$n = d_k \cdot 10^k + d_{k-1} \cdot 10^{k-1} + \cdots + d_2 \cdot 10^2 + d_1 \cdot 10 + 5$$

for some nonnegative integer k and integers d_1, \dots, d_k from 0 to 9. If $k = 0$ then $n = 5$ which is divisible by 5. So we can assume $k \geq 1$. Now

$$n = 10d_k \cdot 10^{k-1} + 10d_{k-1} \cdot 10^{k-2} + \cdots + 10d_2 \cdot 10^{2-1} + 10d_1 + 5$$

Now factoring out a 5 we get

$$n = 5(2d_k \cdot 10^{k-1} + 2d_{k-1} \cdot 10^{k-2} + \cdots + 2d_2 \cdot 10 + 2d_1 + 1)$$

Let $t = 2d_k \cdot 10^{k-1} + 2d_{k-1} \cdot 10^{k-2} + \cdots + 2d_2 \cdot 10 + 2d_1 + 1$. Then t is an integer because it is a sum of products of integers. So $5 \mid n$ by definition of divides. \square

4.46 Exercise 46

Prove that if the decimal representation of a nonnegative integer n ends in d_1d_0 and if $4 \mid (10d_1 + d_0)$, then $4 \mid n$. (Hint: If the decimal representation of a nonnegative integer n ends in d_1d_0 , then there is an integer s such that $n = 100s + 10d_1 + d_0$.)

Proof. Assume the decimal representation of a nonnegative integer n ends in d_1d_0 and assume $4 \mid (10d_1 + d_0)$. Let

$$n = d_k \cdot 10^k + d_{k-1} \cdot 10^{k-1} + \cdots + d_2 \cdot 10^2 + d_1 \cdot 10 + d_0.$$

be the decimal representation of n . Since the representation ends in d_1d_0 , k must be at least 2 or greater. Factoring out 10^2 from the first $k - 1$ terms, and leaving the last 2 terms alone, we get

$$n = 10^2(d_k \cdot 10^{k-2} + d_{k-1} \cdot 10^{k-3} + \cdots + d_2 \cdot 10^{2-2}) + d_1 \cdot 10 + d_0.$$

Let $s = d_k \cdot 10^{k-2} + d_{k-1} \cdot 10^{k-3} + \cdots + d_2 \cdot 10^{2-2}$, which is an integer (being a sum of products of integers).

So $n = 100s + 10d_1 + d_0$. Since $4 \mid (10d_1 + d_0)$, we have $10d_1 + d_0 = 4t$ for some integer t . Therefore $n = 100s + 10d_1 + d_0 = 100s + 4t = 4(25s + t)$.

Let $r = 25s + t$, which is an integer (being a sum of products of integers). So $n = 4r$ where r is an integer. So $4 \mid n$ by definition of divides. \square

4.47 Exercise 47

Observe that

$$\begin{aligned} 7,524 &= 7 \cdot 1,000 + 5 \cdot 100 + 2 \cdot 10 + 4 \\ &= 7(999 + 1) + 5(99 + 1) + 2(9 + 1) + 4 \\ &= (7 \cdot 999 + 7) + (5 \cdot 99 + 5) + (2 \cdot 9 + 2) + 4 \\ &= (7 \cdot 999 + 5 \cdot 99 + 2 \cdot 9) + (7 + 5 + 2 + 4) \\ &= (7 \cdot 111 \cdot 9 + 5 \cdot 11 \cdot 9 + 2 \cdot 9) + (7 + 5 + 2 + 4) \\ &= (7 \cdot 111 + 5 \cdot 11 + 2) \cdot 9 + (7 + 5 + 2 + 4) \\ &= (\text{an integer divisible by } 9) + (\text{the sum of the digits of } 7,524). \end{aligned}$$

Since the sum of the digits of 7,524 is divisible by 9, 7,524 can be written as a sum of two integers each of which is divisible by 9. It follows from exercise 15 that 7,524 is divisible by 9. Generalize the argument given in this example to any nonnegative integer n . In other words, prove that for any nonnegative integer n , if the sum of the digits of n is divisible by 9, then n is divisible by 9.

Hint: You may take it as a fact that for any positive integer k ,

$$10^k = 9 \cdot 10^{k-1} + 9 \cdot 10^{k-2} + \dots + 9 \cdot 10^1 + 9 \cdot 10^0 + 1.$$

Proof. First let's talk about the Hint. It tells us that, for any positive integer k , we have $10^k = 9(10^{k-1} + 10^{k-2} + \dots + 10^1 + 10^0) + 1$. (For example, if $k = 5$ then $10^5 = 100000 = 99999 + 1 = 9 \cdot 11111 + 1$.)

In other words, every power of 10 can be written as: (some integer divisible by 9) + 1, or equivalently as: $(9 \cdot \text{some integer}) + 1$.

Assume n is any nonnegative integer such that the sum of the digits of n is divisible by 9. Let

$$n = d_k \cdot 10^k + d_{k-1} \cdot 10^{k-1} + \dots + d_2 \cdot 10^2 + d_1 \cdot 10 + d_0.$$

be the decimal representation of n . By the Hint,

$$n = d_k \cdot (9a_k + 1) + d_{k-1} \cdot (9a_{k-1} + 1) + \dots + d_2 \cdot (9a_2 + 1) + d_1 \cdot (9a_1 + 1) + d_0$$

for some integers a_1, \dots, a_k . Multiplying out, we get:

$$n = (9d_k a_k + d_k) + (9d_{k-1} a_{k-1} + d_{k-1}) + \dots + (9d_2 a_2 + d_2) + (9d_1 a_1 + d_1) + d_0$$

Reorganizing, we get

$$n = 9(d_k a_k + d_{k-1} a_{k-1} + \dots + d_2 a_2 + d_1 a_1) + (d_k + d_{k-1} + \dots + d_2 + d_1 + d_0)$$

The first term is divisible by 9. Therefore, n is divisible by 9 if, and only if, the second term $(d_k + d_{k-1} + \dots + d_2 + d_1 + d_0)$, in other words the sum of n 's digits, is also divisible by 9. \square

4.48 Exercise 48

Prove that for any nonnegative integer n , if the sum of the digits of n is divisible by 3, then n is divisible by 3.

Proof. This follows from the proof of Exercise 47. We can simply repeat the same proof, and at the end, since $9 = 3 \cdot 3$, we have

$$n = (3 \cdot 3) \cdot (d_k a_k + d_{k-1} a_{k-1} + \cdots + d_2 a_2 + d_1 a_1) + (d_k + d_{k-1} + \cdots + d_2 + d_1 + d_0)$$

so the first term is divisible by 3, therefore n is divisible by 3 if and only if the second term (sum of its digits) is also divisible by 3. \square

4.49 Exercise 49

Given a positive integer n written in decimal form, the alternating sum of the digits of n is obtained by starting with the right-most digit, subtracting the digit immediately to its left, adding the next digit to the left, subtracting the next digit, and so forth. For example, the alternating sum of the digits of 180,928 is $8 - 2 + 9 - 0 + 8 - 1 = 22$. Justify the fact that for any nonnegative integer n , if the alternating sum of the digits of n is divisible by 11, then n is divisible by 11.

Proof. The idea is the same as in Exercises 47 and 48. This time, we have to notice that for even powers of 10, like 100 for example, we can write it as $100 = 99 + 1 = 11 \cdot 9 + 1$; but for odd powers of 10, like 1000 for example, we can write $1000 = 1001 - 1 = 11 \cdot 91 - 1$.

Taking a look at more examples of even powers of 10, we see $10,000 = 9,999 + 1 = 11 \cdot 909 + 1$, and $1,000,000 = 999,999 + 1 = 11 \cdot 90,909 + 1$ and so on.

Taking a look at more examples of odd powers of 10, we see $100,000 = 100,001 - 1 = 11 \cdot 9091 - 1$, and $10,000,000 = 10,000,001 - 1 = 11 \cdot 909,091 - 1$ and so on.

In general, if k is a nonnegative integer, then we have $10^{2k} = 11 \cdot a + 1$ and $10^{2k+1} = 11 \cdot b - 1$ for some integers a, b . Putting these into one formula we get:

$$\forall k \geq 0, \exists a_k \geq 0 \text{ such that } 10^k = 11 \cdot a_k + (-1)^k.$$

Now we can repeat the proof in Exercise 47 but using the above equation instead. We'll end up with:

$$n = 11 \cdot (d_k a_k + \cdots + d_2 a_2 + d_1 a_1) + ((-1)^k d_k + \cdots + (-1)^2 d_2 + (-1)^1 d_1 + (-1)^0 d_0)$$

The first term on the right-hand side is divisible by 11. So n is divisible by 11 if, and only if, the second term (the alternating sum of its digits) is divisible by 11. \square

4.50 Exercise 50

The integer 123,123 has the form abc,abc , where a, b , and c are integers from 0 through 9. Consider all six-digit integers of this form. Which prime numbers divide every one of these integers? Prove your answer.

Proof. 11 divides every one of these integers. Every one of these integers has an alternating sum of digits equal to 0, because: $a - b + c - a + b - c = 0$. Since $11 \mid 0$, by Exercise 49, 11 divides every one of these integers. \square

5 Exercise Set 4.5

5.1 Exercise 1

Proof. \square

5.2 Exercise 2

Proof. \square

5.3 Exercise 3

Proof. \square

5.4 Exercise 4

Proof. \square

5.5 Exercise 5

Proof. \square

5.6 Exercise 6

Proof. \square

5.7 Exercise 7

5.7.1 (a)

Proof. \square

5.7.2 (b)

Proof. \square

5.8 Exercise 8

5.8.1 (a)

Proof. \square

5.8.2 (b)

Proof.



5.9 Exercise 9

5.9.1 (a)

Proof.



5.9.2 (b)

Proof.



5.10 Exercise 10

5.10.1 (a)

Proof.



5.10.2 (b)

Proof.



5.11 Exercise 11

5.11.1 (a)

Proof.



5.11.2 (b)

Proof.



5.11.3 (c)

Proof.



5.12 Exercise 12

Proof.



5.13 Exercise 13

Proof.



5.14 Exercise 14

Proof.



5.15 Exercise 15

Proof.



5.16 Exercise 16

Proof.



5.17 Exercise 17

Proof.



5.18 Exercise 18

5.18.1 (a)

Proof.



5.18.2 (b)

Proof.



5.19 Exercise 19

Proof.



5.20 Exercise 20

Proof.



5.21 Exercise 21

Proof.



5.22 Exercise 22

Proof.



5.23 Exercise 23

Proof.



5.24 Exercise 24

Proof.



5.25 Exercise 25

Proof.



5.26 Exercise 26

Proof.



5.27 Exercise 27

Proof.



5.28 Exercise 28

5.28.1 (a)

Proof.



5.28.2 (b)

Proof.



5.29 Exercise 29

5.29.1 (a)

Proof.



5.29.2 (b)

Proof.



5.30 Exercise 30

5.30.1 (a)

Proof.



5.30.2 (b)

Proof.



5.31 Exercise 31

5.31.1 (a)

Proof.



5.31.2 (b)

Proof.



5.31.3 (c)

Proof.



5.32 Exercise 32

Proof.



5.33 Exercise 33

Proof.



5.34 Exercise 34

Proof.



5.35 Exercise 35

Proof.



5.36 Exercise 36

Proof.



5.37 Exercise 37

Proof.



5.38 Exercise 38

Proof.



5.39 Exercise 39

Proof.



5.40 Exercise 40

Proof.



5.41 Exercise 41

Proof.



5.42 Exercise 42

Proof.



5.43 Exercise 43

Proof.



5.44 Exercise 44

5.44.1 (a)

Proof.



5.44.2 (b)

Proof.



5.44.3 (c)

Proof.



5.45 Exercise 45

Proof.



5.46 Exercise 46

Proof.



5.47 Exercise 47

Proof.



5.48 Exercise 48

Proof.



5.49 Exercise 49

Proof.



5.50 Exercise 50

Proof.



6 Exercise Set 4.6

6.1 Exercise 1

Proof.



6.2 Exercise 2

Proof.



6.3 Exercise 3

Proof.



6.4 Exercise 4

Proof.



6.5 Exercise 5

Proof.



6.6 Exercise 6

Proof.



6.7 Exercise 7

Proof.



6.8 Exercise 8

Proof.



6.9 Exercise 9

Proof.



6.10 Exercise 10

6.10.1 (a)

Proof.



6.10.2 (b)

Proof.



6.11 Exercise 11

Proof.



6.12 Exercise 12

Proof.



6.13 Exercise 13

Proof.



6.14 Exercise 14

Proof.



6.15 Exercise 15

Proof.



6.16 Exercise 16

Proof.



6.17 Exercise 17

Proof.



6.18 Exercise 18

Proof.



6.19 Exercise 19

Proof.



6.20 Exercise 20

Proof.



6.21 Exercise 21

Proof.



6.22 Exercise 22

Proof.



6.23 Exercise 23

Proof.



6.24 Exercise 24

Proof.



6.25 Exercise 25

Proof.



6.26 Exercise 26

Proof.



6.27 Exercise 27

Proof.



6.28 Exercise 28

Proof.



6.29 Exercise 29

Proof.



6.30 Exercise 30

Proof.



6.31 Exercise 31

Proof.



6.32 Exercise 32

Proof.



6.33 Exercise 33

Proof.



7 Exercise Set 4.7

7.1 Exercise 1

Proof.



7.2 Exercise 2

Proof.



7.3 Exercise 3

Proof.



7.4 Exercise 4

Proof.



7.5 Exercise 5

Proof.



7.6 Exercise 6

Proof.



7.7 Exercise 7

Proof.



7.8 Exercise 8

Proof.



7.9 Exercise 9

7.9.1 (a)

Proof.



7.9.2 (b)

Proof.



7.10 Exercise 10

Proof.



7.11 Exercise 11

Proof.



7.12 Exercise 12

7.12.1 (a)

Proof.



7.12.2 (b)

Proof.



7.13 Exercise 13

7.13.1 (a)

Proof.



7.13.2 (b)

Proof.



7.14 Exercise 14

7.14.1 (a)

Proof.



7.14.2 (b)

Proof.



7.15 Exercise 15

Proof.



7.16 Exercise 16

Proof.



7.17 Exercise 17

Proof.



7.18 Exercise 18

Proof.



7.19 Exercise 19

Proof.



7.20 Exercise 20

Proof.



7.21 Exercise 21

7.21.1 (a)

Proof.



7.21.2 (b)

Proof.



7.22 Exercise 22

7.22.1 (a)

Proof.



7.22.2 (b)

Proof.



7.23 Exercise 23

Proof.



7.24 Exercise 24

Proof.



7.25 Exercise 25

Proof.



7.26 Exercise 26

Proof.



7.27 Exercise 27

Proof.



7.28 Exercise 28

Proof.



7.29 Exercise 29

Proof.



7.30 Exercise 30

7.30.1 (a)

Proof.



7.30.2 (b)

Proof.



7.31 Exercise 31

7.31.1 (a)

Proof.



7.31.2 (b)

Proof.



7.31.3 (c)

Proof.



7.32 Exercise 32

Proof.



7.33 Exercise 33

Proof.



7.34 Exercise 34

7.34.1 (a)

Proof.



7.34.2 (b)

Proof.



7.34.3 (c)

Proof.



7.34.4 (d)

Proof.



7.35 Exercise 35

Proof.



7.36 Exercise 36

Proof.



8 Exercise Set 4.8

8.1 Exercise 1

Proof.



8.2 Exercise 2

Proof.



8.3 Exercise 3

Proof.



8.4 Exercise 4

Proof.



8.5 Exercise 5

Proof.



8.6 Exercise 6

Proof.



8.7 Exercise 7

Proof.



8.8 Exercise 8

Proof.



8.9 Exercise 9

Proof.



8.10 Exercise 10

Proof.



8.11 Exercise 11

Proof.



8.12 Exercise 12

Proof.



8.13 Exercise 13

Proof.



8.14 Exercise 14

Proof.



8.15 Exercise 15

Proof.



8.16 Exercise 16

Proof.



8.17 Exercise 17

Proof.



8.18 Exercise 18

8.18.1 (a)

Proof.



8.18.2 (b)

Proof.



8.19 Exercise 19

8.19.1 (a)

Proof.



8.19.2 (b)

Proof.



8.19.3 (c)

Proof.



8.20 Exercise 20

Proof.



8.21 Exercise 21

Proof.



8.22 Exercise 22

Proof.



8.23 Exercise 23

Proof.



8.24 Exercise 24

Proof.



8.25 Exercise 25

Proof.



8.26 Exercise 26

Proof.



8.27 Exercise 27

Proof.



8.28 Exercise 28

Proof.



8.29 Exercise 29

Proof.



8.30 Exercise 30

8.30.1 (a)

Proof.



8.30.2 (b)

Proof.



8.31 Exercise 31

Proof.



8.32 Exercise 32

Proof.



8.33 Exercise 33

Proof.



8.34 Exercise 34

8.34.1 (a)

Proof.



8.34.2 (b)

Proof.



8.35 Exercise 35

Proof.



8.36 Exercise 36

Proof.



8.37 Exercise 37

Proof.



8.38 Exercise 38

Proof.



9 Exercise Set 4.9

9.1 Exercise 1

Proof.



9.2 Exercise 2

Proof.



9.3 Exercise 3

Proof.



9.4 Exercise 4

Proof.



9.5 Exercise 5

Proof.



9.6 Exercise 6

Proof.



9.7 Exercise 7

Proof.



9.8 Exercise 8

Proof.



9.9 Exercise 9

Proof.



9.10 Exercise 10

Proof.



9.11 Exercise 11

Proof.



9.12 Exercise 12

Proof.



9.13 Exercise 13

Proof.



9.14 Exercise 14

9.14.1 (a)

Proof.



9.14.2 (b)

Proof.



9.15 Exercise 15

9.15.1 (a)

Proof.



9.15.2 (b)

Proof.



9.16 Exercise 16

9.16.1 (a)

Proof.



9.16.2 (b)

Proof.



9.17 Exercise 17

Proof.



9.18 Exercise 18

Proof.



9.19 Exercise 19

Proof.



9.20 Exercise 20

9.20.1 (a)

Proof.



9.20.2 (b)

Proof.



9.21 Exercise 21

9.21.1 (a)

Proof.



9.21.2 (b)

Proof.



9.21.3 (c)

Proof.



9.22 Exercise 22

Proof.



9.23 Exercise 23

9.23.1 (a)

Proof.



9.23.2 (b)

Proof.



9.23.3 (c)

Proof.



9.23.4 (d)

Proof.



9.23.5 (e)

Proof.



9.23.6 (f)

Proof.



9.24 Exercise 24

9.24.1 (a)

Proof.



9.24.2 (b)

Proof.



9.24.3 (c)

Proof.



9.24.4 (d)

Proof.



9.24.5 (e)

Proof.



9.24.6 (f)

Proof.



9.25 Exercise 25

Proof.



10 Exercise Set 4.10

10.1 Exercise 1

Proof.



10.2 Exercise 2

Proof.



10.3 Exercise 3

10.3.1 (a)

Proof.



10.3.2 (b)

Proof.



10.4 Exercise 4

Proof.



10.5 Exercise 5

Proof.



10.6 Exercise 6

Proof.



10.7 Exercise 7

Proof.



10.8 Exercise 8

10.8.1 (a)

Proof.



10.8.2 (b)

Proof.



10.9 Exercise 9

Proof.



10.10 Exercise 10

Proof.



10.11 Exercise 11

Proof.



10.12 Exercise 12

Proof.



10.13 Exercise 13

Proof.



10.14 Exercise 14

Proof.



10.15 Exercise 15

Proof.



10.16 Exercise 16

Proof.



10.17 Exercise 17

Proof.



10.18 Exercise 18

Proof.



10.19 Exercise 19

Proof.



10.20 Exercise 20

Proof.



10.21 Exercise 21

Proof.



10.22 Exercise 22

Proof.



10.23 Exercise 23

10.23.1 (a)

Proof.



10.23.2 (b)

Proof.



10.24 Exercise 24

Proof.



10.25 Exercise 25

10.25.1 (a)

Proof.



10.25.2 (b)

Proof.



10.26 Exercise 26

10.26.1 (a)

Proof.



10.26.2 (b)

Proof.



10.27 Exercise 27

10.27.1 (a)

Proof.



10.27.2 (b)

Proof.



10.27.3 (c)

Proof.



10.28 Exercise 28

10.28.1 (a)

Proof.



10.28.2 (b)

Proof.



10.28.3 (c)

Proof.



10.29 Exercise 29

Proof.



10.30 Exercise 30

Proof.



10.31 Exercise 31

Proof.



10.32 Exercise 32

Proof.

