

Chapter 6 Solutions, Susanna Epp Discrete Math 5th Edition

<https://github.com/spamegg1>

August 26, 2023

Contents

1	Exercise Set 6.1	8
1.1	Exercise 1	8
1.1.1	(a)	8
1.1.2	(b)	8
1.1.3	(c)	8
1.1.4	(d)	8
1.1.5	(e)	9
1.1.6	(f)	9
1.2	Exercise 2	9
1.3	Exercise 3	10
1.3.1	(a)	10
1.3.2	(b)	10
1.3.3	(c)	10
1.4	Exercise 4	11
1.4.1	(a)	11
1.4.2	(b)	11
1.5	Exercise 5	11
1.5.1	(a)	11
1.5.2	(b)	11
1.6	Exercise 6	12
1.6.1	(a)	12
1.6.2	(b)	12
1.6.3	(c)	12
1.7	Exercise 7	12
1.7.1	(a)	13
1.7.2	(b)	13
1.7.3	(c)	13
1.8	Exercise 8	13
1.8.1	(a)	13
1.8.2	(b)	14

	1.8.3	(c)	14
	1.8.4	(d)	14
1.9	Exercise 9		14
	1.9.1	(a)	14
	1.9.2	(b)	14
	1.9.3	(c)	14
1.10	Exercise 10		14
	1.10.1	(a)	15
	1.10.2	(b)	15
	1.10.3	(c)	15
	1.10.4	(d)	15
	1.10.5	(e)	15
	1.10.6	(f)	15
	1.10.7	(g)	15
	1.10.8	(h)	15
1.11	Exercise 11		16
	1.11.1	(a)	16
	1.11.2	(b)	16
	1.11.3	(c)	16
	1.11.4	(d)	16
	1.11.5	(e)	16
	1.11.6	(f)	16
	1.11.7	(g)	16
	1.11.8	(h)	17
	1.11.9	(i)	17
	1.11.10	(j)	17
1.12	Exercise 12		17
	1.12.1	(a)	17
	1.12.2	(b)	17
	1.12.3	(c)	17
	1.12.4	(d)	17
	1.12.5	(e)	18
	1.12.6	(f)	18
	1.12.7	(g)	18
	1.12.8	(h)	18
	1.12.9	(i)	18
	1.12.10	(j)	18
1.13	Exercise 13		18
	1.13.1	(a)	18
	1.13.2	(b)	19
	1.13.3	(c)	19
	1.13.4	(d)	19
1.14	Exercise 14		19
	1.14.1	(a)	19
	1.14.2	(b)	19
1.15	Exercise 15		19

1.15.1	(a)	20
1.15.2	(b)	20
1.15.3	(c)	20
1.16	Exercise 16	20
1.16.1	(a)	21
1.16.2	(b)	21
1.16.3	(c)	21
1.17	Exercise 17	21
1.17.1	(a)	21
1.17.2	(b)	22
1.17.3	(c)	22
1.17.4	(d)	22
1.17.5	(e)	23
1.17.6	(f)	23
1.18	Exercise 18	23
1.18.1	(a)	23
1.18.2	(b)	23
1.18.3	(c)	23
1.18.4	(d)	23
1.19	Exercise 19	24
1.19.1	(a)	24
1.19.2	(b)	24
1.19.3	(c)	24
1.20	Exercise 20	24
1.20.1	(a)	24
1.20.2	(b)	24
1.20.3	(c)	24
1.21	Exercise 21	24
1.21.1	(a)	25
1.21.2	(b)	25
1.21.3	(c)	25
1.21.4	(d)	25
1.21.5	(e)	25
1.21.6	(f)	26
1.21.7	(g)	26
1.22	Exercise 22	26
1.22.1	(a)	26
1.22.2	(b)	26
1.22.3	(c)	26
1.22.4	(d)	27
1.22.5	(e)	27
1.22.6	(f)	27
1.22.7	(g)	27
1.23	Exercise 23	27
1.23.1	(a)	27
1.23.2	(b)	28

1.23.3	(c)	28
1.23.4	(d)	28
1.23.5	(e)	28
1.23.6	(f)	28
1.23.7	(g)	28
1.24	Exercise 24	29
1.24.1	(a)	29
1.24.2	(b)	29
1.24.3	(c)	29
1.24.4	(d)	29
1.24.5	(e)	29
1.24.6	(f)	30
1.24.7	(g)	30
1.25	Exercise 25	30
1.25.1	(a)	30
1.25.2	(b)	30
1.25.3	(c)	30
1.25.4	(d)	31
1.25.5	(e)	31
1.25.6	(f)	31
1.25.7	(g)	31
1.26	Exercise 26	31
1.26.1	(a)	31
1.26.2	(b)	32
1.26.3	(c)	32
1.26.4	(d)	32
1.26.5	(e)	32
1.26.6	(f)	32
1.26.7	(g)	32
1.27	Exercise 27	33
1.27.1	(a)	33
1.27.2	(b)	33
1.27.3	(c)	33
1.27.4	(d)	33
1.27.5	(e)	33
1.28	Exercise 28	33
1.29	Exercise 29	34
1.30	Exercise 30	34
1.31	Exercise 31	34
1.31.1	(a)	34
1.31.2	(b)	34
1.31.3	(c)	34
1.31.4	(d)	35
1.32	Exercise 32	35
1.32.1	(a)	35
1.32.2	(b)	35

1.33	Exercise 33	35
1.33.1	(a)	35
1.33.2	(b)	35
1.33.3	(c)	35
1.34	Exercise 34	36
1.34.1	(a)	36
1.34.2	(b)	36
1.35	Exercise 35	36
1.35.1	(a)	36
1.35.2	(b)	36
1.35.3	(c)	36
1.35.4	(d)	36
1.36	Exercise 36	37
1.37	Exercise 37	37
1.38	Exercise 38	37
2	Exercise Set 6.2	38
2.1	Exercise 1	38
2.1.1	(a)	38
2.1.2	(b)	38
2.1.3	(c)	38
2.1.4	(d)	38
2.1.5	(e)	38
2.2	Exercise 2	38
2.2.1	(a)	39
2.2.2	(b)	39
2.3	Exercise 3	39
2.4	Exercise 4	39
2.5	Exercise 5	40
2.6	Exercise 6	40
2.7	Exercise 7	41
2.8	Exercise 8	42
2.9	Exercise 9	42
2.10	Exercise 10	43
2.11	Exercise 11	43
2.12	Exercise 12	43
2.13	Exercise 13	44
2.14	Exercise 14	44
2.15	Exercise 15	45
2.16	Exercise 16	45
2.17	Exercise 17	45
2.18	Exercise 18	46
2.19	Exercise 19	46
2.20	Exercise 20	46
2.21	Exercise 21	47
2.22	Exercise 22	47

2.23	Exercise 23	48
2.24	Exercise 24	48
2.25	Exercise 25	49
2.26	Exercise 26	49
2.26.1	(a)	49
2.26.2	(b)	50
2.26.3	(c)	50
2.26.4	(d)	50
2.27	Exercise 27	51
2.28	Exercise 28	51
2.29	Exercise 29	51
2.30	Exercise 30	52
2.31	Exercise 31	52
2.32	Exercise 32	52
2.33	Exercise 33	52
2.34	Exercise 34	52
2.35	Exercise 35	53
2.36	Exercise 36	53
2.37	Exercise 37	53
2.38	Exercise 38	53
2.39	Exercise 39	54
2.39.1	(a)	54
2.39.2	(b)	54
2.40	Exercise 40	54
2.41	Exercise 41	55
2.42	Exercise 42	56
2.43	Exercise 43	57
2.44	Exercise 44	58
3	Exercise Set 6.3	59
3.1	Exercise 1	59
3.2	Exercise 2	60
3.3	Exercise 3	60
3.4	Exercise 4	60
3.5	Exercise 5	60
3.6	Exercise 6	60
3.7	Exercise 7	61
3.8	Exercise 8	61
3.9	Exercise 9	61
3.10	Exercise 10	61
3.11	Exercise 11	62
3.12	Exercise 12	62
3.13	Exercise 13	63
3.14	Exercise 14	63
3.15	Exercise 15	63
3.16	Exercise 16	64

3.17	Exercise 17	64
3.18	Exercise 18	64
3.19	Exercise 19	64
3.20	Exercise 20	65
3.21	Exercise 21	65
3.22	Exercise 22	65
3.22.1	(a)	65
3.22.2	(b)	66
3.23	Exercise 23	66
3.24	Exercise 24	66
3.24.1	(a)	66
3.24.2	(b)	66
3.24.3	(c)	66
3.24.4	(d)	66
3.24.5	(e)	67
3.24.6	(f)	67
3.25	Exercise 25	67
3.26	Exercise 26	68
3.27	Exercise 27	68
3.28	Exercise 28	69
3.29	Exercise 29	69
3.30	Exercise 30	69
3.31	Exercise 31	70
3.32	Exercise 32	70
3.33	Exercise 33	70
3.34	Exercise 34	71
3.35	Exercise 35	71
3.36	Exercise 36	71
3.37	Exercise 37	72
3.38	Exercise 38	72
3.39	Exercise 39	72
3.40	Exercise 40	73
3.41	Exercise 41	73
3.42	Exercise 42	74
3.43	Exercise 43	74
3.44	Exercise 44	75
3.44.1	(a)	75
3.44.2	(b)	75
3.44.3	(c)	75
3.45	Exercise 45	75
3.45.1	(a)	75
3.45.2	(b)	76
3.45.3	(c)	77
3.46	Exercise 46	77
3.46.1	(a)	78
3.46.2	(b)	78

3.46.3 (c)	78
3.46.4 (d)	78
3.47 Exercise 47	78
3.48 Exercise 48	78
3.49 Exercise 49	79
3.50 Exercise 50	79
3.51 Exercise 51	79
3.52 Exercise 52	80
3.53 Exercise 53	81
3.54 Exercise 54	81

1 Exercise Set 6.1

1.1 Exercise 1

In each of (a)-(f), answer the following questions: Is $A \subseteq B$? Is $B \subseteq A$? Is either A or B a proper subset of the other?

1.1.1 (a)

$$A = \{2, \{2\}, (\sqrt{2})^2\}, B = \{2, \{2\}, \{\{2\}\}\}$$

Proof. $A = \{2, \{2\}, (\sqrt{2})^2\} = \{2, \{2\}, 2\} = \{2, \{2\}\}$, so $A \subseteq B$ because every element of A is in B , but $B \not\subseteq A$ because $\{\{2\}\} \in B$ but $\{\{2\}\} \notin A$. Thus A is a proper subset of B . \square

1.1.2 (b)

$$A = \{3, \sqrt{5^2 - 4^2}, 24 \bmod 7\}, B = \{8 \bmod 5\}$$

Proof. $A = \{3, \sqrt{5^2 - 4^2}, 24 \bmod 7\} = \{3, 3, 3\} = \{3\}$, $B = \{8 \bmod 5\} = \{3\}$. So $A = B$, which means both $A \subseteq B$ and $B \subseteq A$. \square

1.1.3 (c)

$$A = \{\{1, 2\}, \{2, 3\}\}, B = \{1, 2, 3\}$$

Proof. $A \not\subseteq B$ because $\{1, 2\} \in A$ but $\{1, 2\} \notin B$. $B \not\subseteq A$ because $1 \in B$ but $1 \notin A$. \square

1.1.4 (d)

$$A = \{a, b, c\}, B = \{\{a\}, \{b\}, \{c\}\}$$

Proof. $A \not\subseteq B$ because $a \in A$ but $a \notin B$. $B \not\subseteq A$ because $\{a\} \in B$ but $\{a\} \notin A$. \square

1.1.5 (e)

$$A = \{\sqrt{16}, \{4\}\}, B = \{4\}$$

Proof. $A = \{\sqrt{16}, \{4\}\} = \{4, \{4\}\}$. $B \subseteq A$ because 4 is the only element of B and $4 \in A$. $A \not\subseteq B$ because $\{4\} \in A$ but $\{4\} \notin B$. \square

1.1.6 (f)

$$A = \{x \in \mathbb{R} \mid \cos(x) \in \mathbb{Z}\}, B = \{x \in \mathbb{R} \mid \sin(x) \in \mathbb{Z}\}$$

Proof. From trigonometry we know that $\cos(x)$ and $\sin(x)$ have the only integer values $-1, 0$ and 1 . We also know that

$\cos(x) = -1$ if and only if $x = \pi + 2n\pi$ for some integer $n \in \mathbb{Z}$ ($\dots, -3\pi, -\pi, \pi, 3\pi, \dots$),

$\cos(x) = 0$ if and only if $x = \frac{\pi}{2} + n\pi$ for some integer $n \in \mathbb{Z}$ ($\dots, -\frac{3\pi}{2}, -\frac{\pi}{2}, \frac{\pi}{2}, \frac{3\pi}{2}, \dots$),

$\cos(x) = 1$ if and only if $x = 2n\pi$ for some integer $n \in \mathbb{Z}$ ($\dots, -4\pi, -2\pi, 0, 2\pi, 4\pi, \dots$),

$\sin(x) = -1$ if and only if $x = -\frac{\pi}{2} + 2n\pi$ for some integer $n \in \mathbb{Z}$ ($\dots, -\frac{5\pi}{2}, -\frac{\pi}{2}, \frac{3\pi}{2}, \frac{7\pi}{2}, \dots$),

$\sin(x) = 0$ if and only if $x = n\pi$ for some integer $n \in \mathbb{Z}$, ($\dots, -2\pi, -\pi, 0, \pi, 2\pi, \dots$),

$\sin(x) = 1$ if and only if $x = \frac{\pi}{2} + 2n\pi$ for some integer $n \in \mathbb{Z}$ ($\dots, -\frac{7\pi}{2}, -\frac{3\pi}{2}, \frac{\pi}{2}, \frac{5\pi}{2}, \dots$).

So:

$$A = \{\dots, -\frac{5\pi}{2}, -2\pi, -\frac{3\pi}{2}, -\pi, -\frac{\pi}{2}, 0, \frac{\pi}{2}, \pi, \frac{3\pi}{2}, 2\pi, \frac{5\pi}{2}, \dots\}$$

and

$$B = \{\dots, -\frac{5\pi}{2}, -2\pi, -\frac{3\pi}{2}, -\pi, -\frac{\pi}{2}, 0, \frac{\pi}{2}, \pi, \frac{3\pi}{2}, 2\pi, \frac{5\pi}{2}, \dots\}$$

so $A = B$. \square

1.2 Exercise 2

Complete the proof from Example 6.1.3: Prove that $B \subseteq A$ where $A = \{m \in \mathbb{Z} \mid m = 2a \text{ for some integer } a\}$ and $B = \{n \in \mathbb{Z} \mid n = 2b - 2 \text{ for some integer } b\}$

Proof. Proof That $B \subseteq A$: Suppose x is a particular but arbitrarily chosen element of B .

[We must show that $x \in A$. By definition of A , this means we must show that $x = 2 \cdot (\text{some integer})$.]

By definition of B , there is an integer b such that $x = 2b - 2$.

[Given that $x = 2b - 2$, can x also be expressed as $2 \cdot (\text{some integer})$? That is, is there an integer, say, a , such that $2b - 2 = 2a$? Solve for a to obtain $a = b - 1$. Check to see if this works.]

Let $a = b - 1$.

[First check that a is an integer.]

We know that a is an integer because it is a difference of integers.

[Then check that $x = 2a$.]

By substitution, $2a = 2(b - 1) = 2b - 2 = x$. Thus, by definition of A , x is an element of A , [as was to be shown]. \square

1.3 Exercise 3

Let sets R, S , and T be defined as follows:

$$R = \{x \in \mathbb{Z} \mid x \text{ is divisible by } 2\}$$

$$S = \{y \in \mathbb{Z} \mid y \text{ is divisible by } 3\}$$

$$T = \{z \in \mathbb{Z} \mid z \text{ is divisible by } 6\}$$

Prove or disprove each of the following statements.

1.3.1 (a)

$$R \subseteq T$$

Proof. $R \not\subseteq T$ because there are elements in R that are not in T . For example, the number 2 is in R but 2 is not in T since 2 is not divisible by 6. \square

1.3.2 (b)

$$T \subseteq R$$

Proof. $T \subseteq R$ because every element in T is in R since every integer divisible by 6 is divisible by 2. To see why this is so, suppose n is any integer that is divisible by 6. Then $n = 6m$ for some integer m . Since $6m = 2(3m)$ and since $3m$ is an integer (being a product of integers), it follows that $n = 2 \cdot (\text{some integer})$, and, hence, that n is divisible by 2. \square

1.3.3 (c)

$$T \subseteq S$$

Proof. $T \subseteq S$ because every element in T is in S since every integer divisible by 6 is divisible by 3. To see why this is so, suppose n is any integer that is divisible by 6. Then $n = 6m$ for some integer m . Since $6m = 3(2m)$ and since $2m$ is an integer (being a product of integers), it follows that $n = 3 \cdot (\text{some integer})$, and, hence, that n is divisible by 3. \square

1.4 Exercise 4

Let $A = \{n \in \mathbb{Z} \mid n = 5r \text{ for some integer } r\}$ and

$B = \{m \in \mathbb{Z} \mid m = 20s \text{ for some integer } s\}$. Prove or disprove each of the following statements.

1.4.1 (a)

$$A \subseteq B$$

Proof. $A \not\subseteq B$ because $5 \in A$ but $5 \notin B$. □

1.4.2 (b)

$$B \subseteq A$$

Proof. $B \subseteq A$ is true. Suppose $m \in B$. Then $m = 20s$ for some integer s . So $m = 20s = 5(4s)$ where $4s$ is an integer, therefore $m \in A$. □

1.5 Exercise 5

Let $C = \{n \in \mathbb{Z} \mid n = 6r - 5 \text{ for some integer } r\}$ and $D = \{m \in \mathbb{Z} \mid m = 3s + 1 \text{ for some integer } s\}$. Prove or disprove each of the following statements.

1.5.1 (a)

$$C \subseteq D$$

Proof. $C \subseteq D$ because every element in C is in D . To see why this is so, suppose n is any element of C . Then $n = 6r - 5$ for some integer r . Let $s = 2r - 2$. Then s is an integer (because products and differences of integers are integers), and $3s + 1 = 3(2r - 2) + 1 = 6r - 6 + 1 = 6r - 5$, which equals n . Thus n satisfies the condition for being in D . Hence, every element in C is in D . □

1.5.2 (b)

$$D \subseteq C$$

Proof. $D \not\subseteq C$ because there are elements of D that are not in C . For example, 4 is in D because $4 = 3 \cdot 1 + 1$. But 4 is not in C because if it were, then $4 = 6r - 5$ for some integer r , which would imply that $9 = 6r$, or, equivalently, that $r = 3/2$, and this contradicts the fact that r is an integer. □

1.6 Exercise 6

Let $A = \{x \in \mathbb{Z} \mid x = 5a + 2 \text{ for some integer } a\}$,

$B = \{y \in \mathbb{Z} \mid y = 10b - 3 \text{ for some integer } b\}$ and

$D = \{z \in \mathbb{Z} \mid z = 10c + 7 \text{ for some integer } c\}$. Prove or disprove each of the following statements.

1.6.1 (a)

$$A \subseteq B$$

Proof. $A \not\subseteq B$ because $2 \in A$ because $2 = 5 \cdot 0 + 2$, but $2 \notin B$. To see this, argue by contradiction and assume $2 \in B$. So $2 = 10b - 3$ for some integer b . So $5 = 10b$ and $b = 1/2$ is an integer, a contradiction. Therefore our supposition was false and $2 \notin B$. \square

1.6.2 (b)

$$B \subseteq A$$

Proof. Suppose $y \in B$. Then $y = 10b - 3$ for some integer b . Then $y = 10b - 5 + 5 - 3 = 5(b - 2) + 2$ where $b - 2$ is an integer. Let $a = b - 2$. Therefore $y = 5a + 2$ for some integer a , therefore $y \in A$. This proves $B \subseteq A$. \square

1.6.3 (c)

$$B = C$$

Proof. Sketch of proof that $B \subseteq C$: If r is any element of B then there is an integer b such that $r = 10b - 3$. To show that r is in C , you must show that there is an integer c such that $r = 10c + 7$. In scratch work, assume that c exists and use the information that $10b - 3$ would have to equal $10c + 7$ to deduce the only possible value for c . Then show that this value is (1) an integer and (2) satisfies the equation $r = 10c + 7$, which will allow you to conclude that r is an element of C .

Sketch of proof that $C \subseteq B$: If s is any element of C then there is an integer c such that $s = 10c + 7$. To show that s is in B , you must show that there is an integer b such that $s = 10b - 3$. In scratch work, assume that b exists and use the information that $10c + 7$ would have to equal $10b - 3$ to deduce the only possible value for b . Then show that this value is (1) an integer and (2) satisfies the equation $s = 10b - 3$, which will allow you to conclude that s is an element of B . \square

1.7 Exercise 7

Let $A = \{x \in \mathbb{Z} \mid x = 6a + 4 \text{ for some integer } a\}$,

$B = \{y \in \mathbb{Z} \mid y = 18b - 2 \text{ for some integer } b\}$ and

$D = \{z \in \mathbb{Z} \mid z = 18c + 16 \text{ for some integer } c\}$. Prove or disprove each of the following statements.

1.7.1 (a)

$$A \subseteq B$$

Proof. $A \not\subseteq B$ because $10 \in A$ but $10 \notin B$. To see this, argue by contradiction and assume $10 \in B$. Then $10 = 18b - 2$ for some integer b . Then $12 = 18b$ and $b = 12/18 = 2/3$, which is not an integer, contradiction. Therefore our supposition was false and $10 \notin B$. \square

1.7.2 (b)

$$B \subseteq A$$

Proof. Suppose $y \in B$. Then $y = 18b - 2$ for some integer b . Then $y = 18b - 6 + 6 - 2 = 6(3b - 1) + 4$ where $3b - 1$ is an integer. Let $a = 3b - 1$. So $y = 6a + 4$ for some integer a , therefore $y \in A$. This proves $B \subseteq A$. \square

1.7.3 (c)

$$B = C$$

Proof. Suppose $y \in B$. Then $y = 18b - 2$ for some integer b . Then $y = 18b - 18 + 18 - 2 = 18(b - 1) + 16$ where $b - 1$ is an integer. Let $c = b - 1$. So $y = 18c + 16$ for some integer c , therefore $y \in C$. This proves $B \subseteq C$.

Suppose $z \in C$. Then $z = 18c + 16$ for some integer c . Then $z = 18c + 18 - 18 + 16 = 18(c + 1) - 2$ where $c + 1$ is an integer. Let $b = c + 1$. So $z = 18b - 2$ for some integer b , therefore $z \in B$. This proves $C \subseteq B$. \square

1.8 Exercise 8

Write in words how to read each of the following out loud. Then write each set using the symbols for union, intersection, set difference, or set complement.

1.8.1 (a)

$$\{x \in U \mid x \in A \text{ and } x \in B\}$$

Proof. In words: The set of all x in U such that x is in A and x is in B .

In symbolic notation: $A \cap B$. \square

1.8.2 (b)

$$\{x \in U \mid x \in A \text{ or } x \in B\}$$

Proof. In words: The set of all x in U such that x is in A or x is in B .

In symbolic notation: $A \cup B$. □

1.8.3 (c)

$$\{x \in U \mid x \in A \text{ and } x \notin B\}$$

Proof. In words: The set of all x in U such that x is in A and x is not in B .

In symbolic notation: $A - B$. □

1.8.4 (d)

$$\{x \in U \mid x \notin A\}$$

Proof. In words: The set of all x in U such that x is not in A .

In symbolic notation: A^c . □

1.9 Exercise 9

Complete the following sentences without using the symbols \cup , \cap , or $-$.

1.9.1 (a)

$x \notin A \cup B$ if, and only if, ____ .

Proof. $x \notin A$ and $x \notin B$ □

1.9.2 (b)

$x \notin A \cap B$ if, and only if, ____ .

Proof. $x \notin A$ or $x \notin B$ □

1.9.3 (c)

$x \notin A - B$ if, and only if, ____ .

Proof. $x \notin A$ or $x \in B$ □

1.10 Exercise 10

Let $A = \{1, 3, 5, 7, 9\}$, $B = \{3, 6, 9\}$, and $C = \{2, 4, 6, 8\}$. Find each of the following:

1.10.1 (a)

$$A \cup B$$

Proof. $A \cup B = \{1, 3, 5, 6, 7, 9\}$

□

1.10.2 (b)

$$A \cap B$$

Proof. $A \cap B = \{3, 9\}$

□

1.10.3 (c)

$$A \cup C$$

Proof. $A \cup C = \{1, 2, 3, 4, 5, 6, 7, 8, 9\}$

□

1.10.4 (d)

$$A \cap C$$

Proof. $A \cap C = \emptyset$

□

1.10.5 (e)

$$A - B$$

Proof. $A - B = \{1, 5, 7\}$

□

1.10.6 (f)

$$B - A$$

Proof. $B - A = \{6\}$

□

1.10.7 (g)

$$B \cup C$$

Proof. $B \cup C = \{2, 3, 4, 6, 8, 9\}$

□

1.10.8 (h)

$$B \cap C$$

Proof. $B \cap C = \{6\}$

□

1.11 Exercise 11

Let the universal set be \mathbb{R} , the set of all real numbers, and let $A = \{x \in \mathbb{R} \mid 0 < x \leq 2\}$, $B = \{x \in \mathbb{R} \mid 1 \leq x < 4\}$, and $C = \{x \in \mathbb{R} \mid 3 \leq x < 9\}$. Find each of the following:

1.11.1 (a)

$$A \cup B$$

Proof. $A \cup B = \{x \in \mathbb{R} \mid 0 < x < 4\}$

□

1.11.2 (b)

$$A \cap B$$

Proof. $A \cap B = \{x \in \mathbb{R} \mid 1 \leq x \leq 2\}$

□

1.11.3 (c)

$$A^c$$

Proof. $A^c = \{x \in \mathbb{R} \mid x \leq 0 \text{ or } 2 < x\}$

□

1.11.4 (d)

$$A \cup C$$

Proof. $A \cup C = \{x \in \mathbb{R} \mid 0 < x \leq 2 \text{ or } 3 \leq x < 9\}$

□

1.11.5 (e)

$$A \cap C$$

Proof. $A \cap C = \emptyset$

□

1.11.6 (f)

$$B^c$$

Proof. $B^c = \{x \in \mathbb{R} \mid x < 1 \text{ or } 4 \leq x\}$

□

1.11.7 (g)

$$A^c \cap B^c$$

Proof. $A^c \cap B^c = \{x \in \mathbb{R} \mid x \leq 0 \text{ or } 4 \leq x\}$

□

1.11.8 (h)

$$A^c \cup B^c$$

Proof. $A^c \cup B^c = \{x \in \mathbb{R} \mid x < 1 \text{ or } 2 < x\}$

□

1.11.9 (i)

$$(A \cap B)^c$$

Proof. $(A \cap B)^c = \{x \in \mathbb{R} \mid x < 1 \text{ or } 2 < x\}$

□

1.11.10 (j)

$$(A \cup B)^c$$

Proof. $(A \cup B)^c = \{x \in \mathbb{R} \mid x \leq 0 \text{ or } 4 \leq x\}$

□

1.12 Exercise 12

Let the universal set be R , the set of all real numbers, and let $A = \{x \in R \mid -3 \leq x \leq 0\}$, $B = \{x \in R \mid -1 < x < 2\}$, and $C = \{x \in R \mid 6 < x \leq 8\}$. Find each of the following:

1.12.1 (a)

$$A \cup B$$

Proof. $A \cup B = \{x \in \mathbb{R} \mid -3 \leq x < 2\}$

□

1.12.2 (b)

$$A \cap B$$

Proof. $A \cap B = \{x \in \mathbb{R} \mid -1 < x \leq 0\}$

□

1.12.3 (c)

$$A^c$$

Proof. $A^c = \{x \in \mathbb{R} \mid x < -3 \text{ or } 0 < x\}$

□

1.12.4 (d)

$$A \cup C$$

Proof. $A \cup C = \{x \in \mathbb{R} \mid -3 \leq x \leq 0 \text{ or } 6 < x \leq 8\}$

□

1.12.5 (e)

$$A \cap C$$

$$\textit{Proof. } A \cap C = \emptyset$$

□

1.12.6 (f)

$$B^c$$

$$\textit{Proof. } B^c = \{x \in \mathbb{R} \mid x \leq -1 \text{ or } 2 \leq x\}$$

□

1.12.7 (g)

$$A^c \cap B^c$$

$$\textit{Proof. } A^c \cap B^c = \{x \in \mathbb{R} \mid x < -3 \text{ or } 2 \leq x\}$$

□

1.12.8 (h)

$$A^c \cup B^c$$

$$\textit{Proof. } A^c \cup B^c = \{x \in \mathbb{R} \mid x \leq -1 \text{ or } 0 < x\}$$

□

1.12.9 (i)

$$(A \cap B)^c$$

$$\textit{Proof. } (A \cap B)^c = \{x \in \mathbb{R} \mid x \leq -1 \text{ or } 0 < x\}$$

□

1.12.10 (j)

$$(A \cup B)^c$$

$$\textit{Proof. } (A \cup B)^c = \{x \in \mathbb{R} \mid x < -3 \text{ or } 2 \leq x\}$$

□

1.13 Exercise 13

Let S be the set of all strings of 0's and 1's of length 4, and let A and B be the following subsets of S : $A = \{1110, 1111, 1000, 1001\}$ and $B = \{1100, 0100, 1111, 0111\}$. Find each of the following:

1.13.1 (a)

$$A \cap B$$

$$\textit{Proof. } A \cap B = \{1111\}$$

□

1.13.2 (b)

$$A \cup B$$

Proof. $A \cup B = \{1110, 1111, 1000, 1001, 1100, 0100, 0111\}$

□

1.13.3 (c)

$$A - B$$

Proof. $A - B = \{1110, 1000, 1001\}$

□

1.13.4 (d)

$$B - A$$

Proof. $B - A = \{1100, 0100, 0111\}$


□

1.14 Exercise 14

In each of the following, draw a Venn diagram for sets A , B , and C that satisfy the given conditions.

1.14.1 (a)

$$A \subseteq B, C \subseteq B, A \cap C = \emptyset$$



Proof.

□

1.14.2 (b)


$$C \subseteq A, B \cap C = \emptyset$$

Proof.

□


1.15 Exercise 15

In each of the following, draw a Venn diagram for sets A , B , and C that satisfy the given conditions.



1.15.1 (a)

$$A \cap B = \emptyset, A \subseteq C, C \cap B \neq \emptyset$$




Proof.

□

1.15.2 (b)

$$A \subseteq B, C \subseteq B, A \cap C \neq \emptyset$$



Proof.

□

1.15.3 (c)


$$A \cap B \neq \emptyset, B \cap C \neq \emptyset, A \cap C = \emptyset, A \not\subseteq B, C \not\subseteq B$$

Proof.

□

1.16 Exercise 16

Let $A = \{a, b, c\}$, $B = \{b, c, d\}$, $C = \{b, c, e\}$.



1.16.1 (a)

Find $A \cup (B \cap C)$, $(A \cup B) \cap C$, and $(A \cup B) \cap (A \cup C)$. Which of these sets are equal?

Proof. $A \cup (B \cap C) = \{a, b, c\}$, $(A \cup B) \cap C = \{b, c\}$, and $(A \cup B) \cap (A \cup C) = \{a, b, c, d\} \cap \{a, b, c, e\} = \{a, b, c\}$. Hence $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$. \square

1.16.2 (b)

Find $A \cap (B \cup C)$, $(A \cap B) \cap C$, and $(A \cap B) \cup (A \cap C)$. Which of these sets are equal?


Proof. $A \cap (B \cup C) = \{b, c\}$, $(A \cap B) \cap C = \{b, c\}$, and $(A \cap B) \cup (A \cap C) = \{b, c\} \cup \{b, c\} = \{b, c\}$. Hence all three sets are equal. \square

1.16.3 (c)

Find $(A - B) - C$ and $A - (B - C)$. Are these sets equal?

Proof. $(A - B) - C = \{a\} - \{b, c, e\} = \{a\}$ and $A - (B - C) = \{a, b, c\} - \{d\} = \{a, b, c\}$. The sets are not equal. \square

1.17 Exercise 17




Consider the Venn diagram. For each of (a)-(f), copy the diagram and shade the region corresponding to the indicated set.

1.17.1 (a)

$A \cap B$


Proof.

\square



1.17.2 (b)

$$B \cup C$$




Proof.



1.17.3 (c)

$$A^c$$




Proof.



1.17.4 (d)

$$A - (B \cup C)$$




Proof.



1.17.5 (e)

$$(A \cup B)^c$$



Proof.

□

1.17.6 (f)

$$A^c \cap B^c$$

Proof. Same as (e).

□

1.18 Exercise 18

1.18.1 (a)

Is the number 0 in \emptyset ? Why?

Proof. The number 0 is not in \emptyset because \emptyset has no elements.

□

1.18.2 (b)

Is $\emptyset = \{\emptyset\}$? Why?

Proof. No. The left-hand set is the empty set; it does not have any elements. The right-hand set is a set with one element, namely \emptyset .

□

1.18.3 (c)

Is $\emptyset \in \{\emptyset\}$? Why?

Proof. Yes. The left-hand side is the empty set; the right-hand side is a set with one element, namely \emptyset . So the left-hand side is an element of the right-hand side.

□

1.18.4 (d)

Is $\emptyset \in \emptyset$? Why?

Proof. \emptyset is not in \emptyset because \emptyset has no elements.

□

1.19 Exercise 19

Let $A_i = \{i, i^2\}$ for each integer $i = 1, 2, 3, 4$.

1.19.1 (a)

$$A_1 \cup A_2 \cup A_3 \cup A_4 = ?$$

Proof. $A_1 = \{1, 1^2\} = \{1\}$, $A_2 = \{2, 2^2\} = \{2, 4\}$, $A_3 = \{3, 3^2\} = \{3, 9\}$, $A_4 = \{4, 4^2\} = \{4, 16\}$

$$A_1 \cup A_2 \cup A_3 \cup A_4 = \{1\} \cup \{2, 4\} \cup \{3, 9\} \cup \{4, 16\} = \{1, 2, 3, 4, 9, 16\} \quad \square$$

1.19.2 (b)

$$A_1 \cap A_2 \cap A_3 \cap A_4 = ?$$

Proof. $A_1 \cap A_2 \cap A_3 \cap A_4 = \{1\} \cap \{2, 4\} \cap \{3, 9\} \cap \{4, 16\} = \emptyset \quad \square$

1.19.3 (c)

Are A_1, A_2, A_3, A_4 mutually disjoint? Explain.

Proof. A_1, A_2, A_3, A_4 are not mutually disjoint, because $A_2 \cap A_4 = \{4\} \neq \emptyset. \quad \square$

1.20 Exercise 20

Let $B_i = \{x \in \mathbb{R} \mid 0 \leq x \leq i\}$ for each integer $i = 1, 2, 3, 4$.

1.20.1 (a)

$$B_1 \cup B_2 \cup B_3 \cup B_4 = ?$$

Proof. $B_1 \cup B_2 \cup B_3 \cup B_4 = B_4 = [0, 4] \quad \square$

1.20.2 (b)

$$B_1 \cap B_2 \cap B_3 \cap B_4 = ?$$

Proof. $B_1 \cap B_2 \cap B_3 \cap B_4 = B_1 = [0, 1] \quad \square$

1.20.3 (c)

Are B_1, B_2, B_3, B_4 mutually disjoint? Explain.

Proof. No, because for example $B_1 \cap B_2 = B_1 \neq \emptyset. \quad \square$

1.21 Exercise 21

Let $C_i = \{-i, i\}$ for each nonnegative integer i .

1.21.1 (a)

$$\bigcup_{i=0}^4 C_i = ?$$

Proof. $C_0 = \{0, -0\} = \{0\}$, $C_1 = \{1, -1\}$, $C_2 = \{2, -2\}$, $C_3 = \{3, -3\}$, $C_4 = \{4, -4\}$

$$\bigcup_{i=0}^4 C_i = \{-4, -3, -2, -1, 0, 1, 2, 3, 4\}$$

□

1.21.2 (b)

$$\bigcap_{i=0}^4 C_i = ?$$

$$\textit{Proof. } \bigcap_{i=0}^4 C_i = \{0\} \cap \{1, -1\} \cap \{2, -2\} \cap \{3, -3\} \cap \{4, -4\} = \emptyset$$

□

1.21.3 (c)

Are C_0, C_1, C_2, \dots mutually disjoint? Explain.

Proof. C_0, C_1, C_2, \dots are mutually disjoint because no two of the sets have any elements in common.

□

1.21.4 (d)

$$\bigcup_{i=0}^n C_i = ?$$

$$\textit{Proof. } \bigcup_{i=0}^n C_i = \{-n, -(n-1), \dots, -2, -1, 0, 1, 2, \dots, n-1, n\}$$

□

1.21.5 (e)

$$\bigcap_{i=0}^n C_i = ?$$

$$\textit{Proof. } \bigcap_{i=0}^n C_i = \emptyset$$

□

1.21.6 (f)

$$\bigcup_{i=0}^{\infty} C_i = ?$$

Proof. $\bigcup_{i=0}^{\infty} C_i = \mathbb{Z}$, the set of all integers □

1.21.7 (g)

$$\bigcap_{i=0}^{\infty} C_i = ?$$

Proof. $\bigcap_{i=0}^{\infty} C_i = \emptyset$ □

1.22 Exercise 22

Let $D_i = \{x \in \mathbb{R} \mid -i \leq x \leq i\} = [-i, i]$ for each nonnegative integer i .

1.22.1 (a)

$$\bigcup_{i=0}^4 D_i = ?$$

Proof. $D_0 = [-0, 0] = \{0\}$, $D_1 = [-1, 1]$, $D_2 = [-2, 2]$, $D_3 = [-3, 3]$, $D_4 = [-4, 4]$

$$\bigcup_{i=0}^4 D_i = \{0\} \cup [-1, 1] \cup [-2, 2] \cup [-3, 3] \cup [-4, 4] = [-4, 4] \quad \square$$

1.22.2 (b)

$$\bigcap_{i=0}^4 D_i = ?$$

Proof. $\bigcap_{i=0}^4 D_i = \{0\} \cap [-1, 1] \cap [-2, 2] \cap [-3, 3] \cap [-4, 4] = \{0\}$ □

1.22.3 (c)

Are D_0, D_1, D_2, \dots mutually disjoint? Explain.

Proof. D_0, D_1, D_2, \dots , are not mutually disjoint. In fact, each $D_k \subseteq D_{k+1}$. □

1.22.4 (d)

$$\bigcup_{i=0}^n D_i = ?$$

$$\textit{Proof. } \bigcup_{i=0}^n D_i = [-n, n]$$

□

1.22.5 (e)

$$\bigcap_{i=0}^n D_i = ?$$

$$\textit{Proof. } \bigcap_{i=0}^n D_i = \{0\}$$

□

1.22.6 (f)

$$\bigcup_{i=0}^{\infty} D_i = ?$$

$$\textit{Proof. } \bigcup_{i=0}^{\infty} D_i = \mathbb{R}, \text{ the set of all real numbers}$$

□

1.22.7 (g)

$$\bigcap_{i=0}^{\infty} D_i = ?$$

$$\textit{Proof. } \bigcap_{i=0}^{\infty} D_i = \{0\}$$

□

1.23 Exercise 23

Let $V_i = \{x \in \mathbb{R} \mid -\frac{1}{i} \leq x \leq \frac{1}{i}\} = [-\frac{1}{i}, \frac{1}{i}]$ for each positive integer i .

1.23.1 (a)

$$\bigcup_{i=1}^4 V_i = ?$$

$$\textit{Proof. } \bigcup_{i=1}^4 V_i = V_1 = [-1, 1]$$

□

1.23.2 (b)

$$\bigcap_{i=1}^4 V_i = ?$$

$$\textit{Proof. } \bigcap_{i=1}^4 V_i = V_4 = \left[-\frac{1}{4}, \frac{1}{4}\right]$$

□

1.23.3 (c)

Are V_1, V_2, V_3, \dots mutually disjoint? Explain.

Proof. V_1, V_2, V_3, \dots , are not mutually disjoint. In fact, each $V_{k+1} \subseteq V_k$.

□

1.23.4 (d)

$$\bigcup_{i=1}^n V_i = ?$$

$$\textit{Proof. } \bigcup_{i=1}^n V_i = V_1 = [-1, 1]$$

□

1.23.5 (e)

$$\bigcap_{i=1}^n V_i = ?$$

$$\textit{Proof. } \bigcap_{i=1}^n V_i = V_n = \left[-\frac{1}{n}, \frac{1}{n}\right]$$

□

1.23.6 (f)

$$\bigcup_{i=1}^{\infty} V_i = ?$$

$$\textit{Proof. } \bigcup_{i=1}^{\infty} V_i = V_1 = [-1, 1]$$

□

1.23.7 (g)

$$\bigcap_{i=1}^{\infty} V_i = ?$$

$$\textit{Proof. } \bigcap_{i=1}^{\infty} V_i = \{0\}$$

□

1.24 Exercise 24

Let $W_i = \{x \in \mathbb{R} \mid i < x\} = (i, \infty)$ for each nonnegative integer i .

1.24.1 (a)

$$\bigcup_{i=0}^4 W_i = ?$$

$$\textit{Proof. } \bigcup_{i=0}^4 W_i = W_0 = (0, \infty)$$

□

1.24.2 (b)

$$\bigcap_{i=0}^4 W_i = ?$$

$$\textit{Proof. } \bigcap_{i=0}^4 W_i = W_4 = (4, \infty)$$

□

1.24.3 (c)

Are W_0, W_1, W_2, \dots mutually disjoint? Explain.

Proof. W_0, W_1, W_2, \dots , are not mutually disjoint. In fact, each $W_{k+1} \subseteq W_k$.

□

1.24.4 (d)

$$\bigcup_{i=0}^n W_i = ?$$

$$\textit{Proof. } \bigcup_{i=0}^n W_i = W_0 = (0, \infty)$$

□

1.24.5 (e)

$$\bigcap_{i=0}^n W_i = ?$$

$$\textit{Proof. } \bigcap_{i=0}^n W_i = W_n = (n, \infty)$$

□

1.24.6 (f)

$$\bigcup_{i=0}^{\infty} W_i = ?$$

$$\textit{Proof. } \bigcup_{i=0}^{\infty} W_i = W_0 = (0, \infty)$$

□

1.24.7 (g)

$$\bigcap_{i=0}^{\infty} W_i = ?$$

$$\textit{Proof. } \bigcap_{i=0}^{\infty} W_i = \emptyset$$

□

1.25 Exercise 25

Let $R_i = \{x \in \mathbb{R} \mid 1 \leq x \leq 1 + \frac{1}{i}\} = [1, 1 + \frac{1}{i}]$ for each positive integer i .

1.25.1 (a)

$$\bigcup_{i=1}^4 R_i = ?$$

$$\textit{Proof. } \bigcup_{i=1}^4 R_i = R_1 = [1, 2]$$

□

1.25.2 (b)

$$\bigcap_{i=1}^4 R_i = ?$$

$$\textit{Proof. } \bigcap_{i=1}^4 R_i = R_4 = \left[1, \frac{5}{4}\right]$$

□

1.25.3 (c)

Are R_1, R_2, R_3, \dots mutually disjoint? Explain.

Proof. No, in fact each $R_{k+1} \subseteq R_k$.

□

1.25.4 (d)

$$\bigcup_{i=1}^n R_i = ?$$

$$\textit{Proof. } \bigcup_{i=1}^n R_i = R_1 = [1, 2]$$

□

1.25.5 (e)

$$\bigcap_{i=1}^n R_i = ?$$

$$\textit{Proof. } \bigcap_{i=1}^n R_i = R_n = \left[1, 1 + \frac{1}{n}\right]$$

□

1.25.6 (f)

$$\bigcup_{i=1}^{\infty} R_i = ?$$

$$\textit{Proof. } \bigcup_{i=1}^{\infty} R_i = R_1 = [1, 2]$$

□

1.25.7 (g)

$$\bigcap_{i=1}^{\infty} R_i = ?$$

$$\textit{Proof. } \bigcap_{i=1}^{\infty} R_i = \{1\}$$

□

1.26 Exercise 26

Let $S_i = \{x \in \mathbb{R} \mid 1 < x < 1 + \frac{1}{i}\} = (1, 1 + \frac{1}{i})$ for each positive integer i .

1.26.1 (a)

$$\bigcup_{i=1}^4 S_i = ?$$

$$\textit{Proof. } \bigcup_{i=1}^4 S_i = S_1 = (1, 2)$$

□

1.26.2 (b)

$$\bigcap_{i=1}^4 S_i = ?$$

$$\textit{Proof. } \bigcap_{i=1}^4 S_i = S_4 = (1, 5/4)$$

□

1.26.3 (c)

Are S_0, S_1, S_2, \dots mutually disjoint? Explain.

$$\textit{Proof. } \text{No, in fact each } S_{k+1} \subseteq S_k.$$

□

1.26.4 (d)

$$\bigcup_{i=1}^n S_i = ?$$

$$\textit{Proof. } \bigcup_{i=1}^n S_i = S_1 = (1, 2)$$

□

1.26.5 (e)

$$\bigcap_{i=1}^n S_i = ?$$

$$\textit{Proof. } \bigcap_{i=1}^n S_i = S_n = (1, \frac{n+1}{n})$$

□

1.26.6 (f)

$$\bigcup_{i=1}^{\infty} S_i = ?$$

$$\textit{Proof. } \bigcup_{i=1}^{\infty} S_i = S_1 = (1, 2)$$

□

1.26.7 (g)

$$\bigcap_{i=1}^{\infty} S_i = ?$$

Proof. $\bigcap_{i=1}^{\infty} S_i = (1, 1) = \emptyset$

□

1.27 Exercise 27

1.27.1 (a)

Is $\{\{a, d, e\}, \{b, c\}, \{d, f\}\}$ a partition of $\{a, b, c, d, e, f\}$?

Proof. No. The element d is in two of the sets.

□

1.27.2 (b)

Is $\{\{w, x, v\}, \{u, y, q\}, \{p, z\}\}$ a partition of $\{p, q, u, v, w, x, y, z\}$?

Proof. Yes.

□

1.27.3 (c)

Is $\{\{5, 4\}, \{7, 2\}, \{1, 3, 4\}, \{6, 8\}\}$ a partition of $\{1, 2, 3, 4, 5, 6, 7, 8\}$?

Proof. No. The element 4 is in two of the sets.

□

1.27.4 (d)

Is $\{\{3, 7, 8\}, \{2, 9\}, \{1, 4, 5\}\}$ a partition of $\{1, 2, 3, 4, 5, 6, 7, 8, 9\}$?

Proof. No. None of the sets contains 6.

□

1.27.5 (e)

Is $\{\{1, 5\}, \{4, 7\}, \{2, 8, 6, 3\}\}$ a partition of $\{1, 2, 3, 4, 5, 6, 7, 8\}$?

Proof. Yes.

□

1.28 Exercise 28

Let E be the set of all even integers and O the set of all odd integers. Is $\{E, O\}$ a partition of \mathbb{Z} , the set of all integers? Explain your answer.

Proof. Yes. Every integer is either even or odd, and no integer is both even and odd.

□

1.29 Exercise 29

Let \mathbb{R} be the set of all real numbers. Is $\{R^+, R^-, \{0\}\}$ a partition of \mathbb{R} ? Explain your answer.

Proof. Yes. Every real number is either positive or negative or zero, and no real number is both positive and negative, and zero is neither negative nor positive (so the three sets are mutually disjoint). \square

1.30 Exercise 30

Let Z be the set of all integers and let

$$\begin{aligned}A_1 &= \{n \in \mathbb{Z} \mid n = 4k \text{ for some integer } k\} \\A_2 &= \{n \in \mathbb{Z} \mid n = 4k + 1 \text{ for some integer } k\} \\A_3 &= \{n \in \mathbb{Z} \mid n = 4k + 2 \text{ for some integer } k\} \\A_4 &= \{n \in \mathbb{Z} \mid n = 4k + 3 \text{ for some integer } k\}\end{aligned}$$

Is $\{A_0, A_1, A_2, A_3\}$ a partition of \mathbb{Z} ? Explain your answer.

Proof. Yes. These sets are mutually disjoint, and by the quotient-remainder theorem, every integer has exactly one of the forms $n = 4k$ or $n = 4k + 1$ or $n = 4k + 2$ or $n = 4k + 3$. \square

1.31 Exercise 31

Suppose $A = \{1, 2\}$, $B = \{2, 3\}$. Find each of the following:

1.31.1 (a)

$$\mathcal{P}(A \cap B)$$

Proof. $A \cap B = \{2\}$, so $\mathcal{P}(A \cap B) = \{\emptyset, \{2\}\}$. \square

1.31.2 (b)

$$\mathcal{P}(A)$$

Proof. $A = \{1, 2\}$, so $\mathcal{P}(A) = \{\emptyset, \{1\}, \{2\}, \{1, 2\}\}$. \square

1.31.3 (c)

$$\mathcal{P}(A \cup B)$$

Proof. $A \cup B = \{1, 2, 3\}$, so $\mathcal{P}(A \cup B) = \{\emptyset, \{1\}, \{2\}, \{3\}, \{1, 2\}, \{1, 3\}, \{2, 3\}, \{1, 2, 3\}\}$. \square

1.31.4 (d)

$$\mathcal{P}(A \times B)$$

Proof. $A \times B = \{(1, 2), (1, 3), (2, 2), (2, 3)\}$, so $\mathcal{P}(A \times B) = \{\emptyset, \{(1, 2)\}, \{(1, 3)\}, \{(2, 2)\}, \{(2, 3)\}, \{(1, 2), (1, 3)\}, \{(1, 2), (2, 2)\}, \{(1, 2), (2, 3)\}, \{(1, 3), (2, 2)\}, \{(1, 3), (2, 3)\}, \{(2, 2), (2, 3)\}, \{(1, 2), (1, 3), (2, 2)\}, \{(1, 2), (1, 3), (2, 3)\}, \{(1, 2), (2, 2), (2, 3)\}, \{(1, 3), (2, 2), (2, 3)\}, \{(1, 2), (1, 3), (2, 2), (2, 3)\}\}$. □

1.32 Exercise 32

1.32.1 (a)

Suppose $A = \{1\}$, $B = \{u, v\}$. Find $\mathcal{P}(A \times B)$.

Proof. $\mathcal{P}(A \times B) = \{\emptyset, \{(1, u)\}, \{(1, v)\}, \{(1, u), (1, v)\}\}$ □

1.32.2 (b)

Suppose $X = \{a, b\}$, $Y = \{x, y\}$. Find $\mathcal{P}(X \times Y)$.

Proof. $X \times Y = \{(a, x), (a, y), (b, x), (b, y)\}$

$\mathcal{P}(X \times Y) = \{\emptyset, \{(a, x)\}, \{(a, y)\}, \{(b, x)\}, \{(b, y)\}, \{(a, x), (a, y)\}, \{(a, x), (b, x)\}, \{(a, x), (b, y)\}, \{(a, y), (b, x)\}, \{(a, y), (b, y)\}, \{(b, x), (b, y)\}, \{(a, x), (a, y), (b, x)\}, \{(a, x), (a, y), (b, y)\}, \{(a, x), (b, x), (b, y)\}, \{(a, y), (b, x), (b, y)\}, \{(a, x), (a, y), (b, x), (b, y)\}\}$ □

1.33 Exercise 33

1.33.1 (a)

Find $\mathcal{P}(\emptyset)$.

Proof. $\mathcal{P}(\emptyset) = \{\emptyset\}$ □

1.33.2 (b)

Find $\mathcal{P}(\mathcal{P}(\emptyset))$.

Proof. $\mathcal{P}(\mathcal{P}(\emptyset)) = \mathcal{P}(\{\emptyset\}) = \{\emptyset, \{\emptyset\}\}$ □

1.33.3 (c)

Find $\mathcal{P}(\mathcal{P}(\mathcal{P}(\emptyset)))$.

Proof. $\mathcal{P}(\mathcal{P}(\mathcal{P}(\emptyset))) = \mathcal{P}(\{\emptyset, \{\emptyset\}\}) = \{\emptyset, \{\emptyset\}, \{\{\emptyset\}\}, \{\emptyset, \{\emptyset\}\}\}$ □

1.34 Exercise 34

Let $A_1 = \{1\}$, $A_2 = \{u, v\}$, $A_3 = \{m, n\}$. Find each of the following sets:

1.34.1 (a)

$$A_1 \cup (A_2 \times A_3)$$

$$\begin{aligned} \text{Proof. } A_1 \cup (A_2 \times A_3) &= \{1\} \cup \{(u, m), (u, n), (v, m), (v, n)\} \\ &= \{1, (u, m), (u, n), (v, m), (v, n)\} \end{aligned}$$

□

1.34.2 (b)

$$(A_1 \cup A_2) \times A_3$$

$$\text{Proof. } (A_1 \cup A_2) \times A_3 = \{1, u, v\} \times \{m, n\} = \{(1, m), (1, n), (u, m), (u, n), (v, m), (v, n)\}$$

□

1.35 Exercise 35

Let $A = \{a, b\}$, $B = \{1, 2\}$, $C = \{2, 3\}$. Find each of the following sets.

1.35.1 (a)

$$A \times (B \cup C)$$

$$\text{Proof. } A \times (B \cup C) = \{a, b\} \times \{1, 2, 3\} = \{(a, 1), (a, 2), (a, 3), (b, 1), (b, 2), (b, 3)\}$$

□

1.35.2 (b)

$$(A \times B) \cup (A \times C)$$

$$\begin{aligned} \text{Proof. } (A \times B) \cup (A \times C) &= \{(a, 1), (a, 2), (b, 1), (b, 2), (a, 2), (a, 3), (b, 2), (b, 3)\} \\ &= \{(a, 1), (a, 2), (b, 1), (b, 2), (a, 3), (b, 3)\} \end{aligned}$$

□

1.35.3 (c)

$$A \times (B \cap C)$$

$$\text{Proof. } A \times (B \cap C) = \{a, b\} \times \{2\} = \{(a, 2), (b, 2)\}$$

□

1.35.4 (d)

$$(A \times B) \cap (A \times C)$$

$$\begin{aligned} \text{Proof. } (A \times B) \cap (A \times C) &= \{(a, 1), (a, 2), (b, 1), (b, 2)\} \cap \{(a, 2), (a, 3), (b, 2), (b, 3)\} \\ &= \{(a, 2), (b, 2)\} \end{aligned}$$

□

1.36 Exercise 36

Trace the action of Algorithm 6.1.1 on the variables $i, j, found$, and $answer$ for $m = 3, n = 3$, and sets A and B represented as the arrays $a[1] = u, a[2] = v, a[3] = w, b[1] = w, b[2] = u, b[3] = v$.

i	1				2				3			4
j		1	2	3	1	2	3	4	1		2	
$found$		no	yes		no		yes		no	yes		
$answer$	$A \subseteq B$											

Proof. □

1.37 Exercise 37

Trace the action of Algorithm 6.1.1 on the variables $i, j, found$, and $answer$ for $m = 4, n = 4$, and sets A and B represented as the arrays $a[1] = u, a[2] = v, a[3] = w, a[4] = x, b[1] = r, b[2] = u, b[3] = y, b[4] = z$.

Proof.

i	1				2			
j	1	2	3	4	1	2	3	4
found	no	yes			no			
answer	$A \subseteq B$							$A \not\subseteq B$

□

1.38 Exercise 38

Write an algorithm to determine whether a given element x belongs to a given set that is represented as the array $a[1], a[2], \dots, a[n]$.

Algorithm: Testing whether $x \in A$

Input: x (an element), n (a positive integer), $a[1], \dots, a[n]$ (a one-dimensional array representing the set A).

Algorithm Body:

$i := 1, answer := "x \notin A"$

while ($i \leq n$ and $answer = "x \notin A"$)

if $x = a[i]$ **then** $answer = "x \in A"$

$i := i + 1$

end while

Output: $answer$ [a string]

2 Exercise Set 6.2

Fill the blanks.

2.1 Exercise 1

2.1.1 (a)

To say that an element is in $A \cap (B \cup C)$ means that it is in (1) _____ and in (2) _____ .

Proof. (1) A (2) $B \cup C$

□

2.1.2 (b)

To say that an element is in $(A \cap B) \cup C$ means that it is in (1) _____ or in (2) _____ .

Proof. (1) $A \cap B$ (2) C

□

2.1.3 (c)

To say that an element is in $A - (B \cap C)$ means that it is in (1) _____ and not in (2) _____ .

Proof. (1) A (2) $B \cap C$

□

2.1.4 (d)

To prove that $(A \cup B) \cap C \subseteq A \cup (B \cap C)$, we suppose that x is any element in (1) _____ . Then we must show that (2) _____ .

Proof. (1) $(A \cup B) \cap C$ (2) $A \cup (B \cap C)$

□

2.1.5 (e)

If A , B , and C are any sets such that $B \subseteq C$, to prove that $A \cap B \subseteq A \cap C$, we suppose that x is any element in (1) _____ . Then we must show that (2) _____ .

Proof. (1) $A \cap B$ (2) $A \cap C$

□

2.2 Exercise 2

The following are two proofs that for all sets A and B , $A - B \subseteq A$. The first is less formal, and the second is more formal. Fill in the blanks.

2.2.1 (a)

Proof: Suppose A and B are any sets. To show that $A - B \subseteq A$, we must show that every element in (1) _____ is in (2) _____. But any element in $A - B$ is in (3) _____ and not in (4) _____ (by definition of $A - B$). In particular, such an element is in A .

Proof. (1) $A - B$ (2) A (3) A (4) B

□

2.2.2 (b)

Proof: Suppose A and B are any sets and $x \in A - B$. [We must show that (1) _____.] By definition of set difference, $x \in$ (2) _____ and $x \notin$ (3) _____. In particular, $x \in$ (4) _____ [which is what was to be shown].

Proof. (1) $x \in A$ (2) A (3) B (4) A

□

In 3 and 4, supply explanations of the steps in the given proofs.

2.3 Exercise 3

Theorem: For all sets A, B , and C , if $A \subseteq B$ and $B \subseteq C$ then $A \subseteq C$.

Proof:

Statement	Explanation
Suppose A, B, C are any sets such that $A \subseteq B$ and $B \subseteq C$.	starting point
We must show that $A \subseteq C$.	conclusion to be shown
Let x be any element in A .	start of an element proof
Then x is in B .	(a) _____
It follows that x is in C .	(b) _____
Thus every element of A is in C .	since x could be any element of A
Therefore $A \subseteq C$ [as was to be shown].	(c) _____

Proof. (a) by definition of subset (because $A \subseteq B$) (b) by definition of subset (because $B \subseteq C$) (c) by definition of subset

□

2.4 Exercise 4

Theorem: For all sets A, B , if $A \subseteq B$ then $A \cup B \subseteq B$.

Proof:

Statement	Explanation
Suppose A, B are any sets such that $A \subseteq B$.	starting point
We must show that $A \cup B \subseteq B$.	conclusion to be shown
Let x be any element in $A \cup B$.	start of an element proof
Then $x \in A$ or $x \in B$.	(a) _____
In case $x \in A$, then $x \in B$.	(b) _____
In case $x \in B$, then $x \in B$.	tautology ($p \rightarrow p$)
So in either case $x \in B$.	proof by division into cases
Thus every element of $A \cup B$ is in B .	since x could be any element of $A \cup B$
Therefore $A \cup B \subseteq B$ [as was to be shown].	(c) _____

Proof. (a) by definition of union (b) by definition of subset (because $A \subseteq B$) (c) by definition of subset □

2.5 Exercise 5

Prove that for all sets A and B , $(B - A) = B \cap A^c$.

Proof. Suppose A and B are any sets.

Proof that $B - A \subseteq B \cap A^c$: Suppose $x \in B - A$. By definition of set difference, $x \in B$ and $x \notin A$. It follows by definition of complement that $x \in B$ and $x \in A^c$, and so by definition of intersection, $x \in B \cap A^c$. [Thus $B - A \subseteq B \cap A^c$ by definition of subset.]

Proof that $B \cap A^c \subseteq B - A$: Suppose $x \in B \cap A^c$. By definition of intersection, $x \in B$ and $x \in A^c$. It follows by definition of complement that $x \in B$ and $x \notin A$, and so by definition of set difference, $x \in B - A$. [Thus $B \cap A^c \subseteq B - A$ by definition of subset.]

[Since both subset relations have been proved, $B - A = B \cap A^c$ by definition of set equality.] □

2.6 Exercise 6

Let \cap and \cup stand for the words “intersection” and “union,” respectively. Fill in the blanks in the following proof that for all sets A, B , and C , $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$.

Proof: Suppose A, B , and C are any sets.

(1) Proof that $A \cap (B \cup C) \subseteq (A \cap B) \cup (A \cap C)$:

Let $x \in A \cap (B \cup C)$. [We must show that $x \in$ (a) _____].

By definition of \cup , $x \in$ (b) _____ and $x \in B \cup C$. Thus $x \in A$ and, by definition of \cup , $x \in B$ or (c) _____.

Case 1 ($x \in A$ and $x \in B$): In this case, $x \in A \cap B$ by definition of \cap .

Case 2 ($x \in A$ and $x \in C$): In this case, $x \in A \cap C$ by definition of \cap .

By cases 1 and 2, $x \in A \cap B$ or $x \in A \cap C$, and so, by definition of \cup , (d) ____ .

[So $A \cap (B \cup C) \subseteq (A \cap B) \cup (A \cap C)$ by definition of subset.]

(2) Proof that $(A \cap B) \cup (A \cap C) \subseteq A \cap (B \cup C)$:

Let $x \in (A \cap B) \cup (A \cap C)$. [We must show that $x \in A \cap (B \cup C)$.]

By definition of \cup , $x \in A \cap B$ (a) ____ $x \in A \cap C$.

Case 1 ($x \in A \cap B$): In this case, by definition of \cap , $x \in A$ and $x \in B$. Since $x \in B$, then $x \in B \cup C$ by definition of \cup .

Case 2 ($x \in A \cap C$): In this case, by definition of \cap , $x \in A$ (b) ____ $x \in C$. Since $x \in C$, then $x \in B \cup C$ by definition of \cup .

In both cases $x \in A$ and $x \in B \cup C$, and so, by definition of \cap , (c) ____ .

[So $(A \cap B) \cup (A \cap C) \subseteq A \cap (B \cup C)$ by definition of (d) ____ .]

(3) Conclusion: [Since both subset relations have been proved, it follows, by definition of set equality, that (a) ____ .]

Proof. (1) (a) $(A \cap B) \cup (A \cap C)$ (b) A (c) $x \in C$ (d) $x \in (A \cap B) \cup (A \cap C)$

(2) (a) or (b) and (c) $x \in A \cap (B \cup C)$ (d) subset

(3) (a) for all sets A, B , and C , $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$. □

Use an element argument to prove each statement in 7 – 22. Assume that all sets are subsets of a universal set U .

2.7 Exercise 7

For all sets A and B , $(A \cap B)^c = A^c \cup B^c$.

Proof. Suppose A and B are any sets.

Proof that $(A \cap B)^c \subseteq A^c \cup B^c$: Suppose $x \in (A \cap B)^c$. By definition of complement, $x \notin (A \cap B)$. It follows by definition of intersection that $x \notin A$ or $x \notin B$, and so by definition of complement, $x \in A^c$ or $x \in B^c$. So by definition of union, $x \in A^c \cup B^c$. [Thus $(A \cap B)^c \subseteq A^c \cup B^c$ by definition of subset.]

Proof that $A^c \cup B^c \subseteq (A \cap B)^c$: Suppose $x \in A^c \cup B^c$. By definition of union, $x \in A^c$ or $x \in B^c$. By definition of complement, $x \notin A$ or $x \notin B$. It follows by definition of intersection that $x \notin (A \cap B)$. Then by definition of complement, $x \in (A \cap B)^c$. [Thus $A^c \cup B^c \subseteq (A \cap B)^c$ by definition of subset.]

[Since both subset relations have been proved, $(A \cap B)^c = A^c \cup B^c$ by definition of set equality.] □

2.8 Exercise 8

For all sets A and B , $(A \cap B) \cup (A \cap B^c) = A$. (This property is used in Section 9.9.)

Proof. Suppose A and B are any sets.

Proof that $(A \cap B) \cup (A \cap B^c) \subseteq A$: Suppose $x \in (A \cap B) \cup (A \cap B^c)$. [We must show that $x \in A$.]

By definition of union, $x \in A \cap B$ or $x \in (A \cap B^c)$.

Case 1 ($x \in A \cap B$): In this case x is in A and x is in B , and so, in particular, $x \in A$.

Case 2 ($x \in A \cap B^c$): In this case x is in A and x is not in B , and so, in particular, $x \in A$.

Thus, in either case, $x \in A$ [as was to be shown].

So $(A \cap B) \cup (A \cap B^c) \subseteq A$ [by definition of subset].

Proof that $A \subseteq (A \cap B) \cup (A \cap B^c)$: Suppose $x \in A$. [We must show that $x \in (A \cap B) \cup (A \cap B^c)$.]

Either $x \in B$ or $x \notin B$.

Case 1 ($x \in B$): In this case we know that x is in A and we are also assuming that x is in B . Hence, by definition of intersection, $x \in A \cap B$.

Case 2 ($x \in A \cap B^c$): In this case we know that x is in A and we are also assuming that x is in B^c . Hence, by definition of intersection, $x \in A \cap B^c$.

Thus, in either case $x \in A \cap B$ or $x \in A \cap B^c$, and so, by definition of union, $x \in (A \cap B) \cup (A \cap B^c)$ [as was to be shown].

So $A \subseteq (A \cap B) \cup (A \cap B^c)$ [by definition of subset].

Conclusion: Since both subset relations have been proved it follows by definition of set equality that $(A \cap B) \cup (A \cap B^c) = A$. \square

2.9 Exercise 9

For all sets A , B , and C , $(A - B) \cup (C - B) = (A \cup C) - B$.

Proof. Suppose A , B , and C are any sets.

To show that $(A - B) \cup (C - B) = (A \cup C) - B$, we must show that $(A - B) \cup (C - B) \subseteq (A \cup C) - B$ and that $(A \cup C) - B \subseteq (A - B) \cup (C - B)$.

Proof that $(A - B) \cup (C - B) \subseteq (A \cup C) - B$:

Suppose that x is any element in $(A - B) \cup (C - B)$. [We must show that $x \in (A \cup C) - B$.]

By definition of union, $x \in A - B$ or $x \in C - B$.

Case 1 ($x \in A - B$): Then, by definition of set difference, $x \in A$ and $x \notin B$. Now because $x \in A$, we have that $x \in A \cup C$ by definition of union. Hence $x \in A \cup C$ and $x \notin B$, and so, by definition of set difference, $x \in (A \cup C) - B$.

Case 2 ($x \in C - B$): Then, by definition of set difference, $x \in C$ and $x \notin B$. Now because $x \in C$, we have that $x \in A \cup C$ by definition of union. Hence $x \in A \cup C$ and $x \notin B$, and so, by definition of set difference, $x \in (A \cup C) - B$. Thus, in both cases, $x \in (A \cup C) - B$ [as was to be shown].

So $(A - B) \cup (C - B) \subseteq (A \cup C) - B$ [by definition of subset].

To complete the proof that $(A - B) \cup (C - B) = (A \cup C) - B$, you must show that $(A \cup C) - B \subseteq (A - B) \cup (C - B)$. \square

2.10 Exercise 10

For all sets A , B , and C , $(A \cup B) \cap C \subseteq A \cup (B \cap C)$.

Proof. Suppose A , B , and C are any sets. We will show that $(A \cup B) \cap C \subseteq A \cup (B \cap C)$.

Suppose x is any element $(A \cup B) \cap C$. By definition of intersection x is in $A \cup B$ and x is in C . Then by definition of union x is in A or x is in B , and in both cases x is in C . It follows by definition of union that in case x is in A and x is in C , then x is in $A \cup (B \cap C)$ by virtue of being in A . And in case x is in $B \cap C$, then x is in $A \cup (B \cap C)$ by virtue of being in $B \cap C$. Thus in both cases x is in $A \cup (B \cap C)$, which proves that every element in $(A \cup B) \cap C$ is in $A \cup (B \cap C)$. Hence $(A \cup B) \cap C \subseteq A \cup (B \cap C)$ by definition of subset. \square

2.11 Exercise 11

For all sets A , B , and C , $A \cap (B - C) \subseteq (A \cap B) - (A \cap C)$.

Proof. Assume A, B, C are any sets.

1. Assume $x \in A \cap (B - C)$. [Want to show $x \in (A \cap B) - (A \cap C)$].
2. By 1 and definition of intersection, $x \in A$ and $x \in B - C$.
3. By 2 and definition of difference, $x \in A$ and $x \in B$ and $x \notin C$.
4. By 3 and definition of intersection, $x \in A \cap B$.
5. By 3 and definition of intersection, $x \notin A \cap C$.
6. By 4 and 5 and definition of difference, $x \in (A \cap B) - (A \cap C)$.
7. By 1 and 6 and definition of subset, $A \cap (B - C) \subseteq (A \cap B) - (A \cap C)$. \square

2.12 Exercise 12

For all sets A , B , and C , $(A \cup B) - C \subseteq (A - C) \cup (B - C)$.

Proof. Assume A, B, C are any sets.

1. Assume $x \in (A \cup B) - C$. [*Want to show $x \in (A - C) \cup (B - C)$.*]
2. By 1 and definition of difference, $x \in A \cup B$ and $x \notin C$.
3. By 2 and definition of union, $x \in A$ or $x \in B$.
4. By 2, 3 and definition of difference, $x \in A - C$ or $x \in B - C$.
5. By 4 and definition of union, $x \in (A - C) \cup (B - C)$.
6. By 1 and 5 and definition of subset, $(A \cup B) - C \subseteq (A - C) \cup (B - C)$. □

2.13 Exercise 13

For all sets A, B , and C , $(A - B) \cap (C - B) = (A \cap C) - B$.

Proof. Assume A, B, C are any sets.

1. Assume $x \in (A - B) \cap (C - B)$. [*Want to show $x \in (A \cap C) - B$.*]
2. By 1 and definition of intersection, $x \in A - B$ and $x \in C - B$.
3. By 2 and definition of difference, $x \in A$ and $x \in C$ and $x \notin B$.
4. By 3 and definition of intersection, $x \in A \cap C$.
5. By 3 and 4 and definition of difference, $x \in (A \cap C) - B$.
6. By 1 and 5 and definition of subset, $(A - B) \cap (C - B) \subseteq (A \cap C) - B$.

Now the other direction.

7. Assume $x \in (A \cap C) - B$. [*Want to show $x \in (A - B) \cap (C - B)$.*]
8. By 7 and definition of difference, $x \in A \cap C$ and $x \notin B$.
9. By 8 and definition of intersection, $x \in A$ and $x \in C$ and $x \notin B$.
10. By 9 and definition of difference, $x \in A - B$ and $x \in C - B$.
11. By 10 and definition of intersection, $x \in (A - B) \cap (C - B)$.
12. By 7 and 11 and definition of subset, $(A \cap C) - B \subseteq (A - B) \cap (C - B)$.

Conclusion:

13. By 6, 12 and definition of set equality, $(A - B) \cap (C - B) = (A \cap C) - B$. □

2.14 Exercise 14

For all sets A and B , $A \cup (A \cap B) = A$.

Proof. Suppose A and B are any sets. We will show that $A \cup (A \cap B) \subseteq A$. Suppose x is any element in $A \cup (A \cap B)$. [*We must show that $x \in A$.*] By definition of union, $x \in A$ or $x \in A \cap B$. In the case where $x \in A$, clearly $x \in A$. In the case where $x \in A \cap B$,

$x \in A$ and $x \in B$ (by definition of intersection), and so, in particular, $x \in A$. Hence, in both cases $x \in A$ [as was to be shown]. Thus $A \cup (A \cap B) \subseteq A$ by definition of subset.

To complete the proof that $A \cup (A \cap B) = A$, you must show that $A \subseteq A \cup (B \cap A)$. \square

2.15 Exercise 15

For every set A , $A \cup \emptyset = A$.

Proof. Let A be any set. [We must show that $A \cup \emptyset = A$.]

Proof that $A \cup \emptyset \subseteq A$: Suppose $x \in A \cup \emptyset$. Then $x \in A$ or $x \in \emptyset$ by definition of union. But $x \notin \emptyset$ since \emptyset has no elements. Hence $x \in A$.

Proof that $A \subseteq A \cup \emptyset$: Suppose $x \in A$. Then the statement “ $x \in A$ or $x \in \emptyset$ ” is true. Hence $x \in A \cup \emptyset$ by definition of union. [Alternatively, $A \subseteq A \cup \emptyset$ by the inclusion in union property.]

Since $A \cup \emptyset \subseteq A$ and $A \subseteq A \cup \emptyset$, then $A \cup \emptyset = A$ by definition of set equality. \square

2.16 Exercise 16

For all sets A , B , and C , if $A \subseteq B$ then $A \cap C \subseteq B \cap C$.

Proof. Suppose A , B , and C are any sets such that $A \subseteq B$. Let $x \in A \cap C$. By definition of intersection, $x \in A$ and $x \in C$. Now since $A \subseteq B$ and $x \in A$, then $x \in B$. Hence $x \in B$ and $x \in C$, and so, by definition of intersection, $x \in B \cap C$. [Thus $A \cap C \subseteq B \cap C$ by definition of subset.] \square

2.17 Exercise 17

For all sets A , B , and C , if $A \subseteq B$ then $A \cup C \subseteq B \cup C$.

Proof. Assume A, B, C are any sets.

1. Assume $A \subseteq B$. [Want to show $A \cup C \subseteq B \cup C$.]
2. Assume $x \in A \cup C$. [Want to show $x \in B \cup C$.]
3. By 2 and definition of union, $x \in A$ or $x \in C$.
4. **Case 1** ($x \in A$): Then by 1 and definition of subset, $x \in B$. So by definition of union, $x \in B \cup C$.
5. **Case 2** ($x \in C$): Then by definition of union, $x \in B \cup C$.
6. By 4 and 5, ($x \in B \cup C$).
7. By 2 and 6 and definition of subset, ($A \cup C \subseteq B \cup C$, [as was to be shown.] \square)

2.18 Exercise 18

For all sets A and B , if $A \subseteq B$ then $B^c \subseteq A^c$.

Proof. Assume A, B are any sets.

1. Assume $A \subseteq B$. [*Want to show $B^c \subseteq A^c$.*]
2. Assume $x \in B^c$. [*Want to show $x \in A^c$.*]
3. By 2 and definition of complement, $x \notin B$.
4. Argue by contradiction and assume $x \in A$. Then by 1 and definition of subset, $x \in B$, which contradicts 3. So $x \notin A$.
5. By 4 and definition of complement, $x \in A^c$.
6. By 2 and 5 and definition of subset, $B^c \subseteq A^c$. [*as was to be shown.*] □

2.19 Exercise 19

For all sets A , B , and C , if $A \subseteq B$ and $A \subseteq C$ then $A \subseteq B \cap C$.

Proof. Assume A, B, C are any sets.

1. Assume $A \subseteq B$ and $A \subseteq C$. [*Want to show $A \subseteq B \cap C$.*]
2. Assume $x \in A$. [*Want to show $x \in B \cap C$.*]
3. By 1 and 2 and definition of subset, $x \in B$ and $x \in C$.
4. By 3 and definition of intersection, $x \in B \cap C$.
5. By 2 and 4 and definition of subset, $A \subseteq B \cap C$, [*as was to be shown.*] □

2.20 Exercise 20

For all sets A , B , and C , if $A \subseteq C$ and $B \subseteq C$ then $A \cup B \subseteq C$.

Proof. Assume A, B, C are any sets.

1. Assume $A \subseteq C$ and $B \subseteq C$. [*Want to show $A \cup B \subseteq C$.*]
2. Assume $x \in A \cup B$. [*Want to show $x \in C$.*]
3. By 2 and definition of union, $x \in A$ or $x \in B$.
4. **Case 1** ($x \in A$): Then by 1 and 3 and definition of subset, $x \in C$.
5. **Case 2** ($x \in B$): Then by 1 and 3 and definition of subset, $x \in C$.
6. By 4 and 5, $x \in C$.
7. By 2 and 6 and definition of subset, $A \cup B \subseteq C$, [*as was to be shown.*] □

2.21 Exercise 21

For all sets A , B , and C , $A \times (B \cup C) = (A \times B) \cup (A \times C)$.

Proof. Suppose A , B , and C are arbitrarily chosen sets.

$A \times (B \cup C) \subseteq (A \times B) \cup (A \times C)$:

Suppose $(x, y) \in A \times (B \cup C)$. [We must show that $(x, y) \in (A \times B) \cup (A \times C)$.] Then $x \in A$ and $y \in B \cup C$. By definition of union, this means that $y \in B$ or $y \in C$.

Case 1 ($y \in B$): Then, since $x \in A$, $(x, y) \in A \times B$ by definition of Cartesian product. Hence $(x, y) \in (A \times B) \cup (A \times C)$ by definition of union.

Case 2 ($y \in C$): Then, since $x \in A$, $(x, y) \in A \times C$ by definition of Cartesian product. Hence $(x, y) \in (A \times B) \cup (A \times C)$ by definition of union.

Hence, in either case, $(x, y) \in (A \times B) \cup (A \times C)$ [as was to be shown]. Thus $A \times (B \cup C) \subseteq (A \times B) \cup (A \times C)$ by definition of subset.

$(A \times B) \cup (A \times C) \subseteq A \times (B \cup C)$:

Suppose $(x, y) \in (A \times B) \cup (A \times C)$. Then $(x, y) \in A \times B$ or $(x, y) \in A \times C$.

Case 1 ($(x, y) \in A \times B$): In this case, $x \in A$ and $y \in B$. Now since $y \in B$ then $y \in B \cup C$ by definition of union. Hence $x \in A$ and $y \in B \cup C$, and so, by definition of Cartesian product, $(x, y) \in A \times (B \cup C)$.

Case 2 ($(x, y) \in A \times C$): In this case $x \in A$ and $y \in C$.

Now since $y \in C$, then $y \in B \cup C$ by definition of union. Hence $x \in A$ and $y \in B \cup C$, and so, by definition of Cartesian product, $(x, y) \in A \times (B \cup C)$.

Thus, in either case, $(x, y) \in A \times (B \cup C)$. [Hence, by definition of subset, $(A \times B) \cup (A \times C) \subseteq A \times (B \cup C)$.]

[Since both subset relations have been proved, we can conclude that $A \times (B \cup C) = (A \times B) \cup (A \times C)$ by definition of set equality.] \square

2.22 Exercise 22

For all sets A , B , and C , $A \times (B \cap C) = (A \times B) \cap (A \times C)$.

Proof. Assume A, B, C are any sets.

1. Assume $(x, y) \in A \times (B \cap C)$. [Want to show $(x, y) \in (A \times B) \cap (A \times C)$.]
2. By 1 and definition of Cartesian product, $x \in A$ and $y \in B \cap C$.
3. By 2 and definition of intersection, $y \in B$ and $y \in C$.
4. By 2 and 3 and definition of Cartesian product, $(x, y) \in A \times B$ and $(x, y) \in A \times C$.
5. By 4 and definition of intersection, $(x, y) \in (A \times B) \cap (A \times C)$.
6. By 1 and 5 and definition of subset, $A \times (B \cap C) \subseteq (A \times B) \cap (A \times C)$.

Now the reverse part:

7. Assume $(x, y) \in (A \times B) \cap (A \times C)$. [*Want to show $(x, y) \in A \times (B \cap C)$.*]
8. By 7 and definition of intersection, $(x, y) \in A \times B$ and $(x, y) \in A \times C$.
9. By 8 and definition of Cartesian product, $x \in A$ and $y \in B$ and $y \in C$.
10. By 9 and definition of intersection, $y \in B \cap C$.
11. By 9 and 10 and definition of Cartesian product, $(x, y) \in A \times (B \cap C)$.
12. By 6 and 11 and definition of subset, $(A \times B) \cap (A \times C) \subseteq A \times (B \cap C)$.

Conclusion: By 7 and 14 and definition of set equality, $A \times (B \cap C) = (A \times B) \cap (A \times C)$. \square

2.23 Exercise 23

Find the mistake in the following “proof” that for all sets A , B , and C , if $A \subseteq B$ and $B \subseteq C$ then $A \subseteq C$.

“Proof: Suppose A , B , and C are any sets such that $A \subseteq B$ and $B \subseteq C$. Since $A \subseteq B$, there is an element x such that $x \in A$ and $x \in B$, and since $B \subseteq C$, there is an element x such that $x \in B$ and $x \in C$. Hence there is an element x such that $x \in A$ and $x \in C$ and so $A \subseteq C$.”

Proof. There is more than one error in this “proof.” The most serious is the misuse of the definition of subset. To say that A is a subset of B means that for every x , if $x \in A$ then $x \in B$. It does not mean that there exists an element of A that is also an element of B . The second error in the proof occurs in the last sentence. Even if there is an element in A that is in B and an element in B that is in C , it does not follow that there is an element in A that is in C . For instance, suppose $A = \{1, 2\}$, $B = \{2, 3\}$, and $C = \{3, 4\}$. Then there is an element in A that is in B (namely 2) and there is an element in B that is in C (namely 3), but there is no element in A that is in C . \square

2.24 Exercise 24

Find the mistake in the following “proof.”

“Theorem:” For all sets A and B , $A^c \cup B^c \subseteq (A \cup B)^c$.

“Proof: Suppose A and B are any sets, and $x \in A^c \cup B^c$. Then $x \in A^c$ or $x \in B^c$ by definition of union. It follows that $x \notin A$ or $x \notin B$ by definition of complement, and so $x \notin A \cup B$ by definition of union. Thus $x \in (A \cup B)^c$ by definition of complement, and hence $A^c \cup B^c \subseteq (A \cup B)^c$.”

Hint: The words “and so $x \notin A \cup B$ ” do not necessarily follow from “ $x \notin A$ or $x \notin B$.” Try to think of an example of sets A and B and an element x for which “ $x \notin A$ or $x \notin B$ ” is true and “ $x \notin A \cup B$ ” is false.

Proof. (following the Hint)

Let $A = \{x\}$ and $B = \{y\}$. Then $x \notin B$. So “ $x \notin A$ or $x \notin B$ ” is true. Now $A \cup B = \{x, y\}$ so $x \in A \cup B$, therefore “ $x \notin A \cup B$ ” is false. \square

2.25 Exercise 25

Find the mistake in the following “proof” that for all sets A and B , $(A - B) \cup (A \cap B) \subseteq A$.


“Proof: Suppose A and B are any sets, and suppose $x \in (A - B) \cup (A \cap B)$. If $x \in A$ then $x \in A - B$, and so, by definition of difference, $x \in A$ and $x \notin B$. In particular, $x \in A$, and, therefore, $(A - B) \cup (A \cap B) \subseteq A$ by definition of subset.”

Proof. This proof has circular reasoning: it assumes what is to be proved by saying “if $x \in A$ ”.

Another mistake is that “if $x \in A$ then $x \in A - B$ ” is an implication that does not follow. For example, let $A = B = \{x\}$ so $x \in A$ but $A - B = \emptyset$ thus $x \notin A - B$. \square


2.26 Exercise 26

Consider the Venn diagram below.



2.26.1 (a)

Illustrate one of the distributive laws by shading in the region corresponding to $A \cup (B \cap C)$ on one copy of the diagram and $(A \cup B) \cap (A \cup C)$ on another.




The shaded region is $A \cup (B \cap C)$. The most darkly shaded region is $(A \cup B) \cap (A \cup C)$.

Proof.

\square

2.26.2 (b)

Illustrate the other distributive law by shading in the region corresponding to $A \cap (B \cup C)$ on one copy of the diagram and $(A \cap B) \cup (A \cap C)$ on another.




Proof.

□

2.26.3 (c)

Illustrate one of De Morgan's laws by shading in the region corresponding to $(A \cup B)^c$ on one copy of the diagram and $A^c \cap B^c$ on the other. (Leave the set C out of your diagrams.)



Proof.


□

2.26.4 (d)

Illustrate the other De Morgan's law by shading in the region corresponding to $(A \cap B)^c$ on one copy of the diagram and $A^c \cup B^c$ on the other. (Leave the set C out of your diagrams.)

Proof.

□



2.27 Exercise 27

Fill in the blanks in the following proof that for all sets A and B , $(A - B) \cap (B - A) = \emptyset$.

Proof: Let A and B be any sets and suppose $(A - B) \cap (B - A) \neq \emptyset$. That is, suppose there is an element x in (a) _____. By definition of (b) _____, $x \in A - B$ and $x \in$ (c) _____. Then by definition of set difference, $x \in A$ and $x \notin B$ and $x \in$ (d) _____ and $x \notin$ (e) _____. In particular $x \in A$ and $x \notin$ (f) _____, which is a contradiction. Hence [the supposition that $(A - B) \cap (B - A) \neq \emptyset$ is false, and so] (g) _____.

Proof. (a) $(A - B) \cap (B - A)$ (b) intersection (c) $B - A$ (d) B (e) A (f) A (g) $(A - B) \cap (B - A) = \emptyset$ \square

Use the element method for proving a set equals the empty set to prove each statement in 28 – 38. Assume that all sets are subsets of a universal set U .

2.28 Exercise 28

For all sets A and B , $(A \cap B) \cap (A \cap B^c) = \emptyset$. (This property is used in Section 9.9.)

Proof. By contradiction: Suppose not. That is, suppose there exist sets A and B such that $(A \cap B) \cap (A \cap B^c) \neq \emptyset$. Then there is an element x in $(A \cap B) \cap (A \cap B^c)$. By definition of intersection, $x \in (A \cap B)$ and $x \in (A \cap B^c)$. Applying the definition of intersection again, we have that since $x \in (A \cap B)$, $x \in A$ and $x \in B$, and since $x \in (A \cap B^c)$, $x \in A$ and $x \notin B$. Thus, in particular, $x \in B$ and $x \notin B$, which is a contradiction. It follows that the supposition is false, and so $(A \cap B) \cap (A \cap B^c) = \emptyset$. \square

2.29 Exercise 29

For all sets A , B , and C , $(A - C) \cap (B - C) \cap (A - B) = \emptyset$.

Proof. By contradiction: Suppose not. That is, suppose there exist sets A , B and C such that $(A - C) \cap (B - C) \cap (A - B) \neq \emptyset$. Then there is an element x in $(A - C) \cap (B - C) \cap (A - B)$. By definition of intersection, $x \in (A - C)$, $x \in (B - C)$ and $x \in (A - B)$. Applying the definition of difference, we have that since $x \in (B - C)$, $x \in B$ and $x \notin C$, and since $x \in (A - B)$, $x \in A$ and $x \notin B$. Thus, in particular, $x \in B$

and $x \notin B$, which is a contradiction. It follows that the supposition is false, and so $(A - C) \cap (B - C) \cap (A - B) = \emptyset$. \square

2.30 Exercise 30

For every subset A of a universal set U , $A \cap A^c = \emptyset$.

Proof. Let A be a subset of a universal set U . Suppose $A \cap A^c \neq \emptyset$, that is, suppose there is an element x such that $x \in A \cap A^c$. By definition of intersection, $x \in A$ and $x \in A^c$, and so by definition of complement, $x \in A$ and $x \notin A$. This is a contradiction. [Hence the supposition is false, and we conclude that $A \cap A^c = \emptyset$.] \square

2.31 Exercise 31

If U denotes a universal set, then $U^c = \emptyset$.

Proof. Suppose $U^c \neq \emptyset$, that is, suppose there is an element x such that $x \in U^c$. By definition of complement, $x \notin U$. But by definition of universal set, $x \in U$. This is a contradiction. [Hence the supposition is false, and we conclude that $U^c = \emptyset$.] \square

2.32 Exercise 32

For every set A , $A \times \emptyset = \emptyset$.

Proof. Let A be a set. Suppose $A \times \emptyset \neq \emptyset$. Then there would be an element (x, y) in $A \times \emptyset$. By definition of Cartesian product, $x \in A$ and $y \in \emptyset$. But there are no elements y such that $y \in \emptyset$. Hence there are no elements (x, y) in $A \times \emptyset$, which is a contradiction. [Thus the supposition is false, and so $A \times \emptyset = \emptyset$.] \square

2.33 Exercise 33

For all sets A and B , if $A \subseteq B$ then $A \cap B^c = \emptyset$.

Proof. Let A and B be sets such that $A \subseteq B$. [We must show that $A \cap B^c = \emptyset$.] Suppose $A \cap B^c \neq \emptyset$; that is, suppose there were an element x such that $x \in A \cap B^c$. Then $x \in A$ and $x \in B^c$ by definition of intersection. So $x \in A$ and $x \notin B$ by definition of complement. But $A \subseteq B$ by hypothesis, and, since $x \in A$, then $x \in B$ by definition of subset. Thus $x \notin B$ and also $x \in B$, which is a contradiction. Hence the supposition that $A \cap B^c \neq \emptyset$ is false, and so $A \cap B^c = \emptyset$. \square

2.34 Exercise 34

For all sets A and B , if $B \subseteq A^c$ then $A \cap B = \emptyset$.

Proof. Let A and B be sets such that $B \subseteq A^c$. [We must show that $A \cap B = \emptyset$.] Suppose $A \cap B \neq \emptyset$; that is, suppose there were an element x such that $x \in A \cap B$. Then $x \in A$ and $x \in B$ by definition of intersection. But $B \subseteq A^c$ by hypothesis, and, since $x \in B$,

then $x \in A^c$ by definition of subset. Thus $x \notin A$ by definition of complement, and also $x \in A$, which is a contradiction. Hence the supposition that $A \cap B \neq \emptyset$ is false, and so $A \cap B = \emptyset$. \square

2.35 Exercise 35

For all sets A , B , and C , if $A \subseteq B$ and $B \cap C = \emptyset$ then $A \cap C = \emptyset$.

Proof. Let A, B and C be sets such that $A \subseteq B$ and $B \cap C = \emptyset$. [We must show that $A \cap C = \emptyset$.] Suppose $A \cap C \neq \emptyset$; that is, suppose there were an element x such that $x \in A \cap C$. Then $x \in A$ and $x \in C$ by definition of intersection. But $A \subseteq B$ by hypothesis, and, since $x \in A$, then $x \in B$ by definition of subset. Thus $x \in B$ and also $x \in C$, so by definition of intersection $x \in B \cap C$, which is a contradiction to the fact that $B \cap C = \emptyset$. Hence the supposition that $A \cap C \neq \emptyset$ is false, and so $A \cap C = \emptyset$. \square

2.36 Exercise 36

For all sets A , B , and C , if $C \subseteq B - A$ then $A \cap C = \emptyset$.

Proof. Let A , B , and C be any sets such that $C \subseteq B - A$. Suppose $A \cap C \neq \emptyset$. Then there is an element x such that $x \in A \cap C$. By definition of intersection, $x \in A$ and $x \in C$. Now since $x \in C$ and $C \subseteq B - A$, then $x \in B$ and $x \notin A$. So $x \in A$ and $x \notin A$, which is a contradiction. Hence the supposition is false, and thus $A \cap C = \emptyset$. \square

2.37 Exercise 37

For all sets A , B , and C , if $B \cap C \subseteq A$, then $(C - A) \cap (B - A) = \emptyset$.

Proof. Let A , B , and C be any sets such that $B \cap C \subseteq A$. Suppose $(C - A) \cap (B - A) \neq \emptyset$. Then there is an element x such that $x \in (C - A) \cap (B - A)$. By definition of intersection, $x \in C - A$ and $x \in B - A$. Applying the definition of difference, $x \in C$ and $x \notin A$, and $x \in B$ and $x \notin A$. By definition of intersection, $x \in B \cap C$. By definition of subset, $x \in A$. So $x \in A$ and $x \notin A$, which is a contradiction. Hence the supposition is false, and thus $(C - A) \cap (B - A) = \emptyset$. \square

2.38 Exercise 38

For all sets A , B , C , and D , if $A \cap C = \emptyset$ then $(A \times B) \cap (C \times D) = \emptyset$.

Proof. Let A, B, C and D be any sets such that $A \cap C = \emptyset$. Suppose $(A \times B) \cap (C \times D) \neq \emptyset$. Then there is an element (x, y) such that $(x, y) \in (A \times B) \cap (C \times D)$. By definition of intersection, $(x, y) \in A \times B$ and $(x, y) \in C \times D$. By definition of Cartesian product, $x \in A$ and $x \in C$. By definition of intersection, $x \in A \cap C$, which is a contradiction to the fact that $A \cap C = \emptyset$. Hence the supposition is false, and thus $(A \times B) \cap (C \times D) = \emptyset$. \square

Prove each statement in 39 – 44.

2.39 Exercise 39

For all sets A and B ,

2.39.1 (a)

$$(A - B) \cup (B - A) \cup (A \cap B) = A \cup B$$

Proof. Start of proof that $A \cup B \subseteq (A - B) \cup (B - A) \cup (A \cap B)$:

Given any element x in $A \cup B$, by definition of union x is in at least one of A and B . Thus x satisfies exactly one of the following three conditions:

- (1) $x \in A$ and $x \notin B$ (x is in A only)
- (2) $x \in B$ and $x \notin A$ (x is in B only)
- (3) $x \in A$ and $x \in B$ (x is in both A and B)

□

2.39.2 (b)

The sets $(A - B)$, $(B - A)$, $(A \cap B)$ are mutually disjoint.

Proof. To show that $(A - B)$, $(B - A)$, and $(A \cap B)$ are mutually disjoint, we must show that the intersection of any two of them is the empty set. Now, by definition of set difference and set intersection,

saying that $x \in A - B$ means that (1) $x \in A$ and $x \notin B$,

saying that $x \in B - A$ means that (2) $x \in B$ and $x \notin A$, and

saying that $x \in A \cap B$ means that (3) $x \in A$ and $x \in B$.

Conditions (1)–(3) are mutually exclusive: no two of them can be satisfied at the same time. Thus no element can be in the intersection of any two of the sets, and, therefore, the intersection of any two of the sets is the empty set.

Hence, $(A - B)$, $(B - A)$, and $(A \cap B)$ are mutually disjoint.

□

2.40 Exercise 40

For every positive integer n , if A and B_1, B_2, B_3, \dots are any sets, then

$$A \cap \left(\bigcup_{i=1}^n B_i \right) = \bigcup_{i=1}^n (A \cap B_i).$$

Proof. Suppose that n is any positive integer and that A and B_1, B_2, B_3, \dots are any sets.

Proof that $A \cap (\bigcup_{i=1}^n B_i) \subseteq \bigcup_{i=1}^n (A \cap B_i)$:

Suppose x is any element in $A \cap \left(\bigcup_{i=1}^n B_i \right)$. [We must show that $x \in \bigcup_{i=1}^n (A \cap B_i)$.]

By definition of intersection, $x \in A$ and $x \in \bigcup_{i=1}^n B_i$. Since $x \in \bigcup_{i=1}^n B_i$, the definition of general union implies that $x \in B_i$ for some $i = 1, 2, \dots, n$, and so, since $x \in A$, the definition of intersection implies that $x \in A \cap B_i$. Thus, by definition of general union, $x \in \bigcup_{i=1}^n (A \cap B_i)$ [as was to be shown].

Proof that $\bigcup_{i=1}^n (A \cap B_i) \subseteq A \cap \left(\bigcup_{i=1}^n B_i\right)$:

Suppose x is any element in $\bigcup_{i=1}^n (A \cap B_i)$. [We must show that $x \in A \cap \left(\bigcup_{i=1}^n B_i\right)$.]

By definition of general union, $x \in A \cap B_i$ for some $i = 1, 2, \dots, n$. Thus, by definition of intersection, $x \in A$ and $x \in B_i$. Since $x \in B_i$ for some $i = 1, 2, \dots, n$, then by definition of general union, $x \in \bigcup_{i=1}^n B_i$. Thus we have that $x \in A$ and $x \in \bigcup_{i=1}^n B_i$, and so,

by definition of intersection, $x \in A \cap \bigcup_{i=1}^n B_i$, [as was to be shown].

Conclusion: Since both subset relations have been proved, it follows by definition of set equality that

$$A \cap \left(\bigcup_{i=1}^n B_i\right) = \bigcup_{i=1}^n (A \cap B_i).$$

□

2.41 Exercise 41

For every positive integer n , if A_1, A_2, A_3, \dots and B are any sets, then

$$\bigcup_{i=1}^n (A_i - B) = \left(\bigcup_{i=1}^n A_i\right) - B.$$

Proof. Assume n is a positive integer, A_1, A_2, A_3, \dots and B are any sets.

1. Assume $x \in \bigcup_{i=1}^n (A_i - B)$. [Want to show $x \in \left(\bigcup_{i=1}^n A_i\right) - B$].
2. By 1 and definition of \bigcup , there exists a j in $\{1, \dots, n\}$ such that $x \in A_j - B$.
3. By 2 and definition of difference, $x \in A_j$ and $x \notin B$.
4. By 3 and definition of \bigcup , $x \in \bigcup_{i=1}^n A_i$.
5. By 3 and 4 and definition of difference, $x \in \left(\bigcup_{i=1}^n A_i\right) - B$.

6. By 1 and 5 and definition of subset, $\bigcup_{i=1}^n (A_i - B) \subseteq \left(\bigcup_{i=1}^n A_i \right) - B$.

Now the reverse part:

7. Assume $x \in \left(\bigcup_{i=1}^n A_i \right) - B$. [Want to show $x \in \bigcup_{i=1}^n (A_i - B)$].

8. By 7 and definition of difference, $x \in \bigcup_{i=1}^n A_i$ and $x \notin B$.

9. By 8 and definition of \bigcup , there exists j in $\{1, \dots, n\}$ such that $x \in A_j$.

10. By 8 and 9 and definition of difference, $x \in A_j - B$.

11. By 10 and definition of \bigcup , $x \in \bigcup_{i=1}^n (A_i - B)$.

12. By 11 and 7 and definition of subset, $\left(\bigcup_{i=1}^n A_i \right) - B \subseteq \bigcup_{i=1}^n (A_i - B)$.

13. Conclusion: By 6 and 12 and definition of set equality,

$$\bigcup_{i=1}^n (A_i - B) = \left(\bigcup_{i=1}^n A_i \right) - B.$$

□

2.42 Exercise 42

For every positive integer n , if A_1, A_2, A_3, \dots and B are any sets, then

$$\bigcap_{i=1}^n (A_i - B) = \left(\bigcap_{i=1}^n A_i \right) - B.$$

Proof. Assume n is a positive integer, A_1, A_2, A_3, \dots and B are any sets.

1. Assume $x \in \bigcap_{i=1}^n (A_i - B)$. [Want to show $x \in \left(\bigcap_{i=1}^n A_i \right) - B$].

2. By 1 and definition of \bigcap , $x \in A_i - B$ for all i in $\{1, \dots, n\}$.

3. By 2 and definition of difference, $x \in A_i$ for all i in $\{1, \dots, n\}$, and $x \notin B$.

4. By 3 and definition of \bigcap , $x \in \bigcap_{i=1}^n A_i$.

5. By 3 and 4 and definition of difference, $x \in \left(\bigcap_{i=1}^n A_i\right) - B$.

6. By 1 and 5 and definition of subset, $\bigcap_{i=1}^n (A_i - B) \subseteq \left(\bigcap_{i=1}^n A_i\right) - B$.

Now the reverse part:

7. Assume $x \in \left(\bigcap_{i=1}^n A_i\right) - B$. [Want to show $x \in \bigcap_{i=1}^n (A_i - B)$].

8. By 7 and definition of difference, $x \in \bigcap_{i=1}^n A_i$ and $x \notin B$.

9. By 8 and definition of \bigcap , $x \in A_i$ for all i in $\{1, \dots, n\}$.

10. By 8 and 9 and definition of difference, $x \in A_i - B$ for all i in $\{1, \dots, n\}$.

11. By 10 and definition of \bigcap , $x \in \bigcap_{i=1}^n (A_i - B)$.

12. By 11 and 7 and definition of subset, $\left(\bigcap_{i=1}^n A_i\right) - B \subseteq \bigcap_{i=1}^n (A_i - B)$.

13. Conclusion: By 6 and 12 and definition of set equality,

$$\bigcap_{i=1}^n (A_i - B) = \left(\bigcap_{i=1}^n A_i\right) - B.$$

□

2.43 Exercise 43

For every positive integer n , if A and B_1, B_2, B_3, \dots are any sets, then

$$\bigcup_{i=1}^n (A \times B_i) = A \times \left(\bigcup_{i=1}^n B_i\right).$$

Proof. Assume n is a positive integer, B_1, B_2, B_3, \dots and A are any sets.

1. Assume $(x, y) \in \bigcup_{i=1}^n (A \times B_i)$. [Want to show $(x, y) \in A \times \left(\bigcup_{i=1}^n B_i\right)$].

2. By 1 and definition of \bigcup , there exists a j in $\{1, \dots, n\}$ such that $(x, y) \in A \times B_j$.

3. By 2 and definition of Cartesian product, $x \in A$ and $y \in B_j$.

4. By 3 and definition of \bigcup , $y \in \bigcup_{i=1}^n B_i$.

5. By 3 and 4 and definition of Cartesian product, $(x, y) \in A \times \left(\bigcup_{i=1}^n B_i\right)$.

6. By 1 and 5 and definition of subset, $\bigcup_{i=1}^n (A \times B_i) \subseteq A \times \left(\bigcup_{i=1}^n B_i\right)$.

Now the reverse part:

7. Assume $(x, y) \in A \times \left(\bigcup_{i=1}^n B_i\right)$. [*Want to show $(x, y) \in \bigcup_{i=1}^n (A \times B_i)$*].

8. By 7 and definition of Cartesian product, $x \in A$ and $y \in \bigcup_{i=1}^n B_i$.

9. By 8 and definition of \bigcup , there exists j in $\{1, \dots, n\}$ such that $y \in B_j$.

10. By 8 and 9 and definition of Cartesian product, $(x, y) \in A \times B_j$.

11. By 10 and definition of \bigcup , $(x, y) \in \bigcup_{i=1}^n (A \times B_i)$.

12. By 11 and 7 and definition of subset, $A \times \left(\bigcup_{i=1}^n B_i\right) \subseteq \bigcup_{i=1}^n (A \times B_i)$.

13. Conclusion: By 6 and 12 and definition of set equality,

$$A \times \left(\bigcup_{i=1}^n B_i\right) = \bigcup_{i=1}^n (A \times B_i).$$

□

2.44 Exercise 44

For every positive integer n , if A and B_1, B_2, B_3, \dots are any sets, then

$$\bigcap_{i=1}^n (A \times B_i) = A \times \left(\bigcap_{i=1}^n B_i\right).$$

Proof. Assume n is a positive integer, B_1, B_2, B_3, \dots and A are any sets.

1. Assume $(x, y) \in \bigcap_{i=1}^n (A \times B_i)$. [*Want to show $(x, y) \in A \times \left(\bigcap_{i=1}^n B_i\right)$*].

2. By 1 and definition of \bigcap , $(x, y) \in A \times B_i$ for all i in $\{1, \dots, n\}$.

3. By 2 and definition of Cartesian product, $x \in A$ and $y \in B_i$ for all i in $\{1, \dots, n\}$.

4. By 3 and definition of \bigcap , $y \in \bigcap_{i=1}^n B_i$.

5. By 3 and 4 and definition of Cartesian product, $(x, y) \in A \times \left(\bigcap_{i=1}^n B_i\right)$.

6. By 1 and 5 and definition of subset, $\bigcap_{i=1}^n (A \times B_i) \subseteq A \times \left(\bigcap_{i=1}^n B_i\right)$.

Now the reverse part:

7. Assume $(x, y) \in A \times \left(\bigcap_{i=1}^n B_i\right)$. [Want to show $x \in \bigcap_{i=1}^n (A \times B_i)$].

8. By 7 and definition of Cartesian product, $x \in A$ and $y \in \bigcap_{i=1}^n B_i$.

9. By 8 and definition of \bigcap , $y \in B_i$ for all i in $\{1, \dots, n\}$.

10. By 8 and 9 and definition of Cartesian product, $(x, y) \in A \times B_i$ for all i in $\{1, \dots, n\}$.

11. By 10 and definition of \bigcap , $(x, y) \in \bigcap_{i=1}^n (A \times B_i)$.

12. By 11 and 7 and definition of subset, $A \times \left(\bigcap_{i=1}^n B_i\right) \subseteq \bigcap_{i=1}^n (A \times B_i)$.

13. Conclusion: By 6 and 12 and definition of set equality,

$$\bigcap_{i=1}^n (A \times B_i) = A \times \left(\bigcap_{i=1}^n B_i\right).$$

□

3 Exercise Set 6.3

For each of 1 – 4 find a counterexample to show that the statement is false. Assume all sets are subsets of a universal set U .

3.1 Exercise 1

For all sets A , B , and C , $(A \cup B) \cap C = A \cup (B \cap C)$.

Proof. Counterexample: A , B , and C can be any sets where A has an element that is not in C . For instance, let $A = \{1, 2\}$, $B = \{2\}$, and $C = \{2\}$. Then $(A \cup B) \cap C = (\{1, 2\} \cup \{2\}) \cap \{2\} = \{2\}$.

$\{2\}) \cap \{2\} = \{1, 2\} \cap \{2\} = \{2\}$, and $A \cup (B \cap C) = \{1, 2\} \cup (\{2\} \cap \{2\}) = \{1, 2\} \cup \{2\} = \{1, 2\}$. Thus $1 \in A \cup (B \cap C)$ but $1 \notin (A \cup B) \cap C$, and hence $(A \cup B) \cap C \neq A \cup (B \cap C)$ by definition of subset. \square

3.2 Exercise 2

For all sets A, B , $(A \cup B)^c = A^c \cup B^c$.

Proof. Let $U = \{1, 2\}$, $A = \{1\}$, $B = \{2\}$. Then $A \cup B = \{1, 2\}$ so $(A \cup B)^c = \emptyset$. But $A^c = \{2\}$, $B^c = \{1\}$ and so $A^c \cup B^c = \{1, 2\} \neq \emptyset = (A \cup B)^c$. \square

3.3 Exercise 3

For all sets A, B , and C , if $A \not\subseteq B$ and $B \not\subseteq C$ then $A \not\subseteq C$.

Proof. Counterexample: A, B , and C can be any sets where $A \subseteq C$ and B contains at least one element that is not in either A or C . For instance, let $A = \{1\}$, $B = \{2\}$, and $C = \{1, 3\}$. Then $A \not\subseteq B$ and $B \not\subseteq C$ but $A \subseteq C$. \square

3.4 Exercise 4

For all sets A, B , and C , if $B \cup C \subseteq A$ then $(A - B) \cap (A - C) = \emptyset$.

Proof. Let $A = \{1, 2, 3\}$, $B = \{2\}$, $C = \{3\}$. Then $B \cup C = \{2, 3\} \subseteq A$. And $A - B = \{1, 3\}$ and $A - C = \{1, 2\}$, but $(A - B) \cap (A - C) = \{1, 3\} \cap \{1, 2\} = \{1\} \neq \emptyset$. \square

For each of 5–21 prove each statement that is true and find a counterexample for each statement that is false. Assume all sets are subsets of a universal set U .

3.5 Exercise 5

For all sets A, B , and C , $A - (B - C) = (A - B) - C$.

Proof. False. Counterexample: A, B , and C can be any sets where all three sets have an element in common or where A and C have a common element that is not in B . For instance, let $A = \{1, 2, 3\}$, $B = \{2, 3\}$, and $C = \{3\}$. Then $B - C = \{2\}$, and so $A - (B - C) = \{1, 2, 3\} - \{2\} = \{1, 3\}$. On the other hand, $A - B = \{1, 2, 3\} - \{2, 3\} = \{1\}$, and so $(A - B) - C = \{1\} - \{3\} = \{1\}$. Since $\{1, 3\} \neq \{1\}$, $A - (B - C) \neq (A - B) - C$. \square

3.6 Exercise 6

For all sets A, B , $A \cap (A \cup B) = A$.

Proof. True. Let A and B be any sets.

Proof that $A \cap (A \cup B) \subseteq A$: Suppose $x \in A \cap (A \cup B)$. By definition of intersection, $x \in A$ and $x \in A \cup B$. In particular, $x \in A$. Thus, by definition of subset, $A \cap (A \cup B) \subseteq A$.

Proof that $A \subseteq A \cap (A \cup B)$: Suppose $x \in A$. Then by definition of union, $x \in A \cup B$. Hence $x \in A$ and $x \in A \cup B$, and so, by definition of intersection $x \in A \cap (A \cup B)$. Thus, by definition of subset, $A \subseteq A \cap (A \cup B)$.

Because both $A \cap (A \cup B) \subseteq A$ and $A \subseteq A \cap (A \cup B)$ have been proved, we conclude that $A \cap (A \cup B) = A$. \square

3.7 Exercise 7

For all sets A , B , and C , $(A - B) \cap (C - B) = A - (B \cup C)$.

Proof. Counterexample: Let $A = \{1\}$, $B = \{2\}$, $C = \{3\}$. Then $A - B = \{1\}$, $C - B = \{3\}$, $B \cup C = \{2, 3\}$. So $(A - B) \cap (C - B) = \{1\} \cap \{3\} = \emptyset \neq \{1\} = A - (B \cup C)$. \square

3.8 Exercise 8

For all sets A , B , and C , if $A^c \subseteq B$ then $A \cup B = U$.

Proof. Assume $x \in A \cup B$. By definition of universal set, $x \in U$. So by definition of subset, $A \cup B \subseteq U$.

Assume $x \in U$. There are two cases: $x \in A$ or $x \notin A$.

In the first case, $x \in A \cup B$ by definition of union.

In the second case, $x \in A^c$ by definition of complement, and so $x \in B$ because $A^c \subseteq B$. Since $x \in B$, then $x \in A \cup B$ by definition of union.

In both cases $x \in A \cup B$. So by definition of subset, $U \subseteq A \cup B$.

By definition of set equality, $A \cup B = U$. \square

3.9 Exercise 9

For all sets A , B , and C , if $A \subseteq C$ and $B \subseteq C$ then $A \cup B \subseteq C$.

Proof. True. Suppose A , B , and C are any sets such that $A \subseteq C$ and $B \subseteq C$. Let $x \in A \cup B$. By definition of union, $x \in A$ or $x \in B$. But if $x \in A$ then $x \in C$ (because $A \subseteq C$), and if $x \in B$ then $x \in C$ (because $B \subseteq C$). Hence, in either case, $x \in C$. [So, by definition of subset, $A \cup B \subseteq C$.] \square

3.10 Exercise 10

For all sets A , B , if $A \subseteq B$ then $A \cap B^c = \emptyset$.

Proof. Argue by contradiction and assume $A \cap B^c \neq \emptyset$. There exists x such that $x \in A \cap B^c$. By definition of intersection, $x \in A$ and $x \in B^c$. Since $A \subseteq B$ and $x \in A$, $x \in B$. Since $x \in B^c$, by definition of complement $x \notin B$. So both $x \in B$ and $x \notin B$, contradiction. Therefore $A \cap B^c = \emptyset$. \square

3.11 Exercise 11

For all sets A , B , and C , if $A \subseteq B$ then $A \cap (B \cap C)^c = \emptyset$.

Proof. The statement is false. Let $U = \{1, 2, 3, 4\}$, $A = \{1, 2\}$, $B = \{1, 2, 3\}$, $C = \{2\}$.

Notice that $A = \{1, 2\} \subseteq \{1, 2, 3\} = B$. And $B \cap C = \{1, 2, 3\} \cap \{2\} = \{2\}$, so $(B \cap C)^c = \{1, 3, 4\}$ but $A \cap (B \cap C)^c = \{1, 2\} \cap \{1, 3, 4\} = \{1\} \neq \emptyset$. \square

3.12 Exercise 12

For all sets A , B , and C , $A \cap (B - C) = (A \cap B) - (A \cap C)$.

Proof. 1. Assume $x \in A \cap (B - C)$.

2. By 1 and definition of intersection $x \in A$ and $x \in (B - C)$.

3. By 2 and definition of difference, $x \in B$ and $x \notin C$.

4. By 2 and 3 and definition of intersection, $x \in A \cap B$.

5. If $x \in A \cap C$ then by definition of intersection $x \in C$, contradiction to 3. Thus $x \notin A \cap C$.

6. By 4 and 5 and definition of difference, $x \in (A \cap B) - (A \cap C)$.

7. By 1 and 6 and definition of subset, $A \cap (B - C) \subseteq (A \cap B) - (A \cap C)$.

Now the reverse part:

8. Assume $x \in (A \cap B) - (A \cap C)$.

9. By 8 and definition of difference, $x \in (A \cap B)$ and $x \notin (A \cap C)$.

10. By 9 and definition of intersection, $x \in A$ and $x \in B$.

11. If $x \in C$ then by 10 and definition of intersection, $x \in A \cap C$, contradiction to 9. Thus $x \notin C$.

12. By 10 and 11 and definition of difference, $x \in B - C$.

13. By 10 and 12 and definition of intersection, $x \in A \cap (B - C)$.

14. By 8 and 13 and definition of subset, $(A \cap B) - (A \cap C) \subseteq A \cap (B - C)$.

Conclusion: By 7 and 14 and definition of set equality, $(A \cap B) - (A \cap C) = A \cap (B - C)$. \square

3.13 Exercise 13

For all sets A , B , and C , $A \cup (B - C) = (A \cup B) - (A \cup C)$.

Proof. False. Let $A = \{1\}$, $B = \{2\}$, $C = \{3\}$. Then $A \cup (B - C) = \{1\} \cup (\{2\} - \{3\}) = \{1\} \cup \{2\} = \{1, 2\}$, and $(A \cup B) - (A \cup C) = (\{1\} \cup \{2\}) - (\{1\} \cup \{3\}) = \{1, 2\} - \{1, 3\} = \{2\}$. But $\{1, 2\} \neq \{2\}$. \square

3.14 Exercise 14

For all sets A , B , and C , if $A \cap C = B \cap C$ and $A \cup C = B \cup C$, then $A = B$.

Proof. The statement is true.

1. Suppose $x \in A$. [*We must show that $x \in B$.*]
2. Either $x \in C$ or $x \notin C$.
3. In case $x \in C$, by 1 and definition of intersection $x \in A \cap C = B \cap C$. Then by definition of intersection, $x \in B$.
4. In case $x \notin C$, by 1 and definition of intersection $x \in A \cup C = B \cup C$. Then by definition of union $x \in B$ or $x \in C$, therefore $x \in B$.
5. By 1, 4 and definition of subset, $A \subseteq B$.
6. Suppose $x \in B$. [*We must show that $x \in A$.*]
7. Either $x \in C$ or $x \notin C$.
8. In case $x \in C$, by 6 and definition of intersection $x \in B \cap C = A \cap C$. Then by definition of intersection, $x \in A$.
9. In case $x \notin C$, by 6 and definition of intersection $x \in B \cup C = A \cup C$. Then by definition of union $x \in A$ or $x \in C$, therefore $x \in A$.
10. By 6, 9 and definition of subset, $B \subseteq A$.
11. By 5, 10 and definition of set equality, $A = B$. \square

3.15 Exercise 15

For all sets A , B , and C , $(A - B) \cup C \subseteq A \cup (C - B)$.

Proof. False. Counterexample: Let $A = \{1, 2\}$, $B = \{2, 3\}$, $C = \{3, 4\}$.

Then $(A - B) \cup C = (\{1, 2\} - \{2, 3\}) \cup \{3, 4\} = \{1\} \cup \{3, 4\} = \{1, 3, 4\}$.

And $A \cup (C - B) = \{1, 2\} \cup (\{3, 4\} - \{2, 3\}) = \{1, 2\} \cup \{4\} = \{1, 2, 4\}$.

But $\{1, 3, 4\} \not\subseteq \{1, 2, 4\}$. \square

3.16 Exercise 16

For all sets A, B , if $A \cap B = \emptyset$ then $A \times B = \emptyset$.

Proof. False. Counterexample: Let $A = \{1\}, B = \{2\}$. Then $A \cap B = \emptyset$ but $A \times B = \{(1, 2)\} \neq \emptyset$. \square

3.17 Exercise 17

For all sets A, B , if $A \subseteq B$ then $\mathcal{P}(A) \subseteq \mathcal{P}(B)$.

Proof. Suppose A and B are any sets with $A \subseteq B$. [We must show that $\mathcal{P}(A) \subseteq \mathcal{P}(B)$.] So suppose $X \in \mathcal{P}(A)$. Then $X \subseteq A$ by definition of power set. And because $A \subseteq B$, we also have that $X \subseteq B$ by the transitive property for subsets. Thus, by definition of power set, $X \in \mathcal{P}(B)$. This proves that for all X , if $X \in \mathcal{P}(A)$ then $X \in \mathcal{P}(B)$, and so $\mathcal{P}(A) \subseteq \mathcal{P}(B)$, [as was to be shown]. \square

3.18 Exercise 18

For all sets A, B , $\mathcal{P}(A \cup B) \subseteq \mathcal{P}(A) \cup \mathcal{P}(B)$.

Proof. False. Counterexample: For any sets A and B , the only sets in $\mathcal{P}(A) \cup \mathcal{P}(B)$ are subsets of either A or B , whereas a set in $\mathcal{P}(A \cup B)$ can contain elements from both A and B . Thus, if at least one of A or B contains elements that are not in the other set, $\mathcal{P}(A) \cup \mathcal{P}(B)$ and $\mathcal{P}(A \cup B)$ will not be equal. For instance, let $A = \{1\}$ and $B = \{2\}$. Then $\{1, 2\} \in \mathcal{P}(A \cup B)$ but $\{1, 2\} \notin \mathcal{P}(A) \cup \mathcal{P}(B)$. \square

3.19 Exercise 19

For all sets A, B , $\mathcal{P}(A) \cup \mathcal{P}(B) \subseteq \mathcal{P}(A \cup B)$.

Proof. Assume A, B are any sets.

1. Assume $x \in \mathcal{P}(A) \cup \mathcal{P}(B)$. [Want to show $x \in \mathcal{P}(A \cup B)$].
2. By definition of union $x \in \mathcal{P}(A)$ or $x \in \mathcal{P}(B)$.
3. Case 1: $x \in \mathcal{P}(A)$. By definition of power set, $x \subseteq A$. By definition of union, $A \subseteq A \cup B$. By transitivity of subsets, $x \subseteq A \cup B$. By definition of power set, $x \in \mathcal{P}(A \cup B)$.
4. Case 2: $x \in \mathcal{P}(B)$. By definition of power set, $x \subseteq B$. By definition of union, $B \subseteq A \cup B$. By transitivity of subsets, $x \subseteq A \cup B$. By definition of power set, $x \in \mathcal{P}(A \cup B)$.
5. By 2, 3, 4, $x \in \mathcal{P}(A \cup B)$.
6. By 1, 5 and definition of subset, $\mathcal{P}(A) \cup \mathcal{P}(B) \subseteq \mathcal{P}(A \cup B)$. \square

3.20 Exercise 20

For all sets A, B , $\mathcal{P}(A \cap B) = \mathcal{P}(A) \cap \mathcal{P}(B)$.

Proof. 1. Assume $x \in \mathcal{P}(A \cap B)$. [*Want to show $x \in \mathcal{P}(A) \cap \mathcal{P}(B)$.*]

2. By 1 and definition of power set, $x \subseteq A \cap B$.

3. By 2 and definition of intersection, $x \subseteq A$ and $x \subseteq B$.

4. By 3 and definition of power set, $x \in \mathcal{P}(A)$ and $x \in \mathcal{P}(B)$.

5. By 4 and definition of intersection, $x \in \mathcal{P}(A) \cap \mathcal{P}(B)$.

6. By 1, 5 and definition of subset, $\mathcal{P}(A \cap B) \subseteq \mathcal{P}(A) \cap \mathcal{P}(B)$.

7. Assume $x \in \mathcal{P}(A) \cap \mathcal{P}(B)$. [*Want to show $x \in \mathcal{P}(A \cap B)$.*]

8. By 7 and definition of intersection, $x \in \mathcal{P}(A)$ and $x \in \mathcal{P}(B)$.

9. By 8 and definition of power set, $x \subseteq A$ and $x \subseteq B$.

10. By 9 and definition of intersection, $x \subseteq A \cap B$.

11. By 10 and definition of power set, $x \in \mathcal{P}(A \cap B)$.

12. By 7, 11 and definition of subset, $\mathcal{P}(A) \cap \mathcal{P}(B) \subseteq \mathcal{P}(A \cap B)$.

13. By 6, 12 and definition of set equality, $\mathcal{P}(A \cap B) = \mathcal{P}(A) \cap \mathcal{P}(B)$. □

3.21 Exercise 21

For all sets A, B , $\mathcal{P}(A \times B) = \mathcal{P}(A) \times \mathcal{P}(B)$.

Proof. False. Counterexample: Let $A = \{1\}, B = \{2\}$. Then $A \times B = \{(1, 2)\}$.

So $\mathcal{P}(A) = \{\emptyset, \{1\}\}, \mathcal{P}(B) = \{\emptyset, \{2\}\}$ and $\mathcal{P}(A \times B) = \{\emptyset, \{(1, 2)\}\}$.

But $\mathcal{P}(A) \times \mathcal{P}(B) = \{(\emptyset, \emptyset), (\emptyset, \{2\}), (\{1\}, \emptyset), (\{1\}, \{2\})\} \neq \{\emptyset, \{(1, 2)\}\}$. □

3.22 Exercise 22

Write a negation for each of the following statements. Indicate which is true, the statement or its negation. Justify your answers.

3.22.1 (a)

\forall sets S, \exists a set T such that $S \cap T = \emptyset$.

Proof. True. Let $T = S^c$.

Negation: \exists a set S such that \forall sets $T, S \cap T \neq \emptyset$. □

3.22.2 (b)

\exists a set S such that \forall sets T , $S \cup T = \emptyset$.

Proof. False. If T is nonempty, then no choice of S works.

Negation: \forall sets $S \exists$ a set T such that $S \cup T \neq \emptyset$.

Negation is true: let T be any nonempty set. □

3.23 Exercise 23

Let $S = \{a, b, c\}$, and for each integer $i = 0, 1, 2, 3$, let S_i be the set of all subsets of S that have i elements. List the elements in S_0, S_1, S_2 , and S_3 . Is $\{S_0, S_1, S_2, S_3\}$ a partition of $\mathcal{P}(S)$?

Proof. $S_0 = \{\emptyset\}$, $S_1 = \{\{a\}, \{b\}, \{c\}\}$, $S_2 = \{\{a, b\}, \{a, c\}, \{b, c\}\}$, $S_3 = \{\{a, b, c\}\}$.

Yes it is a partition of $\mathcal{P}(S)$. □

3.24 Exercise 24

Let $A = \{t, u, v, w\}$, and let S_1 be the set of all subsets of A that do not contain w and S_2 the set of all subsets of A that contain w .

3.24.1 (a)

Find S_1 .

Proof. $S_1 = \mathcal{P}(\{t, u, v\}) = \{\emptyset, \{t\}, \{u\}, \{v\}, \{t, u\}, \{t, v\}, \{u, v\}, \{t, u, v\}\}$ □

3.24.2 (b)

Find S_2 .

Proof. $S_2 = \{\{w\}, \{t, w\}, \{u, w\}, \{v, w\}, \{t, u, w\}, \{t, v, w\}, \{u, v, w\}, \{t, u, v, w\}\}$ □

3.24.3 (c)

Are S_1 and S_2 disjoint?

Proof. Yes. □

3.24.4 (d)

Compare the sizes of S_1 and S_2 .

Proof. They are equal (8). □

3.24.5 (e)

How many elements are in $S_1 \cup S_2$?

Proof. 16. □

3.24.6 (f)

What is the relation between $S_1 \cup S_2$ and $\mathcal{P}(A)$?

Proof. $S_1 \cup S_2 = \mathcal{P}(A)$. □

3.25 Exercise 25

Use mathematical induction to prove that for every integer $n \geq 2$, if a set S has n elements, then the number of subsets of S with an even number of elements equals the number of subsets of S with an odd number of elements.

Proof. For the sake of simplicity let's introduce some notation. For a set S define $even(S)$ = the set of elements of S with an even number of elements, and $odd(S)$ = the set of elements of S with an odd number of elements. (Notice we said “elements of S ” not “subsets of S ”.)

Let $P(n)$ be the sentence “If a set S has n elements, then $|even(\mathcal{P}(S))| = |odd(\mathcal{P}(S))|$.” (Here $|a|$ denotes the size of the set a .)

Show that $P(2)$ is true:

Assume a set S has two elements: $S = \{a, b\}$. Then $\mathcal{P}(S) = \{\emptyset, \{a\}, \{b\}, \{a, b\}\}$.

There are two subsets with an even number of elements: \emptyset (which has 0 elements) and $\{a, b\}$ (which has 2 elements).

There are two subsets with an odd number of elements: $\{a\}, \{b\}$ (both of which have 1 element).

Since $2 = 2$, $P(2)$ is true.

Show that for any integer $k \geq 2$ if $P(k)$ is true then $P(k+1)$ is true: Assume “If a set S has k elements, $|even(\mathcal{P}(S))| = |odd(\mathcal{P}(S))|$.”

Assume X is a set with $k+1$ elements. [We want to show $|even(\mathcal{P}(X))| = |odd(\mathcal{P}(X))|$.]

Since $k+1 \geq 1$ we may pick an element $z \in X$. Let $S = X - \{z\}$. Then S has k elements, so by inductive hypothesis $|even(\mathcal{P}(S))| = |odd(\mathcal{P}(S))|$.

Observe that any subset of X either contains z or does not. Let A be the set of subsets of X that contain z , let B be the set of subsets of X that do not contain z .

Notice A and B form a partition of $\mathcal{P}(X)$, so $|even(\mathcal{P}(X))| = |even(A)| + |even(B)|$, and $|odd(\mathcal{P}(X))| = |odd(A)| + |odd(B)|$. Also notice that $B = \mathcal{P}(S)$, so $|even(B)| = |odd(B)|$.

Now consider the following one-to-one correspondence between A and B : for any $b \in B$ there is $a \in A$ given by $a = b \cup \{z\}$. Similarly for any $a \in A$ there is $b \in B$ given by $b = a - \{z\}$. Then A and B have the same number of elements.

Notice that under this correspondence, if $b \in B$ has an even number of elements, then $a = b \cup \{z\}$ has an odd number of elements, and vice versa. Therefore: $|even(A)| = |odd(B)|$ and $|odd(A)| = |even(B)|$. Since $|even(B)| = |odd(B)|$, we get $|even(A)| = |odd(A)|$.

Putting these facts together, we get

$$|even(\mathcal{P}(X))| = |even(A)| + |even(B)| = |odd(A)| + |odd(B)| = |odd(\mathcal{P}(X))|,$$

[as was to be shown.] □

3.26 Exercise 26

The following problem, devised by Ginger Bolton, appeared in the January 1989 issue of the College Mathematics Journal (Vol. 20, No. 1, p. 68): Given a positive integer $n \geq 2$, let S be the set of all nonempty subsets of $\{2, 3, \dots, n\}$. For each $S_i \in S$, let P_i be the product of the elements of S_i . Prove or disprove that

$$\sum_{i=1}^{2^{n-1}-1} P_i = \frac{(n+1)!}{2} - 1$$

Proof. Hint: Use mathematical induction. In the inductive step, you will consider the set of all nonempty subsets of $\{2, \dots, k\}$ and the set of all nonempty subsets of $\{2, \dots, k+1\}$. Any subset of $\{2, \dots, k+1\}$ either contains $k+1$ or does not contain $k+1$. Thus the sum of all products of elements of nonempty subsets of $\{2, \dots, k+1\}$ = the sum of all products of elements of nonempty subsets of $\{2, \dots, k+1\}$ that do not contain $k+1$ + the sum of all products of elements of nonempty subsets of $\{2, \dots, k+1\}$ that contain $k+1$. Now any subset of $\{2, \dots, k+1\}$ that does not contain $k+1$ is a subset of $\{2, \dots, k\}$. And any subset of $\{2, \dots, k+1\}$ that contains $k+1$ is the union of a subset of $\{2, \dots, k\}$ and $\{k+1\}$. □

In 27 and 28 supply a reason for each step in the derivation.

3.27 Exercise 27

For all sets A , B , and C , $(A \cup B) \cap C = (A \cap C) \cup (B \cap C)$.

Proof: Suppose A, B, C are any sets. Then

$$\begin{aligned} (A \cup B) \cap C &= C \cap (A \cup B) && \text{by (a) } \underline{\hspace{2cm}} \\ &= (C \cap A) \cup (C \cap B) && \text{by (b) } \underline{\hspace{2cm}} \\ &= (A \cap C) \cup (B \cap C) && \text{by (c) } \underline{\hspace{2cm}} \end{aligned}$$

Proof. (a) commutative law for \cap (b) distributive law (c) commutative law for \cap □

3.28 Exercise 28

For all sets A , B , and C , $(A \cup B) - (C - A) = A \cup (B - C)$.

Proof: Suppose A, B, C are any sets. Then

$$\begin{aligned}(A \cup B) - (C - A) &= (A \cup B) \cap (C - A)^c && \text{by (a) } \underline{\hspace{1cm}} \\ &= (A \cup B) \cap (C \cap A^c)^c && \text{by (b) } \underline{\hspace{1cm}} \\ &= (A \cup B) \cap (A^c \cap C)^c && \text{by (c) } \underline{\hspace{1cm}} \\ &= (A \cup B) \cap ((A^c)^c \cup C^c) && \text{by (d) } \underline{\hspace{1cm}} \\ &= (A \cup B) \cap (A \cup C^c) && \text{by (e) } \underline{\hspace{1cm}} \\ &= A \cup (B \cap C^c) && \text{by (f) } \underline{\hspace{1cm}} \\ &= A \cup (B - C) && \text{by (g) } \underline{\hspace{1cm}}\end{aligned}$$

Proof. (a) set difference law (b) set difference law (c) commutative law for \cap (d) De Morgan's law (e) double complement law (f) distributive law (g) set difference law \square

3.29 Exercise 29

Some steps are missing from the following proof that for all sets A, B , $(A \cup B^c) - B = (A - B) \cup B^c$. Indicate what they are, and then write the proof correctly.

Proof: Suppose A, B are any sets. Then

$$\begin{aligned}(A \cup B^c) - B &= (A \cup B^c) \cap B^c && \text{by set difference law} \\ &= (B^c \cap A) \cup (B^c \cap B^c) && \text{by distributive law} \\ &= (B^c \cap A) \cup B^c && \text{by idempotent law for } \cup \\ &= (A - B) \cup B^c && \text{by set difference law}\end{aligned}$$

Proof. Before using the distributive law, we must first use the commutative law to get it into the right form:

$$(A \cup B^c) \cap B^c = B^c \cap (A \cup B^c) \text{ by commutative law}$$

Then the same thing before the last step:

$$(B^c \cap A) \cup B^c = (A \cap B^c) \cup B^c \text{ by commutative law}$$

\square

In 30 – 40, construct an algebraic proof for the given statement. Cite a property from Theorem 6.2.2 for every step.

3.30 Exercise 30

For all sets A, B, C , $(A \cap B) \cup C = (A \cup C) \cap (B \cup C)$.

Proof. Let sets A, B, C be given. Then

$$\begin{aligned}
(A \cap B) \cup C &= C \cup (A \cap B) && \text{by the commutative law for } \cup \\
&= (C \cup A) \cap (C \cup B) && \text{by the distributive law} \\
&= (A \cup C) \cap (B \cup C) && \text{by the commutative law for } \cup
\end{aligned}$$

□

3.31 Exercise 31

For all sets A, B , $A \cup (B - A) = A \cup B$.

Proof. Suppose A, B are sets. Then

$$\begin{aligned}
A \cup (B - A) &= A \cup (B \cap A^c) && \text{by set difference law} \\
&= (A \cup B) \cap (A \cup A^c) && \text{by distributive law} \\
&= (A \cup B) \cap U && \text{by complement law for } \cup \\
&= A \cup B && \text{by identity law for } \cup
\end{aligned}$$

□

3.32 Exercise 32

For all sets A, B , $(A - B) \cup (A \cap B) = A$.

Proof. Suppose A, B are sets. Then

$$\begin{aligned}
(A - B) \cup (A \cap B) &= (A \cap B^c) \cup (A \cap B) && \text{by set difference law} \\
&= A \cap (B^c \cup B) && \text{by distributive law for } \cap \\
&= A \cap (B \cup B^c) && \text{by commutative law} \\
&= A \cap U && \text{by complement law} \\
&= A && \text{by identity law for } \cap
\end{aligned}$$

□

3.33 Exercise 33

For all sets A, B , $(A - B) \cap (A \cap B) = \emptyset$.

Proof. Suppose A, B are sets. Then

$$\begin{aligned}
(A - B) \cap (A \cap B) &= (A \cap B^c) \cap (A \cap B) && \text{by set difference law} \\
&= A \cap (B^c \cap (A \cap B)) && \text{by associative law} \\
&= A \cap (B^c \cap (B \cap A)) && \text{by commutative law} \\
&= A \cap ((B^c \cap B) \cap A) && \text{by associative law}
\end{aligned}$$

$$\begin{aligned}
&= A \cap ((B \cap B^c) \cap A) && \text{by commutative law} \\
&= A \cap (\emptyset \cap A) && \text{by complement law} \\
&= A \cap (A \cap \emptyset) && \text{by commutative law} \\
&= A \cap \emptyset && \text{by universal bound law} \\
&= \emptyset && \text{by universal bound law}
\end{aligned}$$

□

3.34 Exercise 34

For all sets A, B, C , $(A - B) - C = A - (B \cup C)$.

Proof. Suppose A, B, C are sets. Then

$$\begin{aligned}
(A - B) - C &= (A \cap B^c) - C && \text{by set difference law} \\
&= (A \cap B^c) \cap C^c && \text{by set difference law} \\
&= A \cap (B^c \cap C^c) && \text{by associative law} \\
&= A \cap (B \cup C)^c && \text{by De Morgan's law} \\
&= A - (B \cup C) && \text{by set difference law}
\end{aligned}$$

□

3.35 Exercise 35

For all sets A, B , $A - (A - B) = A \cap B$.

$$\begin{aligned}
A - (A - B) &= A - (A \cap B^c) && \text{by set difference law} \\
&= A \cap (A \cap B^c)^c && \text{by set difference law} \\
&= A \cap (A^c \cup (B^c)^c) && \text{by De Morgan's law} \\
&= A \cap (A^c \cup B) && \text{by double complement law} \\
&= (A \cap A^c) \cup (A \cap B) && \text{by distributive law for } \cap \\
&= \emptyset \cup (A \cap B) && \text{by complement law} \\
&= (A \cap B) \cup \emptyset && \text{by commutative law} \\
&= A \cap B && \text{by identity law}
\end{aligned}$$

Proof.

□

3.36 Exercise 36

For all sets A, B , $((A^c \cup B^c) - A)^c = A$.

Proof. Let A, B be any sets. Then

$$\begin{aligned}
((A^c \cup B^c) - A)^c &= ((A^c \cup B^c) \cap A^c)^c && \text{by set difference law} \\
&= (A^c \cup B^c)^c \cup (A^c)^c && \text{by De Morgan's law} \\
&= ((A^c)^c \cap (B^c)^c) \cup (A^c)^c && \text{by De Morgan's law} \\
&= (A \cap B) \cup A && \text{by double complement law} \\
&= A \cup (A \cap B) && \text{by commutative law} \\
&= A && \text{by absorption law}
\end{aligned}$$

□

3.37 Exercise 37

For all sets A, B , $(B^c \cup (B^c - A))^c = B$.

$$\begin{aligned}
(B^c \cup (B^c - A))^c &= (B^c \cup (B^c \cap A^c))^c && \text{by set difference law} \\
&= (B^c)^c \cap (B^c \cap A^c)^c && \text{by De Morgan's law} \\
\text{Proof.} \quad &= (B^c)^c \cap ((B^c)^c \cup (A^c)^c) && \text{by De Morgan's law} \\
&= B \cap (B \cup A) && \text{by double complement law} \\
&= B && \text{by absorption law}
\end{aligned}$$

□

3.38 Exercise 38

For all sets A, B , $(A \cap B)^c \cap A = A - B$.

$$\begin{aligned}
(A \cap B)^c \cap A &= (A^c \cup B^c) \cap A && \text{by De Morgan's law} \\
&= A \cap (A^c \cup B^c) && \text{by commutative law} \\
&= (A \cap A^c) \cup (A \cap B^c) && \text{by distributive law} \\
\text{Proof.} \quad &= \emptyset \cup (A \cap B^c) && \text{by complement law} \\
&= (A \cap B^c) \cup \emptyset && \text{by commutative law} \\
&= (A \cap B^c) && \text{by identity law} \\
&= A - B && \text{by set difference law}
\end{aligned}$$

□

3.39 Exercise 39

For all sets A, B , $(A - B) \cup (B - A) = (A \cup B) - (A \cap B)$.

Proof. Let A, B be any sets. Then $(A - B) \cup (B - A)$

$$\begin{aligned}
&= (A \cap B^c) \cup (B \cap A^c) && \text{by set difference law} \\
&= [(A \cap B^c) \cup B] \cap [(A \cap B^c) \cup A^c] && \text{by distributive law} \\
&= [B \cup (A \cap B^c)] \cap [A^c \cup (A \cap B^c)] && \text{by commutative law} \\
&= [(B \cup A) \cap (B \cup B^c)] \cap [(A^c \cup A) \cap (A^c \cup B^c)] && \text{by distributive law} \\
&= [(B \cup A) \cap U] \cap [U \cap (A^c \cup B^c)] && \text{by complement law} \\
&= [B \cup A] \cap [A^c \cup B^c] && \text{by identity law} \\
&= [B \cup A] \cap [A \cap B]^c && \text{by De Morgan's law} \\
&= [A \cup B] \cap [A \cap B]^c && \text{by commutative law} \\
&= (A \cup B) - (A \cap B) && \text{by set difference law}
\end{aligned}$$

□

3.40 Exercise 40

For all sets A, B, C , $(A - B) - (B - C) = A - B$.

Proof. Let A, B, C be any sets. Then

$$\begin{aligned}
(A - B) - (B - C) &= (A \cap B^c) \cap (B \cap C^c)^c && \text{by set difference law} \\
&= (A \cap B^c) \cap (B^c \cup (C^c)^c) && \text{by De Morgan's law} \\
&= (A \cap B^c) \cap (B^c \cup C) && \text{by double complement law} \\
&= A \cap (B^c \cap (B^c \cup C)) && \text{by associative law} \\
&= A \cap [(B^c \cap B^c) \cup (B^c \cap C)] && \text{by distributive law} \\
&= A \cap [B^c \cup (B^c \cap C)] && \text{by idempotent law} \\
&= A \cap B^c && \text{by absorption law} \\
&= A - B && \text{by set difference law}
\end{aligned}$$

□

In 41 – 43 simplify the given expression. Cite a property from Theorem 6.2.2 for every step.

3.41 Exercise 41

$$A \cap ((B \cup A^c) \cap B^c)$$

$$\begin{aligned}
A \cap ((B \cup A^c) \cap B^c) &= A \cap (B^c \cap (B \cup A^c)) && \text{by commutative law} \\
&= A \cap [(B^c \cap B) \cup (B^c \cap A^c)] && \text{by distributive law} \\
&= A \cap [\emptyset \cup (B^c \cap A^c)] && \text{by complement law} \\
&= A \cap [(B^c \cap A^c) \cup \emptyset] && \text{by commutative law} \\
&= A \cap (B^c \cap A^c) && \text{by identity law}
\end{aligned}$$

$$\begin{aligned}
&= A \cap (A^c \cap B^c) && \text{by commutative law} \\
&= (A \cap A^c) \cap B^c && \text{by associative law} \\
&= \emptyset \cap B^c && \text{by complement law} \\
&= B^c \cap \emptyset = \emptyset && \text{by commutative \& universal bound law}
\end{aligned}$$

□

3.42 Exercise 42

$$(A - (A \cap B)) \cap (B - (A \cap B))$$

$$\begin{aligned}
&= (A \cap (A \cap B)^c) \cap (B \cap (A \cap B)^c) && \text{by set difference law} \\
&= A \cap [(A \cap B)^c \cap (B \cap (A \cap B)^c)] && \text{by associative law} \\
&= A \cap [((A \cap B)^c \cap B) \cap (A \cap B)^c] && \text{by associative law}
\end{aligned}$$

$$\begin{aligned}
\text{Proof. } &= A \cap [(B \cap (A \cap B)^c) \cap (A \cap B)^c] && \text{by commutative law} \\
&= A \cap [B \cap ((A \cap B)^c \cap (A \cap B)^c)] && \text{by associative law} \\
&= A \cap [B \cap (A \cap B)^c] && \text{by idempotent law} \\
&= (A \cap B) \cap (A \cap B)^c = \emptyset && \text{by associative \& complement law}
\end{aligned}$$

□

3.43 Exercise 43

$$((A \cap (B \cup C)) \cap (A - B)) \cap (B \cup C^c)$$

$$\begin{aligned}
&= (B \cup C^c) \cap ((A \cap (B \cup C)) \cap (A - B)) && \text{by commutative law} \\
&= (B \cup C^c) \cap (((B \cup C) \cap A) \cap (A - B)) && \text{by commutative law} \\
&= (B \cup C^c) \cap ((B \cup C) \cap (A \cap (A - B))) && \text{by associative law} \\
&= [(B \cup C^c) \cap (B \cup C)] \cap (A \cap (A - B)) && \text{by associative law} \\
&= [B \cup (C^c \cap C)] \cap (A \cap (A - B)) && \text{by distributive law} \\
&= [B \cup \emptyset] \cap (A \cap (A - B)) && \text{by complement law} \\
&= B \cap (A \cap (A - B)) && \text{by identity law} \\
\text{Proof. } &= B \cap (A \cap (A \cap B^c)) && \text{by set difference law} \\
&= B \cap ((A \cap A) \cap B^c) && \text{by associative law} \\
&= B \cap (A \cap B^c) && \text{by idempotent law} \\
&= B \cap (B^c \cap A) && \text{by commutative law} \\
&= (B \cap B^c) \cap A && \text{by associative law} \\
&= \emptyset \cap A && \text{by complement law} \\
&= A \cap \emptyset && \text{by commutative law} \\
&= \emptyset && \text{by universal bound law}
\end{aligned}$$

□

3.44 Exercise 44

Consider the following set property: For all sets A and B , $A - B$ and B are disjoint.

3.44.1 (a)

Use an element argument to derive the property.

Proof. We want to show $(A - B) \cap B = \emptyset$. Argue by contradiction and assume $(A - B) \cap B \neq \emptyset$. So there is some $x \in (A - B) \cap B$. By definition of intersection $x \in A - B$ and $x \in B$. By definition of set difference $x \in A$ and $x \notin B$. So $x \in B$ and $x \notin B$, contradiction. Thus the supposition was false, and $(A - B) \cap B = \emptyset$ [as was to be shown.] \square

3.44.2 (b)

Use an algebraic argument to derive the property (by applying properties from Theorem 6.2.2).

$$\begin{aligned} (A - B) \cap B &= (A \cap B^c) \cap B && \text{by set difference law} \\ &= A \cap (B^c \cap B) && \text{by associative law} \\ \text{Proof.} \quad &= A \cap (B \cap B^c) && \text{by commutative law} \\ &= A \cap \emptyset && \text{by complement law} \\ &= \emptyset && \text{by universal bound law} \end{aligned}$$

\square

3.44.3 (c)

Comment on which method you found easier.

Proof. Second method. \square

3.45 Exercise 45

Consider the following set property: For all sets A, B , and C , $(A - B) \cup (B - C) = (A \cup B) - (B \cap C)$.

3.45.1 (a)

Use an element argument to derive the property.

Proof. 1. Assume $x \in (A - B) \cup (B - C)$. [Want to show $x \in (A \cup B) - (B \cap C)$.]

2. By 1 and definition of union, $x \in A - B$ or $x \in B - C$.

3. **Case 1** ($x \in A - B$): By definition of difference, $x \in A$ and $x \notin B$.

By definition of union and $x \in A$, $x \in A \cup B$.

By definition of intersection and $x \notin B$, $x \notin B \cap C$.

So, by definition of difference, $x \in (A \cup B) - (B \cap C)$.

4. **Case 2** ($x \in B - C$): By definition of difference, $x \in B$ and $x \notin C$.

By definition of union and $x \in B$, $x \in A \cup B$.

By definition of intersection and $x \notin C$, $x \notin B \cap C$.

So, by definition of difference, $x \in (A \cup B) - (B \cap C)$.

5. By 3 and 4, $x \in (A \cup B) - (B \cap C)$.

6. By 1, 5 and definition of subset, $(A - B) \cup (B - C) \subseteq (A \cup B) - (B \cap C)$.

Now the reverse:

7. Assume $x \in (A \cup B) - (B \cap C)$. [Want to show $x \in (A - B) \cup (B - C)$.]

8. By 7 and definition of difference, $x \in A \cup B$ and $x \notin B \cap C$.

9. By 8 and definition of union, $x \in A$ or $x \in B$. Also by 8 and definition of intersection, $x \notin B$ or $x \notin C$.

10. **Case 1** ($x \in A$): Either $x \in B$ or $x \notin B$.

11. **Subcase 1.1** ($x \in B$): Then by 9, $x \notin C$. So by definition of difference, $x \in B - C$. Then by definition of union, $x \in (A - B) \cup (B - C)$.

12. **Subcase 1.2** ($x \notin B$): Then by definition of difference, $x \in A - B$. Then by definition of union $x \in (A - B) \cup (B - C)$.

13. **Case 2** ($x \in B$): Then by 9, $x \notin C$. So by definition of difference $x \in B - C$, and then by definition of union, $x \in (A - B) \cup (B - C)$.

14. By 10-13, $x \in (A - B) \cup (B - C)$ in all cases.

15. By 7, 14 and definition of subset, $(A \cup B) - (B \cap C) \subseteq (A - B) \cup (B - C)$.

16. By 6, 15 and definition of set equality, $(A - B) \cup (B - C) = (A \cup B) - (B \cap C)$. \square

3.45.2 (b)

Use an algebraic argument to derive the property (by applying properties from Theorem 6.2.2).

Proof. $(A - B) \cup (B - C)$

$$\begin{aligned} &= (A \cap B^c) \cup (B \cap C^c) && \text{by set difference law} \\ &= [(A \cap B^c) \cup \emptyset] \cup [(B \cap C^c) \cup \emptyset] && \text{by identity law} \\ &= [(A \cap B^c) \cup (B \cap B^c)] \cup [(B \cap C^c) \cup (B \cap B^c)] && \text{by complement law} \\ &= [(B^c \cap A) \cup (B^c \cap B)] \cup [(B \cap C^c) \cup (B \cap B^c)] && \text{by commutative law} \\ &= [B^c \cap (A \cup B)] \cup [B \cap (C^c \cup B^c)] && \text{by distributive law} \end{aligned}$$

$$\begin{aligned}
&= [B^c \cap (A \cup B)] \cup [B \cap (B^c \cup C^c)] && \text{by commutative law} \\
&= ([B^c \cap (A \cup B)] \cup B) \cap ([B^c \cap (A \cup B)] \cup (B^c \cup C^c)) && \text{by distributive law} \\
&= (B \cup [B^c \cap (A \cup B)]) \cap ((B^c \cup C^c) \cup [B^c \cap (A \cup B)]) && \text{by commutative law} \\
&= ((B \cup B^c) \cap (B \cup (A \cup B))) \cap && \\
&\quad ((B^c \cup C^c) \cup B^c) \cap ((B^c \cup C^c) \cup (A \cup B)) && \text{by distributive law} \\
&= (U \cap (B \cup (A \cup B))) \cap && \\
&\quad ((B^c \cup C^c) \cup B^c) \cap ((B^c \cup C^c) \cup (A \cup B)) && \text{by complement law} \\
&= ((B \cup (A \cup B)) \cap U) \cap && \\
&\quad ((B^c \cup C^c) \cup B^c) \cap ((C^c \cup B^c) \cup (B \cup A)) && \text{by commutative law} \\
&= (B \cup (A \cup B)) \cap ((B^c \cup C^c) \cup B^c) \cap ((C^c \cup B^c) \cup (B \cup A)) && \text{by identity law} \\
&= (B \cup (B \cup A)) \cap ((C^c \cup B^c) \cup B^c) \cap ((C^c \cup B^c) \cup (B \cup A)) && \text{by commutative law} \\
&= ((B \cup B) \cup A) \cap [(C^c \cup (B^c \cup B^c)) \cap [C^c \cup (B^c \cup B) \cup A]] && \text{by associative law} \\
&= (B \cup A) \cap ((C^c \cup B^c) \cap [C^c \cup (B^c \cup B) \cup A]) && \text{by idempotent law} \\
&= (B \cup A) \cap ((C^c \cup B^c) \cap [(C^c \cup U) \cup A]) && \text{by complement law} \\
&= (B \cup A) \cap ((C^c \cup B^c) \cap (U \cup A)) && \text{by univ. bound law} \\
&= (A \cup B) \cap ((B^c \cup C^c) \cap (A \cup U)) && \text{by commutative law} \\
&= (A \cup B) \cap ((B^c \cup C^c) \cap U) && \text{by univ. bound law} \\
&= (A \cup B) \cap (B^c \cup C^c) && \text{by identity law} \\
&= (A \cup B) \cap (B \cap C)^c && \text{by De Morgan's law} \\
&= (A \cup B) - (B \cap C) && \text{by set difference law}
\end{aligned}$$

□

3.45.3 (c)

Comment on which method you found easier.

Proof. First method.

□

Definition 1. Given sets A and B , the symmetric difference of A and B , denoted $A \Delta B$, is

$$A \Delta B = (A - B) \cup (B - A).$$

3.46 Exercise 46

Let $A = \{1, 2, 3, 4\}$ and $B = \{3, 4, 5, 6\}$ and $C = \{5, 6, 7, 8\}$. Find each of the following sets:

3.46.1 (a)

$$A\Delta B$$

Proof. $A\Delta B = (A-B)\cup(B-A) = (\{1, 2, 3, 4\}-\{3, 4, 5, 6\})\cup(\{3, 4, 5, 6\}-\{1, 2, 3, 4\}) = \{1, 2\}\cup\{5, 6\} = \{1, 2, 5, 6\}$ \square

3.46.2 (b)

$$B\Delta C$$

Proof. $B\Delta C = (B-C)\cup(C-B) = \{3, 4\}\cup\{7, 8\} = \{3, 4, 7, 8\}$ \square

3.46.3 (c)

$$A\Delta C$$

Proof. $A\Delta C = (A-C)\cup(C-A) = \{1, 2, 3, 4\}\cup\{5, 6, 7, 8\} = \{1, 2, 3, 4, 5, 6, 7, 8\}$ \square

3.46.4 (d)

$$(A\Delta B)\Delta C$$

Proof. $(A\Delta B)\Delta C = \{1, 2, 5, 6\}\Delta\{5, 6, 7, 8\} = \{1, 2\}\cup\{7, 8\} = \{1, 2, 7, 8\}$ \square

Refer to the definition of symmetric difference given above. Prove each of 47 – 52, assuming that A , B , and C are all subsets of a universal set U .

3.47 Exercise 47

$$A\Delta B = B\Delta A$$

Proof. Let A and B be any subsets of a universal set. By definition of Δ , showing that $A\Delta B = B\Delta A$ is equivalent to showing that $(A-B)\cup(B-A) = (B-A)\cup(A-B)$. This follows immediately from the commutative law for \cup . \square

3.48 Exercise 48

$$A\Delta\emptyset = A$$

$$\begin{aligned}
 A\Delta\emptyset &= (A-\emptyset)\cup(\emptyset-A) && \text{by definition of } \Delta \\
 &= (A\cap\emptyset^c)\cup(\emptyset\cap A^c) && \text{by set difference law} \\
 &= (A\cap U)\cup(\emptyset\cap A^c) && \text{by complement of } U \text{ law} \\
 \text{Proof.} \quad &= (A\cap U)\cup(A^c\cap\emptyset) && \text{by commutative law} \\
 &= A\cup(A^c\cap\emptyset) && \text{by identity law} \\
 &= A\cup\emptyset && \text{by universal bound law} \\
 &= A && \text{by identity law}
 \end{aligned}$$

□

3.49 Exercise 49

$$A\Delta A^c = U$$

$$\begin{aligned}
 A\Delta A^c &= (A - A^c) \cup (A^c - A) && \text{by definition of } \Delta \\
 &= (A \cap (A^c)^c) \cup (A^c \cap A^c) && \text{by set difference law} \\
 \text{Proof.} \quad &= (A \cap A) \cup (A^c \cap A^c) && \text{by double complement law} \\
 &= A \cup A^c && \text{by idempotent law} \\
 &= U && \text{by complement law}
 \end{aligned}$$

□

3.50 Exercise 50

$$A\Delta A = \emptyset$$

$$\begin{aligned}
 A\Delta A &= (A - A) \cup (A - A) && \text{by definition of } \Delta \\
 \text{Proof.} \quad &= (A \cap A^c) \cup (A \cap A^c) && \text{by set difference law} \\
 &= \emptyset \cup \emptyset && \text{by complement law} \\
 &= \emptyset && \text{by idempotent law}
 \end{aligned}$$

□

3.51 Exercise 51

If $A\Delta C = B\Delta C$, then $A = B$.

Proof. Assume A, B, C are any sets such that $A\Delta C = B\Delta C$. [Want to show $A = B$.]

Assume $x \in A$. [Want to show $x \in B$.]

Case 1: $x \in C$. Then by definition of difference, $x \notin A - C$ and $x \notin C - A$. So by definition of Δ , $x \notin A\Delta C$. Since $A\Delta C = B\Delta C$, $x \notin B\Delta C$.

By definition of Δ , $x \notin (B - C)$ and $x \notin (C - B)$. Therefore $x \notin (C - B)$.

So by definition of difference, either $x \notin C$ or $x \in B$. Since $x \in C$, $x \notin C$ is impossible. Thus $x \in B$.

Case 2: $x \notin C$. By definition of difference, $x \in A - C$. By definition of Δ , $x \in A\Delta C$. Since $A\Delta C = B\Delta C$, $x \in B\Delta C$.

By definition of Δ , $x \in B - C$ or $x \in C - B$. Since $x \notin C$, $x \in C - B$ is impossible. Thus $x \in B - C$, so by definition of difference $x \in B$.

Therefore $A \subseteq B$.

The reverse direction ($B \subseteq A$) has a similar, symmetric proof. (Swap the roles of A and B in the above proof.)

Therefore $B \subseteq A$ and by definition of set equality, $A = B$. □

3.52 Exercise 52

$$(A \Delta B) \Delta C = A \Delta (B \Delta C)$$

Proof. Assume $x \in (A \Delta B) \Delta C$. [Want to show $x \in A \Delta (B \Delta C)$.]

By definition of Δ , $x \in (A \Delta B) - C$ or $x \in C - (A \Delta B)$.

Case 1 ($x \in (A \Delta B) - C$):

By definition of difference, $x \in A \Delta B$ and $x \notin C$.

By definition of Δ either $x \in A - B$ or $x \in B - A$.

Subcase 1.1 ($x \in A - B$):

By definition of difference, $x \in A$ and $x \notin B$.

Since $x \notin B$, by definition of difference $x \notin B - C$.

Since $x \notin C$, by definition of difference $x \notin C - B$ either.

So by definition of union, $x \notin (B - C) \cup (C - B)$, in other words, $x \notin B \Delta C$.

So $x \in A - (B \Delta C)$, thus by definition of union $x \in [A - (B \Delta C)] \cup [(B \Delta C) - A]$, in other words $x \in A \Delta (B \Delta C)$.

Subcase 1.2 ($x \in B - A$):

By definition of difference, $x \in B$ and $x \notin A$.

Since $x \in B$, by definition of difference $x \in B - C$. Thus by definition of union $x \in (B - C) \cup (C - B)$ or in other words $x \in B \Delta C$.

Since $x \notin A$, by definition of difference $x \in (B \Delta C) - A$. Thus by definition of union, $x \in [(B \Delta C) - A] \cup [A - (B \Delta C)]$, in other words $x \in A \Delta (B \Delta C)$.

Case 2 ($x \in C - (A \Delta B)$):

By definition of difference, $x \in C$ and $x \notin A \Delta B$.

By definition of Δ , $x \notin A - B$ and $x \notin B - A$.

Subcase 2.1 ($x \in A$):

Since $x \notin A - B$, by definition of difference $x \in B$.

Since $x \in B$, by definition of difference $x \notin C - B$.

Since $x \in C$, by definition of difference $x \notin B - C$.

Thus by definition of union $x \notin (B - C) \cup (C - B)$, in other words $x \notin (B \Delta C)$.

Since $x \in A$, by definition of difference $x \in A - (B\Delta C)$.

Then by definition of union $x \in [A - (B\Delta C)] \cup [(B\Delta C) - A]$, in other words, $x \in A\Delta(B\Delta C)$.

Subcase 2.2 ($x \notin A$):

Since $x \notin B - A$, by definition of difference $x \notin B$.

Since $x \in C$, by definition of difference $x \in C - B$. So by definition of union $x \in (B - C) \cup (C - B)$, in other words $x \in (B\Delta C)$.

Since $x \notin A$, by definition of difference $x \in (B\Delta C) - A$.

Then by definition of union, $x \in [A - (B\Delta C)] \cup [(B\Delta C) - A]$, in other words $x \in A\Delta(B\Delta C)$.

Thus in all cases $x \in A\Delta(B\Delta C)$. This proves that $(A\Delta B)\Delta C \subseteq A\Delta(B\Delta C)$.

[The proof of the reverse direction, $A\Delta(B\Delta C) \subseteq (A\Delta B)\Delta C$ is similar.] □

3.53 Exercise 53

Derive the set identity $A \cup (A \cap B) = A$ from the properties listed in Theorem 6.2.2(1)-(9). Start by showing that for every subset B of a universal set U , $U \cup B = U$. Then intersect both sides with A and deduce the identity.

Proof. $U \cup B = B \cup U = U$ by the commutative law and the universal bound law.

Then $A \cap (U \cup B) = A \cap U$. Now the right hand side is $A \cap U = A$ by the identity law.

The left hand side, by the distributive law, is: $A \cap (U \cup B) = (A \cap U) \cup (A \cap B)$, which is, again by the identity law, $= A \cup (A \cap B)$.

Thus the left hand side equals the right hand side, in other words, $A \cup (A \cap B) = A$. □

3.54 Exercise 54

Derive the set identity $A \cap (A \cup B) = A$ from the properties listed in Theorem 6.2.2(1)-(9). Start by showing that for every subset B of a universal set U , $\emptyset = \emptyset \cap B$. Then take the union of both sides with A and deduce the identity.

Proof. $\emptyset = B \cap \emptyset = \emptyset \cap B$ by the universal bound law and the commutative law.

Then $A \cup \emptyset = A \cup (\emptyset \cap B)$. Now the left hand side is $A \cup \emptyset = A$ by the identity law.

The right hand side is, by distributive law, $A \cup (\emptyset \cap B) = (A \cup \emptyset) \cap (A \cup B)$, which is, by the identity law, $= A \cap (A \cup B)$.

Thus the right hand side equals the left hand side, in other words, $A \cap (A \cup B) = A$. □