



✓ **Congratulations! You passed!**

TO PASS 75% or higher

Keep Learning

GRADE
100%

Non-square matrix multiplication

TOTAL POINTS 8

1. In the previous lecture we saw the Einstein summation convention, in which we sum over any indices which are repeated. In traditional notation we might write, for example, $\sum_{j=1}^3 A_{ij} v_j = A_{i1} v_1 + A_{i2} v_2 + A_{i3} v_3$. With the Einstein summation convention we can avoid the big sigma and write this as $A_{ij} v_j$. We know that we sum over j because it appears twice.

1 / 1 point

We saw that thinking about this type of notation helps us to multiply non-square matrices together. For example, consider the matrices

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 0 & 1 \end{bmatrix} \text{ and } B = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 1 & 0 & 1 \end{bmatrix},$$

and remember that in the A_{ij} notation the first index i represents the row number and the second index j represents the column number. For example, $A_{12} = 2$.

Let's define the matrix $C = AB$. Then in Einstein summation convention notation $C_{mn} = A_{mj} B_{jn}$.

Using the Einstein summation convention, calculate $C_{31} = A_{3j} B_{j1}$.