1. The probability of a leap year selected at random contain 53					
Sunday is:					
(a) 53/ 366	(b) 1/7	(c) 2/7	(d) 53/365		
2. A bag contain	s 3 red and 2 b	olue marbles. A r	marble is drawn at		
random. The pro	_	•			
(a) 3/5	(b) 2/5	(c) $0/5$	(d) 1/5		
3. The probability	ty that it will ra	in tomorrow is 0	.85. What is the		
probability that i					
(a) 0.25	(b) 0.145	(c) 3/20	(d) none of these		
•	•		ed from the numbers		
(1, 2, 3,,15					
	1 /	(c) 2/15	• •		
5. What are the					
	1 7	(c) 8	• •		
-	-	e number selecte	d at random from the		
numbers (1,2,3,					
			(d) none of these		
7. The sum of the	_				
) 0 (d) none			
	•	are given; choos	e the correct answer		
for that which is					
			(d) none of these.		
		nultaneously, tha	n the probability of		
getting at least t	wo heads, is:		(1)		
(a) 1/4	(b) 3/8	(c) ½	(d) 1/8		
10. A letter is c					
♦ ASSASSINATI	ON�. The pro	bability that the	letter chosen has:		
(a) 6/13	(b) 7/13	(c) 1	(d) none of these.		
44 4 10 4 11					
	-	• •	ting an even number.		
(A) 2/3	(B) 1	(C) 5/6	D) 1/2		
12. Two coins are thrown at the same time. Find the probability of					
getting both hea (A) 3/4 (B) 1/4		(D) 0			
13. Two dice are thrown simultaneously. The probability of getting a					

1

sum of 9 is:

(A) 1/10	(B) 3/10	(C) 1/9	(D) 4/9				
14. 100 cards are numbered from 1 to 100. Find the probability of getting a prime number.								
(A) 3/4	(B) 27/50	(C) 1/4		(D) 29/100				
15. A bag contains 5 red balls and some blue balls .If the probability of drawing a blue ball is double that of a red ball, then the number of blue balls in a bag is: (A) 5 (B) 10 (C) 15 (D) 20								
16. A box of 600 bulbs contains 12 defective bulbs. One bulb is taken out at random from this box. Then the probability that it is non-defective bulb is: (A) 143/150 (B) 147/150 (C) 1/25 (D) 1/50								
17. Cards marked with numbers 2 to 101 are placed in a box and mixed thoroughly. One card is drawn from this box randomly, then the probability that the number on card is a perfect square. (A) 9/100 (B) 1/10 (C) 3/10 (D) 19/100								
18. What is (A) 1/7	•	ty of getting ((C) 2/7	-	s in a leap year?) 7/366				
19. A card is drawn from a well shuffled deck of 52 cards. Find the probability of getting a king of red suit. (A) 1/26 (B) 3/26 (C) 7/52 (D) 1/13								
20. A game of chance consists of spinning an arrow which is equally likely to come to rest pointing to one of the number 1,2,312 ,then the probability that it will point to an odd number is: (A) 1/6 (B) 1/12 (C) 7/12 (D) 5/12								
21. A game consists of tossing a one rupee coin 3 times and noting its outcome each time. Aryan wins if all the tosses give the same result i.e. three heads or three tails and loses otherwise. Then the probability that Aryan will lose the game. (A) $3/4$ (B) $1/2$ (C) 1 (D) $1/4$								

22. Riya and Kajal are friends. Probability that both will have the same birthday is the same birthday is:							
(A) 364/365	(B) 31/365	(C) 1/365	(D) 1/133225				
2. Then the pro	x is chosen at ran bability that x² < 2 2/5 (C) 3/5	2 is?	ımbers -2, -1, 0 , 1,				
24. A jar contains 24 marbles. Some are red and others are white. If a marble is drawn at random from the jar, the probability that it is red is 2/3, then the number of white marbles in the jar is: (A) 10 (B) 6 (C) 8 (D) 7							
Then the proba	25. A number is selected at random from first 50 natural numbers. Then the probability that it is a multiple of 3 and 4 is: (A) 7/50 (B) 4/25 (C) 1/25 (D) 2/25						
26. Consider a dice with the property that that probability of a face with n dots showing up is proportional to n. The probability of face showing 4 dots is?							
a) $\frac{1}{7}$	b) $\frac{5}{42}$	c) $\frac{1}{21}$	d) $\frac{4}{21}$				
	ed by batsman in	-	es are 50, 70, 82,				
	e standard deviati b) 25.49		d) 25.69				
28. Find median and mode of the messages received on 9 consecutive days 15, 11, 9, 5, 18, 4, 18, 13, 17.							
a) 13, 15	b) 13, 18	c) 18, 15	d) 13, 16				
29. A coin is tossed up 4 times. The probability that tails turn up in 3 cases is							
a) $\frac{1}{2}$ 30. X is a varia		3. The value of I					
31. The random variables X and Y have variances 0.2 and 0.5							

respectively. Let Z= 5X-2Y. The variance of Z is?

32.Out of the probability?	he following valu	ues, which	one is not po	ssible in
a) $P(x) = 1$	b) ∑ x P(d) P(x)	(x) = 3 = - 0.5		
	2 and E(z) = 4, t b) 6	•	•	Insufficient data
34.The cov	ariance of two in	dependen	t random vari	able is
a) 1	b) 0	c) - 1	d)	Undefined
35.If Σ P(x) a) 0	b) = k² – 8 then, th			Insufficient data
, ,	0.5 and x = 4, th b) 0.5	• •	? d)	2
37.In a disc is always?	rete probability	distributio	n, the sum of	all probabilities
a) 0	b) Infinite	c) 1	d) l	Jndefined
38.If the pr	obability of hitti	ng the tar	get is 0.4, find	l mean and
	b) 0.6, 0.5	24	c) 0.4, 0.16	d) 0.6, 0.16
-	% and if 10 bom	bs are dro		lace will strike the ean and variance? d) 4, 1.6
a) 2	,	8	d) 1 or standard no	rmal distribution?

c) 5

d) 7

a) 3

b) 4

•				•	n is 1 and v an is ∞ and				
	ariance ()				given by _ 2) - (E(X))2		d) (E(X))2		
43.Mean of a random variable X is given by a) $E(X)$ b) $E(X2)$ c) $E(X2) - (E(X))2$ d) $E(X)$									
44.Mo a) 0	44.Mean of a constant 'a' is a) 0								
45.Variance of a constant 'a' is . a) 0									
46.Find the mean and variance of X?									
	Х	0	1	2	3	4			
1	f(x)	1/9	2/9	3/9	2/9	1/9			
a) 2, 4 47.Fii		•	, 4/3 on of a ra		c) 2, 2/3 ariable X?	·	d) 3, 2/3		

	Х	0	1	2	3	
	f(x)	1/6	2/6	2/6	1/6	
a) ().5		b) 1.5		c) 2.5	d) 3.5

48. In a Binomial Distribution, if p, q and n are probability of success, failure and number of trials respectively then variance is given by

b) npq

c) np2q

d) npq2

49. If 'X' is a random variable, taking values 'x', probability of success and failure being 'p' and 'q' respectively and 'n' trials being conducted, then what is the probability that 'X' takes values 'x'? Use **Binomial Distribution.**

- a) P(X = x) = nCx px qx
- b) P(X = x) = nCx px q(n-x)
- c) P(X = x) = xCn qx p(n-x)
- d) P(x = x) = xCn pn qx

50. If 'p', 'q' and 'n' are probability pf success, failure and number of trials respectively in a Binomial Distribution, what is its Standard **Deviation?**

- a) \sqrt{np} b) \sqrt{pq} c) (np)2
- d) \sqrt{npq}