

Rob J Hyndman
George Athanasopoulos

FORECASTING

PRINCIPLES AND PRACTICE

A comprehensive introduction to the latest forecasting methods using R. Learn to improve your forecast accuracy using dozens of real data examples.



3RD EDITION

 **OTexts**
Open Texts Publishing

8. Exponential smoothing

8.6 Estimation and model selection

OTexts.org/fpp3/

Estimating ETS models

- Smoothing parameters α , β , γ and ϕ , and the initial states ℓ_0 , b_0 , s_0 , s_{-1} , \dots , s_{-m+1} are estimated by maximising the “likelihood” = the probability of the data arising from the specified model.
- For models with additive errors equivalent to minimising SSE.
- For models with multiplicative errors, **not** equivalent to minimising SSE.

Innovations state space models

Let $\mathbf{x}_t = (\ell_t, b_t, s_t, s_{t-1}, \dots, s_{t-m+1})$ and $\varepsilon_t \stackrel{\text{iid}}{\sim} N(0, \sigma^2)$.

$$y_t = \underbrace{h(\mathbf{x}_{t-1})}_{\mu_t} + \underbrace{k(\mathbf{x}_{t-1})\varepsilon_t}_{e_t}$$

$$\mathbf{x}_t = f(\mathbf{x}_{t-1}) + g(\mathbf{x}_{t-1})\varepsilon_t$$

Additive errors

$$k(x) = 1. \quad y_t = \mu_t + \varepsilon_t.$$

Multiplicative errors

$$k(\mathbf{x}_{t-1}) = \mu_t. \quad y_t = \mu_t(1 + \varepsilon_t).$$

$\varepsilon_t = (y_t - \mu_t)/\mu_t$ is relative error.

Innovations state space models

Estimation

$$\begin{aligned} L^*(\boldsymbol{\theta}, \mathbf{x}_0) &= T \log \left(\sum_{t=1}^T \varepsilon_t^2 \right) + 2 \sum_{t=1}^T \log |k(\mathbf{x}_{t-1})| \\ &= -2 \log(\text{Likelihood}) + \text{constant} \end{aligned}$$

- Estimate parameters $\boldsymbol{\theta} = (\alpha, \beta, \gamma, \phi)$ and initial states $\mathbf{x}_0 = (\ell_0, b_0, s_0, s_{-1}, \dots, s_{-m+1})$ by minimizing L^* .

Parameter restrictions

Usual region

- Traditional restrictions in the methods $0 < \alpha, \beta^*, \gamma^*, \phi < 1$ (equations interpreted as weighted averages).
- In models we set $\beta = \alpha\beta^*$ and $\gamma = (1 - \alpha)\gamma^*$.
- Therefore $0 < \alpha < 1$, $0 < \beta < \alpha$ and $0 < \gamma < 1 - \alpha$.
- $0.8 < \phi < 0.98$ — to prevent numerical difficulties.

Parameter restrictions

Usual region

- Traditional restrictions in the methods $0 < \alpha, \beta^*, \gamma^*, \phi < 1$ (equations interpreted as weighted averages).
- In models we set $\beta = \alpha\beta^*$ and $\gamma = (1 - \alpha)\gamma^*$.
- Therefore $0 < \alpha < 1$, $0 < \beta < \alpha$ and $0 < \gamma < 1 - \alpha$.
- $0.8 < \phi < 0.98$ — to prevent numerical difficulties.

Admissible region

- To prevent observations in the distant past having a continuing effect on current forecasts.
- Usually (but not always) less restrictive than *traditional* region.
- For example for ETS(A,N,N):
traditional $0 < \alpha < 1$ while *admissible* $0 < \alpha < 2$.

Model selection

Akaike's Information Criterion

$$\text{AIC} = -2 \log(L) + 2k$$

where L is the likelihood and k is the number of parameters & initial states estimated in the model.

Model selection

Akaike's Information Criterion

$$\text{AIC} = -2 \log(L) + 2k$$

where L is the likelihood and k is the number of parameters & initial states estimated in the model.

Corrected AIC

$$\text{AIC}_c = \text{AIC} + \frac{2k(k+1)}{T - k - 1}$$

which is the AIC corrected (for small sample bias).

Model selection

Akaike's Information Criterion

$$\text{AIC} = -2 \log(L) + 2k$$

where L is the likelihood and k is the number of parameters & initial states estimated in the model.

Corrected AIC

$$\text{AIC}_c = \text{AIC} + \frac{2k(k+1)}{T-k-1}$$

which is the AIC corrected (for small sample bias).

Bayesian Information Criterion

$$\text{BIC} = \text{AIC} + k[\log(T) - 2].$$

Minimizing the AIC assuming Gaussian residuals is asymptotically equivalent to minimizing one-step time series cross validation MSE.

Automatic forecasting

From Hyndman et al. (IJF, 2002):

- Apply each model that is appropriate to the data. Optimize parameters and initial values using MLE (or some other criterion).
- Select best method using AICc:
- Produce forecasts using best method.
- Obtain forecast intervals using underlying state space model.

Method performed very well in M3 competition.

Some unstable models

- Some of the combinations of (Error, Trend, Seasonal) can lead to numerical difficulties; see equations with **division by a state**.
- These are: $ETS(A,N,M)$, $ETS(A,A,M)$, $ETS(A,A_d,M)$.
- Models with **multiplicative errors** are useful for **strictly positive data**, but are not numerically stable with data containing zeros or negative values. In that case only the **six fully additive models** will be applied.

Example: National populations

```
fit <- global_economy |>
  mutate(Pop = Population / 1e6) |>
  model(ets = ETS(Pop))
fit
```

```
## # A mable: 263 x 2
## # Key:      Country [263]
##   Country                ets
##   <fct>                <model>
## 1 Afghanistan    <ETS(A,A,N)>
## 2 Albania        <ETS(M,A,N)>
## 3 Algeria         <ETS(M,A,N)>
## 4 American Samoa <ETS(M,A,N)>
## 5 Andorra         <ETS(M,A,N)>
## 6 Angola          <ETS(M,A,N)>
## 7 Antigua and Barbuda <ETS(M,A,N)>
## 8 Arab World      <ETS(M,A,N)>
## 9 Argentina       <ETS(A,A,N)>
## 10 Armenia        <ETS(M,A,N)>
```

Example: National populations

```
fit |>  
  forecast(h = 5)
```

```
## # A tibble: 1,315 x 5 [1Y]  
## # Key:      Country, .model [263]  
##   Country      .model Year      Pop .mean  
##   <fct>        <chr>  <dbl>      <dist> <dbl>  
## 1 Afghanistan ets     2018    N(36, 0.012) 36.4  
## 2 Afghanistan ets     2019    N(37, 0.059) 37.3  
## 3 Afghanistan ets     2020    N(38, 0.16) 38.2  
## 4 Afghanistan ets     2021    N(39, 0.35) 39.0  
## 5 Afghanistan ets     2022    N(40, 0.64) 39.9  
## 6 Albania     ets     2018    N(2.9, 0.00012) 2.87  
## 7 Albania     ets     2019    N(2.9, 6e-04) 2.87  
## 8 Albania     ets     2020    N(2.9, 0.0017) 2.87  
## 9 Albania     ets     2021    N(2.9, 0.0036) 2.86  
## 10 Albania    ets     2022    N(2.9, 0.0066) 2.86  
## # ... with 1,305 more rows
```

Residuals

Response residuals

$$\hat{e}_t = y_t - \hat{y}_{t|t-1}$$

Innovation residuals

Additive error model:

$$\hat{\varepsilon}_t = y_t - \hat{y}_{t|t-1}$$

Multiplicative error model:

$$\hat{\varepsilon}_t = \frac{y_t - \hat{y}_{t|t-1}}{\hat{y}_{t|t-1}}$$


Example: Australian holiday tourism

```
aus_holidays <- tourism |>
  filter(Purpose == "Holiday") |>
  summarise(Trips = sum(Trips))
fit <- aus_holidays |>
  model(ets = ETS(Trips)) |>
  report()
```

```
## Series: Trips
## Model: ETS(M,N,M)
## Smoothing parameters:
##   alpha = 0.358
##   gamma = 0.000969
##
## Initial states:
## l[0]  s[0] s[-1] s[-2] s[-3]
## 9667 0.943 0.927 0.968 1.16
##
## sigma^2: 0.0022
##
## AIC AICc BIC
## 1331 1333 1348
```



Example: Australian holiday tourism

```
components(fit) |> autoplot() + labs(title = "ETS(M,N,A) components")
```



Example: Australian holiday tourism

```
residuals(fit)  
residuals(fit, type = "response")
```



Example: Australian holiday tourism

```
fit |>  
  augment()
```

```
## # A tsibble: 80 x 6 [1Q]  
## # Key:           .model [1]  
##   .model Quarter Trips .fitted .resid .innov  
##   <chr>      <qtr>  <dbl>   <dbl>  <dbl>   <dbl>  
## 1 ets      1998 Q1 11806.  11230.  576.    0.0513  
## 2 ets      1998 Q2  9276.   9532. -257.   -0.0269  
## 3 ets      1998 Q3  8642.   9036. -393.   -0.0435  
## 4 ets      1998 Q4  9300.   9050.  249.    0.0275  
## 5 ets      1999 Q1 11172.  11260. -88.0  -0.00781  
## 6 ets      1999 Q2  9608.   9358.  249.    0.0266  
## 7 ets      1999 Q3  8914.   9042. -129.   -0.0142  
## 8 ets      1999 Q4  9026.   9154. -129.   -0.0140  
## 9 ets      2000 Q1 11071.  11221. -150.   -0.0134  
## 10 ets     2000 Q2  9196.   9308. -111.   -0.0120  
## # ... with 70 more rows
```

Example: Australian holiday tourism

```
fit |>
  augment()
```

Innovation residuals (`.innov`) are given by $\hat{\varepsilon}_t$ while regular residuals (`.resid`) are $y_t - \hat{y}_{t-1}$. They are different when the model has multiplicative errors.

```
## # A tsibble: 80 x 6 [1Q]
## # Key:           .model [1]
##   .model Quarter  Trips .fitted .resid  .innov
##   <chr>      <qtr>  <dbl>   <dbl>  <dbl>   <dbl>
## 1 ets       1998 Q1 11806.  11230.  576.    0.0513
## 2 ets       1998 Q2  9276.   9532. -257.   -0.0269
## 3 ets       1998 Q3  8642.   9036. -393.   -0.0435
## 4 ets       1998 Q4  9300.   9050.  249.    0.0275
## 5 ets       1999 Q1 11172.  11260. -88.0  -0.00781
## 6 ets       1999 Q2  9608.   9358.  249.    0.0266
## 7 ets       1999 Q3  8914.   9042. -129.   -0.0142
## 8 ets       1999 Q4  9026.   9154. -129.   -0.0140
## 9 ets       2000 Q1 11071.  11221. -150.   -0.0134
## 10 ets      2000 Q2  9196.   9308. -111.   -0.0120
## # ... with 70 more rows
```