

Rob J Hyndman
George Athanasopoulos

FORECASTING

PRINCIPLES AND PRACTICE

A comprehensive introduction to the latest forecasting methods using R. Learn to improve your forecast accuracy using dozens of real data examples.



3RD EDITION

 **OTexts**
Open Texts Publishing

10. Dynamic regression models

10.1 Estimation

OTexts.org/fpp3/

Regression with ARIMA errors

Regression models

$$y_t = \beta_0 + \beta_1 x_{1,t} + \cdots + \beta_k x_{k,t} + \varepsilon_t,$$

- y_t modeled as function of k explanatory variables $x_{1,t}, \dots, x_{k,t}$.
- In regression, we assume that ε_t is WN.
- Now we want to allow ε_t to be autocorrelated.

Regression with ARIMA errors

Regression models

$$y_t = \beta_0 + \beta_1 x_{1,t} + \cdots + \beta_k x_{k,t} + \varepsilon_t,$$

- y_t modeled as function of k explanatory variables $x_{1,t}, \dots, x_{k,t}$.
- In regression, we assume that ε_t is WN.
- Now we want to allow ε_t to be autocorrelated.

Example: ARIMA(1,1,1) errors

$$y_t = \beta_0 + \beta_1 x_{1,t} + \cdots + \beta_k x_{k,t} + \eta_t,$$

$$(1 - \phi_1 B)(1 - B)\eta_t = (1 + \theta_1 B)\varepsilon_t,$$

where ε_t is white noise.

Residuals and errors

Example: $\eta_t = \text{ARIMA}(1,1,1)$

$$y_t = \beta_0 + \beta_1 x_{1,t} + \cdots + \beta_k x_{k,t} + \eta_t,$$
$$(1 - \phi_1 B)(1 - B)\eta_t = (1 + \theta_1 B)\varepsilon_t,$$

Residuals and errors

Example: $\eta_t = \text{ARIMA}(1,1,1)$

$$y_t = \beta_0 + \beta_1 x_{1,t} + \cdots + \beta_k x_{k,t} + \eta_t,$$
$$(1 - \phi_1 B)(1 - B)\eta_t = (1 + \theta_1 B)\varepsilon_t,$$

- Be careful in distinguishing η_t from ε_t .
- Only the errors ε_t are assumed to be white noise.
- In ordinary regression, η_t is assumed to be white noise and so $\eta_t = \varepsilon_t$.

Estimation

If we minimize $\sum \eta_t^2$ (by using ordinary regression):

- 1 Estimated coefficients $\hat{\beta}_0, \dots, \hat{\beta}_k$ are no longer optimal as some information ignored;
- 2 Statistical tests associated with the model (e.g., t-tests on the coefficients) are incorrect.
- 3 p -values for coefficients usually too small ("spurious regression").
- 4 AIC of fitted models misleading.

Estimation

If we minimize $\sum \eta_t^2$ (by using ordinary regression):

- 1 Estimated coefficients $\hat{\beta}_0, \dots, \hat{\beta}_k$ are no longer optimal as some information ignored;
 - 2 Statistical tests associated with the model (e.g., t-tests on the coefficients) are incorrect.
 - 3 p -values for coefficients usually too small ("spurious regression").
 - 4 AIC of fitted models misleading.
- Minimizing $\sum \varepsilon_t^2$ avoids these problems.
 - Maximizing likelihood similar to minimizing $\sum \varepsilon_t^2$.

Stationarity

Regression with ARMA errors

$$y_t = \beta_0 + \beta_1 x_{1,t} + \cdots + \beta_k x_{k,t} + \eta_t,$$

where η_t is an ARMA process.

- All variables in the model **must be stationary**.
- If we estimate the model while any of these are non-stationary, the estimated coefficients **can be incorrect**.
- **Difference** variables until all stationary.
- If necessary, apply same differencing to all variables.

Regression with ARIMA errors

Model with ARIMA(1,1,1) errors

$$y_t = \beta_0 + \beta_1 x_{1,t} + \cdots + \beta_k x_{k,t} + \eta_t,$$
$$(1 - \phi_1 B)(1 - B)\eta_t = (1 + \theta_1 B)\varepsilon_t,$$

Regression with ARIMA errors

Model with ARIMA(1,1,1) errors

$$y_t = \beta_0 + \beta_1 x_{1,t} + \cdots + \beta_k x_{k,t} + \eta_t,$$
$$(1 - \phi_1 B)(1 - B)\eta_t = (1 + \theta_1 B)\varepsilon_t,$$

Equivalent to model with ARIMA(1,0,1) errors

$$y'_t = \beta_1 x'_{1,t} + \cdots + \beta_k x'_{k,t} + \eta'_t,$$
$$(1 - \phi_1 B)\eta'_t = (1 + \theta_1 B)\varepsilon_t,$$

where $y'_t = y_t - y_{t-1}$, $x'_{t,i} = x_{t,i} - x_{t-1,i}$ and $\eta'_t = \eta_t - \eta_{t-1}$.

Regression with ARIMA errors

Any regression with an ARIMA error can be rewritten as a regression with an ARMA error by differencing all variables with the same differencing operator as in the ARIMA model.

Regression with ARIMA errors

Any regression with an ARIMA error can be rewritten as a regression with an ARMA error by differencing all variables with the same differencing operator as in the ARIMA model.

Original data

$$y_t = \beta_0 + \beta_1 x_{1,t} + \cdots + \beta_k x_{k,t} + \eta_t$$

$$\text{where } \phi(B)(1 - B)^d \eta_t = \theta(B)\varepsilon_t$$

Regression with ARIMA errors

Any regression with an ARIMA error can be rewritten as a regression with an ARMA error by differencing all variables with the same differencing operator as in the ARIMA model.

Original data

$$y_t = \beta_0 + \beta_1 x_{1,t} + \cdots + \beta_k x_{k,t} + \eta_t$$

$$\text{where } \phi(B)(1 - B)^d \eta_t = \theta(B)\varepsilon_t$$

After differencing all variables

$$y'_t = \beta_1 x'_{1,t} + \cdots + \beta_k x'_{k,t} + \eta'_t.$$

where

$$\phi(B)\eta'_t = \theta(B)\varepsilon_t, \quad y'_t = (1 - B)^d y_t, \quad x'_{i,t} = (1 - B)^d x_{i,t}, \quad \eta'_t = (1 - B)^d \eta_t$$