

Rob J Hyndman
George Athanasopoulos

FORECASTING

PRINCIPLES AND PRACTICE

A comprehensive introduction to the latest forecasting methods using R. Learn to improve your forecast accuracy using dozens of real data examples.



3RD EDITION

 **OTexts**
OPEN TEXTS FOR PRACTICE

8. Exponential smoothing

8.2 Methods with trend

OTexts.org/fpp3/

Holt's linear trend

Component form

Forecast

$$\hat{y}_{t+h|t} = \ell_t + hb_t$$

Level

$$\ell_t = \alpha y_t + (1 - \alpha)(\ell_{t-1} + b_{t-1})$$

Trend

$$b_t = \beta^*(\ell_t - \ell_{t-1}) + (1 - \beta^*)b_{t-1},$$

Holt's linear trend

Component form

Forecast	$\hat{y}_{t+h t} = \ell_t + hb_t$
Level	$\ell_t = \alpha y_t + (1 - \alpha)(\ell_{t-1} + b_{t-1})$
Trend	$b_t = \beta^*(\ell_t - \ell_{t-1}) + (1 - \beta^*)b_{t-1},$

- Two smoothing parameters α and β^* ($0 \leq \alpha, \beta^* \leq 1$).
- ℓ_t level: weighted average between y_t and one-step ahead forecast for time t , ($\ell_{t-1} + b_{t-1} = \hat{y}_{t|t-1}$)
- b_t slope: weighted average of $(\ell_t - \ell_{t-1})$ and b_{t-1} , current and previous estimate of slope.
- Choose $\alpha, \beta^*, \ell_0, b_0$ to minimise SSE.

Exponential smoothing: trend/slope


Example: Australian population

```
aus_economy <- global_economy |>
  filter(Code == "AUS") |>
  mutate(Pop = Population / 1e6)
fit <- aus_economy |>
  model(AAN = ETS(Pop ~ error("A") + trend("A") + season("N")))
report(fit)
```

```
## Series: Pop
## Model: ETS(A,A,N)
## Smoothing parameters:
##   alpha = 1
##   beta  = 0.327
##
## Initial states:
## l[0]  b[0]
## 10.1  0.222
##
## sigma^2: 0.0041
##
## AIC  AICc  BIC
## -77.0 -75.8 -66.7
```

Example: Australian population

```
components(fit) |> autoplot()
```




Example: Australian population

```
components(fit) |>  
  left_join(fitted(fit), by = c("Country", ".model", "Year"))
```

```
## # A dable: 59 x 8 [1Y]  
## # Key:      Country, .model [1]  
## # :      Pop = lag(level, 1) + lag(slope, 1) + remainder  
##   Country   .model Year   Pop level slope remainder .fitted  
##   <fct>     <chr>  <dbl> <dbl> <dbl> <dbl>      <dbl>    <dbl>  
## 1 Australia AAN     1959  NA    10.1 0.222 NA         NA  
## 2 Australia AAN     1960  10.3  10.3 0.222 -0.000145  10.3  
## 3 Australia AAN     1961  10.5  10.5 0.217 -0.0159    10.5  
## 4 Australia AAN     1962  10.7  10.7 0.231  0.0418    10.7  
## 5 Australia AAN     1963  11.0  11.0 0.223 -0.0229    11.0  
## 6 Australia AAN     1964  11.2  11.2 0.221 -0.00641   11.2  
## 7 Australia AAN     1965  11.4  11.4 0.221 -0.000314  11.4  
## 8 Australia AAN     1966  11.7  11.7 0.235  0.0418    11.6  
## 9 Australia AAN     1967  11.8  11.8 0.206 -0.0869    11.9  
## 10 Australia AAN     1968  12.0  12.0 0.208  0.00350   12.0  
## # i 49 more rows
```

Example: Australian population

```
fit |>  
  forecast(h = 10) |>  
  autoplot(aus_economy) +  
  labs(y = "Millions", title = "Population: Australia")
```



Damped trend method

Component form

$$\hat{y}_{t+h|t} = \ell_t + (\phi + \phi^2 + \dots + \phi^h)b_t$$

$$\ell_t = \alpha y_t + (1 - \alpha)(\ell_{t-1} + \phi b_{t-1})$$

$$b_t = \beta^*(\ell_t - \ell_{t-1}) + (1 - \beta^*)\phi b_{t-1}.$$

Damped trend method

Component form

$$\hat{y}_{t+h|t} = \ell_t + (\phi + \phi^2 + \dots + \phi^h)b_t$$


$$\ell_t = \alpha y_t + (1 - \alpha)(\ell_{t-1} + \phi b_{t-1})$$

$$b_t = \beta^*(\ell_t - \ell_{t-1}) + (1 - \beta^*)\phi b_{t-1}.$$

- Damping parameter $0 < \phi < 1$.
- If $\phi = 1$, identical to Holt's linear trend.
- As $h \rightarrow \infty$, $\hat{y}_{T+h|T} \rightarrow \ell_T + \phi b_T / (1 - \phi)$.
- Short-run forecasts trended, long-run forecasts constant.

Example: Australian population

```
aus_economy |>  
  model(holt = ETS(Pop ~ error("A") + trend("Ad") + season("N"))) |>  
  forecast(h = 20) |>  
  autoplot(aus_economy)
```



Example: Australian population

```
fit <- aus_economy |>
  filter(Year <= 2010) |>
  model(
    ses = ETS(Pop ~ error("A") + trend("N") + season("N")),
    holt = ETS(Pop ~ error("A") + trend("A") + season("N")),
    damped = ETS(Pop ~ error("A") + trend("Ad") + season("N"))
  )
```

```
tidy(fit)
accuracy(fit)
```

Example: Australian population

term	SES	Linear trend	Damped trend
α	1.00	1.00	1.00
β^*		0.30	0.40
ϕ			0.98
ℓ_0	10.28	10.05	10.04
b_0		0.22	0.25
Training RMSE	0.24	0.06	0.07
Test RMSE	1.63	0.15	0.21
Test MASE	6.18	0.55	0.75
Test MAPE	6.09	0.55	0.74
Test MAE	1.45	0.13	0.18