1. The probability of a leap year selected at random contain 53 Sunday is:					
(a) 53/366 (b) 1/7 (c) 2/7 (d) 53/365					
2. A bag contains 3 red and 2 blue marbles. A marble is drawn at					
random. The probability of drawing a black ball is :					
(a) 3/5 (b) 2/5 (c) 0/5 (d) 1/5					
3. The probability that it will rain tomorrow is 0.85. What is the					
probability that it will not rain tomorrow					
(a) 0.25 (b) 0.145 (c) 3/20 (d) none of these					
4. What is the probability that a number selected from the numbers					
(1, 2, 3,,15) is a multiple of 4?					
(a) 1/5 (b) 4/5 (c) 2/15 (d) 1/3					
5. What are the total outcomes when we throw three coins?					
(a) 4 (b) 5 (c) 8 (d) 7					
6. The probability that a prime number selected at random from the					
numbers (1,2,3,35) is :					
(a) 12/35 (b) 11/35 (c) 13/35 (d) none of these					
7. The sum of the probability of an event and non event is:					
(a) 2 (b) 1 (c) 0 (d) none of these.					
8. The following probabilities are given; choose the correct answer					
for that which is not possible					
for that which is not possible. (a) 0.15 (b) 2/7 (c) 7/5 (d) none of these.					
for that which is not possible. (a) 0.15 (b) 2/7 (c) 7/5 (d) none of these. 9. If three coins are tossed simultaneously, than the probability of					
for that which is not possible. (a) 0.15 (b) 2/7 (c) 7/5 (d) none of these. 9. If three coins are tossed simultaneously, than the probability of getting at least two heads, is:					
for that which is not possible. (a) 0.15 (b) $2/7$ (c) $7/5$ (d) none of these. 9. If three coins are tossed simultaneously, than the probability of getting at least two heads, is: (a) $1/4$ (b) $3/8$ (c) $\frac{1}{2}$ (d) $1/8$					
for that which is not possible. (a) 0.15 (b) $2/7$ (c) $7/5$ (d) none of these. 9. If three coins are tossed simultaneously, than the probability of getting at least two heads, is: (a) $1/4$ (b) $3/8$ (c) $\frac{1}{2}$ (d) $1/8$ 10. A letter is chosen at random from the letters of the word					
for that which is not possible. (a) 0.15 (b) 2/7 (c) 7/5 (d) none of these. 9. If three coins are tossed simultaneously, than the probability of getting at least two heads, is: (a) 1/4 (b) 3/8 (c) ½ (d) 1/8 10. A letter is chosen at random from the letters of the word ASSASSINATION. The probability that the letter chosen has:					
for that which is not possible. (a) 0.15 (b) $2/7$ (c) $7/5$ (d) none of these. 9. If three coins are tossed simultaneously, than the probability of getting at least two heads, is: (a) $1/4$ (b) $3/8$ (c) $\frac{1}{2}$ (d) $1/8$ 10. A letter is chosen at random from the letters of the word					
for that which is not possible. (a) 0.15 (b) 2/7 (c) 7/5 (d) none of these. 9. If three coins are tossed simultaneously, than the probability of getting at least two heads, is: (a) 1/4 (b) 3/8 (c) ½ (d) 1/8 10. A letter is chosen at random from the letters of the word ASSASSINATION. The probability that the letter chosen has: (a) 6/13 (b) 7/13 (c) 1 (d) none of these.					
for that which is not possible. (a) 0.15 (b) 2/7 (c) 7/5 (d) none of these. 9. If three coins are tossed simultaneously, than the probability of getting at least two heads, is: (a) 1/4 (b) 3/8 (c) ½ (d) 1/8 10. A letter is chosen at random from the letters of the word ASSASSINATION. The probability that the letter chosen has: (a) 6/13 (b) 7/13 (c) 1 (d) none of these.					
for that which is not possible. (a) 0.15 (b) 2/7 (c) 7/5 (d) none of these. 9. If three coins are tossed simultaneously, than the probability of getting at least two heads, is: (a) 1/4 (b) 3/8 (c) ½ (d) 1/8 10. A letter is chosen at random from the letters of the word ASSASSINATION. The probability that the letter chosen has: (a) 6/13 (b) 7/13 (c) 1 (d) none of these.					
for that which is not possible. (a) 0.15 (b) 2/7 (c) 7/5 (d) none of these. 9. If three coins are tossed simultaneously, than the probability of getting at least two heads, is: (a) 1/4 (b) 3/8 (c) ½ (d) 1/8 10. A letter is chosen at random from the letters of the word ASSASSINATION. The probability that the letter chosen has: (a) 6/13 (b) 7/13 (c) 1 (d) none of these. 11. A dice is thrown. Find the probability of getting an even number. (A) 2/3 (B) 1 (C) 5/6 (D) 1/2 12. Two coins are thrown at the same time. Find the probability of					
for that which is not possible. (a) 0.15 (b) 2/7 (c) 7/5 (d) none of these. 9. If three coins are tossed simultaneously, than the probability of getting at least two heads, is: (a) 1/4 (b) 3/8 (c) ½ (d) 1/8 10. A letter is chosen at random from the letters of the word ASSASSINATION The probability that the letter chosen has: (a) 6/13 (b) 7/13 (c) 1 (d) none of these. 11. A dice is thrown. Find the probability of getting an even number. (A) 2/3 (B) 1 (C) 5/6 (D) 1/2					

13. Two dice are thrown simultaneously. The probability of getting a sum of 9 is:

(A) 1/10	(B) 3/10	(C) 1/9	(D) 4/9	
	rds are numbere rime number.	ed from 1 to 10	0. Find the p	robability of
(A) 3/4	(B) 27/50	(C) 1/4	(D) 2	9/100
•	contains 5 red ba a blue ball is do n a bag is:			
(A) 5	B) 10 (C) 15	(D) 20	
	of 600 bulbs con t random from t ive bulb is:			
(A) 143/15	0 (B) 147/	(C) 1	/25	(D) 1/50
mixed thor	narked with nun oughly. One card ility that the nun (B) 1/10	d is drawn fron	n this box ra	ndomly, then
18. What is	s the probability (B) 53/366	of getting 53 I (C) 2/7	Mondays in a (D) 7/3	
19. A card	is drawn from a of getting a king	well shuffled	. ,	
(A) 1/26	(B) 3/26 (C)) 7/52 (D)	1/13	
equally like 1,2,312	e of chance conely to come to re then the proba (B) 1/12	est pointing to o	one of the nu	ımber odd number is:
its outcome result i.e. the probability	e consists of tos e each time. Ary hree heads or th that Aryan will I (B) 1/2 (C) 1	ran wins if all the ree tails and lo	ne tosses giv	ve the same

•	ajal are friends. P is the same birtho	•	oth will have the
(A) 364/365			(D) 1/133225
2. Then the pro	x is chosen at ran bability that x² < 2 2/5 (C) 3/5	2 is?	ımbers -2, -1, 0 , 1,
a marble is dra red is 2/3, then	ins 24 marbles. Sown at random fron the number of w	m the jar, the pro	_
Then the proba	s selected at rand ability that it is a n 4/25 (C) 1/25	nultiple of 3 and) natural numbers. 4 is:
with n dots sho showing 4 dots	owing up is propo	rtional to n. The	obability of a face probability of face
a) $\frac{1}{7}$	$\left(0\right)\frac{1}{42}$	C) $\frac{1}{21}$	d) $\frac{4}{21}$
	ed by batsman in e standard deviati		es are 50, 70, 82,
a) 25.79			d) 25.69
consecutive da	n and mode of the nys 15, 11, 9, 5, 18	3, 4, 18, 13, 17.	
a) 13, 15	b) 13, 18	,	d) 13, 16
29. A coin is to 3 cases is	•	The probability	that tails turn up in
a) $^{1}/_{2}$	b) $^{1}/_{3}$	c) $\frac{1}{4}$	(d) $\frac{1}{6}$
30. X is a varia b)	te between 0 and c)	3. The value of I 27 d)	
31.The random	n variables X and '	Y have variances	s 0.2 and 0.5

31. The random variables X and Y have variances 0.2 and 0.5 respectively. Let Z= 5X-2Y. The variance of Z is?

a) 3	
------	--

b) 4



d) 7

32.Out of the following values, which one is not possible in probability?

a)
$$P(x) = 1$$

b)
$$\sum x P(x) = 3$$

c)
$$P(x) = 0.5$$

d)
$$P(x) = -0.5$$

33.If E(x) = 2 and E(z) = 4, then E(z - x) = ?

b) 6

c) 0

d) Insufficient data

34. The covariance of two independent random variable is



c) - 1

d) Undefined

35.If $\Sigma P(x) = k^2 - 8$ then, the value of k is?

b) 1



d) Insufficient data

36.If P(x) = 0.5 and x = 4, then E(x) = ?

- a) 1
- b) 0.5
- c) 4



37.In a discrete probability distribution, the sum of all probabilities is always?

- a) 0
- b) Infinite
- c) 1

d) Undefined

38.If the probability of hitting the target is 0.4, find mean and variance.

- a) 0.4, 0.24
- b) 0.6, 0.24
- c) 0.4, 0.16

d) 0.6, 0.16

39. If the probability that a bomb dropped from a place will strike the target is 60% and if 10 bombs are dropped, find mean and variance?

- a) 0.6, 0.24
- (b) 6, 2.4
- c) 0.4, 0.16

d) 4, 1.6


40. Find the mean of tossing 8 coins.

- a) 2
- b) 4
- c) 8

d) 1

41. What is the mean and variance for standard normal distribution?

						d variance and varianc	
	Variance (X)	e of a rand b) E()			s given by 2) - (E(X)		d) (E(X))2
(a) E	(X)	, ,	2)	c) E(X2	ven by) - (E(X))		d) (E(X))2
44.N a) 0		a constan b) a	nt 'a' is _	c) a/2	- ·	d) 1	
(a) 0		of a cons b) a mean and		c) a/:		d) 1	
	Х	0	1	2	3	4	
	f(x)	1/9	2/9	3/9	2/9	1/9	
	, 4/3	b) 3 expectation	3, 4/3 on of a r		c) 2, 2/3 variable X	(?	d) 3, 2/3
- 3-	1						



48. In a Binomial Distribution, if p, q and n are probability of success, failure and number of trials respectively then variance is given by

5



c) np2q

d) npq2

- 49. If 'X' is a random variable, taking values 'x', probability of success and failure being 'p' and 'q' respectively and 'n' trials being conducted, then what is the probability that 'X' takes values 'x'? Use Binomial Distribution.
- (a) P(X = x) = nCx px qx
- b) P(X = x) = nCx px q(n-x)
- c) P(X = x) = xCn qx p(n-x)
- d) P(x = x) = xCn pn qx
- 50. If 'p', 'q' and 'n' are probability pf success, failure and number of trials respectively in a Binomial Distribution, what is its Standard **Deviation?**
- a) \sqrt{np}
- b) \sqrt{pq} c) (np)2