CROP NUTRITION PLAN · PER ACRE

# Hybrid Rice

May–Jun sowing | N-K programme | 15 products · 5 stages

| | PRODUCT | RATE | HOW | PACK | GROUP |
|---|---|---|---|---|---|
| **Land Preparation** | | | | | |
|  | **Green Phosphate**<br>$P_2O_5$ 32% · N 6% | **1 bag** | Drill at sowing; broadcast if no drill | 50 kg | PHOSPHORUS |
|  | **Fusion Potash**<br>$K_2O$ 50% · N 2% · S 1.5% · B 1% · Mg 1% | **1 bag** | Drill at sowing; broadcast if no drill | 25 kg | POTASH |
|  | **Humi Grow**<br>Humic Acid 40% · $K_2O$ 7% | **1 bag** | Drill at sowing; broadcast if no drill | 8 kg | SOIL AMENDMENT |
| **Germination** | | | | | |
| | **Urea**<br>N 46% | **1 bag** | Fertigation | 50 kg | NITROGEN |
|  | **V-Transform**<br>Zinc 21% | **3 bag** | Side dressing | 3 kg | MICRONUTRIENTS |
| **Early Growth** | | | | | |
|  | **V. Ammonium Phosphate (V-Phosphate)**<br>NPK 10:44:0 + Fe | **1 bag** | Fertigation | 10 kg | PHOSPHORUS |
|  | **Vital Urea**<br>N 32% · S 13% | **1 bag** | Side dressing/Broadcasting | 25 kg | NITROGEN |
|  | **Green Sulfur**<br>Sulfur 70% | **2 bag** | Fertigation | 1 kg | SECONDARY NUTRIENT |
|  | **VL-Micromix**<br>Zn 5% · Fe 2% · Mn 2% · Cu 1% (w/v) | **1 pack** | Spray | 1 L | FOLIAR |
| **Grand Growth** | | | | | |
| | **Urea**<br>N 46% | **1 bag** | Fertigation | 50 kg | NITROGEN |
|  | **Vital Potash**<br>NPK 11:0:44, coated with B | **½ bag** | Fertigation | 20 kg | POTASH |
|  | **Green Sulfur**<br>Sulfur 70% | **2 bag** | Fertigation | 1 kg | SECONDARY NUTRIENT |

| | PRODUCT | RATE | HOW | PACK | GROUP |
|---|---|---|---|---|---|
| **Maturity** | | | | | |
|  | **Vital Potash**<br>NPK 11:0:44, coated with B | **½ bag** | Fertigation | 20 kg | POTASH |
|  | **V-Mag Essential**<br>Mg 8.5% · B 1% | **1 bag** | Fertigation | 5 kg | SECONDARY NUTRIENT |
|  | **VL-Boron**<br>Boron 5% | **1 pack** | Fertigation | 1 L | MICRONUTRIENTS |

**Green Phosphate** **V. Ammonium Phosphate (V-Phosphate)** **Vital Urea** **Fusion Potash** **Vital Potash** **Green Sulfur** **V-Mag Essential** **Humi Grow** **V-Transform** **VL-Boron** **VL-Micromix** Urea

grey: general products, not VAN brands

**Drill at sowing, first**
The seed drill's fertilizer box places the granule about 5 cm beside and below the seed row, never touching the seed. The root meets it first and the whole bag is in the root zone.

**Side-dress, second**
A band beside the row, by hand or planter, at sowing or the first irrigation, covered with soil.

**Broadcast, last**
Spread on the surface before the last ploughing or before an irrigation, and mixed in. Most of the bag is away from the root.

**Indicative programme.** Built on best practice and the nutrient requirement of the crop, per acre. VAN prepares farm-specific plans against your soil analysis, yield target and the other variables that decide a rate: ask for one. Vital Urea is applied by drilling at sowing, side dressing or broadcasting, never by fertigation; the fertigation rows in the nitrogen group are general urea, a different material. Every VAN batch is tested before it ships; the batch number on the bag brings its certificate.

Rows as published on van.com.pk/crops/rice-hybrid.html, 27 Oct 2026. Rates are per acre unless marked. Where a row says "Drill at sowing; broadcast if no drill", the drill is first by VAN's order of methods.