CROP NUTRITION PLAN · PER ACRE

Watermelon

VITAL AGRI NUTRIENTS (PVT) LTD

Apr–May sowing | K-P programme | 17 products · 5 stages

	PRODUCT	RATE	HOW	PACK	GROUP
Land Preparation					
	**Green Phosphate** P_2O_5 32% · N 6%	**1.5 bag**	Drill at sowing; broadcast if no drill	50 kg	PHOSPHORUS
	**Vital Urea** N 32% · S 13%	**1 bag**	Drill at sowing; broadcast if no drill	25 kg	NITROGEN
	**Fusion Potash** K_2O 50% · N 2% · S 1.5% · B 1% · Mg 1%	**1 bag**	Drill at sowing; broadcast if no drill	25 kg	POTASH
	**Humi Grow** Humic Acid 40% · K_2O 7%	**1 bag**	Drill at sowing; broadcast if no drill	8 kg	SOIL AMENDMENT
Germination					
	**Vital Urea** N 32% · S 13%	**1 bag**	Side dressing/Broadcasting	25 kg	NITROGEN
	Urea N 46%	½ bag	Side dressing/Broadcasting	50 kg	NITROGEN
	**V-Transform** Zinc 21%	**2 pack**	Side dressing	3 kg	MICRONUTRIENTS
Early Growth					
	**V. Ammonium Phosphate (V-Phosphate)** NPK 10:44:0 + Fe	**1 bag**	Fertigation	10 kg	PHOSPHORUS
	**Vital Potash** NPK 11:0:44, coated with B	**1 bag**	Fertigation	20 kg	POTASH
	**V-Mag Essential** Mg 8.5% · B 1%	**1 bag**	Fertigation	5 kg	SECONDARY NUTRIENT
	**VL-Boron** Boron 5%	**1 pack**	Fertigation	1 L	MICRONUTRIENTS
Grand Growth					
	**Vital Potash** NPK 11:0:44, coated with B	**1 bag**	Fertigation	20 kg	POTASH
	**V-Mag Essential** Mg 8.5% · B 1%	**1 bag**	Fertigation	5 kg	SECONDARY NUTRIENT

	PRODUCT	RATE	HOW	PACK	GROUP
Grand Growth, continued					
	**Tornado** Bio-stimulant	**1 pack**	Spray	0.5 L	MICRONUTRIENTS
Maturity					
	**V. Ammonium Phosphate (V-Phosphate)** NPK 10:44:0 + Fe	**½ bag**	Fertigation	10 kg	PHOSPHORUS
	**Crop Force** NPK 12:12:18	**1 bag**	Side dressing/Placement	10 kg	NITROGEN
	**Vital Potash** NPK 11:0:44, coated with B	**½ bag**	Fertigation	20 kg	POTASH

Green Phosphate **V. Ammonium Phosphate (V-Phosphate)** **Vital Urea** **Crop Force** **Fusion Potash** **Vital Potash** **V-Mag Essential** **Humi Grow** **V-Transform** **VL-Boron** **Tornado** Urea

grey: general products, not VAN brands

Drill at sowing, first
The seed drill's fertilizer box places the granule about 5 cm beside and below the seed row, never touching the seed. The root meets it first and the whole bag is in the root zone.

Side-dress, second
A band beside the row, by hand or planter, at sowing or the first irrigation, covered with soil.

Broadcast, last
Spread on the surface before the last ploughing or before an irrigation, and mixed in. Most of the bag is away from the root.

Indicative programme. Built on best practice and the nutrient requirement of the crop, per acre. VAN prepares farm-specific plans against your soil analysis, yield target and the other variables that decide a rate: ask for one. Vital Urea is applied by drilling at sowing, side dressing or broadcasting, never by fertigation; the fertigation rows in the nitrogen group are general urea, a different material. Every VAN batch is tested before it ships; the batch number on the bag brings its certificate.

Rows as published on van.com.pk/crops/watermelon.html, 27 Oct 2026. Rates are per acre unless marked. Where a row says "Drill at sowing; broadcast if no drill", the drill is first by VAN's order of methods.